

1 **Original Article**

2 Factors associated with acceptance of a digital contact tracing application for COVID-19
3 in the Japanese working-age population

4
5 Tomohiro Ishimaru¹, MD, MPH, PhD, Koki Ibayashi¹, MD, Masako Nagata², MD, PhD,
6 Seiichiro Tateishi³, MD, PhD, Ayako Hino⁴, MD, PhD, Mayumi Tsuji⁵, MD, PhD, Hajime
7 Ando⁶, MD, MOH, PhD, Keiji Muramatsu⁷, MD, PhD, and Yoshihisa Fujino¹, MD, MPH,
8 PhD, for the CORoNaWork Project

9
10 ¹Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences,
11 University of Occupational and Environmental Health, Japan, Kitakyushu, Japan

12 ²Department of Occupational Health Practice and Management, Institute of Industrial
13 Ecological Sciences, University of Occupational and Environmental Health, Japan,
14 Kitakyushu, Japan

15 ³Department of Mental Health, Institute of Industrial Ecological Sciences, University of
16 Occupational and Environmental Health, Japan, Kitakyushu, Japan

17 ⁴Department of Occupational Medicine, School of Medicine, University of Occupational
18 and Environmental Health, Japan, Kitakyushu, Japan

19 ⁵Department of Environmental Health, School of Medicine, University of Occupational and
20 Environmental Health, Japan, Kitakyushu, Japan

21 ⁶Department of Work Systems and Health, Institute of Industrial Ecological Sciences,
22 University of Occupational and Environmental Health, Japan, Kitakyushu, Japan

23 ⁷Department of Preventive Medicine and Community Health, School of Medicine,
24 University of Occupational and Environmental Health, Japan, Kitakyushu, Japan

25
26 **Corresponding Author:** Tomohiro Ishimaru, MD, MPH, PhD, Department of
27 Environmental Epidemiology, Institute of Industrial Ecological Sciences, University of
28 Occupational and Environmental Health, Japan, 1-1 Iseigaoka, Yahata-nishi-ku,
29 Kitakyushu, Fukuoka 807-8555, Japan
30 Tel: +81-93-603-1611; Fax: +81-93-601-7324; E-mail: ishimaru@med.uoeh-u.ac.jp

32 **Funding Sources**

33 This study was supported and partly funded by the University of Occupational and
34 Environmental Health, Japan (no grant number), the Japanese Ministry of Health, Labour
35 and Welfare (H30-josei-ippan-002, H30-roudou-ippan-007, 19JA1004, 20JA1006,
36 210301-1, and 20HB1004), Anshin Zaidan (no grant number), the Collabo-Health Study
37 Group (no grant number), Hitachi Systems Ltd. (no grant number), and scholarship
38 donations from Chugai Pharmaceutical Co. Ltd. (no grant numbers). The funders were
39 not involved in the design of the study, the collection, analysis, and interpretation of data,
40 the writing of the article, or the decision to submit the manuscript for publication.

42 **Conflict of Interest**

43 None declared.

45 **Acknowledgments**

46 The current members of the CORoNaWork Project, in alphabetical order, are: Prof.
47 Yoshihisa Fujino (present chairperson of the study group), Dr. Hajime Ando, Prof. Hisashi
48 Eguchi, Dr. Kazunori Ikegami, Dr. Arisa Harada, Dr. Ayako Hino, Dr. Tomohiro Ishimaru,

49 Dr. Kyoko Kitagawa, Ms. Ning Liu, Dr. Kosuke Mafune, Prof. Shinya Matsuda, Dr. Ryutaro
50 Matsugaki, Prof. Koji Mori, Dr. Keiji Muramatsu, Dr. Masako Nagata, Dr. Tomohisa Nagata,
51 Prof. Akira Ogami, Dr. Rie Tanaka, Dr. Seiishiro Tateishi, Dr. Kei Tokutsu, and Prof.
52 Mayumi Tsuji. All members are affiliated with the University of Occupational and
53 Environmental Health, Japan.

54

55 **Ethical Considerations & Disclosures**

56 This study was approved by the Ethics Committee of the University of Occupational and
57 Environmental Health, Japan (R2-079).

58

59 **Running Title:** Workers' acceptance of a DCT app for COVID-19

60

61 **Abstract**

62 **Objective:** This study aimed to determine factors associated with acceptance of a Digital
63 Contact Tracing (DCT) app for Coronavirus Disease 2019 (COVID-19) in the Japanese
64 working-age population.

65 **Methods:** A cross-sectional study was performed for 27,036 full-time workers registered
66 with an internet survey company during December 2020 in Japan.

67 **Results:** The rate of downloading the DCT app was 25.1%. The DCT app was more
68 likely to be accepted by people with married status, university graduation or above,
69 higher income, and occupations involving desk work. Fear of COVID-19 transmission,
70 wearing a mask, using hand disinfection, willingness to be vaccinated against COVID-19,
71 and presence of an acquaintance infected with COVID-19 were also associated with a
72 greater likelihood of adopting the app.

73 **Conclusions:** The present findings have important implications for widespread adoption
74 of DCT apps in working-age populations in Japan and elsewhere.

75

76 **Keywords:** Contact tracing, COVID-19, SARS-CoV-2, smartphone, worksite

77

78 **Introduction**

79 Digital Contact Tracing (DCT) has been used in many countries in response to the
80 coronavirus disease 2019 (COVID-19) pandemic. DCT apps implemented in mobile
81 devices (e.g. smartphones) can be employed to detect close contact with infected
82 individuals and prevent the spread of COVID-19.¹ Although the method of contact tracing
83 has been used for a long time, traditional paper-based surveys are complicated and
84 time-consuming, and are affected by loss of questionnaires and huge amounts of data
85 entry.² With the recent development of digital technology and the widespread use of
86 mobile devices, DCT has overcome the shortcomings of traditional paper-based surveys
87 and made it possible to conduct surveys more quickly. In recent years, DCT has been
88 associated with effective control of Ebola hemorrhagic fever in Sierra Leone, tuberculosis
89 in Botswana, and whooping cough in the United States.³

90
91 Ensuring that large numbers of people install DCT apps is essential for effective
92 operation of DCT against COVID-19.⁴⁻⁶ Previous studies revealed that demographic,
93 behavioral, psychological, and COVID-19-related factors all influence the installation of
94 DCT apps.^{1, 3, 7-9} From the demographic aspect, ethnic minorities, migrants, low-income
95 people, and elderly people were less likely to install the apps.³ From the behavioral
96 aspect, non-smoking status was associated with high app acceptance, while lack of
97 information technology skills and not having a smartphone were associated with low app
98 acceptance.^{1, 8} From the psychological aspect, low app acceptance was associated with
99 doubts about the effectiveness of DCT apps, anxiety about the collection of personal
100 information, and distrust of the government and app developers.^{8, 9} From the
101 COVID-19-related aspect, acceptance of new lifestyles under the COVID-19 epidemic

102 such as use of hand sanitizers, avoidance of public transportation, and wearing masks
103 was associated with app use.^{7, 8}

104
105 Because the working-age population is at the epicenter of COVID-19 infections, it is
106 important to increase the acceptance of DCT apps in this population.¹⁰ In other words,
107 the public health benefits of adopting these apps need to be particularly recognized by
108 the working-age population.^{9, 11} In countries where installation of DCT apps is optional,
109 such as North America, Europe, Oceania, and Japan, the installation rates tend to be
110 lower compared with countries where it is mandatory.^{1, 7, 12} However, few studies have
111 evaluated the factors associated with acceptance of DCT apps using sufficient sample
112 sizes of workers.¹³ The purpose of the present study was to evaluate the factors
113 associated with acceptance of a DCT app for COVID-19 in the working-age population of
114 Japan, where app installation is optional. The findings provide insights that can help to
115 facilitate the adoption of DCT apps among workers.

116
117 **Methods**

118 Study design and participants

119 The present study had a cross-sectional design and used baseline questionnaire data
120 from a prospective cohort research project in Japan, the Collaborative Online Research
121 on the Novel-coronavirus and Work (CORoNaWork), between 22 and 26 December
122 2020.¹⁴ The study supplements another study from the CORoNaWork project that
123 evaluated occupational factors associated with DCT adoption.¹⁰ During the research
124 period, COVID-19 vaccines had not yet been introduced in Japan and the third wave of
125 infection was at its peak, with more than 3,000 new infections per day.¹⁵ The inclusion

126 criteria for the study were: full-time worker status, age between 20 and 65 years, and
127 registration with an internet survey company. Invitations were sent by e-mail to 605,381
128 people registered with an internet survey company, of whom 55,045 people accessed the
129 linked website. Of these, 33,302 participants who met the inclusion criteria provided
130 informed consent and completed the baseline survey in an online format. After excluding
131 6,266 participants who gave incorrect answers to questions designed to identify
132 inappropriate responses, 27,036 participants were eligible for the analysis.

133
134 The Contact-Confirming Application (COCOA) DCT app in Japan
135 The COCOA is the only DCT app in Japan and was released by the Ministry of Health,
136 Labour and Welfare in June 2020. The COCOA uses a decentralized approach through
137 Bluetooth and is designed not to store personal information on a central server controlled
138 by the government or the developer. Furthermore, downloading and deleting the COCOA
139 is optional. If a person comes into contact with a COVID-19-positive person within a 1-m
140 radius for more than 15 minutes, the person is notified as a close contact.⁶

141
142 **Dependent variable**
143 The dependent variable in the study was the following question about adoption of the
144 COCOA: “Have you downloaded the Contact-Confirming Application (COCOA)?” The
145 response options were “yes” and “no”.

146
147 **Independent variables**
148 Based on relevant studies, the independent variables were divided into four aspects:
149 demographic, behavioral, psychological, and COVID-19 transmission-related variables.^{1,}

150 ^{3, 7-9} Demographic variables included sex, age, marital status, education, annual
151 household income (in Japanese yen), occupation (desk work, work with interpersonal
152 communication, manual work), and family members living together. Behavioral variables
153 included smoking status, alcohol consumption, and regular doctor's visits for chronic
154 diseases. Psychological variables included fear of COVID-19 transmission. Variables
155 related to COVID-19 transmission included wearing a mask, using hand sanitizer,
156 willingness to be vaccinated against COVID-19, and presence of an acquaintance
157 infected with COVID-19.

158
159 **Statistical analysis**
160 Factors associated with COCOA adoption were evaluated by univariate and multivariate
161 logistic regression analyses, and the results were presented as odds ratios [ORs] and
162 95% confidence intervals [CIs]. The multivariate analyses were adjusted for sex, age,
163 marital status, education, and annual household income, based on relevant studies.^{1, 3, 7-9}
164 The *P*-values were two-sided, and statistical significance was set at $P < 0.05$. All
165 statistical analyses were performed using Stata/SE 16.1 software (StataCorp, College
166 Station, TX, USA).

167
168 **Results**
169 Table 1 shows the general characteristics of the study participants. Of the 27,036
170 participants, around half were men (51.1%), married (55.6%), educated at university or
171 graduate school (48.7%), and engaged in desk work (49.8%). Regarding acceptance of
172 the DCT app for COVID-19, 6,786 respondents (25.1%) reported that they had
173 downloaded the COCOA.

174
175 Table 2 presents the factors associated with acceptance of the DCT app for COVID-19. In
176 the multivariate analyses, married (adjusted OR: 1.29; 95% CI: 1.17–1.43) and
177 divorced/widowed (adjusted OR: 1.16; 95% CI: 1.09–1.25) individuals had significantly
178 higher acceptance of the DCT app than single individuals. Similarly, people with a
179 university or graduate school degree had higher acceptance of the DCT app than people
180 with a junior high school or high school degree (adjusted OR: 1.19; 95% CI: 1.11–1.28).
181 Higher annual income was associated with higher acceptance of the app ($\geq 8,000,000$
182 yen: adjusted OR: 2.08; 95% CI: 1.80–2.39). In contrast, acceptance of the app was
183 lower among people living with family members (adjusted OR: 0.82; 95% CI: 0.76–0.89).

184
185 From the behavioral aspect, past smokers were more accepting of the app than
186 non-smokers (adjusted OR: 1.24; 95% CI: 1.15–1.34), but there was no significant
187 difference for smokers. Patients who regularly visited the hospital for chronic diseases
188 were more likely to accept the app than patients who did not (adjusted OR: 1.39; 95% CI:
189 1.32–1.48). From the psychological aspect, the odds of accepting the app were 2.29
190 times (95% CI: 2.13–2.47) higher in people who feared COVID-19 transmission. From
191 the COVID-19-related aspect, people who always wore a mask (adjusted OR: 2.44; 95%
192 CI: 1.94–3.08) or used hand sanitizer (adjusted OR: 1.88; 95% CI: 1.71–2.07) were more
193 likely to accept the app than people who rarely or never performed these actions.
194 Acceptance of the app had 1.83 times higher odds (95% CI: 1.73–1.94) among people
195 who were willing to receive the COVID-19 vaccine than among people who were not.
196 Similarly, people who had an acquaintance infected with COVID-19 had 1.76 times
197 higher odds (95% CI: 1.61–1.93) of accepting the app than people who did not.

198

199 **Discussion**

200 In the present study, the rate of downloading the COCOA in Japan was not very high at
201 25.1%. The results revealed that there were several factors associated with downloading
202 of the DCT app for each of the demographic, behavioral, psychological, and
203 COVID-19-related variables. Although different countries have differing policies toward
204 DCT apps, the present results suggest that adoption of DCT apps may be influenced by
205 individual characteristics. The findings of the present study have important implications
206 for the widespread adoption of DCT apps in working-age populations in Japan and
207 elsewhere.

208

209 From the demographic aspect, the present study showed that the DCT app was more
210 likely to be accepted by people with married status, university graduation or above,
211 higher income, and desk work. These findings suggest that relatively wealthy people in
212 society are more likely to use DCT apps, consistent with previous studies conducted in
213 many countries including the United States, the United Kingdom, Switzerland, and
214 Japan.^{8, 16-18} In contrast, the present study found that living with family members tended
215 to reduce the use of the DCT app, which was inconsistent with a previous study.¹⁹
216 Although the reason for the discrepancy remains unclear, it is possible that family
217 interactions reduced anxiety about COVID-19, which in turn reduced the need for the
218 DCT app. Conflicting results were also reported in another study. Specifically, Tomczyk et
219 al.²⁰ pointed out that greater perceived usefulness of DCT apps was associated with
220 greater fear of COVID-19 infection and lower adoption of the apps. Therefore, people
221 who are afraid of letting their families know that they are infected with COVID-19 may

222 tend to avoid DCT apps.

223

224 From the behavioral aspect, individuals with regular doctor's visits for chronic diseases
225 and past smokers were more willing to adopt the DCT app, while current smokers
226 showed no association in the present study. The severity of COVID-19 was reported to
227 be higher in people with regular doctor's visits for chronic diseases such as hypertension
228 and respiratory disease, and in smokers.²¹ Therefore, the present results suggest that
229 people with a greater risk of severe COVID-19 are more positive about adopting DCT
230 apps. On the contrary, current smokers generally have limited access to health
231 information, which may have attenuated the result.²² Measures are needed to facilitate
232 adoption of DCT apps by these people. For example, it may be useful for family
233 physicians to encourage their patients to adopt DCT apps.

234

235 From the psychological aspect, fear of COVID-19 transmission was strongly associated
236 with higher acceptance of the DCT app, consistent with a previous study in the United
237 Kingdom that found a positive association between risk perception for COVID-19
238 infection and acceptance of DCT apps.¹⁷ Kawakami et al.¹⁹ found that installation of a
239 DCT app mitigated the psychological distress associated with the risk of COVID-19
240 infection. Taken together, the findings suggest that fear of COVID-19 transmission
241 increases the adoption of DCT apps and that appropriate adoption of DCT apps may
242 reduce anxiety about COVID-19.

243

244 From the COVID-19-related aspect, wearing a mask, using hand disinfection, willingness
245 to be vaccinated against COVID-19, and presence of an acquaintance infected with

246 COVID-19 led to greater likelihood of adopting the DCT app. A previous study in
247 Singapore demonstrated similar findings for mask wearing and hand sanitizer use.⁷ A
248 similar trend was also observed in a study conducted in the United States, wherein
249 people infected with COVID-19 had a higher acceptance rate of DCT apps.¹⁸ Based on
250 these findings, posting Quick Response codes for DCT apps on the packages of masks
251 and hand sanitizers and at vaccination sites may be an effective measure. A function for
252 inviting friends within the apps may also be effective.⁹

253
254 There are several limitations to this study. First, we recruited participants who were
255 registered with an internet research company. Therefore, we may have selected people
256 who actively used the internet, smartphones, and apps. However, we consider this
257 potential selection bias to be negligible because the COCOA downloading rate
258 announced by the Ministry of Health, Labour and Welfare in December 2020 was
259 20.8%,²³ and thus close to the rate in the present study (25.1%). Second, the present
260 study used cross-sectional data for a single period of time. Therefore, the findings can
261 only suggest associations between factors for this given period of time, and cannot
262 capture changes over time within individuals. Third, in the multivariate logistic regression
263 analyses, there was a possibility of unmeasured confounding. Unmeasured personal
264 habits and behavioral patterns may have influenced the adoption of the DCT app.

265
266 **Conclusions**
267 The present study revealed factors associated with adoption of a DCT app among the
268 working-age population in Japan. Use of the COCOA among Japanese workers was not
269 sufficient, and it is important to develop an approach based on the present findings. We

270 found that relatively wealthy people in society were more likely to use the DCT app. In
271 addition, people with a greater risk of severe COVID-19 were more positive about
272 adopting the DCT app. Fear of COVID-19 transmission increased adoption of the DCT
273 app, suggesting that appropriate adoption of DCT apps can reduce anxiety about
274 COVID-19. Because wearing a mask, using hand disinfection, and willingness to be
275 vaccinated against COVID-19 were associated with greater likelihood of adopting the
276 DCT app, posting Quick Response codes for DCT apps on the packages of masks and
277 hand sanitizers and at vaccination sites may be an effective measure.

278

279 **References**

- 280 1. Elkhodr M, Mubin O, Iftikhar Z, et al. Technology, Privacy, and User Opinions of
281 COVID-19 Mobile Apps for Contact Tracing: Systematic Search and Content Analysis. *J*
282 *Med Internet Res.* 2021;23:e23467.
- 283 2. Kouliaridis V, Kambourakis G, Chatzoglou E, Geneiatakis D, Wang H. Dissecting
284 contact tracing apps in the Android platform. *PLoS One.* 2021;16:e0251867.
- 285 3. Anglemyer A, Moore TH, Parker L, et al. Digital contact tracing technologies in
286 epidemics: a rapid review. *Cochrane Database Syst Rev.* 2020;8:CD013699.
- 287 4. Oldeweme A, Martins J, Westmattelmann D, Schewe G. The Role of Transparency,
288 Trust, and Social Influence on Uncertainty Reduction in Times of Pandemics: Empirical
289 Study on the Adoption of COVID-19 Tracing Apps. *J Med Internet Res.* 2021;23:e25893.
- 290 5. Rodriguez P, Grana S, Alvarez-Leon EE, et al. A population-based controlled
291 experiment assessing the epidemiological impact of digital contact tracing. *Nat Commun.*
292 2021;12:587.
- 293 6. Nakamoto I, Jiang M, Zhang J, et al. Evaluation of the Design and Implementation of

- 294 a Peer-To-Peer COVID-19 Contact Tracing Mobile App (COCOA) in Japan. *JMIR*
295 *Mhealth Uhealth*. 2020;8:e22098.
- 296 7. Saw YE, Tan EY, Liu JS, Liu JC. Predicting Public Uptake of Digital Contact Tracing
297 During the COVID-19 Pandemic: Results From a Nationwide Survey in Singapore. *J Med*
298 *Internet Res*. 2021;23:e24730.
- 299 8. von Wyl V, Hoglinger M, Sieber C, et al. Drivers of Acceptance of COVID-19
300 Proximity Tracing Apps in Switzerland: Panel Survey Analysis. *JMIR Public Health*
301 *Surveill*. 2021;7:e25701.
- 302 9. Garrett PM, White JP, Lewandowsky S, et al. The acceptability and uptake of
303 smartphone tracking for COVID-19 in Australia. *PLoS One*. 2021;16:e0244827.
- 304 10. Ishimaru T, Ibayashi K, Nagata M, et al. Industry and workplace characteristics
305 associated with the downloading of a COVID-19 contact tracing app in Japan: a
306 nation-wide cross-sectional study. *Environmental health and preventive medicine*.
307 2021;26:94.
- 308 11. Garrett PM, Wang Y, White JP, et al. Young Adults View Smartphone Tracking
309 Technologies for COVID-19 as Acceptable: The Case of Taiwan. *Int J Environ Res Public*
310 *Health*. 2021;18.
- 311 12. Salathe M, Althaus C, Anderegg N, et al. Early evidence of effectiveness of digital
312 contact tracing for SARS-CoV-2 in Switzerland. *Swiss Med Wkly*. 2020;150:w20457.
- 313 13. Kolasa K, Mazzi F, Leszczuk-Czubkowska E, Zrubka Z, Péntek M. State of the Art in
314 Adoption of Contact Tracing Apps and Recommendations Regarding Privacy Protection
315 and Public Health: Systematic Review. *JMIR Mhealth Uhealth*. 2021;9:e23250.
- 316 14. Fujino Y, Ishimaru T, Eguchi H, et al. Protocol for a Nationwide Internet-based Health
317 Survey of Workers During the COVID-19 Pandemic in 2020. *J uoeh*. 2021;43:217-225.

- 318 15. Ishimaru T, Okawara M, Ando H, et al. Gender differences in the determinants of
319 willingness to get the COVID-19 vaccine among the working-age population in Japan.
320 *Hum Vaccin Immunother.* 2021;in press.
- 321 16. Machida M, Nakamura I, Kojima T, et al. Acceptance of a COVID-19 Vaccine in
322 Japan during the COVID-19 Pandemic. *Vaccines.* 2021;9.
- 323 17. Lewandowsky S, Dennis S, Perfors A, et al. Public acceptance of
324 privacy-encroaching policies to address the COVID-19 pandemic in the United Kingdom.
325 *PLoS One.* 2021;16:e0245740.
- 326 18. Zhang B, Kreps S, McMurry N, McCain RM. Americans' perceptions of privacy and
327 surveillance in the COVID-19 pandemic. *PLoS One.* 2020;15:e0242652.
- 328 19. Kawakami N, Sasaki N, Kuroda R, Tsuno K, Imamura K. The Effects of Downloading
329 a Government-Issued COVID-19 Contact Tracing App on Psychological Distress During
330 the Pandemic Among Employed Adults: Prospective Study. *JMIR Ment Health.*
331 2021;8:e23699.
- 332 20. Tomczyk S, Barth S, Schmidt S, Muehlan H. Utilizing Health Behavior Change and
333 Technology Acceptance Models to Predict the Adoption of COVID-19 Contact Tracing
334 Apps: Cross-sectional Survey Study. *J Med Internet Res.* 2021;23:e25447.
- 335 21. Rahman A, Sathi NJ. Risk factors of the severity of COVID-19: A meta-analysis. *Int J*
336 *Clin Pract.* 2021;75:e13916.
- 337 22. Paul CL, Ross S, Bryant J, Hill W, Bonevski B, Keevy N. The social context of
338 smoking: A qualitative study comparing smokers of high versus low socioeconomic
339 position. *BMC Public Health.* 2010;10:211.
- 340 23. Ministry of Health Labour and Welfare. COVID-19 Contact-Confirming Application.
341 2021; Available at: https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/cocoa_00138.html.

342 Accessed 14 Jul, 2021.

343

344 Table 1. General characteristics of the study participants

	Total N=27,036 n (%)	Acceptance of digital contact tracing application for COVID-19	
		Yes n=6,786 n (%)	No n=20,250 n (%)
Sex			
Women	13,222 (48.9)	3,181 (46.9)	10,041 (49.6)
Men	13,814 (51.1)	3,605 (53.1)	10,209 (50.4)
Age			
20–39 years	6,763 (25.0)	1,669 (24.6)	5,094 (25.2)
40–49 years	8,011 (29.6)	1,886 (27.8)	6,125 (30.2)
50–59 years	9,012 (33.4)	2,359 (34.8)	6,653 (23.9)
60–65 years	3,250 (12.0)	872 (12.8)	2,378 (11.7)
Marital status			
Single	9,164 (33.9)	1,991 (29.4)	7,173 (35.4)
Divorced or widowed	2,843 (10.5)	722 (10.6)	2,121 (10.5)
Married	15,029 (55.6)	4,073 (60.0)	10,956 (54.1)
Education			
Junior high or high school	7,321 (27.1)	1,647 (24.3)	5,674 (28.0)
Vocational school or college	6,544 (24.2)	1,552 (22.9)	4,992 (24.7)
University or graduate school	13,171 (48.7)	3,587 (52.8)	9,584 (47.3)
Annual household income			
< 2,000,000 yen	1,709 (6.3)	277 (4.1)	1,432 (7.1)
2,000,000–3,999,999 yen	5,548 (20.5)	1,188 (17.5)	4,360 (21.5)
4,000,000–7,999,999 yen	11,927 (44.1)	2,925 (43.1)	9,002 (44.5)
≥ 8,000,000 yen	7,852 (29.1)	2,396 (35.3)	5,456 (26.9)
Job type			
Manual work	6,641 (24.6)	1,283 (18.9)	5,358 (26.5)
Desk work	13,468 (49.8)	3,736 (55.1)	9,732 (48.1)
Work involving interpersonal communication	6,927 (25.6)	1,767 (26.0)	5,160 (25.5)
Family living together			
No	5,807 (21.5)	1,400 (20.6)	4,407 (21.8)
Yes	21,229 (78.5)	5,386 (79.4)	15,843 (78.2)
Smoking			
Never	14,587 (54.0)	3,495 (51.5)	11,092 (54.8)
Past	5,445 (20.1)	1,551 (22.9)	3,894 (19.2)
Current	7,004 (25.9)	1,740 (25.6)	5,264 (26.0)

345

346

347 Table 1. General characteristics of the study participants (cont.)

	Total N=27,036 n (%)	Acceptance of digital contact tracing application for COVID-19	
		Yes n=6,786 n (%)	No n=20,250 n (%)
Alcohol intake			
Never	11,472 (42.4)	2,699 (39.8)	8,773 (43.3)
Occasionally	4,547 (16.8)	1,186 (17.5)	3,361 (16.6)
Regular	11,017 (40.8)	2,901 (42.7)	8,116 (40.1)
Regular doctor's visits for chronic diseases			
No	17,526 (64.8)	4,008 (59.1)	13,518 (66.8)
Yes	9,510 (35.2)	2,778 (40.9)	6,732 (33.2)
Fear of COVID-19 transmission			
No	7,019 (26.0)	1,049 (15.5)	5,970 (29.5)
Yes	20,017 (74.0)	5,737 (84.5)	14,280 (70.5)
Wearing a mask			
Rarely/never	682 (2.5)	84 (1.2)	598 (3.0)
Often	3,046 (11.3)	570 (8.4)	2,476 (12.2)
Always	23,308 (86.2)	6,132 (90.4)	17,176 (84.8)
Using hand sanitizer			
Rarely/never	3,740 (13.8)	633 (9.3)	3,107 (15.3)
Often	8,282 (30.6)	1,965 (29.0)	6,317 (31.2)
Always	15,014 (55.4)	4,188 (61.7)	10,826 (53.5)
Willing to receive the COVID-19 vaccine			
No	16,889 (62.5)	3,467 (51.1)	13,422 (66.3)
Yes	10,147 (37.5)	3,319 (48.9)	6,828 (33.7)
Presence of an acquaintance infected with COVID-19			
No	24,760 (91.6)	5,947 (87.6)	18,813 (92.9)
Yes	2,276 (8.4)	839 (12.4)	1,437 (7.1)

348 COVID-19: coronavirus disease 2019.

349

350 Table 2. Factors associated with acceptance of the digital contact tracing application for COVID-19

Category	Variable	Univariate			Multivariate*		
		OR	(95% CI)	P value	OR	(95% CI)	P value
Demographic	Sex						
	Women	1.00	-	-	1.00	-	-
	Men	1.12	(1.06–1.18)	<0.001	1.03	(0.96–1.10)	0.402
	Age						
	20–39 years	1.00	-	-	1.00	-	-
	40–49 years	0.94	(0.87–1.01)	0.108	0.90	(0.83–0.98)	0.011
	50–59 years	1.08	(1.01–1.16)	0.033	0.99	(0.91–1.07)	0.752
	60–65 years	1.12	(1.02–1.23)	0.020	1.01	(0.91–1.12)	0.873
	Marital status						
	Single	1.00	-	-	1.00	-	-
	Divorced or widowed	1.23	(1.11–1.35)	<0.001	1.16	(1.09–1.25)	<0.001
	Married	1.34	(1.26–1.42)	<0.001	1.29	(1.17–1.43)	<0.001
	Education						
	Junior high or high school	1.00	-	-	1.00	-	-
	Vocational school or college	1.07	(0.99–1.16)	0.089	1.08	(0.99–1.17)	0.078
	University or graduate school	1.29	(1.21–1.38)	<0.001	1.19	(1.11–1.28)	<0.001
	Annual household income						
	< 2,000,000 yen	1.00	-	-	1.00	-	-
	2,000,000–3,999,999 yen	1.41	(1.22–1.63)	<0.001	1.40	(1.21–1.61)	<0.001
	4,000,000–7,999,999 yen	1.68	(1.47–1.92)	<0.001	1.60	(1.39–1.83)	<0.001
≥ 8,000,000 yen	2.27	(1.98–2.60)	<0.001	2.08	(1.80–2.39)	<0.001	
Job type							
Manual work	1.00	-	-	1.00	-	-	
Desk work	1.60	(1.49–1.72)	<0.001	1.49	(1.38–1.60)	<0.001	
Work involving interpersonal communication	1.43	(1.32–1.55)	<0.001	1.36	(1.25–1.48)	<0.001	
Family living together							
No	1.00	-	-	1.00	-	-	
Yes	1.07	(1.00–1.15)	0.049	0.82	(0.76–0.89)	<0.001	

351 Table 2. Factors associated with acceptance of the digital contact tracing application for COVID-19 (cont.)

Category	Variable	Univariate			Multivariate*		
		OR	(95% CI)	P value	OR	(95% CI)	P value
Behavioral	Smoking						
	Never smoking	1.00	-	-	1.00	-	-
	Past smoking	1.26	(1.18–1.36)	<0.001	1.24	(1.15–1.34)	<0.001
	Current smoking	1.05	(0.98–1.12)	0.156	1.05	(0.98–1.12)	0.212
	Alcohol intake						
	Never	1.00	-	-	1.00	-	-
	Occasionally	1.15	(1.06–1.24)	0.001	1.12	(1.03–1.21)	0.006
	Regular	1.16	(1.09–1.23)	<0.001	1.08	(1.02–1.15)	0.014
	Regular doctor's visits for chronic diseases						
No	1.00	-	-	1.00	-	-	
Yes	1.39	(1.32–1.47)	<0.001	1.39	(1.32–1.48)	<0.001	
Psychological	Fear of COVID-19 transmission						
	No	1.00	-	-	1.00	-	-
Yes	2.29	(2.13–2.46)	<0.001	2.29	(2.13–2.47)	<0.001	
COVID-19 related	Wearing a mask						
	Rarely/never	1.00	-	-	1.00	-	-
	Often	1.64	(1.28–2.10)	<0.001	1.57	(1.23–2.01)	<0.001
	Always	2.54	(2.02–3.20)	<0.001	2.44	(1.94–3.08)	<0.001
	Using hand sanitizer						
	Rarely/never	1.00	-	-	1.00	-	-
	Often	1.53	(1.38–1.69)	<0.001	1.50	(1.36–1.66)	<0.001
	Always	1.90	(1.73–2.08)	<0.001	1.88	(1.71–2.07)	<0.001
	Willing to receive the COVID-19 vaccine						
	No	1.00	-	-	1.00	-	-
	Yes	1.88	(1.78–1.99)	<0.001	1.83	(1.73–1.94)	<0.001
	Presence of an acquaintance infected with COVID-19						
No	1.00	-	-	1.00	-	-	
Yes	1.85	(1.69–2.02)	<0.001	1.76	(1.61–1.93)	<0.001	

352 COVID-19: coronavirus disease 2019; OR: odds ratio; CI: confidence interval.

353 *Adjusted for sex, age, income, marital status, education, and annual household income.