

1 A neonatal ward strengthening program improves survival for neonates treated with CPAP at
2 district hospitals in Malawi

3

4 Jennifer Carns^{1,2}; Sara Liaghati-Mobarhan¹; Aba Asibon¹; Samuel Ngwala¹; Elizabeth
5 Molyneux³; Maria Oden^{1,2}; Rebecca Richards-Kortum^{1,2*}; Kondwani Kawaza³; Alfred Chalira⁴;
6 Norman Lufesi⁴

7

8 ¹Rice360 Institute for Global Health, Houston, TX

9 ²Department of Bioengineering, Rice University, Houston, TX

10 ³Department of Paediatrics, College of Medicine, Queen Elizabeth Central Hospital, Blantyre,
11 Malawi

12 ⁴Malawi Ministry of Health, Department of Clinical Services, Lilongwe, Malawi

13

14 *Corresponding author

15 E-mail: rkortum@rice.edu (RK)

16 **Abstract**

17 Introduction: From 2013-2015, a CPAP quality improvement program (QIP) was implemented
18 to introduce and monitor CPAP usage and outcomes in the neonatal wards at all government
19 district and central hospitals in Malawi. In 2016 the CPAP QIP was extended into healthcare
20 facilities operated by the Christian Health Association of Malawi. Although clinical outcomes
21 improved, ward assessments indicated that many rural sites lacked other essential equipment and
22 a suitable space to adequately treat sick neonates, which likely limited the impact of improved
23 respiratory care. The aim of this study was to determine if a ward strengthening program
24 improved outcomes for neonates treated with CPAP.

25 Methods: To address the needs identified from ward assessments, a ward strengthening program
26 was implemented from 2017-2018 at rural hospitals in Malawi to improve the care of sick
27 neonates. The ward strengthening program included the distribution of a bundle of equipment,
28 supplemental training, and, in some cases, health facility renovations. Survival to discharge was
29 compared for neonates treated with CPAP at 12 rural hospitals for one year before and for one
30 year immediately after implementation of the ward strengthening program.

31 Results: In the year prior to ward strengthening, 189 neonates were treated with CPAP; in the
32 year after, 232 neonates received CPAP. The overall rate of survival for those treated with CPAP
33 improved from 46.6% to 57.3% after ward strengthening ($p=0.03$). For the subset of neonates
34 with admission weights between 1.00 – 2.49 kg diagnosed with respiratory distress syndrome,
35 survival increased from 39.4% to 60.3% after ward strengthening ($p=0.001$).

36 Conclusion: A ward strengthening program including the distribution of a bundle of equipment,
37 supplemental training, and some health facility renovations, further improved survival among
38 neonates treated with CPAP at district-level hospitals in Malawi.

39 **Introduction**

40 Globally, 85% of neonatal deaths are attributed to complications from prematurity and low
41 birthweight, intrapartum-related hypoxia, and infection [1]. Neonates with these complications
42 are particularly susceptible to respiratory distress, thermal instability, infection, jaundice, and
43 hypoglycaemia [2]. Although delivery in a health facility can reduce the risk of neonatal
44 mortality by 29% in low and middle income countries [3], facilities must have the infrastructure,
45 capacity, and resources to treat small and sick neonates for such reductions in mortality to be
46 realized [4,5]. Healthcare facilities must be able to provide neonatal care (service availability)
47 through the presence of essential infrastructure, functioning equipment, supplies, medicine,
48 trained staff, and current care guidelines (service readiness) [5]

49 Over 91% of deliveries in Malawi occur in a health facility [6]. However, improvements are
50 needed in health facility service availability and service readiness to provide level-2 neonatal
51 care [7], particularly in rural areas where 83% of Malawi's population live [8] and where 85% of
52 healthcare facilities in Malawi are located [5].

53 To address limitations in the capacity to treat neonates with respiratory distress, a quality
54 improvement program (QIP) for neonatal care in Malawi was implemented in 2013-2015 to
55 introduce the use of bubble Continuous Positive Airway Pressure (CPAP) at all government
56 district and central hospitals in Malawi [9]. In 2016 the CPAP QIP was extended to 8 healthcare
57 facilities operated by the Christian Health Association of Malawi (CHAM), the largest non-

58 governmental healthcare provider and trainer of healthcare providers in Malawi. Equipment to
59 support respiratory care (CPAP machines, oxygen concentrators, suction machines, pulse
60 oximeters, all necessary disposable supplies, a storage cabinet, and wall job aids) was installed;
61 training on how to use equipment was conducted and supportive supervision and mentorship
62 were provided [10,11]. This was followed by a reduction in neonatal mortality, especially that
63 caused by Respiratory Distress Syndrome (RDS) [9]. However, after conducting detailed needs
64 assessments at each facility, it became increasingly clear that many neonatal units in rural areas
65 lacked other essential equipment and infrastructure, and that further mortality reductions could
66 only be achieved by taking a broader view of the overall challenges experienced by district-level
67 hospitals. Therefore, in 2017-2018, a ward-strengthening program was implemented to provide a
68 bundle of neonatal care services in a dedicated, well equipped clinical space with training and
69 mentorship extended to all relevant hospital technicians and maintenance staff, as well as clinical
70 staff. The aim of this study was to determine if the ward strengthening program improved
71 outcomes for neonates treated with CPAP at these facilities. Here, we describe the process and
72 outcome of this ward strengthening program.

73 **Methods**

74 **Needs Assessments**

75 In response to the challenges experienced during the CPAP QIP, detailed neonatal ward
76 assessments were done in 2017 at all government district hospitals and CHAM facilities to
77 determine specific challenges and limitations at each site. These assessments documented
78 demand and capacity (number of monthly admissions, number of beds, dimensions of neonatal
79 unit), staffing levels (number of nurses and doctors allocated to the neonatal unit or shared

80 between units), and equipment inventory (functioning and non-functioning equipment allocated
81 to the neonatal unit). Facilities lacking essential equipment and/or suitable space for a neonatal
82 unit were selected for inclusion in a ward strengthening program.

83 We also assessed neonatal training programs at fifteen institutions that train nurses and
84 clinicians to ensure sustainability in the transfer of knowledge and skills. The assessment
85 showed that, despite the presence of neonatal care concepts in training curricula at all
86 institutions, there were gaps in teaching these concepts, particularly due to poorly furnished
87 skills labs. All fifteen training institutions were included in a program to strengthen skills labs
88 for neonatal care.

89 **Ward Strengthening Program**

90 From 2017-2018, a ward strengthening program was implemented at 12 government district
91 hospitals and CHAM facilities to improve the care of sick neonates. Ward strengthening
92 activities included the distribution of a bundle of equipment (Table 1), supplemental training
93 support, and, in some cases, health facility renovations (Table 2).

94

95

96 **Table 1. Equipment bundle and quantity by number of beds**

Item	Quantity
CPAP devices*	2 per unit
Phototherapy lights	1 per 10 cots
Radiant warmer	1 per 10 cots
Glucometer	2 per unit
Glucometer sticks (50/pack)	3 per unit
Ambu Bag	1 per unit + 1 per 10 cots, 2 sizes
Room heater	2 per unit
Room thermometer	1 per unit + 1 in Labour Ward and Theater
Baby thermometer	3 per unit
Oxygen concentrator	1 per unit + 1 per 10 cots
Flow splitter	1 per concentrator
Digital scale	1 per unit
Suction machines	1 per unit
Surge protector	1 for all electrical equipment
Pulse oximeter	1 per ward + 1 spare

* Provided prior to ward strengthening

97

98 **Table 2. Summary of Renovations by Facility**

Hospital	Ward Strengthening Activities	Completion of Ward Strengthening	Bed Capacity of New Unit	No. of Staff trained
A	Renovation and Equipment Installation	11/2017	23	20
B	Renovation and Equipment Installation	10/2017	15	21
C	Renovation and Equipment Installation	7/2017	29	20
D	Renovation and Equipment Installation	11/2017	23	20
E	Renovation and Equipment Installation	11/2017	20	20
F	Renovation and Equipment Installation	1/2018	11	21
G	Renovation and Equipment Installation	11/2017	20	21
H	Renovation and Equipment Installation	1/2018	23	37
I	Equipment installation only	8/2017	-	21
J	Renovation and Equipment Installation	11/2017	23	19
K	Renovation and Equipment Installation	8/2017	14	19
L	Equipment installation only	8/2017	-	15

99

100

101

102 Ten of the 12 facilities lacked a separate or suitable neonatal unit. At these sites, the facility
103 management team worked with the Ministry of Health (MOH) to identify a room or space for
104 renovation, quantify material needs for the renovation, obtain quotations for the materials, and
105 develop a renovation work plan. Renovations varied by facility, but generally included the
106 expansion or partitioning of an existing space, plumbing installation and repairs, installation of
107 lighting and electrical sockets, roofing and ceiling installation and repairs, and painting. For
108 nine of the ten sites where renovations were completed, renovation costs ranged from \$1,300 -
109 \$8,600 USD and were covered by project funds. At the remaining site, the cost of construction of
110 a new ward (\$38k USD) was supplemented by private donor funds.

111 At each site, equipment was distributed according to the bed capacity of the unit (Table 1); for a
112 unit with 20 cots the bundle of equipment was approximately \$14k USD. Before equipment
113 installation, clinical staff and biomedical technicians completed half-day user training courses to
114 ensure familiarity with the operation, troubleshooting and maintenance of the equipment. These
115 trainings were conducted with the Ministry of Health Physical Asset Management (PAM) units
116 to allow biomedical technicians to receive training as well as assist with the instruction and
117 installation process at each site. Costs associated with trainings included travel costs for the
118 instructors and per diems for the trainees. The number of staff trained at each facility is
119 summarized in Table 2.

120 In collaboration with the Paediatric and Child Health Association of Malawi (PACHA), training
121 of the nurses and clinicians was delivered at the facilities using the Care of the Infant and
122 Newborn (COIN) course. The COIN course, which is used nationally, focuses on diagnosis and
123 triage of sick neonates, care at birth for all neonates, neonatal resuscitation, routine postnatal

124 care of the neonate, emergency assessment and treatment of neonates, and identification of
125 neonates in need of referral and safe transport [12].

126 After the completion of the ward strengthening program, follow-up ward assessments were
127 completed at each healthcare facility. The follow-up ward assessments were compared to those
128 conducted prior to the ward strengthening program to evaluate changes in demand, capacity, and
129 staffing levels.

130 **Program to Strengthen Skills Labs for Neonatal Care**

131 Fifteen health training institutions received support to establish skills labs for neonatal care,
132 including equipment and teaching materials to facilitate hands-on teaching of essential skills in
133 the care of neonates. Materials provided included training models, oxygen concentrators,
134 glucometers, CPAP machines, suction machines, phototherapy lights, and teaching aids.
135 Academic staff from these institutions also received training to encourage use of these resources
136 within their institutions.

137 **Study Setting**

138 The impact of the ward strengthening program was evaluated at 12 district and CHAM hospitals
139 located in rural areas (Table 2) where CPAP was being used to treat respiratory distress for one
140 year before, and one year after the implementation of the ward strengthening program. All
141 neonates with known outcomes (i.e. died or survived to discharge) treated with CPAP for
142 respiratory distress at each facility were included in this analysis. Clinical diagnoses were made
143 in accordance with national guidelines for newborn care in Malawi [12]. Because gestational
144 age is often unknown in our setting, a simple, validated algorithm using a combination of vital

145 signs, tone and birth weight was implemented to assist nurses in determining the need for CPAP
146 [13,14]. As previously described [9], patient details including dates of birth and admission,
147 admission weight, discharge diagnosis, and outcome were collected from Acute Respiratory
148 Illness (ARI) forms for each neonate admitted with respiratory illness. An on-site ARI
149 coordinator ensured completion of forms for every qualifying patient. De-identified ARI forms
150 were scanned monthly, and quarterly chart audits were conducted by the MOH ARI team to
151 ensure completion of ARI forms for all patients. Patient data and survival data were compared
152 for neonates treated before and after ward strengthening. Neonates with unknown outcomes,
153 including those that left against medical advice or were transferred to another facility, were
154 excluded from analysis. Rate of survival was defined as the fraction of eligible neonates with
155 known outcomes who survived to discharge.

156 **Statistical Analysis**

157 Differences in transfer rates and survival before and after ward strengthening were compared
158 using a two-sided Fisher's exact test. Differences between continuous variables (i.e., weight and
159 age on admission) were assessed using a two-sided t-test for equality of means (unequal
160 variances assumed). Results were considered significant at the 5% level. Data were analyzed in
161 Excel.

162 **Ethics statement**

163 The protocol was approved by the National Health Sciences Research Committee (NHSRC
164 #1180) of Malawi and the Institutional Review Board of Rice University (13-102X). De-
165 identified patient information was collected from standard Ministry of Health Acute
166 Respiratory Illness forms.

167 **Results**

168 Table 3 summarizes the patient data, outcomes, and survival rates for neonates treated with
169 CPAP one year before and after ward strengthening. A total of 207 neonates were treated with
170 CPAP before ward strengthening, and 254 were treated after ward strengthening. Among
171 neonates with known outcomes, 189 were treated with CPAP before ward strengthening, and
172 232 were treated with CPAP after ward strengthening. There was no significant difference in the
173 number of neonates who transferred out of these facilities before and after ward strengthening,
174 and there was no significant difference in the number of neonates who left against medical
175 advice before and after ward strengthening. There were no significant differences in the rates of
176 diagnosis of birth asphyxia, RDS, pneumonia, and meconium aspiration before and after ward
177 strengthening. The percentage of neonates with known outcomes and a final diagnosis of sepsis
178 significantly decreased after ward strengthening (12.2% vs 4.7%, $p=0.01$).

179

180 **Table 3: Patient data for neonates treated with CPAP for one year before and after ward**
 181 **strengthening**


182
 183
 184

	Before Ward Strengthening	After Ward Strengthening	p-value
Number of Study Participants	207	254	
Outcome			
Died	48.8%	39.0%	0.038
Discharged	42.5%	52.4%	0.039
Transferred	5.3%	3.5%	0.368
Left AMA	3.4%	3.9%	0.809
Unknown	0.0%	1.2%	0.256
Number of Neonates With Known Outcome (Died/Discharged)			
	189	232	
Outcome			
Died	53.4%	42.67%	0.031
Discharged	46.6%	57.33%	0.031
Diagnosis*			
Birth Asphyxia	15.9%	9.9%	0.077
RDS	68.8%	66.8%	0.677
Pneumonia	9.5%	9.1%	0.868
Meconium Aspiration	5.3%	4.7%	0.825
Sepsis	12.2%	4.7%	0.007
No Diagnosis	1.6%	1.7%	1.000
Admission Weight			
<1.00 kg	5.8%	3.45%	0.345
1.00-1.49 kg	37.0%	40.5%	0.483
1.50-1.99 kg	23.3%	23.3%	1.000
2.00-2.49 kg	9.0%	11.2%	0.519
2.50-4.00 kg	21.7%	16.8%	0.214
>4.00 kg	1.1%	1.7%	0.695
Unknown	2.1%	3.0%	1.000
Admission Temperature			
<32.0 °C	0.0%	0.86%	0.504
32.0-34.4 °C	22.8%	24.6%	0.730
34.5-35.4 °C	16.9%	23.7%	0.092
35.5-36.4 °C	27.5%	21.6%	0.171
36.5-37.5 °C	13.8%	13.4%	1.000
>37.5 °C	7.9%	7.8%	1.000
Unknown	11.1%	8.2%	0.321

185 *Some neonates received multiple diagnoses

186

187 Following implementation of the ward strengthening program, there was a significant increase in
188 survival for neonates treated with CPAP (Fig 1A), for the subset of neonates treated with CPAP
189 weighing between 1.00 – 2.49 kg on admission and diagnosed with respiratory distress syndrome
190 (RDS) (Fig 1B), as well as for the subset of neonates treated with CPAP weighing between 1.00
191 – 1.49 kg on admission and diagnosed with respiratory distress syndrome (RDS) (Fig 1C).
192 Before ward strengthening, 46.6% of neonates treated with CPAP survived to discharge; the rate
193 of survival improved to 57.3% after ward strengthening ($p=0.03$). Survival was also compared in
194 the subset of low birth weight neonates diagnosed with RDS and weighing 1.00 – 2.49 kg and
195 1.00 – 1.49 kg on admission; in these groups, survival increased from 39.4% before ward
196 strengthening to 60.3% after ward strengthening ($p=0.001$), and from 25.0% before ward
197 strengthening to 44.3% after ward strengthening ($p=0.03$).



198
199 **Fig 1.** (A) Outcomes for all neonates admitted with respiratory distress and treated with CPAP
200 for one year before and after ward strengthening. There was a significant increase in survival

201 after ward strengthening (46.6% and 57.3%, respectively, $p=0.03$). (B) Outcomes for the subset
 202 of neonates treated with CPAP weighing between 1.00 – 2.49 kg on admission and diagnosed
 203 with respiratory distress syndrome (RDS). There was a significant increase in survival after ward
 204 strengthening (39.4% and 60.3%, respectively, $p=0.001$). (C) Outcomes for the subset of
 205 neonates treated with CPAP weighing between 1.00 – 1.49 kg on admission and diagnosed with
 206 respiratory distress syndrome (RDS). There was a significant increase in survival after ward
 207 strengthening (25.0% and 44.3%, respectively, $p=0.03$).

208 The admission weights, admission temperatures, and age on admission for neonates treated
 209 before and after ward strengthening are compared in Table 4. Results are shown for all neonates
 210 treated with CPAP, and for the subset with admission weights between 1.00 – 2.49 kg and
 211 diagnosed with RDS. There were no significant differences in the distribution of patient
 212 variables before and after ward strengthening, with the exception of a decrease in mean
 213 admission temperature for neonates weighing between 1.00 – 2.49 kg with a diagnosis of RDS (p
 214 = 0.01).

215 **Table 4: Comparison of average weights, temperatures, and age on admission for neonates**
 216 **treated one year before and after ward strengthening.**

	All Admission Weights			Admission Weights 1-2.49 kg, Diagnosed with RDS		
	Before Ward Strengthening	After Ward Strengthening	p- value	Before Ward Strengthening	After Ward Strengthening	p- value
Average Admission Weight (kg)	1.84 +/- 0.81	1.84 +/- 0.96	0.99	1.48 +/-0.34	1.48 +/-0.34	0.87
Average Admission Temperature (°C)	35.5 +/- 1.6	35.4 +/- 1.6	0.48	35.4 +/- 1.5	34.9 +/- 1.3	0.01

Average Age on Admission (day)	2.0 +/- 5.1	2.1 +/- 5.9	0.75	0.7 +/- 3.1	0.4 +/- 2.1	0.48
--------------------------------	-------------	-------------	------	-------------	-------------	------

217
 218 Table 5 compares observations from the initial ward assessments completed in 2017 to the
 219 follow-up ward assessments completed in 2018. These assessments indicated that admissions
 220 increased and staffing levels improved following ward strengthening. Overall, average nursery
 221 admissions at the 12 facilities included in this analysis increased from 39.2 neonates per month
 222 in 2017 to 74.9 neonates per month in 2018. Additionally, the total number of nurses allocated
 223 specifically for neonatal care across these facilities increased for both the day and night shifts.
 224 Overall, the median number of nurses allocated specifically to care for neonates during the day
 225 increased from 1 to 1.5, and the median number of nurses allocated to neonatal care overnight
 226 increased from 0.5 to 1. The number of hospitals with no nurses allocated specifically to care for
 227 neonates during the day remained the same before and after ward strengthening (3 hospitals); the
 228 number of hospitals with no nurses allocated specifically to care for neonates during the night
 229 decreased from 6 to 3. At the conclusion of this program, at least one clinician was assigned to
 230 every neonatal ward included in this analysis.

231
 232 **Table 5: Comparison of ward assessments for average monthly admissions and nurse**
 233 **allocations before and after ward strengthening.**

	Before Ward Strengthening	After Ward Strengthening
Average Monthly Neonatal Admissions (min, max)	39.2 (15, 70)	74.9 (15, 140)
Median Number of Nurses Assigned to Neonatal Care (Day Shift) (min, max)	1 (0, 2)	1.5 (0, 6)
Median Number of Nurses Assigned to Neonatal Care (Night Shift) (min, max)	0.5 (0, 1)	1 (0, 2)

235 **Discussion**

236 We previously reported that implementation of a nurse-led CPAP QIP to treat neonates with
237 respiratory illness in all government district and central hospitals in Malawi led to a significant
238 increase in survival, particularly among neonates of low birth weight [9]. Following the CPAP
239 QIP, survival for neonates with admission weights of 1.00 – 2.49 kg and diagnosed with RDS
240 increased from 39.8% to 48.3% after the implementation of CPAP (p=0.042) [9]. However,
241 improvements in survival fell short of those experienced during a previous quasi-randomized
242 efficacy study of CPAP vs standard nasal oxygen in the neonatal unit of the referral hospital of
243 southern Malawi, where neonates with admission weights of 1.00 – 2.49 kg and diagnosed
244 with RDS experienced survival rates of 64.6% when treated with CPAP [15].

245 During mentorship and supervisory visits to rural and district hospitals, many challenges were
246 identified that likely adversely affected overall survival as well as limited the impact of
247 improved respiratory care. Despite provision of supportive respiratory equipment, training, and
248 peer mentorship along with the CPAP devices, additional equipment, training, and space were
249 necessary to allow more comprehensive care of vulnerable neonates, who often have other
250 complications in addition to respiratory distress. After the ward strengthening program was
251 introduced, survival rates for neonates weighing 1.00-2.49 kg with a diagnosis of RDS
252 improved from 39.4% to 60.3%, approaching the 64.6% survival previously seen in a referral
253 hospital [15].

254 To identify facility-level barriers to improved neonatal care, a needs assessment was undertaken
255 in each hospital by members of their own administrative, clinical and technical/maintenance
256 staff, together with MOH representatives. Together, MOH representatives and research staff

257 determined which parameters to evaluate; only parameters that directly impacted the quality of
258 newborn care were prioritized to avoid burdening hospital staff and prevent disruption of work
259 flow during the assessments. Since hospital staff were familiar with the parameters being
260 evaluated, assessments were completed as part of routine visits to the facilities and no intensive
261 training was required. Building modifications and equipment and staffing needs were thus
262 directed by local requirements. Importantly, this gave ownership and investment to local hospital
263 staff and in itself strengthened interdepartmental relationships.

264 The most common requirement was that of space. Ten of the twelve facilities did not have a
265 separate neonatal ward; if a separate area was available, it was too small or otherwise unsuitable.
266 The absence of a neonatal ward is common in hospitals in many countries [16,17]. As part of the
267 Every Newborn Action Plan process, an analysis of 12 countries in Africa and Asia with the
268 highest burden of neonatal mortality showed that five countries reported limited or no dedicated
269 neonatal space in healthcare facilities [16]. A recent qualitative assessment of health facilities in
270 three regions of Ethiopia from December 2017 to February 2018 found the lack of space for
271 neonates in need of special care across all levels of the health care system [17]. This assessment
272 found that neonates were sometimes relocated to adult and surgical wards due to the lack of
273 space [17]. A dedicated space is essential for the provision of infection control and temperature
274 control for small and sick neonates. Following the implementation of the ward strengthening
275 program presented here, the proportion of neonates receiving CPAP who were diagnosed with
276 sepsis decreased from 12.2% to 4.7%; further investigation is needed to determine if this
277 decrease was a result of improved diagnoses or improved infection control.

278 The care of sick neonates also requires specialized training and skills. Neonatal nursing training
279 programs are rare in low-income countries [16], and a high turnover of trained healthcare

280 providers is a problem in many settings [9,17]. We tried to address these barriers by conducting
281 trainings not only at health facilities where equipment was installed, but also at 15 nursing
282 schools and clinical officer training institutions. Most of the training institutions did not have the
283 basic equipment necessary for practical instruction in neonatal care. Strengthening existing pre-
284 service training programs is necessary to address the high staff turnover due to rotations and
285 attrition.

286 We note that improved space, equipment, and training had important additional beneficial
287 effects. Previously, sick and healthy neonates often remained with their mother in the postnatal
288 ward, but the creation of a separate, well-equipped ward for sick neonates meant that staff were
289 allocated specifically for their care; overall, nurse allocations to the neonatal wards increased and
290 at least one clinician was assigned to the neonatal ward in each facility.

291 As this observational study involved neonates undergoing routine care in a low resource setting,
292 there were some unavoidable limitations. The primary outcome of this study was survival to
293 discharge as follow-up after discharge was not feasible. Few if any neonates had a chest x-ray to
294 diagnose RDS, and gestational age as determined by a first trimester ultrasound was rarely
295 available. Consequently, RDS was diagnosed clinically with the aid of an algorithm to help
296 identify neonates who might benefit the most from CPAP. Although retinopathy of prematurity
297 was addressed during trainings [11], we did not monitor for it during this study. However, in a
298 review of reports from LMICs on the use of CPAP, two studies that looked for retinopathy of
299 prematurity did not find it [18], and we did not encounter it in the early phases of our program
300 [15]. Adverse events such as pneumothoraces and nasal trauma were also not tracked as part of
301 this study, but anecdotally, they were uncommon and limited to nasal bleeds or nasal hyperemia.

302

303 **Learning from our experience**

304 When reflecting on our program, it is clear that efforts to improve the quality of neonatal care
305 must be built on a solid foundation of good basic clinical care that is based on a holistic approach
306 to the needs of a neonate. Improvement in one aspect of need (such as respiratory support) will
307 falter without the provision of warmth, nutrition, infection control and family-centered care
308 given by trained staff in a dedicated space. A needs assessment is essential to identify gaps in
309 overall care so that these can be addressed with an appropriate bundle of interventions to
310 improve care for all infants in the ward. Provision of materials is insufficient without training,
311 supervision, and mentoring to ensure that equipment is used properly and protocols followed
312 correctly.

313 Some challenges are beyond the jurisdiction of hospital management to solve: an unreliable
314 power system and irregular water supply are common and affect many of the facilities included
315 in this program. While power and water outages were not systematically tracked during this
316 audit, power outages were noted on the charts of 16.4% of neonates treated before, and 15.9%
317 treated after ward strengthening. Furthermore, availability of essential commodities and
318 consumables is a challenge if hospitals lack the resources or ability to procure them.

319 It is encouraging to see that the physical changes made to the neonatal ward areas led to
320 improved staff allocation to work in them. In initial discussions about possible space
321 renovations, MOH representatives strongly advocated that each facility dedicate staff solely to
322 planned new units to improve quality of care. Based on our observations, better staffing, more
323 training and mentorship have improved morale, and by planning and creating the changes

324 together, the hospital departments are more interactive. All of these factors have played a part in
325 reducing neonatal mortality.

326 **Conclusion**

327 To improve neonatal outcomes, healthcare facilities need the infrastructure, capacity, and
328 resources to treat small and sick neonates. This means having a family-centred approach to care
329 encompassing the provision of warmth, nutrition and infection control given by trained staff in a
330 dedicated space with functioning equipment. Providing CPAP within such a framework
331 significantly improved survival for neonates with respiratory distress. These improvements in
332 service availability lead to improved staff morale, increased the number of dedicated neonatal
333 staff, and most importantly, improved outcomes for admitted sick and small neonates treated
334 with CPAP.

335 **Acknowledgments**

336 We would like to acknowledge PACHA (Paediatric and Child Health Association) for their
337 leadership in the development and provision of the COIN training course. We would also like to
338 acknowledge all of the staff of the CPAP Project for their contributions to data collection,
339 monitoring, and training associated with the quality improvement initiative, as well as our CPAP
340 trainers and mentors for leading CPAP training sessions and providing mentorship and
341 supportive supervision within the hospitals.

342 **Contributors**

343 NL, AA, KK, RRK, MO, EM, and AA designed the study. JC, SLM, SN, and AA contributed to
344 data collection and/or data and statistical analysis. JC, SLM, AA, and EM prepared the first draft

345 of the manuscript. All authors contributed to critical interpretation of the results and development
346 of the manuscript and approved the final version.

347 **Funding**

348 This work is made possible through the generous support of ELMA Philanthropies through a
349 grant to the University of Malawi College of Medicine (<https://www.elmaphilanthropies.org/>,
350 Grant No. 16-F0007). This manuscript is also made possible through the generous support of the
351 Saving Lives at Birth ([https://www.usaid.gov/global-health/health-areas/maternal-and-child-](https://www.usaid.gov/global-health/health-areas/maternal-and-child-health/projects/saving-lives-birth-a-grand-challenge)
352 [health/projects/saving-lives-birth-a-grand-challenge](https://www.usaid.gov/global-health/health-areas/maternal-and-child-health/projects/saving-lives-birth-a-grand-challenge)) partners: The United States Agency for
353 International Development (USAID), the Government of Norway, the Bill & Melinda Gates
354 Foundation, Grand Challenges Canada, and the UKAID (Grant No. AID-OAA-A-13-00014). It
355 was prepared by Rice University and does not necessarily reflect the views of the Saving Lives at
356 Birth partners. Additionally, this work was supported by the John D. and Catherine T.
357 MacArthur Foundation (<https://www.macfound.org>, Grant No. 17-1709-152484) the Bill &
358 Melinda Gates Foundation (<https://www.gatesfoundation.org>, Grant No. OPP1193531), ELMA
359 Philanthropies (<https://www.elmaphilanthropies.org>, Grant No. 19-F0012), The Children's
360 Investment Fund Foundation UK (<https://ciff.org>, Grant No. R-1810-03159), The Lemelson
361 Foundation (<https://www.lemelson.org>, Grant No. 18-01458), and the Ting Tsung and Wei Fong
362 Chao Foundation under agreements to William Marsh Rice University.

363 References

- 364 1. World Health Organisation. Every Newborn Action Plan. 2014. Available:
365 www.who.int/about/licensing/copyright_form/en/index.html
- 366 2. Maynard KR, Causey L, Kawaza K, Dube Q, Lufesi N, Maria Oden Z, et al. New technologies for
367 essential newborn care in under-resourced areas: what is needed and how to deliver it. *Paediatr*
368 *Int Child Health*. 2015;35: 192–205. doi:10.1179/2046905515Y.0000000034
- 369 3. Tura G, Fantahun M, Worku A. The effect of health facility delivery on neonatal mortality:
370 systematic review and meta-analysis. *BMC Pregnancy Childbirth*. 2013;13: 18. doi:10.1186/1471-
371 2393-13-18
- 372 4. Dickson KE, Simen-Kapeu A, Kinney M V, Huicho L, Vesel L, Lackritz E, et al. Every Newborn:
373 health-systems bottlenecks and strategies to accelerate scale-up in countries. *Lancet*. 2014;384:
374 438–454. doi:[https://doi.org/10.1016/S0140-6736\(14\)60582-1](https://doi.org/10.1016/S0140-6736(14)60582-1)
- 375 5. Winter R, Yourkavitch J, Wang W, Mallick L. Assessment of health facility capacity to provide
376 newborn care in Bangladesh, Haiti, Malawi, Senegal, and Tanzania. *J Glob Health*. 2017;7: 20509.
377 doi:10.7189/jogh.07.020509
- 378 6. National Statistical Office of Malawi. Malawi Demographic and Health Survey. *Natl Stat Off DHS*
379 *Progr*. 2017; 1–658. doi:<http://www.measuredhs.com/>
- 380 7. WHO, UNICEF. Ending Preventable Newborn Deaths and Stillbirths by 2030. 2020.
- 381 8. The World Bank. Rural Population, Malawi. 2018. Available:
382 <https://data.worldbank.org/indicator/SP.RUR.TOTL.ZS?locations=MW>
- 383 9. Carns J, Kawaza K, Liaghati-Mobarhan S, Asibon A, Quinn MK, Chalira A, et al. Neonatal CPAP for
384 Respiratory Distress Across Malawi and Mortality. *Pediatrics*. 2019; e20190668.
385 doi:10.1542/peds.2019-0668
- 386 10. Asibon A, Lufesi N, Choudhury A, Olvera S, Molyneux E, Oden M, et al. Using a peer mentorship
387 approach improved the use of neonatal continuous positive airway pressure and related
388 outcomes in Malawi. *Acta Paediatr*. 2019;n/a. doi:10.1111/apa.15025
- 389 11. NEST360 Clinical Modules. Available: <https://nest360.org/project/clinical-modules/>
- 390 12. Malawi Ministry of Health. Care of the infant and newborn in Malawi (2017): the COIN Course -
391 Participants Manual. 2017. Available: [https://research-repository.st-
392 andrews.ac.uk/handle/10023/12839](https://research-repository.st-andrews.ac.uk/handle/10023/12839)
- 393 13. Crehan C, Colbourn T, Heys M, Molyneux E. Evaluation of 'TRY': an algorithm for neonatal
394 continuous positive airways pressure in low-income settings. *Arch Dis Child*. 2018;103: 732 LP –
395 738. doi:10.1136/archdischild-2017-313867
- 396 14. Hundalani SG, Richards-Kortum R, Oden M, Kawaza K, Gest A, Molyneux E. Development and
397 validation of a simple algorithm for initiation of CPAP in neonates with respiratory distress in
398 Malawi. *Arch Dis Child - Fetal Neonatal Ed*. 2015;100: F332–F336. doi:10.1136/archdischild-2014-
399 308082
- 400 15. Kawaza K, Machen HE, Brown J, Mwanza Z, Iniguez S, Gest A, et al. Efficacy of a low-cost bubble
401 CPAP system in treatment of respiratory distress in a neonatal ward in Malawi. *PLoS One*. 2014;9.
402 doi:10.1371/journal.pone.0086327

- 403 16. Moxon SG, Lawn JE, Dickson KE, Simen-Kapeu A, Gupta G, Deorari A, et al. Inpatient care of small
404 and sick newborns: a multi-country analysis of health system bottlenecks and potential solutions.
405 BMC Pregnancy Childbirth. 2015;15: S7. doi:10.1186/1471-2393-15-S2-S7
- 406 17. Usman AK, Wolka E, Tadesse Y, Tariku A, Yeshidinber A, Teklu AM, et al. Health system readiness
407 to support facilities for care of preterm, low birth weight, and sick newborns in Ethiopia: a
408 qualitative assessment. BMC Health Serv Res. 2019;19: 860. doi:10.1186/s12913-019-4672-2
- 409 18. Thukral A, Sankar MJ, Chandrasekaran A, Agarwal R, Paul VK. Efficacy and safety of CPAP in low-
410 and middle-income countries. J Perinatol. 2016;36: S21. Available:
411 <https://doi.org/10.1038/jp.2016.29>
412