

1 **Title**

2 Relationship between high rates of intestinal parasitic infections and knowledge,
3 attitudes, and practices of Ndelele Health District populations (East Region, Cameroon):
4 a cross-sectional mixed approach

6 **Short Title**

7 Epidemiology of intestinal parasitic infections in the landlocked areas of Eastern
8 Cameroon

10 **Authors and Affiliations**

11 Viviane Ongbassomben¹, Cyrille Ndo^{1,2,3,*}, Ericka A. Lebon⁴, Hugues C. Nana Djeunga⁵,
12 Albert L. Same Ekobo⁶ and Dieudonné Adiogo¹

13
14 ¹Department of Biological Sciences, Faculty of Medicine and Pharmaceutical Sciences,
15 University of Douala, P.O. Box 24157, Douala, Cameroon. ²Department of Parasitology
16 and Micorbiology, Centre for Research in Infectious Disease (CRID), P.O. Box 13591,
17 Yaoundé, Cameroon. ⁴Solutions, Risk Management and Regulatory Framework
18 Information Technology, P.O. Box 5551014, Montréal, Canada. ⁵Molecular Parasitology
19 and Genetic Epidemiology (MPGE) Department, Centre for Research on Filariasis and
20 other Tropical Diseases (CRFiMT), P.O. Box 5797, Yaoundé, Cameroon. ⁶Faculty of
21 Medicine and Biomedical Sciences, University of Yaoundé I, P.O. Box 337, Yaoundé,
22 Cameroon.

23

24 *Correspondence: Email: cyrndo@yahoo.fr; Phone: +237 699262003 / +237 677561088;

25 P.O. Box 13591, Yaoundé, Cameroon.

26

27

28 **Abstract**

29 **Background:** Intestinal parasitic infections remain of public health concern worldwide,
30 especially among rural and poorest populations as a consequence of precariousness,
31 lack of sanitation, non-availability of potable water and poor hygiene conditions. The
32 present study aimed to better understand the epidemiology of intestinal parasitic
33 infections in rural areas of forested Cameroon.

34

35 **Methodology:** A cross-sectional survey was conducted in three Health Areas (Ndelele,
36 Kentzou and Lolo) of the Ndelele Health District (East Region, Cameroon). Information
37 on socio-demographic characteristics, knowledge, attitudes, and practices regarding
38 intestinal parasitic infections were collected using a semi-structured questionnaire. Stool
39 samples were collected and analyzed by the Kato-Katz and formalin-ether concentration
40 techniques to complement simple direct examination.

41

42 **Principal Findings:** A total of 406 individuals belonging to three main groups (Kako or
43 Bantu, Baka or Pygmies and Central African Republic refugees) were enrolled in the
44 study. The overall intestinal parasitic infection rate was 74.9%, including 57.2% cases of
45 polyparasitism. Fourteen parasite species were identified, 89.1% being intestinal

46 protozoa and 41.8% belonging to helminths. Infections with helminths were associated
47 with Baka ($P < 0.0001$). Spring water consumption was associated with hookworm
48 infection (OR = 3.87; $P = 0.008$). Garbage deposited near houses was positively
49 associated with infection with *Giardia lamblia* (OR = 3.41; $P = 0.003$). Polyparasitism was
50 positively linked to washing hand without soap before meal (OR= 11.64; $p= 0.002$).

51
52 **Conclusion/Significance:** Intestinal parasitic infections exhibited high rates in the
53 Ndelele Health District, especially among indigenous and hard-to-reach populations
54 (Pygmies). Hygiene measures appear as the main drivers sustaining transmission, and
55 targeted strategies should be developed to efficiently fight against these debilitating
56 diseases.

57
58 **Keywords:** Intestinal parasitic infection, KAP, Ndelele Health District, East Region,
59 Cameroon.

62 **Author summary**

63 Intestinal parasitic infections such as soil transmitted helminthiasis and schistosomiasis,
64 remain a public health concern in Cameroon. To better understand the epidemiology of
65 these infections in hard-to-reach populations in rural areas, a cross-sectional study was
66 carried out in three health areas of Ndelele health District in the Eastern- Cameroon. The
67 rate of intestinal parasitic infections was high (74.9%) especially in indigenous
68 populations (Baka-Pygmies) which exhibited high prevalence of soil transmitted helminth

69 infections (83.3%). The study confirmed that the lack of sanitation and poor hygiene
70 largely contribute to the endemicity of intestinal parasitic infections, particularly among
71 indigenous populations. Targeted control strategies seem mandatory to reach these
72 populations and offer them appropriate care to interrupt or reduce the transmission of
73 these diseases.

74

75

76 **Introduction**

77 Intestinal parasitosis are infections caused by intestinal helminths (soil transmitted
78 helminth or *Schistosoma spp* infections) and protozoa. They are widespread in
79 developing countries, particularly in the tropics. Absence of sanitation networks, lack of
80 drinking water, poor public and private hygiene standards, and insufficient health
81 education exacerbate infection with eggs, larvae and cysts through contact with
82 contaminated soil, food or water [1-2]. More than 1.5 billion people representing 24% of
83 the world's population are infected with soil-transmitted helminth infections worldwide [3].
84 Estimates show that at least 290.8 million people required preventive treatment for
85 schistosomiasis in 2018, out of which more than 97.2 million have received appropriate
86 treatment [4]. The main species of soil-transmitted helminths that infect humans are
87 roundworm (*Ascaris lumbricoides*), whipworm (*Trichuris trichiura*) and hookworms
88 (*Necator americanus* and *Ancylostoma duodenale*) [5]. The most important intestinal
89 protozoan infections are amoebiasis (*Entamoeba histolytica*), giardiasis (*Giardia lamblia*),
90 cryptosporidiosis (*Cryptosporidium sp*), cyclosporiasis (*Cyclospora cayetanensis*) and
91 isosporiasis (*Isospora belli*) [6]. Children of pre-school and school ages are the most

92 affected by intestinal parasitosis [7-8]. Over 270 million preschool-age children live in
93 areas where these diseases are intensively transmitted [9]. Helminths negatively
94 influence the nutritional status of children and adolescents as they inhibit absorption of
95 nutrients, leading to stunted growth [10]. *Cryptosporidium spp.* is most common in
96 children with diarrhea, followed by *G. lamblia* and *E. histolytica* [11-12]. The strategy
97 deployed by the World Health Organization (WHO) to control soil-transmitted helminth
98 infections is periodic mass administration of Albendazole or Mebendazole to at-risk
99 populations (pre-school age children, school age children, women of reproductive age,
100 adults in certain high-risk occupations such as tea-pickers or miners) living in endemic
101 areas [3-13].

102

103 In Cameroon, intestinal parasitosis remains a public health concern. WHO report on the
104 estimation of helminthic infections in Cameroon illustrates the unequal distribution of
105 helminthiasis across the country. Schistosomiasis is highly endemic in the Northern part
106 of the country. *Ascaris lumbricoides* and *T. trichiura* are widely distributed in the Centre
107 and South Regions of the country, with the highest prevalence found in the South [14].
108 Although studies are carried out in some localities [15,16,17], updated and detailed
109 epidemiological data on intestinal parasitic infections across the country are quite scanty.
110 This is due not only to limited funding but also to the difficulty to access certain zones due
111 to very poor road network. This is especially the case of the East Region of Cameroon
112 which is very landlocked rendering implementation of epidemiological surveys difficult. As
113 consequence, there is lack of data on the epidemiology of several tropical diseases in this
114 region, and people usually feel like they are underlooked. This study was therefore

115 conducted to provide unprecedented detailed data on the epidemiology of intestinal
116 parasitic infections in a hard-to-reach Health District of the East Region of Cameroon.

117

118

119 **Methods**

120 **Ethical considerations**

121 An ethical clearance (N°1750 CEI-UDo/04/2019/M) was granted by the Institutional
122 Review Board (IRB) of the University of Douala. Thereafter, administrative authorizations
123 were delivered by the East Regional Delegate for Public Health, the Ndelele District
124 Medical Officer and the Heads of the Ndelele and Kentzou Sub-Divisions. Prior to
125 enrolment, participants were briefed about the study objectives and design. All adult
126 volunteers who agreed to participate signed a written informed consent form, while formal
127 consent was obtained from the parent or legal guardian for children under the age of 18.
128 A unique identifier was used for each participant for anonymity purpose.

129

130 **Study area and population**

131 The present study was conducted in the Ndelele Health District located in the Kadey
132 Division, East Region, Cameroon. This Health District is bordered to the North by the
133 Batouri Health District, to the South by the Yokadouma Health District, to the West by
134 the Mbang Health District and to the East by the Bombette and the Kadey river which
135 border the Central African Republic. In the study area, prevails a Guinean type tropical
136 climate divided into four seasons: a short rainy season extending from March to June,
137 a short dry season from July to August, a long rainy season from September to

138 November, and a long dry season from December to March. Vegetation is mainly made
139 up of dense forest in the Ndelele Health area, and wooded savannah sprinkled with forest
140 galleries in Kentzou and Lolo Health areas.

141
142 According to the 2019 census, the population of the Ndelele Health District, including
143 Central African refugees, was 63,319 inhabitants. This population is organized in several
144 ethnic groups, of which the most important are the Kako (Bantu), which represents a large
145 majority of the population, the Baka (Pygmies), an indigenous minority groups mostly
146 settled in the forest, and an important community of Central African Bororo refugees.
147 Economic activities of the area are mainly based on agriculture, mining, animal
148 husbandry, logging, hunting, and trading are also practiced in the Ndelele Health District
149 [18-19].

150

151 **Study design**

152 A household-based cross-sectional survey was carried out in the Ndelele Health District
153 to investigate the prevalence, intensity of infection and risk factors associated with
154 intestinal parasitic infections. All volunteers, whatever their age and gender, living in the
155 Ndelele, Kentzou and Lolo Health areas of the Ndelele Health District, were eligible to
156 this study. A semi-structured questionnaire (translated in local language by the
157 community health workers when necessary) was administered to capture their socio-
158 demographic characteristics and potential factors associated with transmission of
159 intestinal parasitic infections (WaSH indicators, gender, age groups, ethics groups and
160 health areas). One part of the questionnaire was only administered to participants over

161 14 years old to know their knowledge about the disease. The WaSH part (hand washing,
162 water supply, defecation place, barefoot walking) of children under 7 years old was
163 answered by parents. All participants with seven years or more were responding to the
164 WaSH part of the questionnaire

165

166 **Sample collection and processing**

167 After the interview, a labelled sterile 60mL plastic screw-cap vial was provided to the
168 participants (or their parents or legal guardian when they were minors), and stool samples
169 were collected the next day in the morning, between 7 and 8 am. Sample were first
170 analyzed by direct observation of fresh stool in saline (NaCl 0.9%) to search for worm
171 eggs, larvae, protozoan trophozoites and cysts. When cyst were present, an iodine wet
172 mount was performed for further their identification by staining their glycogen and nuclei
173 [20]. Two concentration techniques were subsequently performed for the diagnosis of
174 intestinal helminths [21]. Firstly and using the Kato Katz technique thick smears were
175 prepared on microscope slides using standard 41.7 mg templates. Slides were first
176 examined under microscope (magnification x100 or x400) within 30 minutes for the
177 detection of hookworm eggs, then 24 hours later for the identification of other helminth
178 eggs. When present, eggs were identified and counted, and the results expressed as
179 number of eggs per gram of stool (epg). Secondly, stool samples were processed by
180 Formalin-ether concentration technique. Indeed, a suspension of 1g of stool mixed with
181 10 mL of 10% formalin was prepared and strained through sieve in centrifugation tube;
182 formalin was added in centrifugation tube to adjust the total volume to 10 mL. After adding
183 3 mL of ether to the centrifugation tube, the tube was vortexed and then centrifuged for 3

184 min at 500g. From the resulting four layers, the three top layers were discarded, and the
185 resulting sediment was examined using bright field microscopy for detection of helminths
186 eggs and intestinal protozoa cysts [22].

187

188 **Statistical analysis**

189 Data were entered in a purpose-built Microsoft Office Excel sheet and were exported to
190 Statistical Package for the Social Sciences (SPSS) version 23 software (SPSS Inc,
191 Chicago, IL, USA) for analyses. Infection rates were expressed as the number of people
192 infected by at least one parasite in the study population. For STH specifically, intensities
193 of infection were classified according to WHO guidelines, as follows: light, moderate and
194 heavy infections when the epg was 1–1,999, 2,000–3,999 and $\geq 4,000$ for *hookworm*; 1–
195 4,999, 5,000–49,999 and $\geq 50,000$ for *A. lumbricoides*; and 1–999, 1,000–9,999 and
196 $\geq 10,000$ for *T. trichiura* [23]. Chi-square test was used to assess the difference in infection
197 rates and proportions of intensities of infection categories according to age groups,
198 genders, Heath Areas, ethnic groups and WASH indicators (use of latrine, open
199 defecation and water sources). Logistic regression model was used to investigate the
200 association between parasite infections and socio-demographic factors (gender, age,
201 ethnic group and Health Area) as well as WASH indicators adjusted with age, gender,
202 ethnicity, and health area. Threshold for significance was set at 0.05.

203

204

205

206 **Results**

207 **Socio-demographics characteristics**

208 A number of 406 participants were included in the study, (166 males and 240 females).
209 Flighty nine (14.5%) were aged <5 years (preschool-aged children), 98 (24.1%) were
210 aged 5-14 years (school-aged children) and 249 (61.4%) were aged >14 years
211 (adolescents and adults). The enrollees were undereducated, with 7.6% of participants
212 having reached high school, while most of the interviewees have never attended school
213 (34.2%) or have attended primary school (38.4%) or Koranic schools (19.7%). The
214 majority of participants (43.6%) lived in Ndelele Health area, while 26.6% and 29.8% were
215 recruited in Kentzou and Lolo Health areas, respectively. Regarding the ethnic groups,
216 most of them were Kako (47.8%) and refugees from the Central African Republic (35.5%),
217 Baka (pygmies; 11.1%) and other ethnic groups (5.7%) being underrepresented as it is
218 the case in the general population.

219

220 **WASH indicators**

221 All the interviewees declared practicing hand washing before meals (100%), though
222 mostly without soap (96.6%). Although the borehole water was the most used as drinking
223 water (63.1%), spring water consumption was significantly higher in Ndelele Health area
224 (50.3%) ($p < 0.0001$). The number of households without latrine (37.1%) and the proportion
225 of people practicing open defecation (50.3%) were significantly higher in the Ndelele
226 Health Area. As most of Baka households were without latrines (87.5%), this ethnic group
227 practiced open defecation more than other ethnic groups (88.9%) (< 0.0001) (Table 1).
228 Barefoot walking (76.3%) and open defecation (45.8%) were practiced mostly by pre-
229 school aged children ($p = 0.0001$) (Table 2).

230

231 **Knowledge of participants with regards with intestinal parasitic infections**

232 Of the 249 participants interviewed, 145 (79.7%) mentioned food as the main mode of
233 contamination of intestinal parasitic infection, with significantly higher proportion
234 ($p=0.0003$) in the Kentzou Health Area (97.6%), compared to Ndelele (76.5%) and Lolo
235 (72.2%) Health Areas. The most frequent clinical sign associated with intestinal
236 parasitosis by the interviewees were abdominal pain (39.0%), which has been mentioned
237 by 54.8% in Kentzou, 28.1% in Lolo and 23.5% in Ndelele. The other most known
238 symptoms were vomiting (31.3%), and diarrhea (11.5%). The Baka ethnic group
239 (pygmies) exhibited the worst knowledge regarding intestinal parasites.

240

241 **Participants' attitude and practice in case of suspected infection**

242 The majority of the enrollees did not use drugs for intestinal parasitic infections (57.1%).
243 Of the 174 (42.8%) people using medicines for intestinal parasitic infections treatment,
244 89 (51.1%) were using modern medicine (Mebendazole (36.2%), Metronidazole (10.3%),
245 and Albendazole (3.4%) whereas 54 (32.8%) used to use traditional medicine such as
246 bark of trees or leaves of plants. The use of "drugs" was significantly higher ($p<0.0001$)
247 in the Ndélélé Health Area (79.1%) compared to Kentzou (22.2%) and Lolo (6.6%) health
248 Areas. Baka ethnic group (Pygmies) essentially used traditional drugs in case of
249 suspected intestinal parasitic infection.

250

251 **Prevalence of intestinal parasitic infections**

252 A total of 406 enrollees provided stool samples for diagnostic of intestinal parasitic
253 infection. The overall prevalence was 74.9%, and this was significantly higher in Ndelele
254 Health Area (85.3%) compared to the other Health Areas ($p < 0.0001$). Baka ethnic group
255 was significantly more infected (93.3%) compared to the other ethnic groups ($p < 0.0001$).
256 Fourteen parasitic species were found including nine protozoa and five helminth species
257 (Table 3). Intestinal protozoa were found in 89.5% of infected people. The main protozoan
258 species found were *E. coli* (50.0%), *Blastocystis hominis* (48.0%) and *E. histolytica*
259 (26.3%). Helminths were found in 41.8% of infected people, and the main species found
260 were *A. lumbricoides* (31.2%) and *T. trichiura* (11.2%). The distribution of parasitic
261 species was significantly higher in the Ndelele Health Area (74.4%), followed by Kentzou
262 (64.3%) and Lolo (50%) (Fig 1). Prevalence of helminth and protozoa infections was
263 similar between males and females (Fig 2). School-age children were significantly more
264 infected by *A. lumbricoides* (32.7%; $p = 0.045$) whereas pre-school age children were more
265 likely infected by *Strongyloides stercoralis* (13.6%; $p = 0.001$). Regarding the ethnic
266 groups, Baka (Pygmies) were significantly more infected by helminths such as *A.*
267 *lumbricoides* (83.3%; $p < 0.0001$), *T. trichiura* (68.9%; $p < 0.0001$) and hookworm (26.7%;
268 $p < 0.0001$). Although Protozoan infections were the most common among Central African
269 refugees (97.4%; $p = 0.011$), prevalence were lower compare to other ethnics. For
270 example, Kako (autochthones) were most often infected by *E. coli* and *E. histolytica* than
271 Central African Refugees, respectively [(50.0% vs 18.1%; $p < 0.0001$) and (25.3% vs
272 11.1%; $p = 0.012$)]
273

274 Polyparasitism was found in 57.2% of enrollees, including infection with two (44.2%),
275 three (37.9%) and more than four (17.8%) parasites species. Polyparasitism was similar
276 between genders with a decreasing trend with age. It was significantly low among Central
277 African refugees (29.9%) compared to Kako (69.5%) and Baka (66.7%) ethnic groups ($p <$
278 0.0001). Regarding ethnic groups, polyparasitism was significantly higher in Ndelele
279 Health Area (65.6%; $p = 0.012$).

280

281 **Intensity of helminth infections**

282 Intensity of infection was determined only for helminth infections using Kato-Katz
283 technique. This parameter varied from 96 to 76,608 epg [median: 2200; interquartile
284 range (IQR): Q1=1000; Q3=7000] for *A. lumbricoides*, from 96 to 2,520 epg [median: 200;
285 IQR: Q1=96; Q3=475] for *T. trichiura*, and from 96 to 3,500 epg [median: 148; IQR:
286 Q1=96; Q3= 325] for hookworm. According to WHO classification, intensity of helminth
287 infections was light for almost all the enrollees (77.2%) and parasite species, *A.*
288 *lumbricoides* exhibiting only two cases of heavy infections in the Ndelele Health Area
289 (Table 3).

290

291 **Association between parasitic infections, socio-demographic factors, and WASH** 292 **indicators**

293 A positive association was observed between Baka ethnic group (Pygmies) and helminth
294 infections, especially *A. lumbricoides* (OR=4.04; $P < 0.0001$) and *T. trichiura* (OR=2.99;
295 $p = 0.007$). Open defecation was positively associated with *T. trichiura* (OR=5.58;
296 $p < 0.0001$). Spring water consumption was positively associated with hookworm infection

297 (OR=3.87; p=0.008) (Table 4). Infection with *G. lamblia* was positively linked to disposal
298 of garbage near houses (OR=3.41; p= 0.003) and Kentzou Health Area (OR=3.73;
299 p=0.002). Infection with *E. coli* was positively associated with Lolo Health Area (OR=2.61;
300 p=0.011) and open defecation (OR=2.16; p=0.001) (Table 4).

301

302

303 Discussion

304 The present study aimed to understand the epidemiology of intestinal parasitic infections
305 in the Ndelele, Kentzou and Lolo Health Areas of the Ndelele Health District. Overall,
306 74.9% of participants were infected by at least one intestinal parasite species, which is
307 similar to the infection rate (88.2%) in the neighboring areas of Central African Republic
308 [24]. By contrast this prevalence was very high compared to those obtained by Mbuh *et*
309 *al.* (31.0%) and Denecho *et al.* (59.1%) in the South-West Region of Cameroon [25-26]
310 and Saoting *et al.* (30.0%) in the North Region of Cameroon [27]. This very high infection
311 prevalence can be explained by the poor hygiene and sanitation, and reduced access to
312 appropriate treatments in these landlocked areas. Indeed, a KAP survey conducted as
313 part of this study revealed poor knowledge of interviewees about parasitic infection
314 transmission. Those who identified food as the main mode of contamination were making
315 reference to the consumption of sweet fruits (68.1%) such as mango or banana,
316 contaminated food (12.9%) or water (10.7%). In addition, the KAP survey also revealed
317 poor practice of hand hygiene as 96.6% of enrollees was practicing hand washing without
318 soap before each meal. Also, due to absence of latrines in most of the households
319 (20.4%, an important proportion (32.7%) of the population used to practice open

320 defecation that can contribute to the persistence of transmission. It is also worth to
321 mention that most of the enrollees did not use appropriate treatment (drugs known to be
322 effective) against intestinal parasitic infections, likely because of the difficulty to access
323 to modern medicines, and when treatment campaigns are organized, they are only
324 restricted to at-risk groups. Indeed, the use of traditional medicine can represent a viable
325 option but the fact that they have limited knowledge of clinical signs of different infections
326 can jeopardize the efficacy of traditional medicines.

327
328 Likewise, infection by single parasite species, polyparasitism was also important (57.2%)
329 in the framework of this study conducted in rural communities compared to relatively low
330 rates (2.1%-14.7%) of polyparasitism found in urban areas such as in Douala, the
331 Cameroon economic city capital [28-29]. These observations confirm the results of
332 previous studies demonstrating that polyparasitism is more important in rural areas
333 compared to urban settings [28, 30, 31], likely because of poor adherence to WASH.
334 Indeed, a positive association was is often reported between poor practice of hand
335 washing and polyparasitism;, showing that people who never wash their hands with soap
336 before each meal are about 12 times more at risk of intestinal parasitic infections than
337 their counterparts who always wash their hands before meals.

338
339 As part of the present study, two vulnerable/special populations were enrolled, namely
340 the Baka (Pygmies) and the Central African Refugees. Baka's are qualified as
341 "indigenous people" because of their culture which is very different of that of the
342 "dominant" ethnic group (Kako) [32]. Although these two groups are vulnerable, their rate

343 of intestinal parasitic infection were statistically different, the Baka being heavily infested
344 (93.3% vs 53.5%), and highly polyparasitized (66.7% vs 29.9%). Unlike refugees, Baka
345 (Pygmies) live in the forest in extreme poverty conditions. Indeed, the Baka ethnic group
346 exhibited the worst results for WASH indicators while most of their households (87.5%)
347 do not have latrines. As consequence 88.9% of Baka participant practiced open
348 defecation known as a facilitating factor for the dissemination of parasite stages (eggs
349 and cysts) responsible of human infections. This situation is exacerbated by the fact that
350 most may not wash their hands before eating, although they have declared to do so.
351 Previous studies conducted among “indigenous communities” demonstrated that they are
352 most often affected by helminth infections such as *A. lumbricoides*, *T. trichiura* and
353 hookworms [33, 34, 35] as it was found in this study. Contrarily to Baka, Central African
354 refugees found in the Ndelele Health District are, for the vast majority were placed under
355 the protection of the High Commissioner for Refugees benefiting from many facilities from
356 Non-Governmental Organizations, including UNICEF, PRO-ACT Project, *Premiere*
357 *Urgence Internationale*, Care, Red Cross, *Solidarités Internationales*. Central African
358 refugees therefore benefit from donations such as latrines and boreholes construction
359 [36- 37], permanent education on environmental and personal hygiene to improve their
360 living conditions, and appropriate treatments. This can explain the fact that the infection
361 rates of the targeted parasitic infections are quite low among this special population.

362

363

364

365

366

367 **Conclusion**

368 The present study provides useful data on infection rates and risk factors associated with
369 intestinal parasitic infection in the Ndelele Health District. The results highlighted
370 exceptionally high rates of infections in this landlocked. Lack of sanitation and poor
371 hygiene can largely contribute to the endemicity of intestinal parasitic infections,
372 especially among indigenous and hard-to-reach populations (Pygmies). Targeted and
373 complementary control strategies appear compulsory to reach these populations and
374 offer appropriate care to interrupt or at least reduce the transmission of these debilitating
375 diseases.

376

377

378 **Acknowledgements**

379 The authors are thankful to all the population of the Ndelele Health District for their
380 hospitality and their participation. We are grateful to all those who contributed in the
381 realization of this study.

382

383

384 **References**

385 [1] Campbell SJ, Biritwum NK, Woods G, Velleman Y, Fleming F, Stothard JR. Tailoring
386 Water, Sanitation, and Hygiene (WASH) Targets for Soil-Transmitted Helminthiasis and

- 387 Schistosomiasis Control. Trends Parasitol. 2018 Jan;34(1):53-63. doi:
388 10.1016/j.pt.2017.09.004. Epub 2017 Oct 18. PMID: 29055522.
- 389 [2] Crompton, David W. T & Savioli, L. (1993). Intestinal parasitic infections and
390 urbanization / D. W. T. Crompton & L. Savioli. Bulletin of the World Health Organization
391 1993 ; 71(1) : 1-7 <https://apps.who.int/iris/handle/10665/49956>
- 392 [3] World Health Organisation. Soil Transmitted helminths infections. 2 March 2019.
393 Online: [https://www.who.int/news-room/fact-sheets/detail/soil-transmitted-helminth-](https://www.who.int/news-room/fact-sheets/detail/soil-transmitted-helminth-infections)
394 [infections](https://www.who.int/news-room/fact-sheets/detail/soil-transmitted-helminth-infections)
- 395 [4] World Health Organization Global Health Estimates 2016: Deaths by Cause, Age, Sex,
396 by Country and by Region,2000-2016. Geneva, online:
397 https://www.who.int/healthinfo/global_burden_disease/estimates/en/
- 398 [5] Centers for Disease Control and Prevention. Parasites- Soil Transmitted- Helminths.
399 January 10, 2013. Online: <https://www.cdc.gov/parasites/sth/index.html>
- 400 [6] Suna Balkan, Pierre Barel, Marie-Claude Bottineau, Philippa Boule. Médecins Sans
401 Frontières. Online:[https://medicalguidelines.msf.org/viewport/CG/english/intestinal-](https://medicalguidelines.msf.org/viewport/CG/english/intestinal-protozoan-infections-parasitic-diarrhoea-16689774.html#)
402 [protozoan-infections-parasitic-diarrhoea-16689774.html#](https://medicalguidelines.msf.org/viewport/CG/english/intestinal-protozoan-infections-parasitic-diarrhoea-16689774.html#)
- 403 [7] Ibrahim Mohammed, Endalkachew Nibret, Mulugeta Kibret, Bayeh Abera, Melaku
404 Adal. Prevalence of diarrhea causing protozoan infections and associated risk factors in
405 diarrheic under five children in Bahir Dar town, northwest Ethiopia: pediatric clinic based
406 study. Ethiopian Journal of Science and Technology. (2016) doi: 10.4314/ejst.v9i1.2

- 407 [8] WHO Expert Committee on the Control of Schistosomiasis (2001 : Geneva,
408 Switzerland) & World Health Organization. (2002). Prevention and control of
409 schistosomiasis and soil-transmitted helminthiasis : report of a WHO expert
410 committee. World Health Organization. <https://apps.who.int/iris/handle/10665/42588>
- 411 [9] World Health Organization. Integrating neglected tropical diseases into global health
412 and development: fourth WHO report on neglected tropical diseases. World Health
413 Organization (2017). . <https://apps.who.int/iris/handle/10665/255011>. License: CC BY-
414 NC-SA 3.0 IGO
- 415 [10] Ross AG, Papier K, Luceres-Catubig R, Chau TN, Inobaya MT, Ng SK. Poverty,
416 Dietary Intake, Intestinal Parasites, and Nutritional Status among School-Age Children in
417 the Rural Philippines. *Trop Med Infect Dis*. 2017;2(4):49. Published 2017 Sep 21.
418 doi:10.3390/tropicalmed2040049
- 419 [11 Bauhofer AFL, Cossa-Moiane I, Marques S, Guimarães EL, Munlela B, et al.
420 (2020) Intestinal protozoan infections among children 0-168 months with diarrhea in
421 Mozambique: June 2014 - January 2018. *PLOS Neglected Tropical Diseases* 14(4):
422 e0008195. <https://doi.org/10.1371/journal.pntd.0008195>]
- 423 [12] Tellevik MG, Moyo SJ, Blomberg B, Hjøllø T, Maselle SY, Langeland N et al.
424 Prevalence of *Cryptosporidium parvum/hominis*, *Entamoeba histolytica* and *Giardia*
425 *lamblia* among Young Children with and without Diarrhea in Dar es Salaam, Tanzania.
426 *PLoS Negl Trop Dis*. 2015 Oct. doi: 10.1371/journal.pntd.0004125.

- 427 [13] AÅ sbjoÈrnsdoÅttir KH, Ajjampur SSR, Anderson RM, Bailey R, Gardiner I, Halliday
428 KE, et al. (2018) Assessing the feasibility of interrupting the transmission of soil-
429 transmitted helminthes through mass drug administration: The DeWorm3 cluster
430 randomized trial protocol. PLoS Negl Trop Dis 12(1): e0006166. [https://doi.org/10.1371/
431 journal.pntd.0006166](https://doi.org/10.1371/journal.pntd.0006166)
- 432 [14] Simon Brooker, Christl A. Donnelly, Helen L. Guyatt. Estimating the number of
433 helminthic infections in the republic of Cameroon on data on infection prevalence in
434 school children. Bulletin of World Health Organisation, Febuary 2000,
435 DOI: [10.1590/S0042-96862000001200014](https://doi.org/10.1590/S0042-96862000001200014)
- 436 [15] Esther Nana Dankoni et L. A Tchuem Tchuente. Epidémiologie de la schistosomiase
437 et des geohelminthiases dans l'arrondissement de kekem (Ouest-Cameroun).
438 International Journal of innovation and applied studies ISSN 2028-9324 VOL. 8 num 4
439 Oct. 2014, pp. 1782-1790. doi=10.1.1.672.872
- 440 [16] L. Nkengazong, F. Njiokoum, S. Wanji, F. Teukeng, P. Enyong, T. Asonganyi.
441 Prevalence of soil transmitted and impact of Albendazole on parasitic indices in Kotto
442 Barombi and Marumba II villages (South-west Cameroon). Vol. 4 No. 3 (2010)._ doi:
443 [10.5897/ajest09.205](https://doi.org/10.5897/ajest09.205).
- 444 [17] Tchuem Tchuente, Behnke S. M, Gilbert F. S, V. R. Southgate, J. Vercruysse.
445 Polyparasitism with Schistosoma haematobium and soil transmitted helminths infections
446 among school children in Loum, Cameroon. Trop. Med. Int. H., 2003, 8(11), 975-986.
447 <https://doi.org/10.1046/j.1360-2276.2003.01120.x>

- 448 [18] Communes et villes unies du Cameroun: United council and cities of Cameroon.
449 Online:[http://cvuc.cm/national/index.php/fr/carte-communale/region-de-lest/122-](http://cvuc.cm/national/index.php/fr/carte-communale/region-de-lest/122-association/carte-administrative/est/kadey/487-ndelele)
450 [association/carte-administrative/est/kadey/487-ndelele](http://cvuc.cm/national/index.php/fr/carte-communale/region-de-lest/122-association/carte-administrative/est/kadey/487-ndelele)
- 451 [19] Communes et villes unies du Cameroun: United council and cities of Cameroon.
452 Online:[http://cvuc.cm/national/index.php/fr/carte-communale/region-de-lest/122-](http://cvuc.cm/national/index.php/fr/carte-communale/region-de-lest/122-association/carte-administrative/est/kadey/485-kentzou)
453 [association/carte-administrative/est/kadey/485-kentzou](http://cvuc.cm/national/index.php/fr/carte-communale/region-de-lest/122-association/carte-administrative/est/kadey/485-kentzou)
- 454 [20] World Health Organization. Basic laboratory methods in medical parasitology.
455 WHO(1991) . <https://apps.who.int/iris/handle/10665/40793>
- 456 [21] World Health Organization. Planches pour le diagnostic des parasites
457 intestinaux. Genève: Organisation mondiale de la
458 Santé(1994). <https://apps.who.int/iris/handle/10665/37327>
- 459 [22] J. Utzinger, Botero-Kleiven, S. Castelli, P.L. Chiodini, H. Edwards, N. Köhler et al.
460 Microscopic diagnosis of sodium acetate-acetic acid-formalin-fixed stool samples for
461 helminths and intestinal protozoa: a comparison among European reference laboratories.
462 Clinical Microbiology Infectious Diseases. (2010). 16, 267–273. doi: 10.1111/j.1469-
463 0691.2009.02782.x
- 464 [23] E.C. Strunz, David G. Addiss, Meredith E. Stocks, Stephanie Ogden, Jürg Utzinger,
465 Matthew et al. Water, sanitation, hygiene, and soil-transmitted helminth infection: a
466 systematic review and meta-analysis. PLoS Med. 2014 Mar 25;11(3):e1001620.
467 doi:10.1371/journal.pmed.1001620.

468 [24] G. Tékpa, V. Fikouma, E. Gbangba-Ngaï, B.O. Bogning Mejozem, S. Ningatouloum
469 Nazita, B. Koffi, Epidemiological and clinical profile of intestinal parasitosis of children in
470 rural areas in Central African Republic, Archives de Pédiatrie, Volume 26, Issue
471 1, 2019, Pages 34-37, <https://doi.org/10.1016/j.arcped.2018.11.006>.

472 [25] J.V. Mbuh, N.H. Ntonifor and J. Ojong. The epidemiology of soil-transmitted helminth
473 and protozoan infections in south-west Cameroon. Journal of helminthology Volume 86,
474 Issue 1 March 2012, pp. 30-37. doi: <https://doi.org/10.1017/S0022149X1000091X>

475 [26] L. Ndenecho, K. J.N. Ndamukong, M. M. Matute. Soil transmitted nematodes in
476 Children in Buea Health district of Cameroon. East African Medical Journal: 2002 79(8):
477 442-445). doi: 10.4314/eamj.v79i8.8832.

478 [27] Saotoing, R. Djonyang, D. D. Dereng et al. Enquête épidémiologique sur les
479 parasitoses urinaires et intestinales chez les élèves des écoles primaires de
480 l'arrondissement de Maga, Extrême-Nord Cameroun. African Journal Online (2016). doi:
481 10.4314/ijbcs.v10i1.26.

482 [28] L. G. Lehman, L. Koujip, C.F. Bilong Bilong. Diagnostic des parasitoses intestinales
483 à l'aide de la microscopie à fluorescence. Medecine d'Afrique noire. Juillet 2012. Online
484 : [Diagnostic des parasitoses intestinales à l'aide de la microscopie à fluorescence \(ured-
485 douala.com\)](http://www.douala.com)

486 [29] Thomas KUETE, Fomefret Loic, Sorel YEMELI, Emmanuel ESSONO MVOA,
487 Therese NKOA, Roger M. Somo, A. Same Ekobo. Prevalence and Risk Factors of
488 Intestinal Helminth and Protozoa Infections in an Urban Setting of Cameroon: the Case

489 of Douala. *American Journal of Epidemiology and Infectious Disease* (2015). doi:
490 10.12691/ajeid-3-2-4.

491 [30] Al-Delaimy AK, Al-Mekhlafi HM, Nasr NA, Sady H, Atroosh WM, Mohammed Nashiry,
492 Tengku S. Anuar et al. Epidemiology of Intestinal Polyparasitism among Orang Asli
493 School Children in Rural Malaysia. *PLoS Negl Trop Dis* (2014)
494 doi:10.1371/journal.pntd.0003074.

495 [31] Opara KN, Udoidung NI, Opara DC, Okon OE, Edosomwan EU, Udoh AJ. The
496 Impact of Intestinal Parasitic Infections on the Nutritional Status of Rural and Urban
497 School-Aged Children in Nigeria. *Int J MCH AIDS*. (2012) doi:10.21106/ijma.8.

498 [32] Supplementary report submitted in connection with Cameroon's second periodic
499 report. Indigenous peoples' rights in Cameroon. May 2010. Online:
500 [https://www.forestpeoples.org/sites/fpp/files/publication/2010/08/cameroonachprsubmis](https://www.forestpeoples.org/sites/fpp/files/publication/2010/08/cameroonachprsubmissionmay10eng.pdf)
501 [sionmay10eng.pdf](https://www.forestpeoples.org/sites/fpp/files/publication/2010/08/cameroonachprsubmis)

502 [33] Chin, Y.T., Lim, Y.A.L., Chong, C.W. *et al.* Prevalence and risk factors of intestinal
503 parasitism among two indigenous sub-ethnic groups in Peninsular Malaysia. *Infect Dis*
504 *Poverty* (2016). <https://doi.org/10.1186/s40249-016-0168-z>.

505 [34] Rajoo Y, Ambu S, Lim YAL, Rajoo K, Tey SC, Lu CW, et al. Neglected Intestinal
506 Parasites, Malnutrition and Associated Key Factors: A Population Based Cross-Sectional
507 Study among Indigenous Communities in Sarawak, Malaysia. *PLoS ONE* (2017)
508 <https://doi.org/10.1371/journal.pone.0170174>.

509 [35] Carla Scolari, Carlo Torti, Anna Beltrame, Alberto Matteelli, Francesco Castelli,
510 Maurizio Gulletta et al. Prevalence and distribution of soil-transmitted helminth (STH)
511 infections in urban and indigenous schoolchildren in Ortigueira, State of Paraná, Brasil:
512 implications for control. Tropical medicine and international health. December 2001.
513 <https://doi.org/10.1046/j.1365-3156.2000.00549.x>

514 [36] Rapport inter-agences sur la situation des refugiees Centrafricains. Cameroun. 13-
515 19 Octobre 2014. Online : <https://data2.unhcr.org/en/documents/download/32746>

516 [37] Rapport inter-agences sur la situation des refugiees Centrafricains. Cameroun. 1-31
517 Mars 2016. Online:[https://reliefweb.int/report/cameroon/cameroun-rapport-inter-](https://reliefweb.int/report/cameroon/cameroun-rapport-inter-agences-sur-la-situation-des-r-fugi-s-centrafricains-1-31-0)
518 [agences-sur-la-situation-des-r-fugi-s-centrafricains-1-31-0](https://reliefweb.int/report/cameroon/cameroun-rapport-inter-agences-sur-la-situation-des-r-fugi-s-centrafricains-1-31-0)

519

520

521 **Figure legends**

522 **Fig. 1.** Spatial distribution of parasitic infections in Ndelele, Kentzou and Lolo health areas
523 of Ndelele Health District.

524

525 **Fig. 2.** Frequency of parasitic infections according to gender.

526 **Tables**

527 **Table 1: WASH indicators according to Health area and ethnic groups**

	Health area						Ethnic groups							
	Ndelele		Kentzou		Lolo		Kako		Baka		CAR refugees		Other ethnics	
	n	%	n	%	n	%	n	%	n	%	n	%	n	%
Latrine owning*														
Yes	56	61.5	47	92.2	70	95.8	74	78.7	3	12.5	85	98.8	11	100
No	35	38.5	4	7.8	3	4.2	20	21.3	21	87.5	1	1.2	0	0.0
Frequency of hand washing with soap*														
Always	3	1.7	5	4.6	11	9.1	3	1.5	0	0.0	15	10.4	0	0.0
Sometimes	71	40.1	57	52.7	45	37.2	110	56.7	11	24.4	44	30.6	8	34.8
Often	12	6.7	35	34.4	52	43.0	10	5.2	0	0.0	84	58.3	5	21.7
Never	91	51.4	12	11.1	13	10.7	71	36.6	34	75.6	1	0.7	10	43.5
Barefoot walking*														
Always	4	2.3	0	0	2	1.7	2	1.0	3	6.7	1	0.7	0	0.0
Sometimes	83	44.9	49	45.4	50	41.3	87	44.8	19	42.2	68	47.2	8	34.8
Often	78	44.1	39	36.1	17	14.0	99	51.0	22	48.9	6	4.2	7	30.4
Never	12	6.8	20	18.5	52	43.0	6	3.1	1	2.2	69	47.9	8	34.8
Water sources*														
Spring water	89	50.3	51	47.2	1	0.8	72	37.1	45	100	17	11.8	7	30.4
Well water	9	5.1	0	0.0	0	0.0	5	2.6	0	0.0	4	2.8	0	0.0
Borehole water	79	44.6	57	52.8	120	99.2	117	60.3	0	0.0	123	85.4	16	69.6
Defecation place*														
Toilet	89	50.3	78	72.2	106	87.6	107	55.2	3	6.7	142	98.6	21	91.3
Open defecation	72	40.7	17	15.7	11	9.1	58	29.9	40	88.9	1	0.7	1	4.3
Toilet and open defecation	16	9.0	13	12.0	4	3.3	29	14.9	3	4.4	1	0.7	1	4.3
Disposal of garbage in close proximity to houses *														
Yes	98	55.4	24	22.2	17	14.0	83	42.8	39	86.7	137	4.9	10	43.5
No	79	44.6	84	77.8	104	86.0	111	57.2	6	13.3	7	95.1	13	56.5

528 *CAR: Central African Republic; *p-value <0.0001*

529

530

531

532

533

534 **Table 2: WASH indicators according to gender and age groups**

	Gender		Age groups						
	Male n=166 n (%)	Female n=240 n (%)	<5 n=59 n (%)	5-14 n=98 n (%)	15-24 n=66 n (%)	25-34 n=51 n (%)	35-44 n=59 n (%)	45-54 n=32 n (%)	55+ n=41 n (%)
Frequency of hand washing with soap *									
Always	12(7.2)	6(2.5)	0(0.0)	0(0.0)	3(4.5)	5(9.8)	6(10.2)	3(9.4)	1(2.4)
Sometimes	74(44.6)	99(41.3)	25(42.4)	59(60.2)	25(37.9)	21(37.9)	19(32.2)	9(28.1)	15(36.6)
Often	33(19.9)	66(27.5)	7(11.9)	8(8.2)	20(30.3)	16(31.4)	16(27.1)	14(43.8)	18(43.9)
Never	47(28.3)	69(28.8)	27(45.8)	31(31.6)	18(27.3)	9(17.6)	18(30.5)	6(18.8)	7(17.1)
Barefoot walking*									
Always	1(0.6)	5(2.1)	0(0.0)	2(2.0)	0(0.0)	0(0.0)	2(3.4)	1(3.1)	1(2.1)
Sometimes	70(42.2)	112(46.7)	12(20.3)	47(48.0)	42(63.6)	24(47.1)	26(44.1)	14(43.8)	17(41.5)
Often	64(38.6)	70(29.2)	45(76.3)	45(45.9)	12(18.2)	6(11.8)	14(23.7)	5(15.6)	7(17.1)
Never	31(18.7)	53(22.1)	2(3.4)	4(4.1)	12(18.2)	21(41.2)	17(28.8)	12(37.5)	16(39.0)
Water sources									
Spring water	60(36.1)	61(33.8)	23(39.0)	27(27.6)	28(42.4)	16(31.4)	23(39.0)	12(37.5)	12(29.3)
Well water	4(2.4)	5(2.1)	2(3.4)	3(3.1)	0(0.0)	1(2.0)	1(1.7)	0(0.0)	2(4.9)
Borehole water	102(61.4)	154(64.2)	34(57.6)	68(69.4)	38(57.6)	34(66.7)	35(59.3)	20(62.5)	27(65.9)
Defecation place*									
Toilet	108(65.1)	165(68.8)	22(37.3)	53(54.1)	51(77.3)	41(80.4)	43(72.9)	27(84.4)	36(87.8)
Open defecation	44(26.5)	56(23.3)	27(45.8)	27(27.6)	12(18.2)	8(15.7)	16(27.1)	5(15.6)	5(12.2)
Toilet and open defecation	14(8.4)	19(7.9)	10(16.9)	18(18.4)	3(4.5)	2(3.9)	0(0.0)	0(0.0)	0(0.0)
Disposal of garbage in close proximity to houses									
Yes	65(39.2)	74(30.8)	22(37.3)	37(37.8)	24(36.4)	14(27.5)	22(37.3)	9(28.1)	11(26.8)
No	101(60.8)	166(69.2)	37(62.7)	61(62.2)	42(63.6)	37(72.5)	37(62.7)	23(71.9)	30(73.2)

535 * $p < 0.0001$

536

537

538 **Table 3: Parasitic infections in ethnics groups**

Parasitic species	Ethnic groups							
	Kako (n=194)		Baka (n=45)		Central African refugees (n=144)		Other ethnics (n=23)	
	Infected	%	Infected	%	Infected	%	Infected	%
Helminths								
<i>Ascaris lumbricoides</i>	54	27.8	31	68.9**	4	2.8	6	26.1
<i>Trichuris trichiura</i>	21	10.8	12	26.7**	0	0.0	1	4.3
Hookworm	15	7.7**	2	4.4	1	0.7	0	0.0
<i>Strongyloides stercoralis</i>	10	5.2	4	8.9	0	0.0	0	0.0
<i>Enterobius vermicularis</i>	1	0.5	0	0.0	0	0.0	0	0.0
Protozoa								
<i>Entamoeba Coli</i>	97	50.0*	19	42.2	27	18.8	9	39.1
<i>Entamoeba histolytica</i>	49	25.3**	10	22.2	16	11.1	5	21.1
<i>Entamoeba hartmanni</i>	5	2.6	4*	8.9	0	0.0	0	0.0
<i>Endolimax nana</i>	11	5.7	1	2.2	2	1.4	0	0.0
<i>Iodamoeba butshlii</i>	21	10.8*	2	4.4	1	0.7	0	0.0
<i>Giardia lamblia</i>	16	8.2	2	4.4	8	5.6	3	13.0
<i>Balantidium coli</i>	1	0.5	0	0.0	0	0.0	0	0.0
<i>Isospora belli</i>	1	0.5	0	0.0	1	0.7	0	0.0
<i>Blastocystis hominis</i>	80	41.2	18	40.0	42	29.2	6	26.1

539 * $p < 0.05$; ** $p < 0.0001$

540

541

542 **Table 4: Intensity of helminth infections**

	<i>Ascaris lumbricoides</i>			<i>Trichuris trichiura</i>			Hookworm		
	Light (%)	Moderate (%)	Heavy (%)	Light (%)	Moderate (%)	Heavy (%)	Light (%)	Moderate (%)	Heavy (%)
Gender									
Male	23(60.5)	15(39.5)	0(0.0)	11(91.7)	1(8.3)	0(0.0)	6(100)	0(0.0)	0(0.0)
Female	43(75.4)	12(21.1)	2(3.5)	18(90.0)	2(10.0)	0(0.0)	11(91.7)	1(8.3)	0(0.0)
Age groups									
<5	9(60.0)	5(33.3)	1(6.7)	5(83.3)	1(16.7)	0(0.0)	3(75.0)	1(25.0)	0(0.0)
5-14	18(56.3)	14(43.8)	0(0.0)	11(91.7)	1(8.3)	0(0.0)	5(100)	0(0.0)	0(0.0)
15-24	16(84.2)	2(10.5)	1(5.3)	4(80.0)	1(20.0)	0(0.0)	0(0.0)	0(0.0)	0(0.0)
25-34	7(100)	0(0.0)	0(0.0)	2(100)	0(0.0)	0(0.0)	3(100)	0(0.0)	0(0.0)
35-44	5(62.5)	3(37.5)	0(0.0)	2(100)	0(0.0)	0(0.0)	2(100)	0(0.0)	0(0.0)
45-54	4(80.0)	1(20.0)	0(0.0)	1(100)	0(0.0)	0(0.0)	2(100)	0(0.0)	0(0.0)
55+	7(77.8)	2(22.2)	0(0.0)	2(100)	0(0.0)	0(0.0)	2(100)	0(0.0)	0(0.0)
Health area									
Ndelele	47 (63.5)	25(33.8)	2 (2.7)	23 (95.8)	1 (4.2)	0 (0.0)	9 (90.0)	1 (10.0)	0 (0.0)
Kentzou	11 (100)	0 (0.0)	0 (0.0)	6 (75.0)	2 (25.0)	0 (0.0)	6 (100)	0 (0.0)	0 (0.0)
Lolo	8 (80)	2 (20.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (100)	0 (0.0)	0 (0.0)
Ethnic groups									
Kako	40(74.1)	13(24.1)	1(1.9)	16(84.2)	3(15.8)	0(0.0)	14(93.3)	1(6.7)	0 (0.0)
Baka	19(61.3)	11(35.5)	1(3.2)	12(100)	0(0.0)	0(0.0)	2(100)	0(0.0)	0(0.0)
CAR refugees	4(100)	0(0.0)	0(0.0)	0(0.0)	0(0.0)	0(0.0)	1(100)	0(0.0)	0(0.0)
Other ethnics	3(50.0)	3(50.0)	0(0.0)	1(100)	0(0.0)	0(0.0)	0(0.0)	0(0.0)	0(0.0)

543


544

545 **Table 5. Risk factors Parasites**


546

Parasites	Factors	Adjusted Odd ratio (95% CI)	p-value
Helminths			
<i>Ascaris lumbricoides</i>			
Ethnic groups	Kako	1	
	Baka	4.04 (1.926- 8.80)	< 0.0001
	Central African refugees	0.80 (0.02- 0.26)	< 0.0001
	Other Ethnics	0.53 (0.32-0.87)	0.7
Defecation place	Toilet	1	
	Open defecation	3.37 (1.24- 9.10)	0.016
	Toilet and Open defecation	1.162(0.44- 3.05)	0.760
Disposal of garbage close to houses	No	1	
	Yes	2.80 (1.66-4.723))	< 0.0001
<i>Trichuris trichiura</i>			
Ethnic groups	Baka ethnic	2.99 (1.34-6.67)	0.007
	Central African Refugees	000	0.995
	Other Ethnics	0.374 (0.04- 2.92)	0.349
Defecation place	Toilet	1	
	Open defecation	5.58 (2.55- 12.22)	< 0.0001
	Toilet and Open defecation	3.28 (0.983- 10.983)	0.053
Hookworm			
Water sources	Borehole water	1	
	Spring water	3.87 (1.42 - 10.56)	0.008
Protozoa	Well water	000	0.999
<i>Entamoeba coli</i>			
Ethnic groups	Kako	1	
	Baka	0.94 (0.46 - 1.89)	0.869
	Central African refugees	0.13 (0.06 - 0.26)	<0.0001

	Other ethnics	0.72 (0.29 – 1.77)	0.484
Health areas	Ndelele	1	
	Kentzou	1.52 (0.87 – 2.66)	
	Lolo	2.61 (1.24- 5.587)	0.011
Defecation place	Open defecation	2.16 (1.35-3.45)	0.001
	Toilet + open defecation	3.05 (1.46-6.37)	0.003
<i>Entamoeba histolytica</i>			
Ethnic groups	Kako	1	
	Baka	0.84 (0.39 -1.83)	
	Central African refugees	0.37 (0.20 -0.68)	0.001
	Other ethnics	0.822 (0.29- 2.33)	0.712
<i>Giardia lamblia</i>			
Health area	Ndelele	1	
	Kentzou	3.73 (1.61- 8.64)	0.002
	Lolo	0.31 (0.06 -1.47)	1.143
Disposal of garbage near houses	No	1	
	Yes	3.41 (1.52- 7.66)	0.003



Figure



Figure