

Cardiac and aortic size measures in adult echocardiography should be indexed by body surface area regardless of body mass index

Angus Fung^{1,2}, Dhnanjay Soundappan^{1,2}, Daniel E Loewenstein MD³,
David Playford MBBS PhD⁴, Geoffrey Strange PhD⁵, Rebecca Kozor MBBS PhD¹,
James Otton MBBS PhD^{2,6}, Martin Ugander MD PhD^{1,2,3,§}

¹Kolling Institute, Royal North Shore Hospital, and University of Sydney, Sydney, Australia

²St Vincent's Clinical School, University of New South Wales, Sydney, Australia

³Department of Clinical Physiology, Karolinska University Hospital, and Karolinska Institutet, Stockholm, Sweden

⁴School of Medicine, University of Notre Dame, Fremantle, Australia

⁵Faculty of Medicine and Health, University of Sydney, Sydney, Australia

⁶Department of Cardiology, Liverpool Hospital, University of New South Wales, Liverpool, Australia

§ corresponding author:

Professor Martin Ugander MD PhD FCSANZ

Kolling Building, Level 12, Royal North Shore Hospital

St Leonards, Sydney, NSW 2065, Australia

e-mail: martin@ugander.com

Abstract

BACKGROUND: Body size indexation is a foundation of the diagnostic interpretation of cardiac size measures used in imaging assessment of cardiovascular health. Body surface area (BSA) is the most commonly used metric for body size indexation of echocardiographic measures, but its use in patients who are underweight or obese is questioned (body mass index (BMI) <18.5 kg/m² or ≥ 30 kg/m², respectively). We hypothesized that mortality can be used to identify an optimal body size indexation metric for echocardiographic measures that would be a better predictor of survival than BSA regardless of BMI.

METHODS: In this big data, cohort study, adult patients with no prior valve replacement were selected from the National Echo Database Australia. Survival analysis was performed for echocardiographic measures both unindexed and indexed to different body size metrics, with 5-year cardiovascular mortality as the primary endpoint.

FINDINGS: Indexation of echocardiographic measures (left ventricular diameter [n=337,481] and mass [n=330,959], left atrial area [n=136,989], aortic sinus diameter [n=125,130], right atrial area [n=81,699], right ventricular diameter [n=3,575], right ventricular outflow tract diameter [n=2,841]) by BSA had better prognostic performance vs unindexed measures (healthy/overweight: C-statistic 0.656 vs 0.618, average change in Akaike Information Criteria (Δ AIC) 800; underweight: C-statistic 0.669 vs 0.654, Δ AIC 15; obese: C-statistic 0.630 vs 0.612, Δ AIC 113). Indexation by other body size metrics (lean body mass or height and/or weight raised to various powers) did not improve prognostic performance versus BSA by a clinically relevant magnitude (average C-statistic increase ≤ 0.01), with smaller differences in higher BMI subgroups. Similar results were obtained using sex-disaggregated analysis, for indexation of other aortic or cardiac dimension or volume measures, and for all-cause mortality.

INTERPRETATION: Indexing measures of cardiac and aortic size by BSA improves prognostic performance regardless of BMI, and no other body size metric has a clinically meaningful better performance.

FUNDING: This research was supported in part by grants (PI Ugander) from New South Wales Health, Heart Research Australia, and the University of Sydney.

Introduction

Quantification of the dimensions of the heart and great vessels using echocardiography has both diagnostic and prognostic value for the prediction of morbidity and mortality, which also may help guide treatment in patients.¹⁻⁸ Currently, the recommended method for body size indexation of cardiac volumes is body surface area (BSA).⁹ However, there is heterogeneity in the literature as to the best indexation method, and whether or not indexing of cardiac measures improves their predictive value for cardiovascular events.^{6, 10-12} In underweight and overweight patients, correction for BSA can overestimate or underestimate the prevalence of left ventricular hypertrophy, and inaccurately normalise or exaggerate indices of cardiac size.¹⁰ Furthermore, there are concerns regarding the physiological relationships between left ventricular mass and indexation methods. While the relationships between body surface area, height, and weight are non-linear, the indexation of left ventricular mass by these variables often assumes linear relationships.¹³ An argument can also be drawn from the theory of similarity, which reasons that relative geometries are best indexed to body size variables of similar dimensionality. For example, since left ventricular mass is related to cardiac dimensions raised to the third power, and BSA is related to a body dimension raised to the second power, it is logical that left ventricular mass should be proportional to $BSA^{3/2}$ (also expressed as $BSA^{1.5}$).¹⁴

Studies on indexation for prognostic performance have been limited in patient sample size and range of cardiac measures indexed.^{5, 15-18} Using the large-scale data available in the National Echocardiography Database of Australia (NEDA), the aim of the study was to derive one or more formulae based on height and weight to provide a method of body size indexation of cardiac and aortic measures that will be a better predictor of all-cause mortality than current methods based on body surface area.

Methods

Study design. NEDA is a large observational registry that includes routinely recorded echocardiographic data across 30 centres in Australia. Individual data linkage is used to incorporate health outcomes such as all-cause and cardiovascular mortality. The study cohort consists of patients over the age of 18 who have typically been referred clinically for imaging evaluation of known or suspected cardiovascular disease. The study was approved by the lead ethics committee at the Royal Prince Alfred Hospital (2019/ETH06989). NEDA is registered with the Australian New Zealand Clinical Trials Registry (ACTRN12617001387314). Ethical approval has been obtained from the Human Research Ethics Committees at the respective recruiting sites, and the study adheres to the Declaration of Helsinki.

Study cohort. Echocardiographic data and basic patient characteristics were collected from participating centres from 1 January 2000 to 21 May 2019, and were transferred into a central database via an automated data extraction process. All data was cleaned through removal of duplicate, inconsistent, and/or impossible measurements, and transformed into a standardized format. Individuals contributing to NEDA were assigned a unique identifier linked to their echocardiograms and their anonymity protected by stringent security protocols. As shown in Figure 1, 631,824 patients were present in the database. Of these, 77,125 (12%) were excluded for prior valve replacement. Echocardiograms with time from echocardiography to census or death, cause of mortality (cardiovascular and all-cause), height, weight, and the cardiac measure of interest were selected. Different populations were individually filtered for each measure of interest and analysed to maximise the number of patients for analysis. For patients with multiple echocardiograms, the earliest recorded echocardiogram was selected. Patients were grouped according to body mass index (BMI) $<18.5 \text{ kg/m}^2$, BMI $18.5\text{-}30 \text{ kg/m}^2$, or BMI $\geq 30 \text{ kg/m}^2$.

Endpoints. The primary endpoints of interest were all-cause and cardiovascular mortality. Mortality data was obtained by linkage with the National Death Index.¹⁹ A detailed probability matching process involving patient identifiers obtained at echocardiographic recording was used to link survival status of individuals up to the study census date of 21 May 2019. Listed causes of death were described using ICD-10 coding, which allowed for cardiovascular death to be defined (range 100-199 ICD-10AM chapter codes).²⁰

Statistical analysis. NEDA data analyses and reports were generated in agreement with STROBE guidelines.²¹ All data used in analyses were provided and no missing data was imputed. Standard procedures for describing grouped data, such as median [interquartile range (IQR)], and proportions according to patient characteristics were applied.

Cox-proportional hazard models with proportional hazards confirmed by visual inspection and numerical analysis of Schoenfeld residuals were used to derive C-statistics and hazard-ratios (with 95% confidence intervals) for the risk of all-cause and cardiovascular mortality for the entirety of the study follow-up and a five year follow-up duration. The change in Akaike information criterion (Δ AIC) was used to interpret the statistical robustness of the body size indexation metrics. Due to the magnification of Δ AIC by large sample sizes, the C-statistic was chosen to dictate the magnitude of difference between metrics and clinical relevance in a sample size independent fashion. Kaplan-Meier curves were constructed to visually inspect differences between indexation measures. An iterative function was coded to derive 50,000 combinations of body size metrics using different height and weight exponents according to the formula where a given body size metric = $\text{height}^x \cdot \text{weight}^y$. Random combinations of x and y were used as the metric for indexation for the respective echocardiographic measures for subsequent Cox-regression analysis. Cox-regression was performed using the echocardiographic measure indexed by the derived body size metric with five-year cardiovascular mortality as the endpoint to obtain the C-statistic. The C-statistic

was color coded in a scatterplot to present differences in the prognostic strength of different body size metrics including BSA as calculated according to Mosteller²² or Du Bois²³, lean body mass formulas by Hume²⁴, Boer²⁵, or James²⁶, BSA raised to various powers, and height and/or weight raised to various powers. All statistical analyses was performed using R 4.0.4 (R Core Team, R Foundation for Statistical Computing, Vienna, Austria).²⁷ Significance was accepted at the level of $p < 0.05$ (two-sided).

Results

Study Cohort. Subject characteristics and size of the study cohorts for various cardiac and aortic measures are presented in Table 1. Due to the large sample size, differences in baseline characteristics between BMI groups were statistically significant but were not of a clinically meaningful magnitude.

Body size metrics and mortality. Across the echocardiographic measures of left atrial area (n=136,989), right atrial area (n=81,699), left ventricular end-diastolic diameter (n=337,481) and mass (n=330,959), right ventricular end-diastolic diameter (n=3,575), and aortic sinus diameter (n=125,130), indexation by BSA as calculated by Mosteller had an average C-statistic increase from 0.656 to 0.618 for the healthy/overweight cohort (average change in Akaike Information Criteria (Δ AIC) 800), an increase from 0.669 to 0.654 for the underweight cohort (Δ AIC 15), and an increase from 0.630 vs 0.612 for the obese cohort (Δ AIC 113) compared to unindexed measures as shown in Table 2.²² Indexation by other body size metrics yielded a C-statistic increase ≤ 0.01 . Further sex-disaggregated analysis did not differ upon visual inspection, and numerical differences between BSA and the best body size indexation metric were not clinically meaningful (average C-statistic increase ≤ 0.02 , detailed data not shown). Smaller differences in C-statistic between indexation metrics were observed in higher BMI subgroups. Similar results were obtained using long-term cardiovascular mortality not limited to five years, and both long-term and five year all-cause

mortality (Appendix 1-3). Similar trends were observed across indexation of other aortic dimensions (sinotubular junction, ascending, root, arch) and cardiac chamber volumes (left atrial end-systolic diameter and volume, left ventricular end-diastolic volume) (Appendix 4-33). Kaplan Meier curves did not show visually appreciable differences between indexation by BSA compared to indexation by weight across any measure (Appendix 34-48).

Figure 2 shows how indexation by different body size metrics (based on the formula $\text{height}^x \cdot \text{weight}^y$) perform prognostically, averaged across representative cardiac and aortic anatomical measure. The color scale shows increasing C-statistics, where each change in color represents one percentage point of C-statistic. Numerical values for selected measures are presented in Table 2. In summary, in the healthy/overweight cohort, indexation by BSA improves prognostic performance compared to unindexed measures by five percentage points of the C-statistic. Further improvement beyond BSA is limited to <1 percentage point improvement on average. Indexation by BSA yielded smaller improvements in prognostic performance in the underweight and obese cohort, but indexation by any other body size metric did not provide any meaningfully stronger association with survival. Further analyses in obese populations and higher BMI subgroups (BMI 30-35, 35-40 and >40) showed that BSA performed similarly to height raised to various powers (data not shown).

Relative prognostic strength of different cardiac and aortic size measures. Figure 3 shows the prognostic strength of indexing by BSA across all 14 cardiac and aortic measures analysed. In the healthy/overweight group, left atrial size and left ventricular mass had the highest prognostic strength, with a C-statistic ten percentage points higher than left ventricular size and most aortic dimensions. Right atrial and right ventricular sizes had an intermediate prognostic strength. Aortic sinus diameter had the strongest prognostic strength of all aortic measures, and aorta at sinotubular diameter had the weakest prognostic strength.

Discussion

In this study of body size indexation of cardiac and aortic sizes using real-world echocardiographic data from a large-scale nationwide cohort, indexation by BSA is shown to improve prognostic performance compared to unindexed measures regardless of BMI. Furthermore, no other body size indexation metric provided any meaningful improvement in prognostic performance beyond BSA. The current study comprehensively assessed indexation metrics with varying height and weight exponents in different forms (multiplicative, additive/subtractive, both), and in sex-specific cohorts. It was not possible to derive a body size metric with clinically meaningfully better prognostic performance than BSA. This means that using mortality as the arbiter of indexation effectiveness, no other indexation method exists that is clinically meaningfully better than BSA across all investigated cardiac measures. In accordance with current guidelines for echocardiography⁹, cardiac measures should continue to be indexed using any formula for BSA, regardless of BMI or the echocardiographic measure of interest.

Comparison with indexation by height. The current study found that indexation by height raised to various powers does not improve prognostic performance compared to BSA regardless of BMI. It has been suggested that indexation by height improves detection of left ventricular hypertrophy and associations with cardiovascular events and mortality compared to BSA in obese populations.^{12, 18, 28} Recently, it has been shown that indexation by height for left atrial volumes was able to maintain proportionality and avoid overcorrection for body size.²⁹ However, other studies have found no improvement in indexation by height compared to BSA.⁵ The findings of the current study confirm that indexation by height does not provide additional prognostic value compared to indexation by BSA across any BMI group.

Sex differences. The current study found that indexation of cardiac measures disaggregated by sex does not improve prognostic performance. Differences in left ventricular mass have

been found between male and female patients.^{30, 31} A consideration of sex in the indexation of cardiac measures measured by cardiac magnetic resonance imaging has been suggested for improved prediction of incident heart failure.¹¹ The current study analysed sex-specific cohorts and found that body size metrics derived from sex-specific cohorts were interchangeable with negligible differences in prognostic performance. Thus, the findings of the current study suggest that the relationship between cardiac or aortic size and survival in relation to body size does not fundamentally differ between the sexes. Notably, this is still consistent with using sex-specific cut-offs for normality for a given measure.

Relative prognostic association for different aortic or cardiac size measures. The current study found that the prognostic strength of indexed echocardiographic measures varied markedly. Studies comparing prognostic strength of indexed measures are limited with most studies only investigating the prognostic strength of one or two measures.^{5, 6, 32} The current study found that left ventricular mass index and left atrial volume index had the highest prognostic strengths of any measure, with C-statistics broadly similar to those found in previous smaller studies.⁵ Right atrial and right ventricular sizes, and the dimensions of the pulmonic valve expressed as RVOT diameter had moderate prognostic strength that has yet to be evaluated in literature. The current study was the first to compare the prognostic strength of different indexed aortic sizes. Furthermore, when indexed to body size, aortic sinus diameter is the only aortic size measure with good prognostic performance, and other aortic sizes have a considerably weaker association with prognosis. Notably, left ventricular end-diastolic diameter or volume, both measures of left ventricular dilatation, had among the weakest associations with prognosis.

Strengths and limitations. The limitations of applying and interpreting big data in NEDA have been acknowledged previously.^{20, 33} At the time of analysis, NEDA did not include important clinical details of common conditions such as coronary artery disease, ischaemic

heart disease, and clinically diagnosed HF, which may impact mortality. That said, the current study used large-scale, real-world data with relevant clinical outcomes that inevitably have measurement variability between centres and observers. While on one hand this is a limitation as an uncontrolled source of data heterogeneity, it is in fact a strength that reinforces the integrity of observed trends that exist despite sources of variability.

The main endpoint of consideration, cardiovascular mortality, was linked from the Australian National Death Index which has a high sensitivity and specificity (93% and 90%, respectively) validating its use as the primary endpoint.¹⁹ Furthermore, in the current study, similar trends were also observed across all-cause mortality, further reinforcing the validity of the results.

The current study did not include the impact of age on aortic size or mortality. An increase in age has long been established to be related to an increased aortic size, which would affect interpretation of the indexed measure.^{34, 35} The association between age and mortality is both intuitive and widely accepted.^{36, 37} However, a consideration of age in choice of body size indexation metric is impractical clinically, and fails to achieve the goal of indexation, namely, to account for body size. A consideration of age is more appropriate for cut-off values of the indexed cardiac and aortic size measure, but not necessarily for the choice of body size indexation metric. Importantly, similar age distributions existed between the respective cohorts in the current study. Thus, while theoretically attractive, consideration of the effect of age is beyond the scope of this study.

The NEDA cohort comprises patients being investigated for known or suspected cardiovascular disease. Data was largely obtained from specialist centres or clinics in Australia. While the NEDA cohort is representative of the diverse and multiethnic population

in Australia with a largely high-functioning level of health care, applicability may be different in other contexts.

In conclusion, the current study has definitively demonstrated that measures of cardiac and aortic size should continue to be indexed by BSA regardless of BMI. No other existing or derived body size metric (lean body mass or height and/or weight raised to various powers) is clinically meaningfully better.

References

1. Casale PN, Devereux RB, Milner M, et al. Value of echocardiographic measurement of left ventricular mass in predicting cardiovascular morbid events in hypertensive men. *Ann Intern Med.* 1986; **105**: 2:173-8.
2. Verdecchia P, Schillaci G, Borgioni C, et al. Adverse prognostic significance of concentric remodeling of the left ventricle in hypertensive patients with normal left ventricular mass. *J Am Coll Cardiol.* 1995; **25**: 4:871-8.
3. Krumholz HM, Larson M, Levy D. Prognosis of left ventricular geometric patterns in the Framingham Heart Study. *J Am Coll Cardiol.* 1995; **25**: 4:879-84.
4. Koren MJ, Devereux RB, Casale PN, Savage DD, Laragh JH. Relation of left ventricular mass and geometry to morbidity and mortality in uncomplicated essential hypertension. *Ann Intern Med.* 1991; **114**: 5:345-52.
5. Ristow B, Ali S, Na B, Turakhia MP, Whooley MA, Schiller NB. Predicting heart failure hospitalization and mortality by quantitative echocardiography: is body surface area the indexing method of choice? The Heart and Soul Study. *J Am Soc Echocardiogr.* 2010; **23**: 4:406-13.
6. Liao Y, Cooper RS, Durazo-Arvizu R, Mensah GA, Ghali JK. Prediction of mortality risk by different methods of indexation for left ventricular mass. *J Am Coll Cardiol.* 1997; **29**: 3:641-7.
7. Levy D, Garrison RJ, Savage DD, Kannel WB, Castelli WP. Prognostic implications of echocardiographically determined left ventricular mass in the Framingham Heart Study. *N Engl J Med.* 1990; **322**: 22:1561-6.
8. Benjamin EJ, D'Agostino RB, Belanger AJ, Wolf PA, Levy D. Left atrial size and the risk of stroke and death: the Framingham Heart Study. *Circulation.* 1995; **92**: 4:835-41.
9. Lang RM, Badano LP, Mor-Avi V, et al. Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American Society of Echocardiography and the European association of Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging.* 2015; **16**: 3:233-71.
10. Lang RM, Bierig M, Devereux RB, et al. Recommendations for chamber quantification. *Eur Heart J Cardiovasc Imaging.* 2006; **7**: 2:79-108.
11. Bluemke DA, Kronmal RA, Lima JA, et al. The relationship of left ventricular mass and geometry to incident cardiovascular events: the MESA (Multi-Ethnic Study of Atherosclerosis) study. *J Am Coll Cardiol.* 2008; **52**: 25:2148-55.
12. Chirinos JA, Segers P, De Buyzere ML, et al. Left ventricular mass: allometric scaling, normative values, effect of obesity, and prognostic performance. *Hypertension.* 2010; **56**: 1:91-8.
13. De Simone G, Daniels SR, Devereux RB, et al. Left ventricular mass and body size in normotensive children and adults: assessment of allometric relations and impact of overweight. *J Am Coll Cardiol.* 1992; **20**: 5:1251-60.
14. Dewey FE, Rosenthal D, Murphy Jr DJ, Froelicher VF, Ashley EA. Does size matter? Clinical applications of scaling cardiac size and function for body size. *Circulation.* 2008; **117**: 17:2279-87.
15. Devereux RB, Lutas EM, Casale PN, et al. Standardization of M-mode echocardiographic left ventricular anatomic measurements. *J Am Coll Cardiol.* 1984; **4**: 6:1222-30.
16. Airale L, Painsi A, Ianniello E, et al. Left atrial volume indexed for height² is a new sensitive marker for subclinical cardiac organ damage in female hypertensive patients. *Hypertens Res.* 2021:1-8.

17. Cuspidi C, Carugo S, Tadic M. Looking at the best indexing method of left atrial volume in the hypertensive setting. *Hypertens Res.* 2021;1-3.
18. De Simone G, Kizer JR, Chinali M, et al. Normalization for body size and population-attributable risk of left ventricular hypertrophy: the Strong Heart Study. *Am J Hypertens.* 2005; **18**: 2:191-6.
19. Magliano D, Liew D, Pater H, et al. Accuracy of the Australian National Death Index: comparison with adjudicated fatal outcomes among Australian participants in the Long-term Intervention with Pravastatin in Ischaemic Disease (LIPID) study. *Aust N Z J Public Health.* 2003; **27**: 6:649-53.
20. Playford D, Strange G, Celermajer DS, et al. Diastolic dysfunction and mortality in 436 360 men and women: the National Echo Database Australia (NEDA). *Eur Heart J Cardiovasc Imaging.* 2021; **22**: 5:505-15.
21. Sharp MK, Tokalić R, Gómez G, Wager E, Altman DG, Hren D. A cross-sectional bibliometric study showed suboptimal journal endorsement rates of STROBE and its extensions. *J Clin Epidemiol.* 2019; **107**:42-50.
22. Mosteller R. Simplified calculation of body-surface area. *N Engl J Med.* 1987; **317**: 17:1098-.
23. DuBois D. A formula to estimate the approximate surface area if height and body mass be known. *Arch Intern Med.* 1916; **17**:863-71.
24. Hume R. Prediction of lean body mass from height and weight. *J Clin Pathol.* 1966; **19**: 4:389-91.
25. Boer P. Estimated lean body mass as an index for normalization of body fluid volumes in humans. *Am J Physiol Renal Physiol.* 1984; **247**: 4:F632-F6.
26. Research DMGoO, James WPT, Waterlow JC. Research on Obesity: a Report of the DHSS/MRC Group; Compiled by WPT James: HM Stationery Office; 1976.
27. Team RC. R: A language and environment for statistical computing. 2013.
28. De Simone G, Devereux RB, Roman MJ, Alderman MH, Laragh JH. Relation of obesity and gender to left ventricular hypertrophy in normotensive and hypertensive adults. *Hypertension.* 1994; **23**: 5:600-6.
29. Jeyaprakash P, Moussad A, Pathan S, et al. A Systematic Review of Scaling Left Atrial Size: Are Alternative Indexation Methods Required for an Increasingly Obese Population? *J Am Soc Echocardiogr.* 2021; **34**: 10:1067-76.e3.
30. Hammond IW, Devereux RB, Alderman MH, Laragh JH. Relation of blood pressure and body build to left ventricular mass in normotensive and hypertensive employed adults. *J Am Coll Cardiol.* 1988; **12**: 4:996-1004.
31. Liebson PR, Grandits G, Prineas R, et al. Echocardiographic correlates of left ventricular structure among 844 mildly hypertensive men and women in the Treatment of Mild Hypertension Study (TOMHS). *Circulation.* 1993; **87**: 2:476-86.
32. Ramu B, Elwan AM, Coleman CI, Silverman DI, Gluck JA. The association between baseline left atrial volume index and all-cause mortality in patients with heart failure: a meta-analysis. *Conn Med.* 2015; **79**: 8.
33. Strange G, Stewart S, Celermajer D, et al. Poor long-term survival in patients with moderate aortic stenosis. *J Am Coll Cardiol.* 2019; **74**: 15:1851-63.
34. Lederle FA, Johnson GR, Wilson SE, et al. Relationship of age, gender, race, and body size to infrarenal aortic diameter. *J Vasc Surg.* 1997; **26**: 4:595-601.
35. Wolak A, Gransar H, Thomson LE, et al. Aortic size assessment by noncontrast cardiac computed tomography: normal limits by age, gender, and body surface area. *JACC Cardiovasc Imaging.* 2008; **1**: 2:200-9.
36. Vaupel JW. How change in age-specific mortality affects life expectancy. *Population Studies.* 1986; **40**: 1:147-57.

37. Romero-Ortuno R, Kenny RA. The frailty index in Europeans: association with age and mortality. *Age Ageing*. 2012; **41**: 5:684-9.

Figure texts

Figure 1. Flowchart describing patient inclusion. Exact numbers for the respective cardiac measure specific populations are given in Table 1.

Figure 2. Average prognostic strength for predicting 5-year cardiovascular mortality when indexing for body size for RA area, RV diameter, LA area, LV diameter, LV mass, and aortic sinus diameter. The horizontal axis represents the height exponent and the vertical axis represents the weight exponent of a body size indexation metric of the format $\text{height}^x * \text{weight}^y$. The color scale shows the prognostic strength which ranges from a C-statistic of 0.6 to 0.7, each increment representing a 1%-point improvement. The figure is composed of 50,000 points each representing an individual Cox regression. Existing body size metrics were plotted: h = height, w = weight, hw = height * weight, BMI = body mass index, BSA = body surface area, M = Mosteller, D = DuBois. In the healthy/overweight group, from unindexed to body surface area by Mosteller there is a 5%-point improvement, but further improvement is limited to <1%-point. Similar trends can be observed in the underweight and obese group.

Figure 3. The prognostic strength of indexing by body surface area across cardiac measures for the prediction of 5-year cardiovascular mortality. In the healthy/overweight group, left atrial size and left ventricular mass had the highest prognostic strength, with a C-statistic 10%-points higher than left ventricular size and most aortic dimensions. Right atrial and right ventricular sizes had an intermediate prognostic strength. Aortic sinus diameter had the strongest prognostic strength of all aortic measures, and aorta at sinotubular diameter had the weakest prognostic strength.

Table 1. Subject characteristics for patients with available follow-up data for the respective cardiac or aortic measures. RA = right atrial, RV = right ventricular, RVOT = right

ventricular outflow tract, LA = left atrial, LV = left ventricular, IVS = interventricular septum.

Table 2. C-statistic for 5-year cardiovascular mortality both unindexed and indexed by different body size metrics for representative anatomical measures, and their average. BMI = body mass index, BSA = body surface area, LBM = lean body mass, RA = right atrial, RV = right ventricular, RVOT = right ventricular outflow tract, LA = left atrial, LV = left ventricular, IVS = interventricular septum.

Appendix 1. C-statistic for long-term cardiovascular mortality both unindexed and indexed by different body size metrics for representative anatomical measures, and their average. BMI = body mass index, BSA = body surface area, LBM = lean body mass, RA = right atrial, RV = right ventricular, RVOT = right ventricular outflow tract, LA = left atrial, LV = left ventricular, IVS = interventricular septum.

Appendix 2. C-statistic for 5-year all-cause mortality both unindexed and indexed by different body size metrics for representative anatomical measures, and their average. BMI = body mass index, BSA = body surface area, LBM = lean body mass, RA = right atrial, RV = right ventricular, RVOT = right ventricular outflow tract, LA = left atrial, LV = left ventricular, IVS = interventricular septum.

Appendix 3. C-statistic for long-term all-cause mortality both unindexed and indexed by different body size metrics for representative anatomical measures, and their average. BMI = body mass index, BSA = body surface area, LBM = lean body mass, RA = right atrial, RV = right ventricular, RVOT = right ventricular outflow tract, LA = left atrial, LV = left ventricular, IVS = interventricular septum.

Appendix 4. Average prognostic strength of indexing for body size in aorta at sinotubular diameter.

Appendix 5. Average prognostic strength of indexing for body size in aortic sinus diameter.

Appendix 6. Average prognostic strength of indexing for body size in aortic arch diameter.

Appendix 7. Average prognostic strength of indexing for body size in aortic root diameter.

Appendix 8. Average prognostic strength of indexing for body size in ascending aorta diameter.

Appendix 9. Average prognostic strength of indexing for body size in LA area.

Appendix 10. Average prognostic strength of indexing for body size in LA diameter.

Appendix 11. Average prognostic strength of indexing for body size in LA volume.

Appendix 12. Average prognostic strength of indexing for body size in LV diameter.

Appendix 13. Average prognostic strength of indexing for body size in LV volume.

Appendix 14. Average prognostic strength of indexing for body size in IVS thickness.

Appendix 15. Average prognostic strength of indexing for body size in LV mass.

Appendix 16. Average prognostic strength of indexing for body size in RA area.

Appendix 17. Average prognostic strength of indexing for body size in RV diameter. Visual artifact present in BMI <18.5 kg/m² due to small population size (n=96).

Appendix 18. Average prognostic strength of indexing for body size in RVOT diameter.

Visual artifact present in BMI <18.5 kg/m² due to small population size (n=78).

Appendix 19. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for aorta at sinotubular diameter.

Appendix 20. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for aortic sinus diameter.

Appendix 21. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for aortic arch diameter.

Appendix 22. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for aortic root diameter.

Appendix 23. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for ascending aorta diameter.

Appendix 24. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for LA area.

Appendix 25. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for LA diameter.

Appendix 26. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for LA volume.

Appendix 27. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for LV diameter.

Appendix 28. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for LV volume.

Appendix 29. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for IVS diameter.

Appendix 30. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for LV mass.

Appendix 31. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for RA area.

Appendix 32. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for RV diameter.

Appendix 33. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for RVOT diameter.

Appendix 34. Long-term cardiovascular mortality for aorta at sinotubular diameter unindexed, indexed by BSA and indexed by weight

Appendix 35. Long-term cardiovascular mortality for aortic sinus diameter unindexed, indexed by BSA and indexed by weight.

Appendix 36. Long-term cardiovascular mortality for aortic arch diameter unindexed, indexed by BSA and indexed by weight.

Appendix 37. Long-term cardiovascular mortality for aortic root diameter unindexed, indexed by BSA and indexed by weight.

Appendix 38. Long-term cardiovascular mortality for ascending aorta diameter unindexed, indexed by BSA and indexed by weight.

Appendix 39. Long-term cardiovascular mortality for LA area unindexed, indexed by BSA and indexed by weight.

Appendix 40. Long-term cardiovascular mortality for LA diameter unindexed, indexed by BSA and indexed by weight.

Appendix 41. Long-term cardiovascular mortality for LA volume unindexed, indexed by BSA and indexed by weight.

Appendix 42. Long-term cardiovascular mortality for LV diameter unindexed, indexed by BSA and indexed by weight.

Appendix 43. Long-term cardiovascular mortality for LV volume unindexed, indexed by BSA and indexed by weight.

Appendix 44. Long-term cardiovascular mortality for IVS thickness unindexed, indexed by BSA and indexed by weight.

Appendix 45. Long-term cardiovascular mortality for LV mass unindexed, indexed by BSA and indexed by weight.

Appendix 46. Long-term cardiovascular mortality for RA area unindexed, indexed by BSA and indexed by weight.

Appendix 47. Long-term cardiovascular mortality for RV diameter unindexed, indexed by BSA and indexed by weight.

Appendix 48. Long-term cardiovascular mortality for RVOT diameter unindexed, indexed by BSA and indexed by weight.




Figure 1. Flowchart describing patient inclusion. Exact numbers for the respective cardiac measure specific populations are given in Table 1.

	RA area				RV diameter				RVOT diameter			
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	All	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	All	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	All
Subjects, n (%)	1730 (2)	53277 (64)	27797 (34)	82804 (100)	96 (3)	2412 (67)	1067 (30)	3575 (100)	78 (3)	1913 (67)	850 (30)	2841 (100)
Events, n (%)	199 (11.5)	3190 (6.0)	1129 (4.1)	4518 (5.5)	15 (15.6)	211 (8.7)	82 (7.7)	308 (8.6)	6 (7.7)	111 (5.8)	34 (4.0)	151 (5.3)
Female sex, n (%)	1185 (69)	22642 (43)	12572 (45)	36399 (44)	49 (51)	838 (35)	422 (40)	1309 (37)	49 (63)	894 (47)	451 (53)	1394 (49)
Age, years	69 [35-82]	67 [53-78]	64 [53-73]	66 [53-76]	60 [39-78]	63 [46-75]	60 [48-70]	62 [47-74]	39 [22-68]	55 [36-73]	59 [44-71]	56 [37-72]
Follow-up, years	4.1 [1.9-7.6]	6 [3.1-9.5]	5.9 [3.2-9.1]	5.9 [3.1-9.3]	6.6 [3-9.1]	6.7 [3.4-9.2]	6.9 [4.5-9.4]	6.8 [3.7-9.3]	3.8 [2.3-5.6]	3.9 [2.1-6.2]	3.7 [1.7-5.8]	3.8 [1.9-6.0]
	LA area				LA diameter				LA volume			
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	All	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	All	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	All
Subjects, n (%)	3163 (2)	86455 (63)	47371 (35)	136989 (100)	2711 (2)	73144 (65)	36366 (32)	112221 (100)	4290 (2)	138410 (67)	65093 (31)	207793 (100)
Events, n (%)	302 (9.5)	4903 (5.7)	1997 (4.2)	7202 (5.3)	227 (8.4)	4138 (5.7)	1543 (4.2)	5908 (5.3)	455 (10.6)	6928 (5.0)	2215 (3.4)	9598 (4.6)
Female sex, n (%)	2191 (69)	39157 (45)	22982 (49)	64330 (47)	1850 (68)	35702 (49)	19054 (52)	56606 (50)	3136 (73)	65665 (47)	32574 (50)	101375 (49)
Age, years	63 [34-79]	65 [49-76]	63 [52-72]	64 [50-75]	61 [34-77]	63 [47-75]	62 [51-71]	63 [48-74]	68 [40-82]	65 [50-76]	62 [51-72]	64 [51-75]
Follow-up, years	3.9 [1.9-7.3]	5.1 [2.8-9]	5.0 [2.9-8.7]	5.0 [2.8-8.8]	5.6 [2.6-9.9]	6.8 [3.7-11.5]	6.4 [3.7-10.5]	6.6 [3.7-11.1]	4.4 [2.3-7.7]	6.0 [3.5-9.2]	6.0 [3.5-9.2]	5.9 [3.5-9.2]
	LV diameter				LV volume				IVS thickness			
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	All	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	All	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	All
Subjects, n (%)	8410 (2)	222977 (66)	106094 (31)	337481 (100)	1856 (2)	67279 (71)	25284 (27)	94419 (100)	8513 (3)	225193 (66)	106764 (31)	340470 (100)
Events, n (%)	673 (8.0)	10941 (4.9)	3685 (3.5)	15299 (4.5)	177 (9.5)	2890 (4.3)	668 (2.6)	3735 (4.0)	677 (8.0)	11163 (5.0)	3723 (3.5)	15563 (4.6)
Female sex, n (%)	5705 (68)	104928 (47)	53137 (50)	163770 (49)	1406 (76)	31877 (47)	12855 (51)	46138 (49)	5785 (68)	105785 (47)	53476 (50)	165046 (49)
Age, years	63 [38-79]	63 [48-75]	62 [50-71]	63 [49-74]	70 [44-82]	64 [50-76]	62 [51-71]	63 [50-75]	63 [38-79]	64 [48-75]	62 [50-71]	63 [49-74]
Follow-up, years	4.6 [2.2-8.5]	5.5 [3.1-8.9]	5.3 [3.1-8.6]	5.4 [3.1-8.8]	4.5 [2.7-7.1]	5.5 [3.6-8.2]	5.5 [3.7-8.1]	5.5 [3.6-8.2]	4.6 [2.2-8.5]	5.5 [3.1-8.9]	5.3 [3.1-8.5]	5.4 [3.1-8.8]

Table 1. Subject characteristics for patients with available follow-up data for the respective cardiac or aortic measures. RA = right atrial, RV = right ventricular, RVOT = right ventricular outflow tract, LA = left atrial, LV = left ventricular, IVS = interventricular septum.

	LV mass				Aortic sinus diameter				Aorta at sinotubular diameter			
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	All	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	All	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	All
Subjects, n (%)	8263 (2)	218970 (66)	103726 (31)	330959 (100)	2829 (2)	84509 (68)	37792 (30)	125130 (100)	1730 (2)	53277 (64)	27797 (34)	82804 (100)
Events, n (%)	650 (7.9)	10586 (4.8)	3523 (3.4)	14759 (4.5)	284 (10.0)	4238 (5.0)	1244 (3.3)	5766 (4.6)	199 (11.5)	3190 (6.0)	1129 (4.1)	4518 (5.5)
Female sex, n (%)	5619 (68)	103367 (47)	52210 (50)	161196 (49)	2031 (72)	39146 (46)	18168 (48)	59345 (47)	1185 (69)	22642 (43)	12572 (45)	36399 (44)
Age, years	63 [38-79]	63 [48-75]	62 [50-71]	63 [49-74]	68 [39-82]	65 [50-77]	62 [51-72]	64 [50-75]	69 [35-82]	67 [53-78]	64 [53-73]	66 [53-76]
Follow-up, years	4.6 [2.2-8.5]	5.5 [3.1-8.9]	5.3 [3.1-8.5]	5.4 [3-8.7]	4.3 [2.1-7.4]	5.9 [3.4-8.8]	5.9 [3.5-8.7]	5.9 [3.4-8.8]	4.1 [1.9-7.6]	6.0 [3.1-9.5]	5.9 [3.2-9.1]	5.9 [3.1-9.3]
	Aortic root diameter				Ascending aorta diameter				Aortic arch diameter			
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	All	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	All	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	All
Subjects, n (%)	6028 (2)	180514 (67)	83013 (31)	269555 (100)	3689 (2)	102736 (63)	57160 (35)	163585 (100)	761 (2)	21122 (63)	11531 (35)	33414 (100)
Events, n (%)	541 (9.0)	8973 (5.0)	2846 (3.4)	12360 (4.6)	295 (8.0)	5588 (5.4)	2253 (3.9)	8136 (5.0)	43 (5.7)	803 (3.8)	312 (2.7)	1158 (3.5)
Female sex, n (%)	4171 (69)	85985 (48)	42230 (51)	132386 (49)	2442 (66)	46046 (45)	27458 (48)	75946 (46)	509 (67)	9575 (45)	5511 (48)	15595 (47)
Age, years	65 [38-80]	64 [49-76]	62 [51-71]	63 [49-74]	63 [37-79]	65 [50-76]	63 [52-72]	64 [51-75]	50 [24-75]	63 [46-75]	61 [49-71]	62 [47-73]
Follow-up, years	4.8 [2.3-8.5]	5.9 [3.4-9.6]	5.8 [3.5-9.2]	5.9 [3.4-9.4]	4.2 [2-8.1]	4.7 [2.5-7.7]	4.7 [2.6-7.6]	4.7 [2.6-7.7]	3.6 [1.8-6.6]	3.9 [2.1-6.8]	3.7 [2.1-6.6]	3.8 [2.1-6.7]

Table 1. Subject characteristics for patients with available follow-up data for the respective cardiac or aortic measures. RA = right atrial, RV = right ventricular, RVOT = right ventricular outflow tract, LA = left atrial, LV = left ventricular, IVS = interventricular septum.






Figure 2. Average prognostic strength for predicting 5-year cardiovascular mortality when indexing for body size for RA area, RV diameter, LA area, LV diameter, LV mass, and aortic sinus diameter. The horizontal axis represents the height exponent and the vertical axis represents the weight exponent of a body size indexation metric of the format $\text{height}^x * \text{weight}^y$. The color scale shows the prognostic strength which ranges from a C-statistic of 0.6 to 0.7, each increment representing a 1%-point improvement. The figure is composed of 50,000 points each representing an individual Cox regression. Existing body size metrics were plotted: h = height, w = weight, hw = height * weight, BMI = body mass index, BSA = body surface area, M = Mosteller, D = DuBois. In the healthy/overweight group, from unindexed to body surface area by Mosteller there is a 5%-point improvement, but further improvement is limited to <1%-point. Similar trends can be observed in the underweight and obese group.

	LV mass			LA area			RA area			Aortic sinus diameter			RV diameter			LV diameter			RVOT diameter			Average		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.726	0.741	0.681	0.761	0.725	0.647	0.723	0.680	0.602	0.652	0.646	0.567	0.608	0.646	0.650	0.563	0.618	0.592	0.630	0.569	0.607	0.666	0.661	0.621
Weight ^{1.5}	0.718	0.745	0.677	0.758	0.715	0.638	0.726	0.682	0.601	0.638	0.635	0.559	0.417	0.638	0.639	0.566	0.615	0.581	0.598	0.565	0.592	0.632	0.656	0.612
Height*Weight	0.729	0.745	0.687	0.758	0.721	0.647	0.723	0.684	0.604	0.638	0.638	0.569	0.601	0.644	0.646	0.568	0.617	0.591	0.601	0.566	0.625	0.660	0.659	0.624
BSA[Mosteller] ^{1.5}	0.733	0.740	0.687	0.760	0.725	0.651	0.721	0.680	0.604	0.648	0.646	0.575	0.617	0.648	0.654	0.567	0.618	0.598	0.627	0.570	0.631	0.667	0.661	0.628
BSA[Du Bois] ^{1.5}	0.736	0.739	0.688	0.759	0.725	0.653	0.719	0.680	0.604	0.646	0.645	0.578	0.619	0.647	0.654	0.569	0.617	0.600	0.621	0.570	0.641	0.667	0.660	0.631
BSA[Mosteller]	0.734	0.727	0.681	0.759	0.724	0.652	0.714	0.673	0.601	0.657	0.647	0.580	0.620	0.644	0.660	0.562	0.610	0.602	0.638	0.569	0.635	0.669	0.656	0.630
BSA[Du Bois]	0.736	0.727	0.682	0.759	0.723	0.653	0.714	0.673	0.602	0.655	0.646	0.583	0.625	0.644	0.662	0.565	0.609	0.604	0.641	0.569	0.642	0.671	0.656	0.632
LBM[Hume]	0.741	0.732	0.686	0.753	0.718	0.647	0.714	0.674	0.598	0.637	0.628	0.568	0.630	0.638	0.649	0.572	0.603	0.590	0.601	0.560	0.645	0.664	0.650	0.626
LBM[Boer]	0.741	0.730	0.679	0.754	0.712	0.634	0.714	0.671	0.589	0.639	0.616	0.546	0.629	0.630	0.629	0.572	0.593	0.569	0.601	0.553	0.641	0.664	0.644	0.612
LBM[James]	0.730	0.731	0.673	0.754	0.714	0.632	0.716	0.671	0.585	0.639	0.618	0.547	0.612	0.631	0.628	0.555	0.594	0.566	0.593	0.555	0.652	0.657	0.645	0.612
BMI	0.706	0.710	0.652	0.751	0.711	0.631	0.705	0.659	0.590	0.633	0.617	0.535	0.576	0.627	0.642	0.527	0.588	0.566	0.604	0.562	0.537	0.643	0.639	0.593
Height ^{2.7}	0.747	0.723	0.687	0.753	0.716	0.653	0.710	0.673	0.603	0.633	0.618	0.582	0.663	0.636	0.656	0.585	0.600	0.602	0.632	0.561	0.674	0.675	0.647	0.637
Height ^{2.13}	0.745	0.719	0.684	0.755	0.717	0.655	0.709	0.670	0.602	0.642	0.622	0.586	0.673	0.636	0.660	0.581	0.598	0.606	0.652	0.564	0.669	0.680	0.647	0.637
Height ²	0.744	0.718	0.683	0.755	0.717	0.655	0.709	0.669	0.602	0.644	0.622	0.586	0.674	0.636	0.662	0.580	0.598	0.606	0.658	0.565	0.668	0.681	0.646	0.637
Height ^{1.5}	0.740	0.712	0.678	0.755	0.715	0.653	0.706	0.665	0.600	0.650	0.619	0.586	0.675	0.634	0.664	0.574	0.592	0.605	0.644	0.565	0.663	0.678	0.643	0.636
Height	0.736	0.705	0.672	0.753	0.712	0.650	0.702	0.660	0.597	0.651	0.608	0.579	0.672	0.629	0.665	0.565	0.583	0.600	0.647	0.563	0.647	0.675	0.637	0.630
<i>Unindexed</i>	0.723	0.687	0.655	0.745	0.699	0.640	0.690	0.646	0.590	0.625	0.569	0.551	0.659	0.612	0.662	0.541	0.558	0.577	0.595	0.553	0.607	0.654	0.618	0.612

Table 2. C-statistic for 5-year cardiovascular mortality both unindexed and indexed by different body size metrics for representative anatomical measures, and their average. BMI = body mass index, BSA = body surface area, LBM = lean body mass, RA = right atrial, RV = right ventricular, RVOT = right ventricular outflow tract, LA = left atrial, LV = left ventricular, IVS = interventricular septum.




Figure 3. The prognostic strength of indexing by body surface area across cardiac measures for the prediction of 5-year cardiovascular mortality. In the healthy/overweight group, left atrial size and left ventricular mass had the highest prognostic strength, with a C-statistic 10%-points higher than left ventricular size and most aortic dimensions. Right atrial and right ventricular sizes had an intermediate prognostic strength. Aortic sinus diameter had the strongest prognostic strength of all aortic measures, and aorta at sinotubular diameter had the weakest prognostic strength. RA = right atrial, RV = right ventricular, RVOT = right ventricular outflow tract, LA = left atrial, LV = left ventricular, IVS = interventricular septum.

	LV mass			LA area			RA area			Aortic sinus diameter			RV diameter			LV diameter			RVOT diameter			Average		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.721	0.737	0.681	0.764	0.724	0.651	0.729	0.684	0.610	0.655	0.644	0.570	0.428	0.641	0.641	0.560	0.612	0.591	0.630	0.570	0.606	0.641	0.659	0.621
Weight ^{1.5}	0.712	0.741	0.677	0.761	0.714	0.642	0.731	0.684	0.608	0.642	0.632	0.563	0.449	0.632	0.630	0.564	0.610	0.582	0.598	0.566	0.590	0.637	0.654	0.613
Height*Weight	0.725	0.742	0.686	0.761	0.721	0.651	0.729	0.687	0.611	0.643	0.637	0.573	0.434	0.638	0.638	0.566	0.613	0.591	0.601	0.567	0.623	0.637	0.658	0.625
BSA[Mosteller] ^{1.5}	0.729	0.737	0.686	0.763	0.725	0.656	0.727	0.685	0.611	0.652	0.646	0.579	0.579	0.643	0.645	0.564	0.613	0.598	0.627	0.571	0.630	0.663	0.660	0.629
BSA[Du Bois] ^{1.5}	0.732	0.736	0.687	0.762	0.725	0.657	0.726	0.685	0.612	0.650	0.645	0.582	0.581	0.642	0.646	0.567	0.613	0.600	0.621	0.571	0.640	0.663	0.659	0.632
BSA[Mosteller]	0.730	0.724	0.680	0.762	0.724	0.657	0.721	0.678	0.609	0.659	0.647	0.583	0.580	0.641	0.651	0.560	0.606	0.601	0.638	0.571	0.636	0.664	0.656	0.631
BSA[Du Bois]	0.732	0.723	0.681	0.762	0.723	0.658	0.721	0.678	0.610	0.659	0.646	0.586	0.583	0.641	0.653	0.563	0.605	0.603	0.641	0.571	0.643	0.666	0.655	0.633
LBM[Hume]	0.738	0.730	0.685	0.757	0.718	0.652	0.720	0.679	0.607	0.642	0.630	0.573	0.591	0.632	0.642	0.571	0.600	0.590	0.601	0.562	0.644	0.660	0.650	0.627
LBM[Boer]	0.738	0.728	0.678	0.758	0.713	0.639	0.721	0.676	0.597	0.645	0.618	0.552	0.592	0.624	0.624	0.572	0.591	0.568	0.601	0.554	0.639	0.661	0.643	0.614
LBM[James]	0.725	0.729	0.672	0.757	0.714	0.636	0.723	0.676	0.593	0.643	0.620	0.552	0.579	0.625	0.624	0.553	0.591	0.565	0.593	0.556	0.652	0.653	0.644	0.613
BMI	0.700	0.705	0.651	0.754	0.709	0.635	0.711	0.662	0.597	0.630	0.611	0.535	0.459	0.625	0.631	0.477	0.581	0.564	0.604	0.564	0.540	0.619	0.637	0.593
Height ^{2.7}	0.745	0.721	0.686	0.757	0.717	0.657	0.718	0.678	0.610	0.639	0.622	0.587	0.619	0.632	0.650	0.586	0.599	0.601	0.632	0.563	0.674	0.671	0.647	0.638
Height ^{2.13}	0.742	0.717	0.683	0.759	0.718	0.659	0.717	0.675	0.610	0.647	0.626	0.591	0.629	0.632	0.653	0.582	0.597	0.605	0.652	0.566	0.670	0.675	0.647	0.639
Height ²	0.742	0.716	0.682	0.759	0.718	0.659	0.716	0.674	0.609	0.649	0.626	0.591	0.631	0.633	0.654	0.580	0.596	0.605	0.658	0.567	0.669	0.676	0.647	0.639
Height ^{1.5}	0.738	0.710	0.677	0.758	0.716	0.658	0.714	0.670	0.608	0.654	0.622	0.590	0.631	0.632	0.656	0.574	0.590	0.604	0.644	0.567	0.665	0.673	0.644	0.637
Height	0.732	0.702	0.670	0.757	0.712	0.655	0.710	0.665	0.605	0.653	0.611	0.582	0.625	0.627	0.656	0.565	0.581	0.598	0.647	0.565	0.650	0.670	0.638	0.631
<i>Unindexed</i>	0.719	0.683	0.654	0.748	0.699	0.644	0.698	0.651	0.597	0.623	0.570	0.551	0.612	0.613	0.652	0.540	0.555	0.574	0.595	0.555	0.611	0.648	0.618	0.612




Appendix 1. C-statistic for long-term cardiovascular mortality both unindexed and indexed by different body size metrics for representative anatomical measures, and their average. BMI = body mass index, BSA = body surface area, LBM = lean body mass, RA = right atrial, RV = right ventricular, RVOT = right ventricular outflow tract, LA = left atrial, LV = left ventricular, IVS = interventricular septum.

	LV mass			LA area			RA area			Aortic sinus diameter			RV diameter			LV diameter			RVOT diameter			Average		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.622	0.654	0.606	0.653	0.644	0.602	0.660	0.631	0.588	0.661	0.633	0.569	0.619	0.631	0.620	0.497	0.554	0.535	0.527	0.585	0.576	0.606	0.619	0.585
Weight ^{1.5}	0.620	0.660	0.603	0.655	0.641	0.596	0.663	0.633	0.587	0.648	0.619	0.561	0.617	0.626	0.605	0.484	0.561	0.535	0.533	0.585	0.573	0.603	0.618	0.580
Height*Weight	0.626	0.657	0.610	0.652	0.642	0.602	0.660	0.633	0.590	0.648	0.623	0.569	0.626	0.628	0.611	0.487	0.556	0.538	0.538	0.586	0.592	0.605	0.618	0.588
BSA[Mosteller] ^{1.5}	0.627	0.652	0.609	0.652	0.643	0.605	0.658	0.630	0.590	0.658	0.634	0.575	0.630	0.630	0.621	0.495	0.550	0.539	0.533	0.586	0.592	0.608	0.618	0.590
BSA[Du Bois] ^{1.5}	0.629	0.651	0.611	0.651	0.642	0.606	0.657	0.629	0.591	0.656	0.632	0.576	0.632	0.629	0.621	0.506	0.548	0.540	0.530	0.585	0.600	0.609	0.617	0.592
BSA[Mosteller]	0.625	0.642	0.605	0.648	0.639	0.605	0.654	0.624	0.588	0.665	0.639	0.578	0.628	0.627	0.629	0.507	0.539	0.537	0.522	0.579	0.589	0.607	0.613	0.590
BSA[Du Bois]	0.627	0.641	0.606	0.648	0.639	0.606	0.653	0.624	0.589	0.664	0.638	0.580	0.633	0.625	0.630	0.506	0.537	0.538	0.525	0.578	0.596	0.608	0.612	0.592
LBM[Hume]	0.631	0.643	0.608	0.644	0.634	0.601	0.651	0.622	0.586	0.644	0.616	0.567	0.653	0.620	0.615	0.503	0.533	0.532	0.532	0.575	0.605	0.608	0.606	0.588
LBM[Boer]	0.631	0.640	0.602	0.645	0.629	0.591	0.651	0.618	0.579	0.647	0.604	0.549	0.656	0.614	0.600	0.503	0.525	0.518	0.532	0.568	0.604	0.609	0.600	0.578
LBM[James]	0.620	0.641	0.600	0.644	0.631	0.590	0.653	0.619	0.578	0.645	0.606	0.547	0.628	0.616	0.603	0.509	0.526	0.518	0.538	0.570	0.622	0.605	0.601	0.580
BMI	0.604	0.633	0.586	0.642	0.632	0.589	0.648	0.617	0.577	0.640	0.612	0.541	0.575	0.617	0.621	0.529	0.533	0.517	0.507	0.563	0.517	0.592	0.601	0.564
Height ^{2.7}	0.638	0.636	0.611	0.644	0.632	0.606	0.650	0.621	0.591	0.642	0.607	0.576	0.680	0.613	0.622	0.518	0.531	0.543	0.549	0.575	0.627	0.617	0.602	0.597
Height ^{2.13}	0.635	0.633	0.609	0.644	0.632	0.607	0.649	0.619	0.590	0.650	0.613	0.581	0.676	0.614	0.628	0.510	0.527	0.543	0.538	0.573	0.622	0.615	0.602	0.597
Height ²	0.635	0.633	0.608	0.644	0.632	0.607	0.649	0.619	0.590	0.652	0.614	0.581	0.677	0.614	0.629	0.508	0.525	0.542	0.537	0.572	0.619	0.615	0.601	0.597
Height ^{1.5}	0.631	0.629	0.605	0.643	0.631	0.606	0.647	0.616	0.588	0.657	0.615	0.581	0.668	0.613	0.632	0.499	0.520	0.540	0.524	0.567	0.612	0.610	0.599	0.595
Height	0.626	0.624	0.600	0.640	0.628	0.604	0.645	0.613	0.585	0.656	0.609	0.575	0.658	0.610	0.634	0.511	0.512	0.535	0.493	0.559	0.597	0.604	0.593	0.590
<i>Unindexed</i>	<i>0.615</i>	<i>0.611</i>	<i>0.589</i>	<i>0.632</i>	<i>0.618</i>	<i>0.595</i>	<i>0.637</i>	<i>0.604</i>	<i>0.578</i>	<i>0.627</i>	<i>0.578</i>	<i>0.549</i>	<i>0.626</i>	<i>0.597</i>	<i>0.631</i>	<i>0.531</i>	<i>0.498</i>	<i>0.520</i>	<i>0.516</i>	<i>0.539</i>	<i>0.559</i>	<i>0.598</i>	<i>0.578</i>	<i>0.574</i>




Appendix 2. C-statistic for 5-year all-cause mortality both unindexed and indexed by different body size metrics for representative anatomical measures, and their average. BMI = body mass index, BSA = body surface area, LBM = lean body mass, RA = right atrial, RV = right ventricular, RVOT = right ventricular outflow tract, LA = left atrial, LV = left ventricular, IVS = interventricular septum.

	LV mass			LA area			RA area			Aortic sinus diameter			RV diameter			LV diameter			RVOT diameter			Average		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.621	0.654	0.608	0.657	0.645	0.604	0.665	0.634	0.592	0.665	0.631	0.570	0.433	0.628	0.603	0.495	0.552	0.537	0.527	0.584	0.577	0.581	0.618	0.584
Weight ^{1.5}	0.619	0.659	0.606	0.658	0.642	0.598	0.669	0.635	0.589	0.653	0.616	0.563	0.441	0.620	0.590	0.483	0.559	0.536	0.533	0.582	0.576	0.580	0.616	0.580
Height*Weight	0.626	0.658	0.612	0.656	0.644	0.605	0.666	0.636	0.593	0.653	0.622	0.571	0.569	0.624	0.597	0.515	0.555	0.540	0.538	0.585	0.594	0.603	0.618	0.587
BSA[Mosteller] ^{1.5}	0.627	0.653	0.612	0.655	0.645	0.607	0.664	0.634	0.594	0.662	0.632	0.576	0.576	0.628	0.605	0.507	0.550	0.541	0.533	0.585	0.593	0.603	0.618	0.590
BSA[Du Bois] ^{1.5}	0.629	0.652	0.613	0.654	0.644	0.608	0.663	0.634	0.594	0.660	0.631	0.579	0.578	0.627	0.606	0.508	0.548	0.542	0.530	0.585	0.600	0.603	0.617	0.592
BSA[Mosteller]	0.625	0.643	0.608	0.651	0.641	0.608	0.659	0.629	0.592	0.668	0.638	0.579	0.578	0.626	0.613	0.505	0.538	0.539	0.522	0.579	0.589	0.601	0.613	0.590
BSA[Du Bois]	0.627	0.642	0.608	0.651	0.641	0.608	0.658	0.628	0.593	0.667	0.637	0.582	0.583	0.625	0.614	0.496	0.537	0.540	0.525	0.579	0.596	0.601	0.613	0.592
LBM[Hume]	0.632	0.644	0.610	0.648	0.637	0.603	0.657	0.627	0.590	0.650	0.617	0.571	0.593	0.618	0.600	0.506	0.534	0.534	0.532	0.575	0.605	0.603	0.607	0.588
LBM[Boer]	0.632	0.642	0.605	0.648	0.632	0.594	0.657	0.623	0.582	0.652	0.605	0.553	0.598	0.613	0.587	0.506	0.527	0.520	0.532	0.567	0.604	0.604	0.601	0.578
LBM[James]	0.621	0.643	0.603	0.648	0.633	0.592	0.658	0.624	0.580	0.650	0.607	0.551	0.571	0.614	0.591	0.507	0.527	0.520	0.538	0.569	0.621	0.599	0.602	0.580
BMI	0.602	0.632	0.587	0.644	0.632	0.591	0.651	0.620	0.580	0.640	0.608	0.540	0.463	0.615	0.605	0.530	0.529	0.517	0.507	0.563	0.519	0.577	0.600	0.563
Height ^{2.7}	0.640	0.639	0.613	0.648	0.635	0.609	0.655	0.626	0.594	0.648	0.610	0.580	0.618	0.613	0.609	0.521	0.534	0.545	0.549	0.575	0.626	0.611	0.605	0.597
Height ^{2.13}	0.636	0.635	0.611	0.648	0.635	0.609	0.655	0.625	0.594	0.655	0.616	0.585	0.617	0.615	0.613	0.513	0.529	0.545	0.538	0.574	0.621	0.609	0.604	0.597
Height ²	0.636	0.635	0.610	0.647	0.635	0.609	0.654	0.624	0.593	0.657	0.617	0.585	0.619	0.616	0.614	0.511	0.528	0.544	0.537	0.573	0.619	0.609	0.604	0.596
Height ^{1.5}	0.631	0.630	0.607	0.646	0.633	0.608	0.652	0.622	0.592	0.660	0.618	0.584	0.613	0.615	0.616	0.502	0.522	0.542	0.524	0.569	0.611	0.604	0.601	0.594
Height	0.626	0.625	0.602	0.643	0.630	0.606	0.649	0.618	0.589	0.658	0.611	0.577	0.605	0.612	0.618	0.491	0.514	0.536	0.493	0.561	0.596	0.595	0.596	0.589
<i>Unindexed</i>	<i>0.614</i>	<i>0.612</i>	<i>0.591</i>	<i>0.634</i>	<i>0.621</i>	<i>0.597</i>	<i>0.640</i>	<i>0.609</i>	<i>0.582</i>	<i>0.627</i>	<i>0.579</i>	<i>0.549</i>	<i>0.583</i>	<i>0.601</i>	<i>0.614</i>	<i>0.530</i>	<i>0.499</i>	<i>0.521</i>	<i>0.516</i>	<i>0.541</i>	<i>0.558</i>	<i>0.592</i>	<i>0.580</i>	<i>0.573</i>


Appendix 3. C-statistic for long-term all-cause mortality both unindexed and indexed by different body size metrics for representative anatomical measures, and their average. BMI = body mass index, BSA = body surface area, LBM = lean body mass, RA = right atrial, RV = right ventricular, RVOT = right ventricular outflow tract, LA = left atrial, LV = left ventricular, IVS = interventricular septum.






Appendix 4. Average prognostic strength of indexing for body size in aorta at sinotubular diameter.






Appendix 5. Average prognostic strength of indexing for body size in aortic sinus diameter.






Appendix 6. Average prognostic strength of indexing for body size in aortic arch diameter.






Appendix 7. Average prognostic strength of indexing for body size in aortic root diameter.






Appendix 8. Average prognostic strength of indexing for body size in ascending aorta diameter.






Appendix 9. Average prognostic strength of indexing for body size in LA area.






Appendix 10. Average prognostic strength of indexing for body size in LA diameter.






Appendix 11. Average prognostic strength of indexing for body size in LA volume.






Appendix 12. Average prognostic strength of indexing for body size in LV diameter.






Appendix 13. Average prognostic strength of indexing for body size in LV volume.




Appendix 14. Average prognostic strength of indexing for body size in IVS thickness.




Appendix 15. Average prognostic strength of indexing for body size in LV mass.



Appendix 16. Average prognostic strength of indexing for body size in RA area.



Appendix 17. Average prognostic strength of indexing for body size in RV diameter. Visual artifact present in BMI <math><18.5</math> (kg/m²) due to small population size (n=96).



Appendix 18. Average prognostic strength of indexing for body size in RVOT diameter. Visual artifact present in BMI <18.5 (kg/m²) due to small population size (n=78).

	C statistic			Hazard Ratio per 1 SD increment			Chi-square			p value			ΔAIC			Akaike Weight		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.598	0.573	0.519	1.40 [1.24-1.59]	1.32 [1.28-1.37]	1.10 [1.04-1.17]	28	298	11	<0.0001	<0.0001	0.0012	3	55	7	0.060	1.15E-12	0.017
Weight^1.5	0.611	0.585	0.528	1.40 [1.24-1.58]	1.34 [1.30-1.38]	1.13 [1.06-1.19]	31	372	17	<0.0001	<0.0001	<0.0001	2	0	1	0.107	1.000	0.317
Height*Weight	0.609	0.579	0.528	1.43 [1.26-1.62]	1.33 [1.29-1.37]	1.13 [1.07-1.19]	31	328	18	<0.0001	<0.0001	<0.0001	0	32	0	0.261	9.75E-08	0.489
BSA[Mosteller]^1.5	0.600	0.569	0.521	1.42 [1.25-1.62]	1.31 [1.27-1.35]	1.11 [1.05-1.17]	29	273	12	<0.0001	<0.0001	0.00042	2	76	5	0.109	2.45E-17	0.044
BSA[Du Bois]^1.5	0.600	0.567	0.522	1.43 [1.25-1.62]	1.30 [1.26-1.34]	1.11 [1.05-1.18]	29	258	13	<0.0001	<0.0001	0.00029	1	90	4	0.124	3.00E-20	0.062
BSA[Mosteller]	0.585	0.553	0.513	1.39 [1.22-1.58]	1.26 [1.22-1.30]	1.08 [1.02-1.14]	24	182	6	<0.0001	<0.0001	0.012	6	157	11	0.013	7.94E-35	2.18E-03
BSA[Du Bois]	0.586	0.552	0.514	1.39 [1.22-1.58]	1.25 [1.21-1.29]	1.08 [1.02-1.14]	24	173	7	<0.0001	<0.0001	0.0097	6	166	10	0.015	7.94E-37	2.65E-03
LBM[Hume]	0.596	0.554	0.516	1.42 [1.25-1.61]	1.25 [1.21-1.29]	1.08 [1.02-1.15]	28	180	8	<0.0001	<0.0001	0.0056	2	163	10	0.079	5.01E-36	4.13E-03
LBM[Boer]	0.596	0.550	0.508	1.42 [1.25-1.61]	1.23 [1.19-1.27]	1.05 [0.99-1.12]	29	152	3	<0.0001	<0.0001	0.079	2	189	14	0.097	1.00E-41	4.37E-04
LBM[James]	0.592	0.551	0.503	1.40 [1.23-1.60]	1.23 [1.19-1.27]	1.05 [1.00-1.10]	26	159	5	<0.0001	<0.0001	0.030	4	181	13	0.032	4.41E-40	5.76E-04
BMI	0.551	0.544	0.495	1.25 [1.11-1.42]	1.21 [1.17-1.25]	1.01 [0.95-1.07]	12	124	0	0.00042	<0.0001	0.77	18	213	17	4.01E-05	6.21E-47	9.85E-05
Height^2.7	0.595	0.547	0.523	1.41 [1.24-1.60]	1.22 [1.18-1.26]	1.11 [1.05-1.17]	27	144	13	<0.0001	<0.0001	0.00036	3	196	5	0.058	3.42E-43	0.047
Height^2.13	0.589	0.541	0.518	1.39 [1.22-1.59]	1.20 [1.16-1.24]	1.09 [1.03-1.16]	25	118	9	<0.0001	<0.0001	0.0028	5	218	8	0.022	3.70E-48	7.90E-03
Height^2	0.587	0.539	0.517	1.39 [1.22-1.58]	1.20 [1.16-1.24]	1.09 [1.03-1.15]	24	111	8	<0.0001	<0.0001	0.0045	6	225	9	0.016	1.59E-49	5.10E-03
Height^1.5	0.579	0.532	0.513	1.36 [1.20-1.56]	1.17 [1.13-1.21]	1.07 [1.01-1.13]	21	83	5	<0.0001	<0.0001	0.032	8	252	13	3.98E-03	1.89E-55	9.43E-04
Height	0.568	0.525	0.508	1.33 [1.16-1.52]	1.14 [1.10-1.18]	1.04 [0.98-1.10]	18	52	2	<0.0001	<0.0001	0.19	12	281	15	6.39E-04	7.86E-62	2.27E-04
<i>Unindexed</i>	<i>0.541</i>	<i>0.507</i>	<i>0.506</i>	<i>1.22 [1.07-1.40]</i>	<i>1.05 [1.02-1.09]</i>	<i>0.99 [0.93-1.04]</i>	<i>9</i>	<i>8</i>	<i>0</i>	<i>0.0032</i>	<i>0.0049</i>	<i>0.62</i>	<i>20</i>	<i>325</i>	<i>17</i>	<i>9.32E-06</i>	<i>2.26E-71</i>	<i>1.07E-04</i>

Appendix 19. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for aorta at sinotubular diameter.

	C statistic			Hazard Ratio per 1 SD increment			Chi-square			p value			ΔAIC			Akaike Weight		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.652	0.646	0.567	1.47 [1.35-1.61]	1.59 [1.55-1.63]	1.28 [1.21-1.35]	72	1332	79	<0.0001	<0.0001	<0.0001	11	0	48	2.18E-03	1.000	1.94E-11
Weight^1.5	0.638	0.635	0.559	1.35 [1.25-1.45]	1.53 [1.50-1.57]	1.25 [1.18-1.32]	57	1172	66	<0.0001	<0.0001	<0.0001	27	140	61	5.25E-07	4.35E-31	3.60E-14
Height*Weight	0.638	0.638	0.569	1.45 [1.33-1.60]	1.55 [1.51-1.59]	1.28 [1.22-1.35]	62	1200	94	<0.0001	<0.0001	<0.0001	16	110	39	1.30E-04	1.10E-24	1.80E-09
BSA[Mosteller]^1.5	0.648	0.646	0.575	1.53 [1.39-1.69]	1.59 [1.55-1.63]	1.31 [1.24-1.38]	71	1307	103	<0.0001	<0.0001	<0.0001	6	22	28	0.021	1.61E-05	5.13E-07
BSA[Du Bois]^1.5	0.646	0.645	0.578	1.54 [1.39-1.71]	1.58 [1.54-1.62]	1.32 [1.25-1.38]	70	1260	114	<0.0001	<0.0001	<0.0001	6	57	20	0.027	3.84E-13	2.66E-05
BSA[Mosteller]	0.657	0.647	0.580	1.58 [1.42-1.75]	1.57 [1.53-1.61]	1.31 [1.25-1.38]	76	1263	106	<0.0001	<0.0001	<0.0001	0	76	25	0.376	2.46E-17	2.03E-06
BSA[Du Bois]	0.655	0.646	0.583	1.58 [1.43-1.76]	1.56 [1.53-1.60]	1.32 [1.26-1.39]	76	1222	116	<0.0001	<0.0001	<0.0001	0	108	17	0.452	4.32E-24	1.07E-04
LBM[Hume]	0.637	0.628	0.568	1.52 [1.37-1.69]	1.50 [1.46-1.54]	1.27 [1.21-1.34]	64	987	92	<0.0001	<0.0001	<0.0001	10	287	42	2.77E-03	4.30E-63	4.49E-10
LBM[Boer]	0.639	0.616	0.546	1.53 [1.38-1.70]	1.46 [1.42-1.50]	1.20 [1.14-1.27]	65	823	50	<0.0001	<0.0001	<0.0001	9	413	78	5.10E-03	1.80E-90	8.80E-18
LBM[James]	0.639	0.618	0.547	1.50 [1.36-1.66]	1.49 [1.45-1.53]	1.08 [1.06-1.09]	64	874	66	<0.0001	<0.0001	<0.0001	12	359	97	1.13E-03	1.33E-78	6.59E-22
BMI	0.633	0.617	0.535	1.34 [1.23-1.45]	1.46 [1.42-1.50]	1.13 [1.07-1.20]	51	794	19	<0.0001	<0.0001	<0.0001	32	430	106	5.68E-08	4.01E-94	5.88E-24
Height^2.7	0.633	0.618	0.582	1.53 [1.37-1.70]	1.45 [1.42-1.49]	1.28 [1.23-1.33]	61	758	152	<0.0001	<0.0001	<0.0001	11	454	13	1.65E-03	2.30E-99	9.38E-04
Height^2.13	0.642	0.622	0.586	1.56 [1.40-1.74]	1.47 [1.43-1.51]	1.32 [1.26-1.38]	66	792	151	<0.0001	<0.0001	<0.0001	6	427	1	0.021	1.73E-93	0.317
Height^2	0.644	0.622	0.586	1.57 [1.41-1.75]	1.47 [1.43-1.51]	1.33 [1.27-1.39]	67	792	149	<0.0001	<0.0001	<0.0001	5	429	0	0.032	8.01E-94	0.603
Height^1.5	0.650	0.619	0.586	1.57 [1.41-1.75]	1.45 [1.41-1.49]	1.33 [1.27-1.40]	69	742	136	<0.0001	<0.0001	<0.0001	4	475	4	0.052	6.20E-104	0.079
Height	0.651	0.608	0.579	1.54 [1.39-1.71]	1.41 [1.37-1.44]	1.31 [1.24-1.37]	66	604	109	<0.0001	<0.0001	<0.0001	9	592	25	6.12E-03	3.22E-129	2.19E-06
<i>Unindexed</i>	<i>0.625</i>	<i>0.569</i>	<i>0.551</i>	<i>1.40 [1.26-1.55]</i>	<i>1.26 [1.22-1.30]</i>	<i>1.19 [1.13-1.26]</i>	<i>41</i>	<i>249</i>	<i>41</i>	<i><0.0001</i>	<i><0.0001</i>	<i><0.0001</i>	<i>32</i>	<i>901</i>	<i>85</i>	<i>5.72E-08</i>	<i>1.93E-196</i>	<i>2.05E-19</i>

Appendix 20. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for aortic sinus diameter.

	C statistic			Hazard Ratio per 1 SD increment			Chi-square			p value			ΔAIC			Akaike Weight		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.701	0.621	0.522	1.54 [1.25-1.88]	1.50 [1.42-1.59]	1.14 [1.03-1.27]	17	211	6	<0.0001	<0.0001	0.015	2	0	17	0.040	0.423	1.38E-04
Weight^1.5	0.691	0.624	0.527	1.44 [1.18-1.76]	1.49 [1.41-1.57]	1.15 [1.03-1.28]	13	210	6	0.00028	<0.0001	0.011	5	4	17	9.15E-03	0.066	1.78E-04
Height*Weight	0.691	0.625	0.532	1.50 [1.22-1.84]	1.50 [1.42-1.58]	1.19 [1.08-1.32]	15	212	11	0.00013	<0.0001	0.00082	4	0	12	0.017	0.370	1.60E-03
BSA[Mosteller]^1.5	0.698	0.621	0.527	1.55 [1.26-1.92]	1.50 [1.42-1.58]	1.18 [1.07-1.31]	17	207	10	<0.0001	<0.0001	0.0016	2	2	13	0.044	0.121	9.45E-04
BSA[Du Bois]^1.5	0.696	0.620	0.529	1.56 [1.26-1.93]	1.49 [1.41-1.58]	1.20 [1.08-1.33]	17	202	12	<0.0001	<0.0001	0.00044	2	6	11	0.043	0.019	2.71E-03
BSA[Mosteller]	0.695	0.608	0.519	1.61 [1.30-2.00]	1.47 [1.39-1.56]	1.16 [1.05-1.29]	19	177	8	<0.0001	<0.0001	0.0053	1	24	15	0.104	2.26E-06	3.36E-04
BSA[Du Bois]	0.694	0.607	0.521	1.61 [1.30-2.00]	1.47 [1.39-1.55]	1.17 [1.06-1.30]	19	173	9	<0.0001	<0.0001	0.0023	1	27	14	0.102	4.89E-07	6.83E-04
LBM[Hume]	0.689	0.608	0.523	1.55 [1.25-1.92]	1.44 [1.36-1.53]	1.19 [1.08-1.32]	16	167	12	<0.0001	<0.0001	0.00058	3	36	12	0.033	7.78E-09	2.05E-03
LBM[Boer]	0.692	0.601	0.517	1.56 [1.26-1.93]	1.42 [1.34-1.50]	1.17 [1.06-1.30]	16	147	9	<0.0001	<0.0001	0.0022	3	51	14	0.038	2.97E-12	6.58E-04
LBM[James]	0.691	0.602	0.520	1.56 [1.26-1.93]	1.43 [1.35-1.52]	1.10 [1.07-1.13]	16	152	53	<0.0001	<0.0001	<0.0001	3	45	5	0.039	5.57E-11	0.074
BMI	0.659	0.583	0.490	1.52 [1.24-1.87]	1.39 [1.31-1.47]	1.00 [0.90-1.12]	16	116	0	<0.0001	<0.0001	0.96	3	74	23	0.032	4.13E-17	7.95E-06
Height^2.7	0.682	0.604	0.536	1.58 [1.25-1.98]	1.43 [1.35-1.51]	1.27 [1.16-1.39]	15	148	28	<0.0001	<0.0001	<0.0001	3	49	0	0.034	9.27E-12	0.716
Height^2.13	0.684	0.599	0.530	1.61 [1.28-2.02]	1.42 [1.34-1.51]	1.26 [1.14-1.38]	17	141	22	<0.0001	<0.0001	<0.0001	2	54	4	0.062	1.02E-12	0.121
Height^2	0.685	0.598	0.529	1.61 [1.29-2.02]	1.42 [1.34-1.51]	1.25 [1.13-1.38]	17	138	20	<0.0001	<0.0001	<0.0001	1	55	5	0.070	3.90E-13	0.072
Height^1.5	0.685	0.590	0.522	1.63 [1.30-2.05]	1.40 [1.32-1.49]	1.22 [1.10-1.35]	18	124	15	<0.0001	<0.0001	0.00014	1	67	9	0.107	1.52E-15	7.40E-03
Height	0.678	0.579	0.516	1.65 [1.32-2.07]	1.37 [1.29-1.46]	1.17 [1.06-1.30]	19	103	9	<0.0001	<0.0001	0.0026	0	84	14	0.138	2.90E-19	6.26E-04
<i>Unindexed</i>	<i>0.647</i>	<i>0.552</i>	<i>0.502</i>	<i>1.66 [1.30-2.11]</i>	<i>1.27 [1.19-1.35]</i>	<i>1.07 [0.96-1.19]</i>	<i>17</i>	<i>53</i>	<i>2</i>	<i><0.0001</i>	<i><0.0001</i>	<i>0.22</i>	<i>1</i>	<i>128</i>	<i>21</i>	<i>0.089</i>	<i>8.48E-29</i>	<i>1.70E-05</i>

Appendix 21. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for aortic arch diameter.

	C statistic			Hazard Ratio per 1 SD increment			Chi-square			p value			ΔAIC			Akaike Weight		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.639	0.631	0.575	1.34 [1.26-1.43]	1.54 [1.51-1.57]	1.32 [1.27-1.36]	86	2327	238	<0.0001	<0.0001	<0.0001	63	9	50	1.11E-14	9.88E-03	6.12E-12
Weight^1.5	0.629	0.622	0.569	1.25 [1.18-1.32]	1.48 [1.46-1.51]	1.27 [1.23-1.31]	57	2048	231	<0.0001	<0.0001	<0.0001	88	276	87	3.09E-20	1.13E-60	4.36E-20
Height*Weight	0.632	0.626	0.576	1.37 [1.28-1.46]	1.50 [1.48-1.53]	1.27 [1.24-1.31]	92	2182	313	<0.0001	<0.0001	<0.0001	58	165	54	1.54E-13	1.78E-36	8.47E-13
BSA[Mosteller]^1.5	0.640	0.633	0.580	1.43 [1.34-1.53]	1.54 [1.51-1.57]	1.33 [1.28-1.37]	109	2358	294	<0.0001	<0.0001	<0.0001	40	0	19	1.14E-09	0.990	3.30E-05
BSA[Du Bois]^1.5	0.640	0.632	0.582	1.47 [1.37-1.58]	1.53 [1.51-1.56]	1.32 [1.28-1.36]	117	2320	320	<0.0001	<0.0001	<0.0001	30	39	10	1.21E-07	3.10E-09	2.21E-03
BSA[Mosteller]	0.643	0.632	0.581	1.49 [1.39-1.61]	1.52 [1.50-1.55]	1.34 [1.30-1.39]	121	2396	279	<0.0001	<0.0001	<0.0001	26	95	11	1.21E-06	2.01E-21	1.61E-03
BSA[Du Bois]	0.644	0.631	0.583	1.52 [1.41-1.64]	1.52 [1.49-1.54]	1.35 [1.30-1.40]	127	2369	294	<0.0001	<0.0001	<0.0001	19	133	0	4.02E-05	1.23E-29	0.403
LBM[Hume]	0.633	0.616	0.571	1.48 [1.38-1.58]	1.43 [1.41-1.46]	1.21 [1.19-1.24]	124	2068	339	<0.0001	<0.0001	<0.0001	29	505	97	3.19E-07	1.79E-110	3.97E-22
LBM[Boer]	0.635	0.604	0.551	1.47 [1.38-1.57]	1.40 [1.37-1.42]	1.17 [1.14-1.20]	127	1667	175	<0.0001	<0.0001	<0.0001	28	758	170	3.50E-07	2.34E-165	4.17E-38
LBM[James]	0.624	0.605	0.547	1.39 [1.29-1.48]	1.44 [1.41-1.46]	1.06 [1.05-1.08]	88	1531	80	<0.0001	<0.0001	<0.0001	57	668	241	1.84E-13	7.96E-146	1.63E-53
BMI	0.604	0.602	0.548	1.22 [1.14-1.31]	1.41 [1.39-1.44]	1.19 [1.15-1.23]	35	1353	87	<0.0001	<0.0001	<0.0001	102	823	190	3.73E-23	2.42E-179	2.37E-42
Height^2.7	0.640	0.611	0.583	1.57 [1.45-1.69]	1.44 [1.42-1.47]	1.28 [1.24-1.31]	136	1605	342	<0.0001	<0.0001	<0.0001	7	627	32	0.014	6.92E-137	3.75E-08
Height^2.13	0.645	0.613	0.585	1.59 [1.47-1.72]	1.45 [1.42-1.47]	1.32 [1.28-1.36]	142	1699	330	<0.0001	<0.0001	<0.0001	1	590	4	0.381	8.09E-129	0.061
Height^2	0.646	0.613	0.586	1.59 [1.48-1.72]	1.45 [1.42-1.47]	1.33 [1.29-1.37]	143	1708	326	<0.0001	<0.0001	<0.0001	0	596	0	0.499	3.02E-130	0.408
Height^1.5	0.646	0.610	0.584	1.58 [1.47-1.71]	1.43 [1.41-1.45]	1.34 [1.30-1.39]	141	1629	299	<0.0001	<0.0001	<0.0001	3	695	2	0.106	1.46E-151	0.125
Height	0.641	0.601	0.578	1.54 [1.43-1.66]	1.39 [1.37-1.42]	1.32 [1.28-1.37]	128	1322	251	<0.0001	<0.0001	<0.0001	16	915	38	1.34E-04	2.33E-199	2.25E-09
Unindexed	0.609	0.569	0.555	1.39 [1.29-1.50]	1.27 [1.24-1.29]	1.22 [1.17-1.26]	72	556	116	<0.0001	<0.0001	<0.0001	66	1511	162	2.01E-15	0.00E+00	2.63E-36

Appendix 22. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for aortic root diameter.

	C statistic			Hazard Ratio per 1 SD increment			Chi-square			p value			ΔAIC			Akaike Weight		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.588	0.615	0.565	1.20 [1.09-1.32]	1.48 [1.44-1.52]	1.25 [1.20-1.31]	13	938	108	0.00029	<0.0001	<0.0001	52	41	20	2.13E-12	1.26E-09	3.87E-05
Weight^1.5	0.585	0.613	0.561	1.12 [1.01-1.23]	1.46 [1.42-1.49]	1.25 [1.20-1.30]	5	1053	115	0.024	<0.0001	<0.0001	60	0	17	4.90E-14	1.000	1.77E-04
Height*Weight	0.597	0.613	0.566	1.23 [1.12-1.36]	1.46 [1.43-1.50]	1.27 [1.21-1.32]	18	1005	126	<0.0001	<0.0001	<0.0001	48	21	6	2.05E-11	2.62E-05	0.053
BSA[Mosteller]^1.5	0.599	0.615	0.569	1.29 [1.17-1.42]	1.47 [1.44-1.51]	1.27 [1.21-1.32]	25	908	117	<0.0001	<0.0001	<0.0001	41	66	10	5.50E-10	5.62E-15	4.81E-03
BSA[Du Bois]^1.5	0.605	0.614	0.571	1.34 [1.21-1.48]	1.47 [1.43-1.50]	1.27 [1.22-1.33]	32	888	121	<0.0001	<0.0001	<0.0001	35	84	7	1.27E-08	5.04E-19	0.032
BSA[Mosteller]	0.601	0.615	0.572	1.35 [1.22-1.50]	1.45 [1.42-1.49]	1.26 [1.20-1.31]	33	750	105	<0.0001	<0.0001	<0.0001	34	179	21	2.40E-08	1.08E-39	2.21E-05
BSA[Du Bois]	0.606	0.614	0.574	1.39 [1.25-1.54]	1.45 [1.41-1.49]	1.26 [1.21-1.32]	38	737	107	<0.0001	<0.0001	<0.0001	29	193	18	2.96E-07	1.28E-42	8.49E-05
LBM[Hume]	0.622	0.606	0.565	1.45 [1.31-1.60]	1.42 [1.39-1.46]	1.24 [1.19-1.29]	52	778	114	<0.0001	<0.0001	<0.0001	17	203	24	8.83E-05	1.05E-44	4.44E-06
LBM[Boer]	0.623	0.600	0.554	1.46 [1.32-1.61]	1.40 [1.36-1.43]	1.21 [1.16-1.25]	54	723	94	<0.0001	<0.0001	<0.0001	15	260	47	2.30E-04	2.99E-57	5.61E-11
LBM[James]	0.594	0.601	0.555	1.27 [1.15-1.40]	1.41 [1.38-1.45]	1.05 [1.04-1.06]	22	721	73	<0.0001	<0.0001	<0.0001	44	240	103	1.33E-10	9.20E-53	4.25E-23
BMI	0.558	0.600	0.550	1.10 [0.99-1.22]	1.40 [1.36-1.44]	1.19 [1.14-1.24]	3	612	63	0.072	<0.0001	<0.0001	61	317	66	2.23E-14	1.87E-69	3.11E-15
Height^2.7	0.640	0.603	0.575	1.54 [1.39-1.69]	1.40 [1.37-1.44]	1.27 [1.22-1.32]	74	703	134	<0.0001	<0.0001	<0.0001	0	260	0	0.495	2.78E-57	0.855
Height^2.13	0.637	0.604	0.577	1.54 [1.39-1.70]	1.40 [1.37-1.44]	1.27 [1.22-1.33]	71	652	122	<0.0001	<0.0001	<0.0001	1	288	6	0.270	2.95E-63	0.041
Height^2	0.636	0.605	0.577	1.54 [1.39-1.70]	1.40 [1.37-1.44]	1.27 [1.22-1.33]	71	639	119	<0.0001	<0.0001	<0.0001	2	297	8	0.201	2.77E-65	0.015
Height^1.5	0.631	0.604	0.577	1.52 [1.38-1.69]	1.39 [1.36-1.43]	1.26 [1.21-1.32]	66	579	108	<0.0001	<0.0001	<0.0001	5	344	18	0.033	1.97E-75	1.12E-04
Height	0.624	0.601	0.575	1.50 [1.35-1.66]	1.37 [1.33-1.41]	1.25 [1.19-1.30]	59	510	96	<0.0001	<0.0001	<0.0001	11	407	30	1.71E-03	5.14E-89	2.65E-07
Unindexed	0.602	0.585	0.563	1.39 [1.26-1.54]	1.30 [1.27-1.34]	1.21 [1.15-1.26]	40	361	70	<0.0001	<0.0001	<0.0001	29	560	57	2.95E-07	3.09E-122	2.94E-13

Appendix 23. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for ascending aorta diameter.

	C statistic			Hazard Ratio per 1 SD increment			Chi-square			p value			ΔAIC			Akaike Weight		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.761	0.725	0.647	1.44 [1.39-1.50]	1.44 [1.43-1.45]	1.44 [1.41-1.47]	373	6294	1205	<0.0001	<0.0001	<0.0001	15	110	18	2.00E-04	1.45E-24	1.04E-04
Weight^1.5	0.758	0.715	0.638	1.45 [1.40-1.51]	1.50 [1.49-1.52]	1.46 [1.43-1.50]	358	5836	982	<0.0001	<0.0001	<0.0001	17	0	51	9.46E-05	0.999	5.94E-12
Height*Weight	0.758	0.721	0.647	1.48 [1.42-1.54]	1.48 [1.46-1.49]	1.45 [1.42-1.48]	373	6136	1147	<0.0001	<0.0001	<0.0001	8	15	19	7.05E-03	6.97E-04	7.05E-05
BSA[Mosteller]^1.5	0.760	0.725	0.651	1.47 [1.41-1.52]	1.43 [1.42-1.44]	1.44 [1.41-1.47]	380	6324	1275	<0.0001	<0.0001	<0.0001	10	153	3	2.79E-03	6.31E-34	0.158
BSA[Du Bois]^1.5	0.759	0.725	0.653	1.48 [1.42-1.53]	1.42 [1.41-1.44]	1.44 [1.41-1.47]	383	6295	1285	<0.0001	<0.0001	<0.0001	8	180	0	8.45E-03	8.39E-40	0.755
BSA[Mosteller]	0.759	0.724	0.652	1.45 [1.40-1.50]	1.38 [1.37-1.39]	1.43 [1.40-1.46]	384	5986	1289	<0.0001	<0.0001	<0.0001	16	376	7	1.66E-04	2.19E-82	0.019
BSA[Du Bois]	0.759	0.723	0.653	1.45 [1.40-1.51]	1.38 [1.37-1.39]	1.43 [1.40-1.46]	386	5950	1300	<0.0001	<0.0001	<0.0001	14	395	5	3.35E-04	2.01E-86	0.068
LBM[Hume]	0.753	0.718	0.647	1.52 [1.46-1.59]	1.42 [1.41-1.44]	1.42 [1.39-1.45]	393	5894	1169	<0.0001	<0.0001	<0.0001	1	273	45	0.325	6.68E-60	1.00E-10
LBM[Boer]	0.754	0.712	0.634	1.51 [1.45-1.58]	1.44 [1.43-1.46]	1.39 [1.36-1.42]	393	5596	941	<0.0001	<0.0001	<0.0001	2	284	123	0.162	2.45E-62	1.53E-27
LBM[James]	0.754	0.714	0.632	1.49 [1.43-1.55]	1.44 [1.43-1.45]	1.08 [1.07-1.09]	382	5695	328	<0.0001	<0.0001	<0.0001	7	272	469	0.013	1.08E-59	1.15E-102
BMI	0.751	0.711	0.631	1.39 [1.34-1.44]	1.38 [1.37-1.39]	1.44 [1.41-1.48]	346	5328	839	<0.0001	<0.0001	<0.0001	40	550	105	9.59E-10	3.20E-120	1.15E-23
Height^2.7	0.753	0.716	0.653	1.53 [1.47-1.59]	1.39 [1.38-1.41]	1.33 [1.31-1.35]	390	5699	1180	<0.0001	<0.0001	<0.0001	0	428	110	0.442	1.03E-93	1.14E-24
Height^2.13	0.755	0.717	0.655	1.50 [1.44-1.56]	1.37 [1.36-1.38]	1.41 [1.38-1.44]	392	5620	1180	<0.0001	<0.0001	<0.0001	6	504	37	0.026	4.23E-110	6.62E-09
Height^2	0.755	0.717	0.655	1.49 [1.43-1.55]	1.37 [1.36-1.38]	1.42 [1.39-1.45]	392	5582	1186	<0.0001	<0.0001	<0.0001	7	523	29	0.012	2.26E-114	3.75E-07
Height^1.5	0.755	0.715	0.653	1.47 [1.41-1.52]	1.35 [1.34-1.36]	1.43 [1.40-1.46]	388	5379	1208	<0.0001	<0.0001	<0.0001	14	606	21	4.07E-04	2.17E-132	2.09E-05
Height	0.753	0.712	0.650	1.44 [1.39-1.50]	1.34 [1.33-1.35]	1.43 [1.40-1.46]	378	5097	1164	<0.0001	<0.0001	<0.0001	22	699	33	7.38E-06	1.45E-152	6.17E-08
Unindexed	0.745	0.699	0.640	1.40 [1.35-1.45]	1.32 [1.31-1.33]	1.44 [1.40-1.47]	341	4372	924	<0.0001	<0.0001	<0.0001	41	907	87	6.51E-10	1.22E-197	1.23E-19

Appendix 24. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for LA area.

	C statistic			Hazard Ratio per 1 SD increment			Chi-square			p value			ΔAIC			Akaike Weight		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.731	0.713	0.662	1.74 [1.60-1.90]	1.77 [1.73-1.80]	1.75 [1.68-1.83]	169	3625	654	<0.0001	<0.0001	<0.0001	41	234	117	6.51E-10	1.93E-51	2.08E-26
Weight ^{1.5}	0.722	0.688	0.638	1.50 [1.40-1.61]	1.64 [1.61-1.67]	1.62 [1.55-1.69]	124	2915	479	<0.0001	<0.0001	<0.0001	82	714	269	8.50E-19	1.01E-155	2.74E-59
Height*Weight	0.728	0.699	0.650	1.74 [1.60-1.89]	1.69 [1.66-1.72]	1.64 [1.57-1.71]	170	3253	552	<0.0001	<0.0001	<0.0001	41	535	217	6.67E-10	7.00E-117	3.83E-48
BSA[Mosteller] ^{1.5}	0.735	0.714	0.666	1.86 [1.71-2.03]	1.77 [1.74-1.81]	1.74 [1.67-1.82]	196	3651	680	<0.0001	<0.0001	<0.0001	20	222	105	1.94E-05	7.53E-49	9.34E-24
BSA[Du Bois] ^{1.5}	0.737	0.713	0.666	1.88 [1.73-2.05]	1.77 [1.74-1.81]	1.73 [1.66-1.80]	209	3587	681	<0.0001	<0.0001	<0.0001	14	241	109	3.45E-04	3.98E-53	9.80E-25
BSA[Mosteller]	0.738	0.723	0.681	1.94 [1.78-2.12]	1.86 [1.83-1.90]	1.82 [1.75-1.90]	217	3714	811	<0.0001	<0.0001	<0.0001	7	0	2	0.015	1.000	0.224
BSA[Du Bois]	0.739	0.722	0.681	1.94 [1.78-2.12]	1.86 [1.83-1.90]	1.82 [1.75-1.90]	226	3655	812	<0.0001	<0.0001	<0.0001	4	20	2	0.057	3.62E-05	0.216
LBM[Hume]	0.734	0.701	0.655	1.78 [1.65-1.92]	1.71 [1.68-1.74]	1.66 [1.59-1.73]	223	3136	585	<0.0001	<0.0001	<0.0001	21	526	193	1.23E-05	5.97E-115	6.55E-43
LBM[Boer]	0.734	0.690	0.630	1.76 [1.64-1.90]	1.68 [1.64-1.71]	1.55 [1.49-1.62]	224	2718	417	<0.0001	<0.0001	<0.0001	23	722	328	5.08E-06	1.66E-157	3.25E-72
LBM[James]	0.727	0.691	0.625	1.83 [1.68-2.00]	1.74 [1.70-1.77]	1.12 [1.11-1.14]	183	2732	398	<0.0001	<0.0001	<0.0001	28	594	582	2.97E-07	9.62E-130	2.04E-127
BMI	0.707	0.706	0.655	1.66 [1.52-1.82]	1.80 [1.77-1.84]	1.71 [1.64-1.79]	119	3050	580	<0.0001	<0.0001	<0.0001	69	324	179	4.74E-16	3.68E-71	7.55E-40
Height ^{2.7}	0.737	0.699	0.660	1.85 [1.72-2.00]	1.77 [1.73-1.80]	1.52 [1.48-1.57]	244	2767	748	<0.0001	<0.0001	<0.0001	8	517	243	6.69E-03	4.33E-113	8.68E-54
Height ^{2.13}	0.739	0.707	0.671	1.89 [1.75-2.04]	1.82 [1.78-1.86]	1.66 [1.60-1.72]	251	2975	772	<0.0001	<0.0001	<0.0001	3	340	134	0.118	1.59E-74	3.57E-30
Height ²	0.739	0.708	0.674	1.90 [1.75-2.05]	1.84 [1.80-1.88]	1.69 [1.63-1.75]	252	3008	773	<0.0001	<0.0001	<0.0001	2	306	111	0.186	3.41E-67	5.05E-25
Height ^{1.5}	0.738	0.711	0.681	1.93 [1.78-2.09]	1.87 [1.83-1.92]	1.79 [1.72-1.86]	247	3060	792	<0.0001	<0.0001	<0.0001	0	216	35	0.444	1.21E-47	1.26E-08
Height	0.733	0.711	0.685	1.95 [1.79-2.13]	1.89 [1.85-1.94]	1.83 [1.76-1.91]	233	2978	816	<0.0001	<0.0001	<0.0001	2	215	0	0.173	2.23E-47	0.559
<i>Unindexed</i>	<i>0.713</i>	<i>0.695</i>	<i>0.675</i>	<i>1.95 [1.77-2.15]</i>	<i>1.84 [1.79-1.88]</i>	<i>1.77 [1.70-1.85]</i>	<i>181</i>	<i>2456</i>	<i>714</i>	<i><0.0001</i>	<i><0.0001</i>	<i><0.0001</i>	<i>21</i>	<i>532</i>	<i>80</i>	<i>1.44E-05</i>	<i>3.07E-116</i>	<i>2.80E-18</i>

Appendix 25. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for LA diameter.

	C statistic			Hazard Ratio per 1 SD increment			Chi-square			p value			ΔAIC			Akaike Weight		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.741	0.740	0.677	1.64 [1.56-1.73]	1.60 [1.58-1.62]	1.51 [1.46-1.55]	361	5473	783	<0.0001	<0.0001	<0.0001	0	80	21	0.477	5.26E-18	2.50E-05
Weight^1.5	0.741	0.742	0.673	1.65 [1.56-1.74]	1.57 [1.56-1.59]	1.50 [1.46-1.54]	355	6089	812	<0.0001	<0.0001	<0.0001	2	0	14	0.199	1.000	8.87E-04
Height*Weight	0.740	0.742	0.679	1.64 [1.55-1.72]	1.59 [1.57-1.61]	1.51 [1.47-1.55]	354	5800	823	<0.0001	<0.0001	<0.0001	4	42	0	0.065	7.67E-10	0.985
BSA[Mosteller]^1.5	0.739	0.739	0.680	1.63 [1.55-1.72]	1.60 [1.58-1.62]	1.51 [1.47-1.55]	357	5358	794	<0.0001	<0.0001	<0.0001	3	124	12	0.100	1.10E-27	3.07E-03
BSA[Du Bois]^1.5	0.739	0.739	0.681	1.63 [1.55-1.71]	1.60 [1.58-1.62]	1.51 [1.47-1.56]	355	5303	796	<0.0001	<0.0001	<0.0001	5	155	9	0.044	2.71E-34	0.011
BSA[Mosteller]	0.737	0.734	0.678	1.63 [1.55-1.71]	1.60 [1.58-1.62]	1.50 [1.46-1.55]	356	4858	752	<0.0001	<0.0001	<0.0001	5	306	37	0.032	3.14E-67	1.04E-08
BSA[Du Bois]	0.737	0.734	0.679	1.63 [1.54-1.71]	1.60 [1.58-1.62]	1.50 [1.46-1.55]	355	4819	753	<0.0001	<0.0001	<0.0001	6	327	35	0.019	8.46E-72	2.22E-08
LBM[Hume]	0.734	0.734	0.678	1.62 [1.54-1.71]	1.59 [1.57-1.61]	1.50 [1.46-1.55]	348	5083	767	<0.0001	<0.0001	<0.0001	10	307	30	4.04E-03	1.82E-67	2.75E-07
LBM[Boer]	0.735	0.732	0.672	1.62 [1.54-1.71]	1.58 [1.56-1.60]	1.49 [1.44-1.53]	349	5029	724	<0.0001	<0.0001	<0.0001	9	374	66	6.07E-03	5.72E-82	5.59E-15
LBM[James]	0.735	0.732	0.671	1.63 [1.55-1.72]	1.58 [1.56-1.60]	1.16 [1.15-1.17]	349	5013	905	<0.0001	<0.0001	<0.0001	7	331	359	0.012	1.44E-72	8.63E-79
BMI	0.734	0.727	0.663	1.64 [1.56-1.73]	1.58 [1.56-1.60]	1.46 [1.42-1.51]	354	4663	660	<0.0001	<0.0001	<0.0001	5	482	111	0.038	1.78E-105	6.15E-25
Height^2.7	0.734	0.731	0.682	1.61 [1.53-1.69]	1.58 [1.56-1.60]	1.50 [1.46-1.55]	343	4834	779	<0.0001	<0.0001	<0.0001	15	436	27	3.14E-04	2.39E-95	1.48E-06
Height^2.13	0.734	0.730	0.681	1.61 [1.53-1.69]	1.58 [1.56-1.60]	1.50 [1.46-1.55]	346	4626	756	<0.0001	<0.0001	<0.0001	13	475	36	6.76E-04	8.78E-104	1.24E-08
Height^2	0.734	0.729	0.681	1.61 [1.53-1.69]	1.58 [1.56-1.60]	1.50 [1.46-1.55]	346	4579	751	<0.0001	<0.0001	<0.0001	13	489	40	7.40E-04	8.25E-107	2.58E-09
Height^1.5	0.733	0.727	0.679	1.61 [1.53-1.69]	1.58 [1.56-1.60]	1.50 [1.45-1.54]	347	4395	727	<0.0001	<0.0001	<0.0001	13	559	55	7.72E-04	4.50E-122	1.40E-12
Height	0.732	0.724	0.676	1.61 [1.53-1.69]	1.58 [1.55-1.60]	1.49 [1.44-1.53]	346	4207	701	<0.0001	<0.0001	<0.0001	14	653	74	4.93E-04	1.35E-142	8.94E-17
Unindexed	0.727	0.715	0.668	1.61 [1.53-1.69]	1.55 [1.53-1.57]	1.46 [1.42-1.51]	339	3812	639	<0.0001	<0.0001	<0.0001	19	899	123	4.42E-05	4.98E-196	2.40E-27

Appendix 26. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for LA volume.

	C statistic			Hazard Ratio per 1 SD increment			Chi-square			p value			ΔAIC			Akaike Weight		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.563	0.618	0.592	1.13 [1.06-1.21]	1.54 [1.52-1.57]	1.40 [1.37-1.44]	13	3034	700	0.00025	<0.0001	<0.0001	74	38	158	3.76E-17	5.88E-09	5.07E-35
Weight^1.5	0.566	0.615	0.581	1.07 [1.00-1.15]	1.48 [1.46-1.50]	1.30 [1.27-1.32]	4	2540	572	0.035	<0.0001	<0.0001	83	446	321	5.82E-19	1.11E-97	1.38E-70
Height*Weight	0.568	0.617	0.591	1.17 [1.09-1.25]	1.51 [1.48-1.53]	1.27 [1.25-1.29]	22	2790	822	<0.0001	<0.0001	<0.0001	67	267	283	1.60E-15	1.18E-58	2.40E-62
BSA[Mosteller]^1.5	0.567	0.618	0.598	1.21 [1.13-1.29]	1.55 [1.53-1.57]	1.35 [1.33-1.38]	31	3102	975	<0.0001	<0.0001	<0.0001	59	0	144	1.02E-13	0.892	4.06E-32
BSA[Du Bois]^1.5	0.569	0.617	0.600	1.24 [1.16-1.33]	1.55 [1.52-1.57]	1.33 [1.30-1.35]	41	3065	1057	<0.0001	<0.0001	<0.0001	49	37	155	1.11E-11	6.91E-09	1.51E-34
BSA[Mosteller]	0.562	0.610	0.602	1.25 [1.16-1.34]	1.56 [1.53-1.58]	1.46 [1.42-1.49]	40	3097	929	<0.0001	<0.0001	<0.0001	50	4	4	7.48E-12	0.108	0.096
BSA[Du Bois]	0.565	0.609	0.604	1.28 [1.19-1.37]	1.55 [1.53-1.58]	1.44 [1.41-1.48]	48	3050	1020	<0.0001	<0.0001	<0.0001	42	39	0	3.46E-10	2.76E-09	0.859
LBM[Hume]	0.572	0.603	0.590	1.28 [1.21-1.36]	1.46 [1.44-1.48]	1.26 [1.24-1.28]	73	2503	767	<0.0001	<0.0001	<0.0001	31	639	297	1.03E-07	1.55E-139	2.46E-65
LBM[Boer]	0.572	0.593	0.569	1.28 [1.21-1.36]	1.41 [1.39-1.44]	1.22 [1.20-1.25]	78	1977	420	<0.0001	<0.0001	<0.0001	29	996	438	2.30E-07	5.18E-217	6.98E-96
LBM[James]	0.555	0.594	0.566	1.17 [1.09-1.25]	1.45 [1.42-1.47]	1.06 [1.05-1.06]	21	1884	183	<0.0001	<0.0001	<0.0001	67	912	613	1.37E-15	9.14E-199	5.92E-134
BMI	0.527	0.588	0.566	1.03 [0.96-1.10]	1.44 [1.42-1.47]	1.35 [1.30-1.39]	1	1884	347	0.45	<0.0001	<0.0001	86	932	344	9.51E-20	3.39E-203	1.48E-75
Height^2.7	0.585	0.600	0.602	1.39 [1.30-1.48]	1.46 [1.44-1.48]	1.25 [1.23-1.27]	100	2267	900	<0.0001	<0.0001	<0.0001	0	714	269	0.538	7.01E-156	3.92E-59
Height^2.13	0.581	0.598	0.606	1.40 [1.31-1.49]	1.48 [1.46-1.51]	1.32 [1.30-1.34]	97	2333	1044	<0.0001	<0.0001	<0.0001	1	599	151	0.295	7.25E-131	1.16E-33
Height^2	0.580	0.598	0.606	1.39 [1.30-1.49]	1.49 [1.46-1.51]	1.34 [1.31-1.36]	95	2334	1060	<0.0001	<0.0001	<0.0001	2	591	123	0.164	4.32E-129	1.61E-27
Height^1.5	0.574	0.592	0.605	1.38 [1.29-1.47]	1.48 [1.46-1.51]	1.42 [1.39-1.45]	85	2238	994	<0.0001	<0.0001	<0.0001	11	654	28	2.75E-03	7.15E-143	7.42E-07
Height	0.565	0.583	0.600	1.34 [1.25-1.44]	1.45 [1.43-1.48]	1.48 [1.44-1.52]	69	1942	803	<0.0001	<0.0001	<0.0001	24	890	6	2.85E-06	4.04E-194	0.045
Unindexed	0.541	0.558	0.577	1.23 [1.14-1.32]	1.33 [1.31-1.36]	1.40 [1.36-1.45]	30	1017	508	<0.0001	<0.0001	<0.0001	58	1663	211	1.21E-13	0.00E+00	1.12E-46

Appendix 27. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for LV diameter.

	C statistic			Hazard Ratio per 1 SD increment			Chi-square			p value			ΔAIC			Akaike Weight		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.539	0.589	0.614	1.30 [1.15-1.47]	1.42 [1.39-1.45]	1.47 [1.41-1.54]	19	962	300	<0.0001	<0.0001	<0.0001	3	114	18	0.066	1.48E-25	1.36E-04
Weight ^{1.5}	0.562	0.607	0.622	1.32 [1.18-1.47]	1.45 [1.42-1.48]	1.48 [1.42-1.54]	23	1183	328	<0.0001	<0.0001	<0.0001	0	0	8	0.259	1.000	0.021
Height*Weight	0.554	0.598	0.623	1.33 [1.18-1.50]	1.44 [1.41-1.47]	1.49 [1.42-1.55]	22	1097	338	<0.0001	<0.0001	<0.0001	0	37	0	0.307	1.18E-08	0.956
BSA[Mosteller] ^{1.5}	0.539	0.586	0.617	1.31 [1.16-1.47]	1.42 [1.39-1.45]	1.48 [1.42-1.55]	19	929	312	<0.0001	<0.0001	<0.0001	3	126	11	0.079	3.43E-28	4.97E-03
BSA[Du Bois] ^{1.5}	0.540	0.584	0.617	1.31 [1.16-1.48]	1.42 [1.38-1.45]	1.48 [1.42-1.55]	19	910	316	<0.0001	<0.0001	<0.0001	3	135	8	0.081	4.17E-30	0.016
BSA[Mosteller]	0.524	0.573	0.608	1.27 [1.13-1.44]	1.39 [1.35-1.42]	1.47 [1.40-1.53]	15	724	275	0.00011	<0.0001	<0.0001	6	239	29	0.016	1.29E-52	4.28E-07
BSA[Du Bois]	0.525	0.572	0.609	1.27 [1.13-1.44]	1.38 [1.35-1.42]	1.47 [1.40-1.54]	15	712	277	0.00011	<0.0001	<0.0001	6	245	28	0.016	5.06E-54	9.20E-07
LBM[Hume]	0.536	0.576	0.615	1.29 [1.14-1.46]	1.40 [1.37-1.43]	1.48 [1.41-1.55]	17	781	298	<0.0001	<0.0001	<0.0001	5	206	16	0.032	2.26E-45	2.66E-04
LBM[Boer]	0.534	0.573	0.612	1.29 [1.14-1.46]	1.39 [1.36-1.42]	1.47 [1.40-1.54]	17	741	275	<0.0001	<0.0001	<0.0001	5	229	29	0.032	2.02E-50	5.60E-07
LBM[James]	0.535	0.574	0.606	1.28 [1.13-1.45]	1.39 [1.36-1.43]	1.40 [1.35-1.46]	16	752	252	<0.0001	<0.0001	<0.0001	5	224	59	0.023	2.38E-49	1.84E-13
BMI	0.513	0.569	0.592	1.22 [1.08-1.39]	1.36 [1.33-1.39]	1.42 [1.35-1.49]	10	595	196	0.0017	<0.0001	<0.0001	10	319	74	2.06E-03	4.78E-70	8.11E-17
Height ^{2.7}	0.537	0.571	0.617	1.30 [1.15-1.46]	1.39 [1.35-1.42]	1.49 [1.42-1.56]	17	715	298	<0.0001	<0.0001	<0.0001	4	241	13	0.043	4.48E-53	1.29E-03
Height ^{2.13}	0.528	0.566	0.612	1.28 [1.13-1.45]	1.37 [1.34-1.41]	1.48 [1.41-1.55]	15	643	282	<0.0001	<0.0001	<0.0001	6	284	22	0.019	2.19E-62	1.33E-05
Height ²	0.526	0.565	0.611	1.27 [1.13-1.44]	1.37 [1.34-1.40]	1.48 [1.41-1.55]	15	625	278	0.00012	<0.0001	<0.0001	6	295	25	0.015	1.09E-64	3.48E-06
Height ^{1.5}	0.519	0.560	0.606	1.26 [1.11-1.42]	1.35 [1.32-1.39]	1.46 [1.40-1.53]	13	555	258	0.00034	<0.0001	<0.0001	8	337	37	6.98E-03	5.93E-74	9.51E-09
Height	0.513	0.555	0.600	1.24 [1.09-1.41]	1.34 [1.30-1.37]	1.45 [1.38-1.52]	11	484	235	0.00097	<0.0001	<0.0001	9	381	50	3.19E-03	1.61E-83	1.15E-11
<i>Unindexed</i>	<i>0.501</i>	<i>0.547</i>	<i>0.589</i>	<i>1.20 [1.05-1.37]</i>	<i>1.30 [1.27-1.34]</i>	<i>1.42 [1.35-1.49]</i>	<i>7</i>	<i>352</i>	<i>187</i>	<i>0.0069</i>	<i><0.0001</i>	<i><0.0001</i>	<i>12</i>	<i>467</i>	<i>79</i>	<i>7.45E-04</i>	<i>4.67E-102</i>	<i>7.13E-18</i>

Appendix 28. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for LV volume.

	C statistic			Hazard Ratio per 1 SD increment			Chi-square			p value			ΔAIC			Akaike Weight		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.703	0.705	0.628	1.61 [1.52-1.69]	1.63 [1.61-1.64]	1.47 [1.43-1.50]	318	8608	999	<0.0001	<0.0001	<0.0001	123	0	77	1.27E-27	1.000	1.56E-17
Weight^1.5	0.681	0.687	0.612	1.42 [1.35-1.49]	1.60 [1.58-1.62]	1.41 [1.38-1.44]	186	7518	923	<0.0001	<0.0001	<0.0001	226	565	215	4.64E-50	2.22E-123	1.84E-47
Height*Weight	0.695	0.692	0.621	1.60 [1.51-1.68]	1.61 [1.60-1.63]	1.41 [1.38-1.44]	308	7652	1083	<0.0001	<0.0001	<0.0001	132	415	158	1.40E-29	9.28E-91	4.12E-35
BSA[Mosteller]^1.5	0.710	0.704	0.631	1.67 [1.59-1.76]	1.63 [1.61-1.64]	1.47 [1.43-1.50]	397	8257	1060	<0.0001	<0.0001	<0.0001	76	105	53	2.06E-17	1.23E-23	2.58E-12
BSA[Du Bois]^1.5	0.712	0.702	0.632	1.68 [1.60-1.77]	1.62 [1.61-1.64]	1.46 [1.43-1.50]	430	8032	1077	<0.0001	<0.0001	<0.0001	61	200	53	3.71E-14	4.73E-44	3.49E-12
BSA[Mosteller]	0.723	0.710	0.640	1.70 [1.62-1.78]	1.62 [1.60-1.64]	1.48 [1.45-1.52]	488	8295	1117	<0.0001	<0.0001	<0.0001	32	101	4	6.11E-08	1.16E-22	0.105
BSA[Du Bois]	0.725	0.708	0.641	1.70 [1.63-1.78]	1.62 [1.60-1.64]	1.48 [1.45-1.52]	509	8117	1124	<0.0001	<0.0001	<0.0001	25	177	0	2.74E-06	3.67E-39	0.895
LBM[Hume]	0.716	0.690	0.623	1.67 [1.59-1.74]	1.58 [1.57-1.60]	1.28 [1.26-1.30]	483	7228	1515	<0.0001	<0.0001	<0.0001	46	749	281	5.25E-11	1.96E-163	8.09E-62
LBM[Boer]	0.718	0.680	0.604	1.66 [1.59-1.74]	1.56 [1.55-1.58]	1.24 [1.22-1.25]	492	6669	1163	<0.0001	<0.0001	<0.0001	44	1070	419	2.11E-10	3.61E-233	7.88E-92
LBM[James]	0.701	0.683	0.603	1.65 [1.57-1.73]	1.59 [1.57-1.61]	1.05 [1.04-1.06]	371	6829	312	<0.0001	<0.0001	<0.0001	94	843	764	3.06E-21	9.10E-184	1.35E-166
BMI	0.697	0.703	0.617	1.55 [1.47-1.63]	1.59 [1.58-1.61]	1.43 [1.39-1.46]	277	7992	754	<0.0001	<0.0001	<0.0001	156	424	241	7.99E-35	9.29E-93	5.24E-53
Height^2.7	0.724	0.685	0.630	1.68 [1.60-1.75]	1.59 [1.57-1.60]	1.41 [1.38-1.43]	526	6434	1057	<0.0001	<0.0001	<0.0001	28	1035	152	5.08E-07	2.17E-225	1.09E-33
Height^2.13	0.730	0.691	0.637	1.69 [1.62-1.76]	1.59 [1.58-1.61]	1.45 [1.42-1.49]	559	6756	1051	<0.0001	<0.0001	<0.0001	12	856	71	1.54E-03	1.53E-186	3.00E-16
Height^2	0.732	0.692	0.638	1.69 [1.62-1.76]	1.60 [1.58-1.61]	1.46 [1.43-1.49]	564	6804	1055	<0.0001	<0.0001	<0.0001	9	830	58	6.64E-03	6.63E-181	1.81E-13
Height^1.5	0.735	0.696	0.641	1.69 [1.62-1.77]	1.60 [1.58-1.61]	1.47 [1.44-1.51]	580	6868	1074	<0.0001	<0.0001	<0.0001	1	796	31	0.347	1.81E-173	1.57E-07
Height	0.737	0.695	0.642	1.69 [1.62-1.77]	1.59 [1.57-1.61]	1.47 [1.44-1.51]	582	6706	1060	<0.0001	<0.0001	<0.0001	0	882	41	0.645	3.05E-192	1.41E-09
Unindexed	0.728	0.682	0.631	1.66 [1.59-1.74]	1.56 [1.54-1.58]	1.44 [1.40-1.47]	545	5685	874	<0.0001	<0.0001	<0.0001	24	1435	166	3.45E-06	0.00E+00	9.17E-37

Appendix 29. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for IVS diameter.

	C statistic			Hazard Ratio per 1 SD increment			Chi-square			p value			ΔAIC			Akaike Weight		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.726	0.741	0.681	1.79 [1.70-1.88]	1.82 [1.80-1.84]	1.71 [1.67-1.75]	485	11151	1849	<0.0001	<0.0001	<0.0001	116	14	39	2.78E-26	7.57E-04	1.85E-09
Weight ^{1.5}	0.718	0.745	0.677	1.60 [1.53-1.68]	1.74 [1.73-1.76]	1.61 [1.58-1.64]	358	12472	2178	<0.0001	<0.0001	<0.0001	223	77	151	1.97E-49	1.87E-17	1.32E-33
Height*Weight	0.729	0.745	0.687	1.78 [1.69-1.87]	1.78 [1.76-1.80]	1.58 [1.55-1.61]	508	11995	2625	<0.0001	<0.0001	<0.0001	109	0	103	1.10E-24	0.999	2.65E-23
BSA[Mosteller] ^{1.5}	0.733	0.740	0.687	1.86 [1.76-1.95]	1.82 [1.80-1.85]	1.70 [1.66-1.74]	558	10834	1979	<0.0001	<0.0001	<0.0001	65	112	0	3.97E-15	5.29E-25	0.655
BSA[Du Bois] ^{1.5}	0.736	0.739	0.688	1.87 [1.78-1.97]	1.82 [1.80-1.84]	1.69 [1.65-1.73]	588	10636	2061	<0.0001	<0.0001	<0.0001	48	204	1	2.41E-11	5.46E-45	0.345
BSA[Mosteller]	0.734	0.727	0.681	1.89 [1.80-1.99]	1.83 [1.81-1.85]	1.71 [1.67-1.76]	584	9227	1759	<0.0001	<0.0001	<0.0001	43	714	67	2.54E-10	1.00E-155	2.02E-15
BSA[Du Bois]	0.736	0.727	0.682	1.90 [1.80-2.00]	1.83 [1.80-1.85]	1.71 [1.67-1.76]	602	9102	1771	<0.0001	<0.0001	<0.0001	33	782	61	3.57E-08	1.51E-170	3.71E-14
LBM[Hume]	0.741	0.732	0.686	1.88 [1.79-1.97]	1.80 [1.78-1.82]	1.64 [1.61-1.68]	650	10025	2138	<0.0001	<0.0001	<0.0001	21	620	90	1.09E-05	2.40E-135	2.04E-20
LBM[Boer]	0.741	0.730	0.679	1.88 [1.79-1.97]	1.78 [1.76-1.80]	1.58 [1.55-1.61]	649	9903	2130	<0.0001	<0.0001	<0.0001	21	794	235	1.62E-05	4.14E-173	6.21E-52
LBM[James]	0.730	0.731	0.673	1.84 [1.75-1.94]	1.79 [1.77-1.81]	1.10 [1.09-1.10]	540	10031	1081	<0.0001	<0.0001	<0.0001	77	632	1234	8.05E-18	6.50E-138	7.06E-269
BMI	0.706	0.710	0.652	1.68 [1.60-1.77]	1.77 [1.75-1.79]	1.62 [1.57-1.66]	391	8135	1285	<0.0001	<0.0001	<0.0001	190	1533	435	3.14E-42	0.00E+00	1.93E-95
Height ^{2.7}	0.747	0.723	0.687	1.88 [1.80-1.98]	1.78 [1.76-1.80]	1.53 [1.50-1.55]	688	8864	2401	<0.0001	<0.0001	<0.0001	3	1192	249	0.129	1.52E-259	5.15E-55
Height ^{2.13}	0.745	0.719	0.684	1.91 [1.82-2.00]	1.79 [1.77-1.81]	1.63 [1.60-1.67]	675	8373	1987	<0.0001	<0.0001	<0.0001	0	1311	155	0.505	2.14E-285	1.23E-34
Height ²	0.744	0.718	0.683	1.91 [1.82-2.00]	1.79 [1.77-1.81]	1.65 [1.61-1.68]	671	8241	1900	<0.0001	<0.0001	<0.0001	1	1360	149	0.358	5.57E-296	2.82E-33
Height ^{1.5}	0.740	0.712	0.678	1.91 [1.82-2.01]	1.79 [1.76-1.81]	1.68 [1.63-1.72]	651	7686	1672	<0.0001	<0.0001	<0.0001	8	1617	170	8.27E-03	0.00E+00	7.55E-38
Height	0.736	0.705	0.672	1.90 [1.81-2.00]	1.77 [1.75-1.79]	1.67 [1.63-1.71]	627	7071	1529	<0.0001	<0.0001	<0.0001	24	1972	242	3.87E-06	0.00E+00	2.15E-53
Unindexed	0.723	0.687	0.655	1.84 [1.75-1.93]	1.71 [1.69-1.73]	1.61 [1.57-1.65]	562	5756	1250	<0.0001	<0.0001	<0.0001	74	2863	458	4.00E-17	0.00E+00	2.24E-100

Appendix 30. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for LV mass.

	C statistic			Hazard Ratio per 1 SD increment			Chi-square			p value			ΔAIC			Akaike Weight		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.723	0.680	0.602	1.73 [1.59-1.89]	1.69 [1.65-1.72]	1.50 [1.43-1.57]	159	2324	299	<0.0001	<0.0001	<0.0001	1	5	6	0.253	0.075	0.019
Weight ^{1.5}	0.726	0.682	0.601	1.73 [1.59-1.89]	1.64 [1.61-1.68]	1.47 [1.40-1.53]	160	2404	282	<0.0001	<0.0001	<0.0001	0	44	25	0.372	2.68E-10	1.80E-06
Height*Weight	0.723	0.684	0.604	1.71 [1.57-1.85]	1.68 [1.64-1.71]	1.48 [1.42-1.55]	161	2374	305	<0.0001	<0.0001	<0.0001	2	0	10	0.116	0.873	3.24E-03
BSA[Mosteller] ^{1.5}	0.721	0.680	0.604	1.71 [1.58-1.86]	1.69 [1.66-1.73]	1.50 [1.44-1.57]	159	2297	309	<0.0001	<0.0001	<0.0001	3	6	0	0.100	0.051	0.456
BSA[Du Bois] ^{1.5}	0.719	0.680	0.604	1.70 [1.57-1.85]	1.69 [1.66-1.73]	1.50 [1.43-1.57]	158	2272	315	<0.0001	<0.0001	<0.0001	4	15	1	0.060	5.43E-04	0.334
BSA[Mosteller]	0.714	0.673	0.601	1.71 [1.57-1.86]	1.69 [1.65-1.73]	1.51 [1.44-1.58]	153	2096	297	<0.0001	<0.0001	<0.0001	5	94	3	0.027	3.38E-21	0.084
BSA[Du Bois]	0.714	0.673	0.602	1.71 [1.57-1.86]	1.69 [1.65-1.73]	1.51 [1.44-1.58]	153	2084	298	<0.0001	<0.0001	<0.0001	6	102	3	0.020	5.53E-23	0.103
LBM[Hume]	0.714	0.674	0.598	1.67 [1.54-1.82]	1.68 [1.64-1.72]	1.45 [1.39-1.51]	155	2073	306	<0.0001	<0.0001	<0.0001	8	107	30	7.37E-03	5.54E-24	1.11E-07
LBM[Boer]	0.714	0.671	0.589	1.67 [1.54-1.81]	1.67 [1.63-1.71]	1.37 [1.32-1.43]	155	1960	273	<0.0001	<0.0001	<0.0001	8	163	77	7.10E-03	3.20E-36	8.18E-18
LBM[James]	0.716	0.671	0.585	1.70 [1.56-1.85]	1.67 [1.63-1.71]	1.17 [1.14-1.19]	155	2018	220	<0.0001	<0.0001	<0.0001	6	142	170	0.023	1.60E-31	7.06E-38
BMI	0.705	0.659	0.590	1.74 [1.58-1.90]	1.65 [1.61-1.69]	1.44 [1.38-1.51]	138	1745	235	<0.0001	<0.0001	<0.0001	10	271	56	2.80E-03	1.04E-59	3.90E-13
Height ^{2.7}	0.710	0.673	0.603	1.66 [1.53-1.80]	1.67 [1.63-1.71]	1.27 [1.24-1.30]	151	2007	417	<0.0001	<0.0001	<0.0001	10	161	81	2.29E-03	1.05E-35	9.86E-19
Height ^{2.13}	0.709	0.670	0.602	1.67 [1.54-1.82]	1.67 [1.63-1.71]	1.37 [1.33-1.41]	150	1972	399	<0.0001	<0.0001	<0.0001	10	180	44	3.06E-03	6.03E-40	1.15E-10
Height ²	0.709	0.669	0.602	1.67 [1.54-1.82]	1.67 [1.63-1.71]	1.39 [1.35-1.44]	150	1956	382	<0.0001	<0.0001	<0.0001	10	189	37	3.04E-03	9.01E-42	3.31E-09
Height ^{1.5}	0.706	0.665	0.600	1.68 [1.55-1.83]	1.66 [1.62-1.70]	1.47 [1.40-1.53]	148	1864	306	<0.0001	<0.0001	<0.0001	10	235	23	2.23E-03	9.83E-52	4.66E-06
Height	0.702	0.660	0.597	1.68 [1.55-1.83]	1.65 [1.61-1.69]	1.49 [1.42-1.56]	144	1731	268	<0.0001	<0.0001	<0.0001	12	301	26	1.03E-03	4.47E-66	1.24E-06
<i>Unindexed</i>	<i>0.690</i>	<i>0.646</i>	<i>0.590</i>	<i>1.68 [1.54-1.84]</i>	<i>1.61 [1.57-1.65]</i>	<i>1.45 [1.38-1.52]</i>	<i>132</i>	<i>1397</i>	<i>231</i>	<i><0.0001</i>	<i><0.0001</i>	<i><0.0001</i>	<i>18</i>	<i>479</i>	<i>54</i>	<i>5.34E-05</i>	<i>6.87E-105</i>	<i>7.24E-13</i>




Appendix 31. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for RA area.

	C statistic			Hazard Ratio per 1 SD increment			Chi-square			p value			ΔAIC			Akaike Weight		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.608	0.646	0.650	1.09 [0.68-1.75]	1.55 [1.41-1.70]	1.67 [1.38-2.01]	0	82	29	0.72	<0.0001	<0.0001	5	0	3	0.013	0.251	0.058
Weight ^{1.5}	0.417	0.638	0.639	0.93 [0.56-1.56]	1.51 [1.37-1.67]	1.57 [1.29-1.90]	0	71	21	0.80	<0.0001	<0.0001	5	6	9	0.013	0.011	2.49E-03
Height*Weight	0.601	0.644	0.646	1.10 [0.68-1.76]	1.53 [1.39-1.69]	1.60 [1.32-1.93]	0	73	24	0.71	<0.0001	<0.0001	5	4	7	0.014	0.030	7.43E-03
BSA[Mosteller] ^{1.5}	0.617	0.648	0.654	1.21 [0.76-1.93]	1.55 [1.41-1.70]	1.67 [1.38-2.01]	1	80	29	0.42	<0.0001	<0.0001	4	1	3	0.017	0.193	0.062
BSA[Du Bois] ^{1.5}	0.619	0.647	0.654	1.28 [0.80-2.04]	1.55 [1.41-1.71]	1.66 [1.38-2.00]	1	78	29	0.31	<0.0001	<0.0001	4	1	3	0.021	0.131	0.054
BSA[Mosteller]	0.620	0.644	0.660	1.36 [0.85-2.16]	1.55 [1.41-1.70]	1.70 [1.42-2.04]	2	82	33	0.20	<0.0001	<0.0001	3	0	0	0.027	0.201	0.249
BSA[Du Bois]	0.625	0.644	0.662	1.41 [0.89-2.24]	1.55 [1.41-1.70]	1.70 [1.41-2.03]	2	80	33	0.15	<0.0001	<0.0001	3	1	0	0.034	0.148	0.228
LBM[Hume]	0.630	0.638	0.649	1.49 [0.93-2.37]	1.53 [1.38-1.69]	1.61 [1.34-1.95]	3	66	25	0.095	<0.0001	<0.0001	2	8	6	0.046	5.69E-03	0.012
LBM[Boer]	0.629	0.630	0.629	1.50 [0.94-2.39]	1.50 [1.35-1.67]	1.51 [1.25-1.83]	3	58	18	0.087	<0.0001	<0.0001	2	13	12	0.049	4.02E-04	5.73E-04
LBM[James]	0.612	0.631	0.628	1.21 [0.76-1.93]	1.51 [1.36-1.68]	1.37 [1.16-1.62]	1	61	14	0.42	<0.0001	0.0002	4	11	17	0.017	9.63E-04	4.37E-05
BMI	0.576	0.627	0.642	1.06 [0.66-1.71]	1.53 [1.38-1.68]	1.63 [1.36-1.96]	0	72	27	0.81	<0.0001	<0.0001	5	6	5	0.013	0.011	0.025
Height ^{2.7}	0.663	0.636	0.656	1.67 [1.07-2.59]	1.52 [1.37-1.69]	1.57 [1.31-1.87]	5	61	25	0.024	<0.0001	<0.0001	0	10	7	0.122	1.37E-03	6.36E-03
Height ^{2.13}	0.673	0.636	0.660	1.67 [1.08-2.58]	1.53 [1.38-1.70]	1.61 [1.35-1.93]	5	65	28	0.020	<0.0001	<0.0001	0	8	5	0.132	4.25E-03	0.026
Height ²	0.674	0.636	0.662	1.67 [1.08-2.57]	1.53 [1.38-1.70]	1.62 [1.36-1.94]	5	66	29	0.020	<0.0001	<0.0001	0	8	4	0.133	4.89E-03	0.035
Height ^{1.5}	0.675	0.634	0.664	1.66 [1.08-2.54]	1.53 [1.38-1.70]	1.65 [1.38-1.97]	5	67	31	0.020	<0.0001	<0.0001	0	8	2	0.132	5.17E-03	0.082
Height	0.672	0.629	0.665	1.64 [1.08-2.50]	1.52 [1.38-1.69]	1.66 [1.39-1.97]	5	65	32	0.021	<0.0001	<0.0001	0	9	2	0.123	2.43E-03	0.113
<i>Unindexed</i>	<i>0.659</i>	<i>0.612</i>	<i>0.662</i>	<i>1.59 [1.04-2.42]</i>	<i>1.49 [1.34-1.66]</i>	<i>1.62 [1.36-1.92]</i>	<i>5</i>	<i>53</i>	<i>30</i>	<i>0.031</i>	<i><0.0001</i>	<i><0.0001</i>	<i>1</i>	<i>17</i>	<i>4</i>	<i>0.093</i>	<i>5.93E-05</i>	<i>0.039</i>

Appendix 32. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for RV diameter.

	C statistic			Hazard Ratio per 1 SD increment			Chi-square			p value			ΔAIC			Akaike Weight		
	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²	BMI <18.5 kg/m ²	BMI 18.5-30 kg/m ²	BMI ≥30 kg/m ²
Weight	0.630	0.569	0.607	1.52 [0.69-3.32]	1.21 [1.06-1.39]	1.29 [1.00-1.68]	1	8	4	0.30	0.0055	0.052	0	1	4	0.064	0.126	0.022
Weight ^{1.5}	0.598	0.565	0.592	1.55 [0.76-3.18]	1.23 [1.07-1.41]	1.22 [0.96-1.55]	1	8	3	0.23	0.0036	0.11	0	0	5	0.073	0.174	0.013
Height*Weight	0.601	0.566	0.625	1.55 [0.75-3.19]	1.22 [1.06-1.41]	1.24 [1.01-1.52]	1	7	4	0.24	0.0069	0.036	0	1	4	0.072	0.123	0.021
BSA[Mosteller] ^{1.5}	0.627	0.570	0.631	1.54 [0.70-3.38]	1.21 [1.05-1.39]	1.32 [1.04-1.66]	1	7	5	0.28	0.0093	0.02	0	1	3	0.066	0.096	0.036
BSA[Du Bois] ^{1.5}	0.621	0.570	0.641	1.54 [0.70-3.38]	1.20 [1.04-1.39]	1.32 [1.06-1.64]	1	6	6	0.28	0.012	0.013	0	1	2	0.067	0.082	0.043
BSA[Mosteller]	0.638	0.569	0.635	1.45 [0.64-3.27]	1.18 [1.03-1.37]	1.34 [1.06-1.70]	1	5	6	0.37	0.021	0.014	1	2	2	0.057	0.056	0.046
BSA[Du Bois]	0.641	0.569	0.642	1.46 [0.65-3.31]	1.18 [1.02-1.37]	1.36 [1.08-1.71]	1	5	7	0.36	0.025	0.0093	0	2	2	0.057	0.050	0.058
LBM[Hume]	0.601	0.560	0.645	1.45 [0.68-3.09]	1.18 [1.01-1.37]	1.31 [1.06-1.62]	1	4	6	0.33	0.038	0.012	0	3	2	0.059	0.041	0.043
LBM[Boer]	0.601	0.553	0.641	1.44 [0.67-3.07]	1.17 [0.99-1.37]	1.39 [1.08-1.78]	1	4	7	0.35	0.059	0.011	1	3	2	0.057	0.031	0.063
LBM[James]	0.593	0.555	0.652	1.44 [0.66-3.15]	1.17 [1.01-1.36]	1.26 [1.06-1.50]	1	4	7	0.37	0.042	0.0086	1	3	3	0.056	0.038	0.040
BMI	0.604	0.562	0.537	1.20 [0.56-2.54]	1.18 [1.02-1.37]	1.14 [0.84-1.53]	0	5	1	0.64	0.027	0.41	1	3	6	0.042	0.049	6.27E-03
Height ^{2.7}	0.632	0.561	0.674	1.52 [0.70-3.31]	1.16 [0.99-1.36]	1.29 [1.10-1.53]	1	3	9	0.29	0.071	0.0025	0	4	1	0.064	0.027	0.074
Height ^{2.13}	0.652	0.564	0.669	1.51 [0.66-3.44]	1.15 [0.99-1.35]	1.38 [1.13-1.68]	1	3	10	0.33	0.074	0.0016	0	4	0	0.061	0.026	0.136
Height ²	0.658	0.565	0.668	1.50 [0.65-3.44]	1.15 [0.99-1.35]	1.39 [1.13-1.71]	1	3	10	0.34	0.077	0.0016	0	4	0	0.060	0.025	0.148
Height ^{1.5}	0.644	0.565	0.663	1.44 [0.62-3.33]	1.14 [0.98-1.33]	1.41 [1.13-1.75]	1	3	10	0.40	0.091	0.002	1	4	0	0.054	0.022	0.141
Height	0.647	0.563	0.647	1.35 [0.59-3.08]	1.13 [0.97-1.32]	1.37 [1.11-1.70]	1	2	8	0.48	0.12	0.0038	1	4	1	0.049	0.019	0.086
<i>Unindexed</i>	<i>0.595</i>	<i>0.553</i>	<i>0.607</i>	<i>1.17 [0.53-2.58]</i>	<i>1.10 [0.93-1.30]</i>	<i>1.28 [1.02-1.60]</i>	<i>0</i>	<i>1</i>	<i>4</i>	<i>0.69</i>	<i>0.25</i>	<i>0.034</i>	<i>1</i>	<i>5</i>	<i>4</i>	<i>0.041</i>	<i>0.012</i>	<i>0.024</i>

Appendix 33. Statistical analysis for 5-year cardiovascular mortality by different indexation metrics for RVOT diameter.area



Strata




9266 (1)	4622 (4243)	994 (7725)	19 (8689)
9266 (1)	5475 (3495)	2020 (6786)	139 (8634)
9265 (2)	5939 (2904)	2505 (6049)	149 (8336)

Strata

9266 (1)	4550 (4354)	987 (7795)	28 (8744)
9266 (2)	5515 (3426)	1966 (6812)	120 (8625)
9265 (1)	5971 (2862)	2566 (5953)	159 (8290)

Strata

9266 (1)	4628 (4293)	1104 (7694)	44 (8739)
9266 (2)	5438 (3474)	1895 (6851)	96 (8625)
9265 (1)	5970 (2875)	2520 (6015)	167 (8295)



Strata




17759 (2)	8743 (7912)	2178 (14145)	82 (16208)
17759 (1)	10857 (6066)	4401 (12150)	348 (16130)
17759 (0)	11284 (5243)	5037 (10825)	327 (15383)

Strata

17759 (2)	9090 (7763)	2383 (14179)	72 (16459)
17759 (1)	10686 (6165)	4254 (12227)	358 (16059)
17759 (0)	11108 (5293)	4979 (10714)	327 (15203)

Strata

17759 (2)	9506 (7437)	2678 (13970)	101 (16511)
17759 (0)	10579 (6218)	4259 (12135)	320 (15997)
17759 (1)	10790 (5566)	4679 (11015)	336 (15213)



Strata

577 (0)	214 (304)	44 (473)	2 (515)
577 (0)	263 (252)	85 (417)	9 (491)
576 (1)	261 (237)	91 (370)	6 (451)


Strata

577 (0)	233 (297)	51 (478)	1 (528)
577 (0)	258 (258)	88 (416)	9 (492)
576 (1)	247 (238)	81 (366)	7 (437)




Strata

577 (0)	247 (288)	61 (472)	1 (532)
577 (0)	252 (257)	83 (412)	10 (481)
576 (1)	239 (248)	76 (376)	6 (444)

Appendix 34. Long-term cardiovascular mortality for aorta at sinotubular diameter unindexed, indexed by BSA and indexed by weight.






Appendix 35. Long-term cardiovascular mortality for aortic sinus diameter unindexed, indexed by BSA and indexed by weight.



Strata	3844 (3)	1559 (2175)	255 (3445)	0 (3699)
—	3844 (1)	1406 (2341)	276 (3438)	0 (3711)
—	3843 (1)	1447 (2293)	352 (3334)	1 (3681)

Strata	3844 (1)	1555 (2185)	245 (3471)	0 (3715)
—	3844 (2)	1431 (2317)	293 (3415)	0 (3704)
—	3843 (2)	1426 (2307)	345 (3331)	1 (3672)




Strata	3844 (1)	1489 (2252)	234 (3482)	0 (3715)
—	3844 (3)	1490 (2262)	307 (3408)	0 (3712)
—	3843 (1)	1433 (2295)	342 (3327)	1 (3664)



Strata	7041 (2)	2979 (3828)	554 (6197)	0 (6745)
—	7041 (1)	2721 (4088)	612 (6122)	0 (6731)
—	7040 (3)	2798 (3911)	795 (5815)	0 (6585)

Strata	7041 (2)	3182 (3670)	673 (6122)	0 (6790)
—	7041 (2)	2679 (4140)	609 (6137)	0 (6740)
—	7040 (2)	2637 (4017)	679 (5875)	0 (6531)

Strata	7041 (3)	3160 (3712)	676 (6140)	0 (6809)
—	7041 (1)	2754 (4068)	649 (6090)	0 (6733)
—	7040 (2)	2584 (4047)	636 (5904)	0 (6519)




Strata	254 (0)	108 (138)	25 (220)	0 (245)
—	254 (0)	87 (151)	24 (212)	0 (236)
—	253 (0)	82 (152)	21 (211)	0 (232)


Strata	254 (0)	117 (132)	30 (219)	0 (249)
—	254 (0)	95 (148)	25 (216)	0 (241)
—	253 (0)	65 (161)	15 (208)	0 (223)

Strata	254 (0)	119 (130)	31 (218)	0 (249)
—	254 (0)	96 (147)	26 (215)	0 (241)
—	253 (0)	62 (164)	13 (210)	0 (223)


Appendix 36. Long-term cardiovascular mortality for aortic arch diameter unindexed, indexed by BSA and indexed by weight.




—	27671 (5)	16226 (10888)	5318 (21249)	1172 (25286)	228 (26202)	9 (26416)	1 (26424)
- - -	27671 (5)	16021 (10759)	5745 (20573)	1312 (24853)	177 (25956)	8 (26122)	0 (26130)
...	27671 (5)	15764 (10750)	5928 (19954)	1273 (24395)	154 (25484)	6 (25632)	0 (25638)




—	27671 (4)	16037 (10934)	5142 (21519)	1062 (25517)	171 (26387)	6 (26551)	1 (26556)
- - -	27671 (7)	16168 (10673)	5732 (20684)	1296 (24978)	181 (26065)	6 (26236)	0 (26242)
...	27671 (4)	15806 (10590)	6117 (19573)	1399 (24039)	207 (25190)	11 (25383)	0 (25394)




—	27671 (4)	15944 (11019)	5119 (21520)	1049 (25495)	144 (26381)	5 (26519)	1 (26523)
- - -	27671 (8)	16108 (10721)	5743 (20668)	1313 (24962)	201 (26044)	8 (26234)	0 (26242)
...	27671 (3)	15959 (10457)	6129 (19586)	1395 (24077)	214 (25217)	10 (25417)	0 (25427)




—	30172 (18)	36699 (21213)	13852 (43218)	3683 (53130)	1009 (55749)	38 (56716)	2 (56751)
- - -	30171 (8)	35070 (22123)	13408 (42637)	3534 (52133)	730 (54857)	18 (55560)	2 (55576)
...	30171 (5)	34128 (22361)	13652 (41215)	3523 (50808)	597 (53631)	17 (54195)	2 (54209)




—	30172 (11)	37805 (20657)	14026 (43802)	3626 (53977)	915 (56635)	20 (57527)	3 (57544)
- - -	30171 (13)	35386 (22184)	13853 (42624)	3790 (52360)	769 (55301)	27 (56029)	1 (56055)
...	30171 (7)	32706 (22856)	13033 (40644)	3324 (49734)	652 (52301)	26 (52915)	2 (52937)




—	30172 (10)	37439 (21010)	13780 (43974)	3449 (54065)	784 (56674)	17 (57435)	2 (57450)
- - -	30171 (15)	35572 (21977)	13986 (44246)	3765 (52287)	808 (55157)	24 (55933)	2 (55955)
...	30171 (6)	32886 (22710)	13146 (40670)	3526 (49719)	744 (52406)	32 (53103)	2 (53131)



—	2010 (0)	1103 (787)	426 (1430)	148 (1705)	36 (1814)	1 (1849)
- - -	2009 (0)	980 (858)	349 (1435)	120 (1654)	32 (1740)	0 (1770)
...	2009 (0)	809 (952)	284 (1407)	87 (1587)	18 (1652)	1 (1669)




—	2010 (0)	1180 (741)	488 (1407)	192 (1701)	54 (1837)	1 (1889)
- - -	2009 (0)	929 (895)	329 (1450)	104 (1662)	20 (1742)	0 (1761)
...	2009 (0)	783 (961)	242 (1415)	59 (1583)	12 (1627)	1 (1638)




—	2010 (0)	1182 (739)	499 (1397)	198 (1696)	55 (1837)	1 (1890)
- - -	2009 (0)	934 (896)	320 (1459)	97 (1667)	20 (1739)	0 (1758)
...	2009 (0)	776 (962)	240 (1416)	60 (1583)	11 (1630)	1 (1640)


Appendix 37. Long-term cardiovascular mortality for aortic root diameter unindexed, indexed by BSA and indexed by weight.




Appendix 38. Long-term cardiovascular mortality for ascending aorta diameter unindexed, indexed by BSA and indexed by weight.




Strata	15791 (6)	7481 (7976)	2255 (13062)	295 (14996)	92 (15194)	5 (15278)
—	15790 (2)	8074 (7149)	3060 (11911)	481 (14412)	81 (14804)	7 (14895)
—	15790 (2)	8032 (6099)	3220 (10975)	483 (13595)	44 (13990)	5 (14027)




Strata	15791 (4)	7405 (8050)	2133 (13224)	215 (15122)	68 (15264)	4 (15327)
—	15790 (4)	8073 (7165)	3174 (11817)	557 (14362)	104 (14804)	10 (14895)
—	15790 (2)	8089 (6609)	3228 (10907)	467 (13519)	45 (13920)	3 (13959)




Strata	15791 (4)	7390 (8044)	2117 (13216)	217 (15093)	63 (15240)	5 (15298)
—	15790 (3)	8009 (7239)	3109 (11888)	527 (14401)	106 (14814)	9 (14907)
—	15790 (3)	8168 (6541)	3309 (10844)	495 (13509)	48 (13934)	3 (13976)




Strata	8819 (13)	14446 (13785)	5097 (22912)	955 (27003)	302 (27644)	25 (27916)
—	28818 (5)	14908 (12651)	6110 (20987)	1175 (25792)	250 (26688)	16 (26917)
—	28818 (5)	14561 (11234)	6307 (18271)	1146 (23086)	165 (24011)	10 (24160)




Strata	8819 (13)	14481 (13809)	4871 (23221)	755 (27294)	249 (27794)	16 (28024)
—	28818 (6)	15258 (12457)	6621 (20662)	1479 (25675)	327 (26782)	24 (27078)
—	28818 (4)	14176 (11404)	6022 (18287)	1042 (22912)	141 (23767)	11 (23891)




Strata	8819 (14)	14479 (13790)	4875 (23181)	743 (27264)	239 (27761)	15 (27982)
—	28818 (4)	15407 (12354)	6709 (20620)	1485 (25708)	315 (26831)	21 (27118)
—	28818 (5)	14029 (11526)	5930 (18369)	1048 (22909)	163 (23751)	15 (23893)



Strata	1055 (0)	436 (584)	143 (873)	18 (996)	7 (1007)	0 (1014)
—	1054 (0)	457 (539)	168 (814)	36 (938)	8 (965)	1 (972)
—	1054 (0)	359 (486)	129 (665)	26 (758)	4 (776)	0 (779)






Strata	1055 (0)	454 (571)	151 (870)	23 (996)	9 (1010)	0 (1019)
—	1054 (0)	458 (541)	181 (807)	35 (945)	8 (971)	1 (978)
—	1054 (0)	340 (497)	108 (675)	22 (751)	2 (767)	0 (768)



Strata	1055 (0)	470 (558)	163 (861)	28 (993)	8 (1013)	0 (1021)
—	1054 (0)	451 (543)	176 (807)	33 (942)	10 (964)	1 (972)
—	1054 (0)	331 (508)	101 (684)	19 (757)	1 (771)	0 (772)

Appendix 39. Long-term cardiovascular mortality for LA area unindexed, indexed by BSA and indexed by weight.



Strata




12122 (0)	7730 (4154)	2922 (8824)	852 (10835)	194 (11477)	6 (11664)	0 (11670)
12122 (0)	7936 (3799)	3456 (8059)	1004 (10415)	194 (11204)	8 (11385)	1 (11392)
12122 (4)	7282 (3935)	3595 (7138)	974 (9560)	155 (10337)	9 (10480)	0 (10489)

Strata

12122 (0)	7610 (4277)	2669 (9102)	715 (11016)	133 (11586)	4 (11715)	0 (11719)
12122 (1)	7970 (3771)	3596 (7926)	1059 (10385)	223 (11180)	9 (11392)	1 (11400)
12122 (3)	7368 (3840)	3708 (6993)	1056 (9429)	187 (10252)	10 (10422)	0 (10432)

Strata

12122 (0)	7562 (4301)	2676 (9052)	694 (10992)	119 (11555)	2 (11671)	0 (11673)
12122 (1)	7889 (3834)	3558 (7957)	1049 (10354)	223 (11158)	12 (11368)	1 (11379)
12122 (3)	7497 (3753)	3739 (7012)	1087 (9464)	201 (10305)	9 (10490)	0 (10499)



Strata




24382 (6)	16310 (7491)	7214 (16338)	2759 (20693)	899 (22510)	31 (23371)	2 (23400)
24381 (1)	16302 (7060)	8326 (14596)	3057 (19642)	835 (21798)	22 (22603)	1 (22624)
24381 (5)	14489 (7369)	7692 (12962)	2480 (17689)	555 (19517)	20 (20038)	3 (20054)

Strata

24382 (6)	16411 (7472)	7061 (16596)	2509 (21056)	749 (22778)	21 (23503)	2 (23522)
24381 (2)	16492 (6968)	8668 (14346)	3295 (19512)	925 (21818)	30 (22700)	1 (22729)
24381 (4)	14198 (7480)	7503 (12954)	2492 (17456)	615 (19229)	22 (19809)	3 (19827)

Strata

24382 (6)	16477 (7386)	7132 (16489)	2491 (21025)	690 (22778)	20 (23444)	2 (23462)
24381 (3)	16319 (7073)	8539 (14364)	3173 (19509)	905 (21716)	24 (22586)	1 (22609)
24381 (3)	14305 (7461)	7561 (13043)	2632 (17490)	694 (19331)	29 (19982)	3 (20007)



Strata

904 (0)	530 (346)	226 (641)	100 (785)	34 (831)	0 (865)
904 (1)	544 (307)	274 (556)	114 (711)	30 (793)	0 (823)
903 (0)	421 (338)	168 (533)	69 (614)	19 (659)	1 (675)




Strata

904 (0)	560 (320)	264 (612)	120 (755)	41 (834)	0 (875)
904 (1)	520 (333)	253 (578)	109 (715)	28 (793)	1 (820)
903 (0)	415 (338)	151 (540)	54 (620)	14 (656)	0 (668)

Strata

904 (0)	568 (312)	271 (602)	122 (748)	40 (830)	0 (870)
904 (1)	514 (343)	252 (582)	110 (719)	29 (797)	1 (825)
903 (0)	413 (336)	145 (546)	51 (623)	14 (656)	0 (668)




Appendix 40. Long-term cardiovascular mortality for LA diameter unindexed, indexed by BSA and indexed by weight.



21698 (6)	13827 (7560)	4233 (16980)	503 (20664)
21698 (1)	12588 (8504)	4154 (16605)	382 (20314)
21697 (5)	11949 (8477)	4446 (15320)	212 (19440)

21698 (6)	13845 (7563)	4247 (17002)	499 (20708)
21698 (1)	12580 (8543)	4094 (16704)	387 (20345)
21697 (5)	11939 (8435)	4492 (15199)	211 (19365)




21698 (6)	13772 (7643)	4189 (17070)	472 (20748)
21698 (1)	12599 (8514)	4110 (16674)	408 (20306)
21697 (5)	11993 (8384)	4534 (15161)	217 (19364)



46137 (6)	29774 (15570)	9687 (35221)	1085 (43720)
46137 (8)	27334 (17258)	9359 (34503)	932 (42780)
46136 (4)	24371 (17225)	9135 (30793)	562 (39084)

46137 (8)	30004 (15448)	9727 (35347)	1122 (43855)
46137 (5)	27442 (17282)	9450 (34562)	925 (42951)
46136 (5)	24033 (17323)	9004 (30608)	532 (38778)

46137 (5)	30083 (15397)	9732 (35374)	1135 (43870)
46137 (7)	27457 (17262)	9546 (34466)	916 (42967)
46136 (6)	23939 (17394)	8903 (30677)	528 (38747)






1430 (0)	748 (630)	228 (1137)	25 (1340)
1430 (0)	675 (649)	209 (1085)	14 (1277)
1430 (0)	487 (646)	135 (936)	13 (1054)

1430 (0)	762 (624)	232 (1142)	26 (1348)
1430 (0)	683 (641)	217 (1080)	13 (1281)
1430 (0)	465 (660)	123 (936)	13 (1042)

1430 (0)	780 (608)	242 (1135)	26 (1351)
1430 (0)	667 (655)	207 (1087)	14 (1277)
1430 (0)	463 (662)	123 (936)	12 (1043)

Appendix 41. Long-term cardiovascular mortality for LA volume unindexed, indexed by BSA and indexed by weight.



Strata




35365 (9)	18089 (16218)	5957 (27910)	1421 (32294)	217 (33472)	4 (33681)	0 (33685)
35365 (7)	19365 (15121)	6247 (27817)	1135 (32804)	125 (33796)	7 (33912)	1 (33918)
35364 (3)	18957 (14703)	6561 (26359)	1177 (31533)	145 (32531)	9 (32665)	0 (32674)

Strata

35365 (10)	18043 (16399)	5781 (26309)	1328 (32654)	175 (33785)	2 (33957)	0 (33959)
35365 (5)	19129 (15332)	6203 (27800)	1157 (32703)	132 (33711)	8 (33834)	1 (33841)
35364 (4)	19239 (14311)	6781 (25977)	1248 (31274)	180 (32303)	10 (32467)	0 (32477)

Strata

35365 (9)	18092 (16359)	5747 (26340)	1258 (32714)	144 (33809)	3 (33949)	0 (33952)
35365 (6)	19021 (15360)	6259 (27676)	1180 (32610)	149 (33620)	6 (33762)	1 (33767)
35364 (4)	19298 (14323)	6759 (26070)	1295 (31307)	194 (32370)	11 (32547)	0 (32558)



Strata




4326 (27)	38219 (32626)	14306 (55296)	4186 (65027)	983 (68154)	24 (69107)	3 (69127)
4326 (19)	41993 (29758)	14706 (55936)	3097 (67267)	503 (69811)	16 (70293)	0 (70309)
4325 (9)	40842 (28670)	15088 (52793)	3336 (64048)	639 (66652)	32 (67244)	3 (67272)

Strata

4326 (28)	39513 (32044)	14692 (55824)	4189 (66013)	904 (69227)	16 (70109)	3 (70122)
4326 (16)	41510 (30197)	14566 (56040)	3092 (67206)	527 (69712)	18 (70215)	2 (70231)
4325 (11)	40031 (28813)	14842 (52161)	3338 (63123)	694 (65678)	38 (66320)	1 (66355)

Strata

4326 (27)	40226 (31648)	14617 (56262)	3824 (66745)	719 (69779)	11 (70484)	2 (70493)
4326 (13)	41319 (30066)	14890 (55295)	3396 (66464)	688 (69107)	23 (69760)	3 (69779)
4325 (15)	39509 (29340)	14593 (52468)	3399 (63133)	718 (65731)	38 (66400)	1 (66436)



Strata

2804 (0)	1117 (1462)	426 (2100)	152 (2368)	32 (2485)	1 (2516)
2803 (0)	1364 (1289)	501 (2097)	100 (2486)	19 (2564)	0 (2583)
2803 (0)	1418 (1091)	593 (1848)	98 (2330)	27 (2396)	1 (2422)


Strata

2804 (0)	1188 (1431)	466 (2111)	185 (2380)	41 (2521)	1 (2561)
2803 (0)	1293 (1335)	429 (2141)	99 (2461)	19 (2537)	0 (2566)
2803 (0)	1418 (1076)	625 (1793)	66 (2343)	18 (2389)	1 (2404)



















Strata

2804 (0)	1242 (1395)	492 (2102)	189 (2393)	43 (2535)	1 (2577)
2803 (0)	1258 (1344)	403 (2146)	104 (2434)	20 (2515)	0 (2535)
2803 (0)	1399 (1103)	625 (1797)	57 (2357)	15 (2397)	1 (2409)




Appendix 42. Long-term cardiovascular mortality for LV diameter unindexed, indexed by BSA and indexed by weight.



Appendix 43. Long-term cardiovascular mortality for LV volume unindexed, indexed by BSA and indexed by weight.



Appendix 44. Long-term cardiovascular mortality for IVS thickness unindexed, indexed by BSA and indexed by weight.



Strata




4576 (10)	17642 (16376)	5391 (28370)	1180 (32494)	239 (33412)	5 (33642)	0 (33647)
34575 (6)	18568 (15073)	5954 (27276)	1126 (31974)	123 (32955)	9 (33067)	1 (33075)
34575 (2)	18569 (14020)	6734 (25005)	1218 (30279)	98 (31370)	6 (31460)	0 (31466)

Strata

4576 (10)	17445 (16630)	5268 (26594)	1161 (32632)	230 (33545)	5 (33767)	0 (33772)
34575 (5)	18812 (14939)	5858 (27526)	1072 (32195)	114 (33131)	8 (33235)	1 (33242)
34575 (3)	18522 (13900)	6953 (24531)	1291 (29920)	116 (31061)	7 (31167)	0 (31174)

Strata

4576 (10)	17334 (16742)	5122 (28750)	1109 (32697)	205 (33584)	5 (33782)	0 (33787)
34575 (5)	18828 (14920)	5939 (27456)	1095 (32187)	130 (33131)	7 (33253)	1 (33259)
34575 (3)	18617 (13807)	7018 (24445)	1320 (29883)	125 (31022)	8 (31134)	0 (31142)



Strata




2990 (34)	39132 (32331)	13759 (57105)	3862 (66805)	1106 (69507)	28 (70581)	3 (70605)
2990 (14)	40372 (29994)	14055 (55249)	3031 (65975)	544 (68396)	22 (68904)	0 (68926)
2990 (4)	38901 (27725)	15098 (49347)	3314 (60508)	399 (63331)	19 (63704)	2 (63721)

Strata

2990 (31)	39216 (32546)	13652 (57624)	3874 (67233)	1101 (69553)	24 (71027)	3 (71048)
2990 (17)	40969 (29820)	14143 (55741)	2942 (66675)	509 (69056)	20 (69534)	0 (69553)
2990 (4)	38220 (27684)	15117 (48336)	3391 (59380)	439 (62225)	25 (62628)	2 (62651)

Strata

2990 (29)	39313 (32558)	13564 (57850)	3822 (67428)	1079 (70117)	19 (71174)	3 (71190)
2990 (18)	41271 (29641)	14312 (55729)	2941 (66838)	527 (69203)	22 (69698)	0 (69719)
2990 (5)	37821 (27851)	15036 (48122)	3444 (59022)	443 (61914)	28 (62317)	2 (62343)



Strata

2755 (0)	1299 (1377)	461 (2192)	144 (2505)	34 (2614)	0 (2648)
2754 (0)	1352 (1255)	562 (1992)	122 (2425)	22 (2521)	2 (2540)
2754 (0)	1182 (1152)	473 (1763)	73 (2143)	16 (2196)	0 (2211)




Strata

2755 (0)	1284 (1393)	444 (2213)	167 (2485)	37 (2614)	0 (2651)
2754 (0)	1356 (1264)	544 (2021)	113 (2445)	25 (2529)	2 (2551)
2754 (0)	1193 (1127)	508 (1713)	59 (2143)	10 (2188)	0 (2197)

Strata

2755 (0)	1292 (1394)	451 (2217)	172 (2491)	39 (2622)	0 (2661)
2754 (0)	1291 (1306)	482 (2058)	114 (2419)	24 (2505)	2 (2526)
2754 (0)	1250 (1084)	563 (1672)	53 (2163)	9 (2204)	0 (2212)

Appendix 45. Long-term cardiovascular mortality for LV mass unindexed, indexed by BSA and indexed by weight.



Strata




9503 (1)	5123 (4079)	1119 (8020)	435 (8662)	111 (8974)	6 (9074)
9503 (1)	5838 (3269)	2463 (6450)	644 (8175)	98 (8705)	11 (8790)
9503 (2)	5023 (3900)	1817 (8829)	159 (8407)	8 (8549)	0 (8557)

Strata

9503 (1)	5137 (4075)	1040 (8111)	403 (8718)	109 (9000)	6 (9099)
9503 (2)	5788 (3333)	2526 (8420)	699 (8141)	100 (8725)	11 (8811)
9503 (1)	5059 (3840)	1833 (6768)	136 (8365)	8 (8503)	0 (8511)

Strata

9503 (1)	5106 (4094)	1040 (8106)	393 (8716)	94 (9004)	6 (9091)
9503 (1)	5764 (3377)	2489 (6478)	683 (8187)	108 (8748)	10 (8840)
9503 (2)	5114 (3777)	1870 (6715)	162 (8341)	15 (8476)	1 (8490)



Strata




17118 (9)	9451 (7075)	2704 (13679)	1099 (15226)	325 (15979)	29 (16266)
17118 (1)	10728 (5537)	4875 (11056)	1610 (14117)	333 (15345)	18 (15656)
17117 (2)	9112 (6340)	3857 (10903)	565 (13974)	58 (14454)	3 (14506)

Strata

17118 (8)	9665 (6946)	2748 (13738)	1161 (15256)	337 (16055)	25 (16363)
17118 (7)	10667 (5637)	4995 (11011)	1659 (14159)	336 (15436)	22 (15739)
17117 (1)	8959 (6369)	3695 (10889)	454 (13902)	43 (14287)	3 (14326)

Strata

17118 (8)	9702 (6897)	2772 (13701)	1155 (15244)	323 (16046)	19 (16347)
17118 (7)	10740 (5639)	5038 (11027)	1670 (14220)	352 (15497)	28 (15809)
17117 (1)	8849 (6416)	3626 (10910)	449 (13853)	41 (14235)	3 (14272)



Strata

613 (0)	285 (295)	85 (493)	22 (555)	8 (569)	0 (577)
612 (0)	319 (250)	122 (427)	44 (497)	10 (528)	1 (537)
612 (0)	216 (286)	72 (396)	14 (444)	1 (455)	0 (455)


Strata

613 (0)	299 (293)	96 (495)	28 (562)	9 (581)	0 (590)
612 (0)	309 (250)	115 (426)	39 (493)	8 (521)	1 (528)
612 (0)	212 (288)	68 (395)	13 (441)	2 (450)	0 (451)



















Strata

613 (0)	305 (290)	101 (493)	32 (560)	10 (582)	0 (592)
612 (0)	308 (249)	114 (424)	35 (495)	7 (520)	1 (526)
612 (0)	207 (292)	64 (399)	13 (441)	2 (450)	0 (451)

Appendix 46. Long-term cardiovascular mortality for RA area unindexed, indexed by BSA and indexed by weight.



Appendix 47. Long-term cardiovascular mortality for RV diameter unindexed, indexed by BSA and indexed by weight.



Appendix 48. Long-term cardiovascular mortality for RVOT diameter unindexed, indexed by BSA and indexed by weight.