

Changing of therapeutic trends between the 1st and 2nd wave did not reduce COVID-19 related mortality of renal transplant recipients: a national registry study.

Bastien Berger¹, Marc Hazzan², Nassim Kamar³, H el ene Francois⁴, Marie Matignon⁵, Clarisse Greze⁶, Philippe Gatault⁷, Luc Frimat⁸, Pierre F Westeel⁹, Valentin Goutaudier¹⁰, Renaud Snanoudj¹¹, Charlotte Colosio¹², Antoine Sicard¹³, Dominique Bertrand¹⁴, Christiane Mousson¹⁵, Jamal Bamoulid¹⁶, Antoine Thierry¹⁷, Dany Anglicheau¹⁸, Lionel Couzi¹⁹, Jonathan M Chemouny²⁰, Agnes Duveau²¹, Valerie Moal²², Yannick Le Meur²³, Gilles Blancho²⁴, J er ome Turret²⁵, Paolo Malvezzi²⁶, Christophe Mariat²⁷, Jean-Philippe Rerolle²⁸, Nicolas Bouvier²⁹, the French SOT COVID Registry, Sophie Caillard^{30,31}, and Olivier Thauvat^{1,32,33,#}

1) Department of Transplantation, Nephrology and Clinical Immunology, Edouard Herriot Hospital, Hospices civils de Lyon, Lyon, France.

2) Department of Nephrology and Transplantation, University of Lille, Lille, France.

3) Department of Nephrology and Transplantation, University of Toulouse, Toulouse, France.

4) Nephrology and Renal Transplantation Department, AP-HP (Assistance Publique-H opitaux de Paris), H opital Tenon, Paris, France.

5) Nephrology and Renal Transplantation Department, AP-HP (Assistance Publique-H opitaux de Paris), Institut Francilien de Recherche en N ephrologie et Transplantation (IFRNT), Groupe Hospitalier Henri-Mondor/Albert-Chenevier, Universit  Paris-Est-

Créteil (UPEC), DHU (Département Hospitalo-Universitaire) VIC (Virus-Immunité-Cancer), IMRB (Institut Mondor de Recherche Biomédicale), Equipe 21, INSERM U 955, Créteil, France.

6) Department of Nephrology and Transplantation, Hôpital Bichat, Paris, France.

7) Department of Nephrology and Transplantation, University of Tours, Tours, France.

8) Department of Nephrology, University of Lorraine, CHRU-Nancy, Vandoeuvre, France; INSERM CIC-EC CIE6, Nancy, France.

9) Department of Nephrology and Transplantation, University of Amiens, Amiens, France.

10) Department of Nephrology and Transplantation, University of Montpellier, Montpellier, France.

11) Nephrology and Renal Transplantation Department, Hôpital Foch, Paris, France.

12) Department of Nephrology and Transplantation, University of Reims, Reims, France.

13) Service de Néphrologie-Dialyse-Transplantation, Hôpital Pasteur 2, CHU de Nice, Unité de Recherche Clinique Côte d'Azur (UR2CA), Université Côte d'Azur, Nice, France.

14) Department of Nephrology and Transplantation, University of Rouen, Rouen, France.

15) Department of Nephrology and Transplantation, University of Dijon, Dijon, France.

16) Department of Nephrology, University of Besançon, Besançon, France.

17) Department of Nephrology and Transplantation, University of Poitiers, Poitiers, France.

18) Service de Néphrologie et Transplantation Adultes, Hôpital Universitaire Necker - APHP Centre - Université de Paris INEM INSERM U 1151 - CNRS UMR 8253, Paris, France.

19) Service de Néphrologie-Transplantation-Dialyse-Aphérèse, Hôpital Pellegrin, CHU de Bordeaux Pellegrin, Unité Mixte de Recherche "ImmunoConcEpT" 5164 - Université de Bordeaux, Bordeaux, France.

20) University of Rennes, CHU Rennes, Inserm, EHESP, Irset (Institut de Recherche en Santé, Environnement et Travail) - UMR_S 1085, CIC-P 1414, Rennes, France.

21) Department of Nephrology and Transplantation, University of Angers, Angers, France.

22) Centre de Néphrologie et Transplantation Rénale, Aix Marseille Université, Hôpitaux Universitaires de Marseille, Hôpital Conception, Marseille, France.

23) Department of Nephrology, CHU de Brest, UMR1227, Lymphocytes B et Autoimmunité, Université de Brest, Inserm, Labex IGO, Brest, France.

24) Department of Nephrology and Transplantation, Centre Hospitalier Universitaire de Nantes, Nantes, France.

25) Nephrology and Renal Transplantation Department, AP-HP (Assistance Publique-Hôpitaux de Paris), Hôpital de la Pitié Salpêtrière, Paris, France.

26) Department of Nephrology, University of Grenoble, Grenoble, France.

27) Department of Nephrology and Transplantation, University of St Etienne, St Etienne, France.

28) Department of Nephrology and Transplantation, University of Limoges, Limoges, France.

29) Department of Nephrology and Transplantation, University of Caen, Caen, France.

30) Department of Nephrology and Transplantation, Strasbourg University Hospital, Strasbourg, France.

31) INSERM, IRM UMR-S 1109, University of Strasbourg, Strasbourg, France.

32) CIRI, INSERM U1111, University Claude Bernard Lyon I, Lyon, France.

33) Claude Bernard University (Lyon 1), Villeurbanne, France.

Correspondence to:

Olivier Thauvat MD, PhD

Service de Transplantation, Néphrologie et Immunologie Clinique,

Hôpital Edouard Herriot, 5 Place d'Arsonval, 69003 Lyon, France

Phone: +33 472110170

Email: olivier.thauvat@chu-lyon.fr

Keywords : Transplantation, SARS-CoV2, COVID-19.

Content: 3036 words, 1 table, 4 figures

Abstract (246 words)

SARS-CoV-2 pandemic evolved in two consecutive waves over 2020 (for France: 1st wave from March 1 to July 31; and 2nd wave from August 1 to December 31).

Improvements in the management of COVID-19 led to a reduction of mortality rates in hospitalized patients during the second wave. Whether this progress also benefited to kidney transplant recipients (KTR), a population particularly vulnerable to severe COVID-19, remained unclear.

In France, 957 KTR were hospitalized for COVID-19 in 2020 and their data were prospectively collected in the French SOT COVID registry. The presentation, management, and outcomes of the 359 KTR diagnosed during the 1st wave were compared to those of the 598 of the 2nd wave. Baseline comorbidities were largely similar between KTR of the 2 waves. Maintenance immunosuppression was reduced in most patients but withdrawal of antimetabolite (73.7% vs 58.4%, $p<0.001$) or CNI (32.1% vs 16.6%, $p<0.001$) was less frequent during the 2nd wave. Hydroxychloroquine and azithromycin that were commonly used during the 1st wave (21.7% and 30.9%, respectively) were almost abandoned during the 2nd. In contrast, the use of high dose corticosteroids doubled (19.5% vs. 41.6%, $p<0.001$). Despite these changing trends in COVID-19 management, 60-day mortality was not statistically different between the 2 waves (25.3% vs. 23.9%; Log Rank, $p=0.48$).

We conclude that changing of therapeutic trends during 2020 did not reduce COVID-19 related mortality in KTR. Our data indirectly support the importance of vaccination and monoclonal neutralizing anti-SARS-CoV-2 antibodies to protect KTR from severe COVID-19.

Introduction

After the initial outbreak in China in late 2019, coronavirus disease (COVID-19) spread globally¹. As on October 14, 2021, the pandemic had affected more than 238 million people causing more than 4.8 million deaths worldwide².

Like in the rest of the world^{3,4}, the viral pandemic evolved during 2020 in two consecutive waves in France. The first hit France in spring, only three months after SARS-CoV-2 discovery⁵, in a context of limited knowledge about COVID-19, absence of proven specific treatment, and shortage of essential equipment such as face masks and diagnostic tests^{6,7}. The government imposed a national lockdown from March 17, 2020 to May 10, 2020, which successfully reduced the spread of the virus and led to the resolution of this first wave⁸. However, in the absence of available vaccine, SARS-CoV-2 resurged following the ease of social distancing rules during the summer. As a result, a second pandemic wave started in fall 2020. However, in contrast with the first wave, enhanced testing capacities allowed diagnosis of asymptomatic cases during this second wave. Additionally, intensivists had better experience of the stereotypical course of severe COVID-19, including the prolonged mechanical ventilation and Intensive Care Unit (ICU) stay⁹, the increased risk of thrombotic events¹⁰, and the high rates of acute kidney injury¹¹. More importantly, RECOVERY trial¹² had been published, providing evidence that Dexamethasone reduces mortality in hospitalized patients requiring oxygen therapy by one fifth. These changes in medical care resulted in a 10% reduction of mortality rates in French hospitalized patients during the second wave compared to the first one^{13,14}.

Whether kidney transplant recipients (KTR), a particularly vulnerable population to COVID-19¹⁵⁻¹⁷, also benefited from the progresses made along 2020 in COVID-19 management remained unclear. Aiming at addressing this question, we retrospectively analyzed the prospectively collected data of the French Solid Organ Transplant (SOT) COVID registry and compare the course, management and outcomes of COVID-19 diagnosed in the first versus second waves in 957 hospitalized French KTR.

Methods

Data Collection

KTR hospitalized for COVID-19 in France between March 1 and December 31 2020 were identified by the interrogation of the French Solid Organ Transplant (SOT) COVID registry.

This prospective registry was approved by the Institutional Review Board of Strasbourg University (approval number 02.26) and registered at clinicaltrials.gov (NCT04360707). Of note, all patients were informed about their inclusion in the registry but the need for informed consent was waived.

Study Design and Patients

Inclusion criteria were age > 18 years at the diagnosis of COVID-19 and presence of a functioning kidney graft. The diagnostic criteria for COVID-19 was based on: (i) a positive RT-PCR for SARS-CoV-2 in nasopharyngeal swab or (ii) the presence of typical respiratory symptoms accompanied by evocative pulmonary lesions on low-dose chest CT when RT-PCR yielded negative results. Cases were considered to have occurred during the 1st wave if they were diagnosed between March 1 to July 31, 2020 and during 2nd wave if they were diagnosed between August 1 to December 31, 2020. We used the time cutoff of December 31, 2020 for the end of the second wave to have an equal length of time compared to the first wave and to avoid the effect of the vaccination to increase baseline comparability. Cardiovascular disease included heart failure, coronary vascular disease and dysrhythmia. Respiratory disease included chronic respiratory failure, asthma and chronic obstructive pulmonary disease.

Statistical Analysis

Categorical variables are reported as counts and percentages. Continuous variables are presented as medians and interquartile ranges. Differences between groups were assessed with the χ^2 test or 2-sided Fisher's exact test for categorical variables and with Student's t-test or Wilcoxon's rank-sum test for continuous variables. Survival curves were represented using the Kaplan-Meier method and compared with the log-rank test. The primary outcome is 60-day mortality. Secondary outcomes are: admission to the ICU, 60-day mortality in ICU, initiation of renal replacement therapy (RRT), use of mechanical ventilation, use of vasopressor support, occurrence of bacterial pulmonary superinfection, or thrombo-embolic event. The multiple imputations method¹⁸ was used to handle missing data on relevant covariates. Five imputed data sets were generated and analyses were performed on each of them. Then, the results were combined using the Rubin rules¹⁹ to obtain average values. To assess risk factors for mortality, Cox proportional hazard univariable and multivariable models were built. All the variables with a univariable threshold $p < 0.1$ were selected as covariates for the initial multivariable model. The covariates in the final multivariable model were selected using a backward conditional procedure with a threshold $p < 0.05$. Results are expressed as hazard ratios (HRs) with their 95% confidence intervals. All analyses were conducted in the R environment (R Foundation for Statistical Computing, Vienne, Austria) version 4.1.2²⁰ using the "survival" and "mice" packages. All tests were 2-sided, and $p < 0.05$ was considered statistically significant.

Results

Baseline patient characteristics:

Shortage in diagnosis assays during the first pandemic wave resulted in the fact that only symptomatic patients were tested to confirm clinically or radiologically suspected COVID-19^{21,22}. As the result of enhanced availability of these assays along the year 2020, asymptomatic COVID-19 were identified during the second wave²¹. Furthermore, from January 2021 onward anti-SARS-CoV-2 vaccines became available, reducing the risk of severe COVID-19 and contributing to the resolution of the second pandemic wave. Since the criteria for hospitalization of KTR with symptomatic COVID-19 little evolved over time and given the fact that our aim was to compare the two pandemic waves, the present study focused on the 957 cases [n=359 (37.5%) from the first and n=598 (62.5%) from the second wave] of COVID-19 diagnosed in KTR that require hospitalization and occurred before January 1st 2021.

The characteristics of enrolled patients, which were prospectively collected in the French SOT COVID registry, are presented in **Table 1**. Briefly, a little less than 10% of the cohort received a graft from a living donor. The median recipient age was 63.0 [52.0-70.0] years and males represented 68.1% of the cohort. The majority of patients (537/864, 62.1%) were overweight and the median BMI of the cohort was 26.0 [23.0-29.4] kg/m². The most common comorbidity was hypertension (798/918, 86.9%), followed by diabetes (371/914, 40.6%) and cardiovascular disease (352/908, 38.8%). The median baseline estimated glomerular filtration rate (eGFR) was 41.0 [30.0-54.0] mL/min/1.73m². Regarding therapeutic immunosuppression, the vast majority of patients received an induction therapy, either with anti-interleukin-2 (385/931, 41.4%) or with

antithymocyte globulin (508/931, 54.6%). At diagnosis of COVID-19, maintenance regimen of most patients consisted in a combination of calcineurin inhibitor (807/957, 84%, either tacrolimus 65.3% or cyclosporine 19%), an antimetabolite (722/957, 75.4% on mycophenolic acid) and corticosteroids (726/957, 75.9%). Only 4.0% of the cohort were on belatacept.

The patients of the two pandemic waves were largely similar except for diabetes, the prevalence of which was slightly lower in patients of the second wave (37.3% versus 45.7% respectively; $p=0.014$). Difference in immunosuppression regimen were also minor with only less patients on corticosteroids (72.7% vs 81.1%, $p=0.005$) and mTOR inhibitors (8.9% vs 13.1%, $p=0.050$) in the second pandemic wave.

Clinical and biological presentation of COVID-19 at admission

Almost all diagnoses of COVID-19 (919/957, 96%) were confirmed by reverse transcriptase-polymerase chain reaction. SARS-CoV-2 infection occurred after a median of 67.6 [28.2-134.2] months after kidney transplantation. Of note, despite the fact that KT activity in France was interrupted during the first wave but maintained during the 2nd there was no difference in the median delay from transplantation to COVID-19 diagnosis between the two pandemic waves (71.1 [31.0-144.5] vs 65.6 [27.3-129.9] months, $p=0.215$).

Considering the whole cohort (**Figure 1A**), the most frequent symptom on admission was fever (585/957, 67.2%), followed by cough (494/957, 56.8%), dyspnea (466/957, 52.3%) and diarrhea (317/957, 36.2%). Median levels of C-reactive protein and procalcitonin were 67 [28-121] mg/L and 0.22 [0.12-0.70] ng/mL respectively. At admission, most (580/653, 89%) patients had low lymphocyte count (median lymphocyte

count of the cohort $0.65 \cdot 10^9$ [0.40-1.00]/L) and median creatinemia was 174 [129-256] $\mu\text{mol/L}$.

KTR from the second wave differed from those of the first in that they less frequently exhibited fever, cough and myalgias, which could indicate earlier diagnosis during the second wave (**Figure 1B**). This hypothesis is coherent with the increased availability of diagnosis assays during the second half of 2020. However, no significant differences in CRP and PCT levels, nor in lymphocyte count could be observed between the 2 pandemic waves (data not shown). Furthermore, chest CT scan severity at presentation was also similar between the two waves with ~45%, 30% and 25% of KTR presented with mild, moderate and severe degree of involvement respectively (**Figure 1C**; $p=0.921$).

Management of immunosuppression and anti-viral therapies

The exact impact of maintenance immunosuppression during COVID-19 is unclear²³. On one hand these drugs could be protective against the overproduction of proinflammatory cytokines during critical COVID-19^{24,25}, on the other solid organ transplant recipients have been found to have delayed SARS-CoV-2 clearance^{26,27}.

Maintenance immunosuppression was tapered in KTR hospitalized for symptomatic COVID-19, particularly antimetabolites and mTOR inhibitors, which were discontinued in the majority of patients of both pandemic waves (**Figure 2A**). However, if modifications of maintenance immunosuppression did not differ in nature between the two waves, they were made in a smaller proportion of patients during the second wave, particularly regarding withdrawal of CNI (32.1% vs 16.6%, $p<0.001$) and of antimetabolites (73.7% vs 58.4%, $p<0.001$; **Figure 2A**), which is in line with a previous report from the USA²⁸.

Contrasting with the global stability of immunosuppression management, anti-SARS-CoV-2 therapies differed in many respects between the two waves (**Figure 2B**). KTR with COVID-19 from the second wave received less frequently probabilistic antibiotics compared to those of the first wave (75.8% vs 49.2%, $p<0.001$). Hydroxychloroquine and azithromycin, which were commonly used during the first wave were almost completely abandoned during the second (21.7% vs 1.7% and 30.9% vs 5.0%, $p<0.001$, respectively). Tocilizumab use declined between the first and second waves (7.5% vs 2.2%, $p<0.001$). Conversely, the use of high dose corticosteroids doubled (19.5% vs. 41.6%, $p<0.001$).

Risk factors associated with death due to COVID-19 in KTR

Univariate analysis conducted on the whole cohort identified: age, hypertension, preexisting cardiovascular disease, history of cancer, diabetes, dyspnea at admission, CRP > 60 mg/L at admission, baseline eGFR as significantly associated with mortality (data not shown). In contrast, diarrhea, anosmia and headaches were associated with reduced risk of death.

In multivariable analysis, only age>50 years, history of cancer, dyspnea or CRP > 60 mg/L at admission, and baseline eGFR<30ml/min/m² remained independently associated with a higher risk of death in KTR hospitalized for COVID-19 (**Figure 3**), while anosmia at admission was associated with a better prognosis (**Figure 3**).

Comparison of first vs second wave outcomes

The incidence of thromboembolic events (9.5% vs 6.4%, $p=0.135$) and bacterial superinfection (27.0% vs 30.7%, $p=0.304$) was similar between the 2 pandemic waves. A non-significant trend for lesser use of RRT (15.9% vs 12.9%, $p=0.230$), mechanical ventilation (26.5% vs 22.1%, $p=0.152$) and vasopressor support (20.5% vs 15.9%,

p=0.304) was observed during the 2nd wave but mortality at 60 days from admission (24.5%) was in the range of what previously reported^{29,30}, with no significant difference between the first and second wave (**Figure 4A**; Log rank test, p=0.48).

A slight difference in dynamic between the two waves could however be observed on Kaplan Meier curves (**Figure 4A**), with shorter delay between admission and death due to COVID-19 in KTR of the first wave (death at 14 days: 16.7% vs 10.1%, p=0.002). This difference is to be interpreted together with a faster and higher incidence of transfer in ICU for patients of the first wave (**Figure 4B**), without difference on the mortality for patients transferred in ICU (**Figure 4C**). Altogether, these findings could indicate that patients of the first wave were diagnosed (and therefore hospitalized) later in the course of COVID-19, a hypothesis in line with the difference in clinical presentation between the 2 waves reported above (**Figure 1B**).

In contrast with the second wave that impacted the entire French territory, the first pandemic wave had a heterogeneous geographic distribution³¹ that could have introduce a "learning-curve" bias. Physicians from the geographic area impacted by the first wave could have accumulated knowledge and skills useful to better manage patients from the second wave. To test this hypothesis, we compared the survival of KTR hospitalized for COVID-19 during the second wave in geographic area impacted (in red on the map **Figure 4D**) vs preserved (in green on the map **Figure 4D**) during the first pandemic wave. The similarity in survival for patients of the second wave hospitalized in either of these two areas strongly argue against the theory of the learning curve bias (**Figure 4E**).

Discussion

Kidney transplant recipients (KTR), who are characterized by a highly comorbid profile and receive therapeutic immunosuppression to prevent graft rejection, were very early identified as particularly vulnerable to COVID-19¹⁵⁻¹⁷. An excess of mortality, integrally explained by COVID-19, was indeed reported in this population during the first wave of the pandemic in France³¹ and several large multicenter cohorts KTR estimated short-term intra-hospital mortality around 20-32%^{29,32,33}. Among the risk factors identified in previous publications for death due to COVID-19 in KTR are age, eGFR and presence of comorbidities, including cardiovascular disease, diabetes, and/or obesity^{31,32,34,35}. Additionally, dyspnea and elevations of biochemical markers of inflammation at diagnosis of COVID-19 were also associated with less favorable survival figures³⁶⁻³⁸. Our study largely confirms these data. In addition, it provides original additional information regarding the stability of the risk of death due to COVID-19 in KTR, despite the impressive accumulation of knowledge regarding the disease, which translated in better outcomes in the general population^{13,14,39}. Indeed, despite a more homogeneous COVID-19 management with wider prescription of Dexamethasone and important decrease in the use of treatments deemed inefficient such as azithromycin⁴⁰, hydroxychloroquine^{40,41}, lopinavir/ritonavir⁴², survival of hospitalized KTR during the second wave remained similar to that observed during the first wave. Our findings concur with a meta-analysis including 5559 KTR with COVID-19 that reported a mean mortality rate of 23% (similar to what we observed) without significant difference between “early” (studies submitted before July 2020) and “late” (studies submitted from July 2020 onwards) phases of the pandemic⁴³. These conclusions conflict with a recent

study showing a better prognosis in “late” (from June 20 to December 31, 2020) compared to “early” 2020 (from March 1 to June 19, 2020) among 973 solid organ transplant recipients (SOTR) hospitalized in USA for COVID-19²⁸. In their report, crude mortality by 28 days indeed declined from 19.6% in early period to 13.7% in late period and after adjusting for differences in baseline comorbidities between both periods, the odds of death remained lower in the late period (aOR 0.67, 95% CI 0.46 – 0.98, $p = 0.04$). Instead of the changing trends in management of COVID-19 patients, we believe that the observations made by Heldman et al, could be explained by the numerous differences in the baseline comorbid profiles of SOTR between the early and late period (SOTR in late period presented with less hypertension, diabetes, heart failure, coronary artery disease, chronic lung disease) and/or by the short follow-up period of the study. Indeed, when we assessed 14-days mortality in our own cohort, we also found a significant difference between the first and second wave (16.7% vs. 10.1%, p -value = 0.002, respectively) that progressively disappeared by the end of the 60-days follow-up period. Whether this effect is attributable to earlier diagnosis of COVID-19 in KTR during the second wave is possible and supported by some clues discussed above but remains to be formally demonstrated.

Among the strengths of our study are the relative high number of patients enrolled and the prospective collection of data. Our study has however also some limitations. First, we compared two periods (first and second wave) but did not take into account COVID-19 ICU occupancy rates, a factor thought to impact on mortality rates¹³. Second, our study was not designed to capture the impact of vaccines, which only became available early 2021.

Accumulating evidence suggests however that KTR have an impaired response to the “standard” 2-dose of mRNA vaccine^{44–47}, which leaves them at high risk of severe COVID-19^{46,48}. Despite intensified scheme of vaccination (with third and even a fourth vaccine dose now recommended in weak responders), up to 20% of KTR will not develop sufficient protection against COVID-19^{49–51}. In this regard, the development of monoclonal neutralizing anti-SARS-CoV-2 Spike Protein Antibodies represent an interesting therapeutic option. The latter are already available in high-risk patients diagnosed with mild to moderate COVID-19⁵² (post-exposition therapy) and first reports about their use for prophylaxis (pre-exposition therapy) are promising⁵³. In conclusion, changing of therapeutic trends during 2020 did not reduce COVID-19 related mortality in KTR. Our data thus indirectly stress the importance of therapeutic progresses made during 2021, including vaccination and monoclonal neutralizing anti-SARS-CoV-2 spike protein antibodies, to protect this vulnerable population from death due to COVID-19.

Disclosure

All the authors declared no competing interests.

References

1. Hu B, Guo H, Zhou P, et al. Characteristics of SARS-CoV-2 and COVID-19. *Nat Rev Microbiol.* 2021;19(3):141-154. doi:10.1038/s41579-020-00459-7
2. Coronavirus Disease (COVID-19) Situation Reports. Accessed October 14, 2021. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>
3. Vahidy FS, Drews AL, Masud FN, et al. Characteristics and Outcomes of COVID-19 Patients During Initial Peak and Resurgence in the Houston Metropolitan Area. *JAMA.* 2020;324(10):998. doi:10.1001/jama.2020.15301
4. Soriano V, Ganado-Pinilla P, Sanchez-Santos M, et al. Main differences between the first and second waves of COVID-19 in Madrid, Spain. *Int J Infect Dis.* 2021;105:374-376. doi:10.1016/j.ijid.2021.02.115
5. Zhu N, Zhang D, Wang W, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med.* 2020;382(8):727-733. doi:10.1056/NEJMoa2001017
6. Vandenberg O, Martiny D, Rochas O, et al. Considerations for diagnostic COVID-19 tests. *Nat Rev Microbiol.* 2021;19(3):171-183. doi:10.1038/s41579-020-00461-z
7. COVID19-APHP Group. Electronic address: martin.hirsch@aphp.fr. Assistance Publique-Hôpitaux de Paris' response to the COVID-19 pandemic. *Lancet.* 2020;395(10239):1760-1761. doi:10.1016/S0140-6736(20)31210-1
8. Roques L, Klein EK, Papaix J, et al. Impact of Lockdown on the Epidemic Dynamics of COVID-19 in France. *Front Med (Lausanne).* 2020;7:274. doi:10.3389/fmed.2020.00274
9. Torres Acosta MA, Singer BD, et al. Pathogenesis of COVID-19-induced ARDS: implications for an ageing population. *Eur Respir J.* 2020;56(3):2002049. doi:10.1183/13993003.02049-2020
10. Bikdeli B, Madhavan MV, Jimenez D, et al. COVID-19 and Thrombotic or Thromboembolic Disease: Implications for Prevention, Antithrombotic Therapy, and Follow-Up: JACC State-of-the-Art Review. *J Am Coll Cardiol.* 2020;75(23):2950-2973. doi:10.1016/j.jacc.2020.04.031
11. Gupta S, Coca SG, Chan L, et al. AKI Treated with Renal Replacement Therapy in Critically Ill Patients with COVID-19. *J Am Soc Nephrol.* 2021;32(1):161-176. doi:10.1681/ASN.2020060897
12. RECOVERY Collaborative Group, Horby P, Lim WS, et al. Dexamethasone in Hospitalized Patients with Covid-19. *N Engl J Med.* 2021;384(8):693-704. doi:10.1056/NEJMoa2021436

13. Lefrancq N, Paireau J, Hozé N, et al. Evolution of outcomes for patients hospitalised during the first 9 months of the SARS-CoV-2 pandemic in France: A retrospective national surveillance data analysis. *Lancet Reg Health Eur.* 2021;5:100087. doi:10.1016/j.lanepe.2021.100087
14. Covid-19: facteurs de risques hospitalisation décès - 2ème vague. EPI-PHARE. Published July 23, 2021. Accessed October 14, 2021. <https://www.epi-phare.fr/actualites/communiqué-presse-20210723/>
15. Alberici F, Delbarba E, Manenti C, et al. A single center observational study of the clinical characteristics and short-term outcome of 20 kidney transplant patients admitted for SARS-CoV2 pneumonia. *Kidney Int.* 2020;97(6):1083-1088. doi:10.1016/j.kint.2020.04.002
16. Akalin E, Azzi Y, Bartash R, et al. Covid-19 and Kidney Transplantation. *New England Journal of Medicine.* Published online April 24, 2020. doi:10.1056/NEJMc2011117
17. Williamson EJ, Walker AJ, Bhaskaran K, et al. Factors associated with COVID-19-related death using OpenSAFELY. *Nature.* 2020;584(7821):430-436. doi:10.1038/s41586-020-2521-4
18. Buuren S van, Groothuis-Oudshoorn K. mice: Multivariate Imputation by Chained Equations in R. *Journal of Statistical Software.* 2011;45:1-67. doi:10.18637/jss.v045.i03
19. Multiple Imputation for Nonresponse in Surveys | Wiley. Wiley.com. Accessed November 27, 2021. <https://www.wiley.com/en-us/Multiple+Imputation+for+Nonresponse+in+Surveys-p-9780471655749>
20. R: The R Project for Statistical Computing. Accessed November 27, 2021. <https://www.r-project.org/>
21. Gitman MR, Shaban MV, Paniz-Mondolfi AE, et al. Laboratory Diagnosis of SARS-CoV-2 Pneumonia. *Diagnostics (Basel).* 2021;11(7):1270. doi:10.3390/diagnostics11071270
22. HCSP. *Signes cliniques d'orientation diagnostique du Covid-19.* Haut Conseil de la Santé Publique; 2020. Accessed November 20, 2021. <https://www.hcsp.fr/explore.cgi/avisrapportsdomaine?clefr=812>
23. Caillard S, Chavarot N, Francois H, et al. Is COVID-19 infection more severe in kidney transplant recipients? *Am J Transplant.* 2021;21(3):1295-1303. doi:10.1111/ajt.16424
24. Chavarot N, Gueguen J, Bonnet G, et al. COVID-19 severity in kidney transplant recipients is similar to nontransplant patients with similar comorbidities. *Am J Transplant.* 2021;21(3):1285-1294. doi:10.1111/ajt.16416

25. Dęborska-Materkowska D, Kamińska D, et al. The Immunology of SARS-CoV-2 Infection and Vaccines in Solid Organ Transplant Recipients. *Viruses*. 2021;13(9):1879. doi:10.3390/v13091879
26. Epstein RL, Sperring H, Hofman M, et al. Time to SARS-CoV-2 PCR Clearance in Immunocompromising Conditions: Is Test-Based Removal From Isolation Necessary in Severely Immunocompromised Individuals? *Open Forum Infect Dis*. 2021;8(6):ofab164. doi:10.1093/ofid/ofab164
27. Benotmane I, Risch S, Doderer-Lang C, et al. Long-term shedding of viable SARS-CoV-2 in kidney transplant recipients with COVID-19. *Am J Transplant*. 2021;21(8):2871-2875. doi:10.1111/ajt.16636
28. Heldman MR, Kates OS, Safa K, et al. Changing trends in mortality among solid organ transplant recipients hospitalized for COVID-19 during the course of the pandemic. *Am J Transplant*. Published online September 12, 2021. doi:10.1111/ajt.16840
29. Cravedi P, Mothi SS, Azzi Y, et al. COVID-19 and kidney transplantation: Results from the TANGO International Transplant Consortium. *Am J Transplant*. 2020;20(11):3140-3148. doi:10.1111/ajt.16185
30. Coll E, Fernández-Ruiz M, Sánchez-Álvarez JE, et al. COVID-19 in transplant recipients: The Spanish experience. *Am J Transplant*. 2021;21(5):1825-1837. doi:10.1111/ajt.16369
31. Thauinat O, Legeai C, Anglicheau D, et al. IMPact of the COVID-19 epidemic on the moRTality of kidney transplant recipients and candidates in a French Nationwide registry sTudy (IMPORTANT). *Kidney Int*. 2020;98(6):1568-1577. doi:10.1016/j.kint.2020.10.008
32. Caillard S, Anglicheau D, Matignon M, et al. An initial report from the French SOT COVID Registry suggests high mortality due to COVID-19 in recipients of kidney transplants. *Kidney Int*. 2020;98(6):1549-1558. doi:10.1016/j.kint.2020.08.005
33. Jager KJ, Kramer A, Chesnaye NC, et al. Results from the ERA-EDTA Registry indicate a high mortality due to COVID-19 in dialysis patients and kidney transplant recipients across Europe. *Kidney Int*. 2020;98(6):1540-1548. doi:10.1016/j.kint.2020.09.006
34. Udomkarnjananun S, Kerr SJ, Townamchai N, et al. Mortality risk factors of COVID-19 infection in kidney transplantation recipients: a systematic review and meta-analysis of cohorts and clinical registries. *Sci Rep*. 2021;11(1):20073. doi:10.1038/s41598-021-99713-y
35. Kates OS, Haydel BM, Florman SS, et al. COVID-19 in solid organ transplant: A multi-center cohort study. *Clin Infect Dis*. Published online August 7, 2020:ciaa1097. doi:10.1093/cid/ciaa1097

36. Caillard S, Chavarot N, Francois H, et al. Clinical Utility of Biochemical Markers for the Prediction of COVID-19–Related Mortality in Kidney Transplant Recipients. *Kidney International Reports*. 2021;6(10):2689-2693. doi:10.1016/j.ekir.2021.06.034
37. Requião-Moura LR, Sandes-Freitas TV de, Viana LA, et al. High mortality among kidney transplant recipients diagnosed with coronavirus disease 2019: Results from the Brazilian multicenter cohort study. *PLoS One*. 2021;16(7):e0254822. doi:10.1371/journal.pone.0254822
38. Benotmane I, Perrin P, Vargas GG, et al. Biomarkers of Cytokine Release Syndrome Predict Disease Severity and Mortality From COVID-19 in Kidney Transplant Recipients. *Transplantation*. 2021;105(1):158-169. doi:10.1097/TP.0000000000003480
39. Ioannou GN, O'Hare AM, Berry K, et al. Trends Over Time in the Risk of Adverse Outcomes Among Patients With Severe Acute Respiratory Syndrome Coronavirus 2 Infection. *Clinical Infectious Diseases*. 2021;(ciab419). doi:10.1093/cid/ciab419
40. Cavalcanti AB, Zampieri FG, Rosa RG, et al. Hydroxychloroquine with or without Azithromycin in Mild-to-Moderate Covid-19. *N Engl J Med*. 2020;383(21):2041-2052. doi:10.1056/NEJMoa2019014
41. Martins-Filho PR, Ferreira LC, Heimfarth L, et al. Efficacy and safety of hydroxychloroquine as pre-and post-exposure prophylaxis and treatment of COVID-19: A systematic review and meta-analysis of blinded, placebo-controlled, randomized clinical trials. *Lancet Reg Health Am*. 2021;2:100062. doi:10.1016/j.lana.2021.100062
42. RECOVERY Collaborative Group. Lopinavir-ritonavir in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. *Lancet*. 2020;396(10259):1345-1352. doi:10.1016/S0140-6736(20)32013-4
43. Kremer D, Pieters TT, Verhaar MC, et al. A systematic review and meta-analysis of COVID-19 in kidney transplant recipients: Lessons to be learned. *Am J Transplant*. Published online July 1, 2021. doi:10.1111/ajt.16742
44. Boyarsky BJ, Werbel WA, Avery RK, et al. Antibody Response to 2-Dose SARS-CoV-2 mRNA Vaccine Series in Solid Organ Transplant Recipients. *JAMA*. 2021;325(21):2204-2206. doi:10.1001/jama.2021.7489
45. Benotmane I, Gautier-Vargas G, Cognard N, et al. Low immunization rates among kidney transplant recipients who received 2 doses of the mRNA-1273 SARS-CoV-2 vaccine. *Kidney International*. 2021;99(6):1498-1500. doi:10.1016/j.kint.2021.04.005
46. Charmetant X, Espi M, Benotmane I, et al. *Comparison of Infected and Vaccinated Transplant Recipients Highlights the Role of Tfh and Neutralizing IgG in*

COVID-19 Protection.; 2021:2021.07.22.21260852.
doi:10.1101/2021.07.22.21260852

47. Caillard S, Thaunat O, et al. COVID-19 vaccination in kidney transplant recipients. *Nat Rev Nephrol.* 2021;17(12):785-787. doi:10.1038/s41581-021-00491-7
48. Caillard S, Chavarot N, Bertrand D, et al. Occurrence of severe COVID-19 in vaccinated transplant patients. *Kidney Int.* 2021;100(2):477-479. doi:10.1016/j.kint.2021.05.011
49. Benotmane I, Gautier G, Perrin P, et al. Antibody Response After a Third Dose of the mRNA-1273 SARS-CoV-2 Vaccine in Kidney Transplant Recipients With Minimal Serologic Response to 2 Doses. *JAMA.* 2021;326(11):1063-1065. doi:10.1001/jama.2021.12339
50. Werbel WA, Boyarsky BJ, Ou MT, et al. Safety and Immunogenicity of a Third Dose of SARS-CoV-2 Vaccine in Solid Organ Transplant Recipients: A Case Series. *Ann Intern Med.* 2021;174(9):1330-1332. doi:10.7326/L21-0282
51. Caillard S, Thaunat O, Benotmane I, on behalf of the French-speaking Society of Transplantation. Antibody response to a fourth mRNA Covid-19 vaccine boost in weak responder kidney transplant recipients. *medRxiv.* Published online January 1, 2021:2021.09.03.21262691. doi:10.1101/2021.09.03.21262691
52. Dougan M, Nirula A, Azizad M, et al. Bamlanivimab plus Etesevimab in Mild or Moderate Covid-19. *N Engl J Med.* 2021;385(15):1382-1392. doi:10.1056/NEJMoa2102685
53. Subcutaneous REGEN-COV Antibody Combination to Prevent Covid-19 | NEJM. Accessed November 21, 2021. https://www.nejm.org/doi/10.1056/NEJMoa2109682?url_ver=Z39.88-2003&rfr_id=ori:rid:crossref.org&rfr_dat=cr_pub%20%20pubmed

Acknowledgements

OT is supported by the Etablissement Français du Sang and the Fondation pour la Recherche Médicale (PME20180639518).

TABLE 1 | Baseline characteristics of kidney transplant patients at admission for COVID-19

Variables median [IQR] or n (%)	All cohort (n=957)	Missing data	1 st Wave (n=359)	2 nd Wave (n=598)	p
Clinical characteristics					
Age (yr)	63.0 [52.0-70.0]	0 (0.0%)	63.0 [54.0-70.0]	62.0 [51.2-70.0]	0.298
Male	652 (68.1%)	0 (0.0%)	243 (67.7%)	409 (68.4%)	0.876
BMI (kg/m ²)	26.0 [23.0-29.4]	93 (9.7%)	26.0 [23.0-29.0]	26.0 [23.2-29.6]	0.564
Blood group		30 (3.1%)			0.472
A	395 (42.6%)		144 (40.4%)	251 (44.0%)	
AB	59 (6.4%)		21 (5.9%)	38 (6.7%)	
B	107 (11.5%)		39 (11.0%)	68 (11.9%)	
O	366 (39.5%)		152 (42.7%)	214 (37.5%)	
Retransplantation	104 (11.5%)	50 (5.2%)	45 (12.6%)	59 (10.7%)	0.462
Living donor	90 (9.5%)	13 (1.3%)	27 (7.5%)	63 (10.8%)	0.125
Delay Tx-COVID (mo)	67.6 [28.2-134.2]	0 (0.0%)	71.1 [31.0-144.5]	65.6 [27.3-129.9]	0.215
Hypertension	798 (86.9%)	39 (4.1%)	320 (89.4%)	478 (85.4%)	0.096
CV disease	352 (38.8%)	49 (5.1%)	148 (41.2%)	204 (37.2%)	0.246
Respiratory disease	122 (13.4%)	45 (4.7%)	43 (12.0%)	79 (14.3%)	0.368
Diabetes	371 (40.6%)	43 (4.5%)	164 (45.7%)	207 (37.3%)	0.014
Cancer	144 (15.8%)	47 (4.9%)	63 (17.5%)	81 (14.7%)	0.290
Smoking	126 (15.0%)	115 (12.0%)	40 (12.1%)	86 (16.8%)	0.079
Statin	307 (46.2%)	292 (30.5%)	154 (49.7%)	153 (43.1%)	0.105
RAS blockers	371 (44.8%)	129 (13.5%)	155 (48.1%)	216 (42.7%)	0.143
eGFR (ml/min/1.73m ²)	41.0 [30.0-54.0]	36 (3.8%)	40.0 [29.0-55.0]	42.0 [30.0-54.0]	0.336
Immunosuppression					
Induction		26 (2.7%)			0.140
No induction	38 (4.1%)		10 (2.9%)	28 (4.8%)	
anti-IL2R	385 (41.4%)		137 (39.1%)	248 (42.7%)	
ATG	508 (54.6%)		203 (58.0%)	305 (52.5%)	
Maintenance					
CNI		0 (0.0%)			0.234
No CNI	150 (15.7%)		47 (13.1%)	103 (17.2%)	
Tacrolimus	625 (65.3%)		242 (67.4%)	383 (64.0%)	
Cyclosporine	182 (19.0%)		70 (19.5%)	112 (18.7%)	
Mycophenolate	722 (75.4%)	0 (0.0%)	278 (77.4%)	444 (74.2%)	0.302
Azathioprin	32 (3.3%)	0 (0.0%)	12 (3.3%)	20 (3.3%)	1.000
mTOR inhibitor	100 (10.4%)	0 (0.0%)	47 (13.1%)	53 (8.9%)	0.050
Steroids	726 (75.9%)	0 (0.0%)	291 (81.1%)	435 (72.7%)	0.005
Belatacept	38 (4.0%)	0 (0.0%)	20 (5.6%)	18 (3.0%)	0.073

The p values are for the comparisons of 1st wave 1 vs. 2nd wave. Bold indicates $p < 0.05$.

Abbreviations are: yr, year; BMI, body mass index; Tx, transplantation; mo, months; CV, cardiovascular; RAS, renin-angiotensin-system; eGFR, estimated glomerular filtration rate; anti-IL2R, anti-interleukin-2 receptor; ATG, anti-thymocyte globulin; CNI, calcineurin inhibitor; mTor, mechanistic target of rapamycin. Baseline eGFR is determined with the MDRD equation.

Figure legends

FIGURE 1 | Clinical and biological presentation of COVID-19 at admission

A Summary of the main clinical and biological characteristics of the entire cohort (n=957 KTR), Median [IQR] or n (%), at hospital admission for COVID-19. **B** Comparison of characteristics at hospital admission for COVID-19 of patients from the 1st vs 2nd pandemic wave. **C** Comparison of chest CT scan severity between the 1st vs 2nd pandemic wave.

Abbreviations are: CRP, C-reactive protein; PCT, procalcitonin; SCr, serum creatinine. χ^2 test; p>0.05, ns.

FIGURE 2 | Changing of therapeutic trends between the 1st and 2nd pandemic COVID-19 wave

Comparison of the management of immunosuppression (**A**) and the use of COVID-19 specific treatments (**B**) between the 1st (blue) vs 2nd (red) pandemic wave.

Abbreviations are: CNI, calcineurin inhibitor; mTor, mechanistic target of rapamycin; ATB, antibiotics. χ^2 test; p>0.05, ns.

FIGURE 3 | Variables associated with the risk of death due to COVID-19 in KTR

This forest plot shows the variable independently associated with the risk of death in multivariate analysis for the 957 KTR diagnosed with COVID-19 during the 1st or the 2nd pandemic wave.

FIGURE 4 | Comparison of COVID-19 outcomes between the 1st and 2nd wave

A In-hospital survival of KTR diagnosed with COVID-19 during the 1st and 2nd wave.

B Cumulative incidence of Intensive Care Unit (ICU) admission of KTR diagnosed with COVID-19 during the 1st and 2nd wave.

C Survival of KTR diagnosed with COVID-19 transferred in ICU.

D Map of the geographic distribution of the cases of COVID-19 in France during the 1st wave. Area in which the incidence of COVID-19 was the highest are in red.


E In-hospital survival of KTR diagnosed with COVID-19 during the 2nd wave according to their geographic location (in the red or green area defined in panel D).

Comparison were made using the Log Rank test.


Appendix

The French SOT COVID Registry Collaborators are as follows: Sophie Caillard, Bruno Moulin, Service de Néphrologie et Transplantation, Hôpitaux Universitaires de Strasbourg, Strasbourg; Samira Fafi-Kremer, Laboratoire de Virologie, Hôpitaux Universitaires de Strasbourg, Strasbourg; Marc Hazzan, Service de Néphrologie, Hôpital Huriez, Lille; Dany Anglicheau, Service de Néphrologie et Transplantation Adultes, AP-HP, Hôpital Necker, Paris; Alexandre Hertig, Jérôme Turret, Benoit Barrou, Service de Néphrologie, AP-HP, Hôpital La Pitié Salpêtrière, Paris; Emmanuel Morelon, Olivier Thauinat, Service de Néphrologie, Hôpital Edouard Herriot, Lyon; Lionel Couzi, Pierre Merville, Service de Néphrologie–Transplantation–Dialyse, Hôpital Pellegrin, Bordeaux; Valérie Moal, Tristan Legris, Service de Néphrologie et Transplantation, AP-HM, Hôpital de la Conception, Marseille; Pierre-François Westeel, Maïté Jaureguy, Service de Néphrologie, CHU Amiens Picardie, Amiens; Luc Frimat, Service de Néphrologie, CHRU Nancy, Vandoeuvre; Didier Ducloux, Jamal Bamoulid, Service de Néphrologie, Hôpital Jean-Minjoz, Besancon; Dominique Bertrand, Service de Néphrologie, CHU de Rouen, Rouen; Michel Tsimaratos, Florentine Garaix-Gilardo, Service de Pédiatrie Multidisciplinaire, Hôpital La Timone, Marseille; Jérôme Dumortier, Service d'Hépatogastroentérologie, Hôpital Edouard Herriot, Lyon; Sacha Mussot, Antoine Roux, Centre Chirurgical Marie Lannelongue, Le Plessis Robinson; Laurent Sebbag, Service d'Insuffisance Cardiaque, Hôpital Louis Pradel, Bron; Yannick Le Meur, Service de Néphrologie, Hôpital de la Cavale Blanche, Brest; Gilles Blancho, Christophe Masset, Service de Néphrologie–Transplantation, Hôtel Dieu, Nantes; Nassim Kamar, Service de Néphrologie et Transplantation, Hôpital Rangueil, Toulouse; Hélène Francois, Eric Rondeau, Service de Néphrologie, Dialyse et Transplantation, AP-HP, Hôpital Tenon, Paris; Nicolas Bouvier, Service de Néphrologie, Dialyse, Transplantation Rénale, CHU, Caen; Christiane Mousson, Service de Néphrologie, Dijon; Matthias Buchler, Philippe Gatault, Service de Néphrologie, Tours; Jean-François Augusto, Agnès Dureau, Service de Néphrologie, Dialyse, Transplantation, CHU Angers, Angers; Cécile Vigneau, Marie-Christine Morin, Jonathan Chemouny, Leonard Golbin, Service de Néphrologie, CHU de Rennes, Rennes; Philippe Grimbert, Marie Matignon, Antoine Durrbach, Service de Néphrologie, Hôpital Henri-Mondor, Creteil; Clarisse Greze, Service de Néphrologie, AP-HP, Hôpital Bichat Claude Bernard, Paris; Renaud Snanoudj, Service de Néphrologie, Hôpital Foch, Service de Néphrologie et Transplantation Hôpital du Kremlin Bicêtre, Le Kremlin Bicêtre; Charlotte Colosio, Betoul Schwartz, Service de Néphrologie, Hôpital Maison Blanche, Reims; Paolo Malvezzi, Service de Néphrologie, Hémodialyse, Transplantation Rénale, Hôpital La Tronche, Grenoble; Christophe Mariat, Service de Néphrologie, CHU de Saint Etienne, Saint Etienne; Antoine Thierry, Service de Néphrologie, Hémodialyse et Transplantation Rénale, Hôpital Jean Bernard, Poitiers; Moglie Le Quintrec, Service de Néphrologie–Transplantation–Dialyse, CHU Lapeyronie, Montpellier; Antoine Sicard, Service de Néphrologie, Hôpital Pasteur, Nice; Jean Philippe Rerolle, Service de Néphrologie, CHU Dupuytren, Limoges; Anne-Élisabeth Heng, Cyril Garrouste, Service de Néphrologie, CHU Gabriel Montpied, Clermont-Ferrand; Henri Vacher Coponat, Service de Néphrologie, CHU de La Réunion, Saint Denis; Éric Epailly, Service de Cardiologie, Hôpitaux Universitaires de Strasbourg, Strasbourg; Olivier Brugiere, Service d'Hépatologie, Hôpital Foch, Suresnes; Sébastien Dharancy, Service d'Hépatologie, Hôpital Huriez, Lille; Éphrem Salame, Service de Chirurgie Hépatique, Hôpital Universitaire de Tours, Tours; Faouzi Saliba, Service d'Hépatologie, Centre hépato-biliaire Paul Brousse, Villejuif, France.


A




B



C



A



B

