

## Evaluation of the Implementation of the 4C Mortality Score in United Kingdom hospitals during the second pandemic wave

Andrew E. Blunsum<sup>1\*</sup>, Jonathan S. Perkins<sup>1, 2\*</sup>, Areeb Arshad<sup>3</sup>, Sukrit Bajpai<sup>4</sup>, Karen Barclay-Elliott<sup>5</sup>, Sanjita Brito-Mutunayagam<sup>6, 7</sup>, Rebecca Brooks<sup>8</sup>, Terrence Chan<sup>9</sup>, Dominic J. G. Coates<sup>10</sup>, Alina Corobana<sup>11</sup>, Tim Crocker-Buqué<sup>12</sup>, Terry J. Evans<sup>11</sup>, Jasmine Gordon-Brown<sup>13</sup>, Berkin Hack<sup>14</sup>, Heather Hiles<sup>15</sup>, Aakash Khanijau<sup>16</sup>, Salina Lalwani<sup>6, 17</sup>, Clare Leong<sup>8</sup>, Kirsty MacKay<sup>18, 19, 20, 21</sup>, Catriona Macrae<sup>22</sup>, Bryony Martin<sup>23</sup>, Christopher A. Martin<sup>24, 25, 27</sup>, Emily McKemey<sup>4</sup>, Joshua Nazareth<sup>24, 25, 27</sup>, Daniel Pan<sup>25, 26, 27</sup>, Marcello Scopazzini<sup>28</sup>, David Simons<sup>29</sup>, Sophie Swinhoe<sup>16</sup>, Julia Thomas<sup>30, 31</sup>, Fiona Thorburn<sup>1</sup>, Sarah Walpole<sup>32</sup>, Esmie Warne<sup>33</sup>, Rory Wilson<sup>34</sup>, Alisdair MacConnachie<sup>1</sup>, Antonia Ho<sup>1, 35</sup>

\*Joint first authors

### Affiliations

1. Queen Elizabeth University Hospital, Glasgow
2. Glasgow Royal Infirmary, Glasgow
3. Forth Valley Royal Hospital, Larbert
4. Queen Elizabeth Hospital, Birmingham
5. University Hospital Coventry, Coventry
6. St. John's Hospital, Livingston
7. Western General Hospital, Edinburgh
8. Addenbrooke's Hospital, Cambridge
9. Kingston Hospital, Kingston upon Thames
10. University Hospital Wishaw, Wishaw
11. St. George's Hospital, London
12. Royal Free Hospital, London
13. University Hospital Hairmyres, East Kilbride
14. Newham University Hospital, London
15. Royal Hallamshire Hospital, Sheffield
16. Warrington Hospital, Warrington
17. Edinburgh Royal Infirmary, Edinburgh
18. Lady Margaret Hospital, Millport
19. University Hospital Crosshouse, Kilmarnock
20. Arran War Memorial Hospital, Lamlash
21. University Hospital Ayr, Ayr
22. University Hospital Monklands, Airdrie
23. Royal Alexandra Hospital, Paisley
24. Leicester General Hospital, Leicester
25. Glenfield Hospital, Leicester
26. Leicester Royal Infirmary, Leicester
27. Department of Respiratory Sciences, University of Leicester
28. Victoria Hospital, Kirkcaldy
29. University College Hospital, London
30. Dumfries and Galloway Royal Infirmary, Dumfries
31. Galloway Community Hospital, Stranraer
32. Royal Victoria Infirmary, Newcastle
33. John Radcliffe Hospital, Oxford

34. Ninewells Hospital, Dundee

35. MRC-University of Glasgow Centre for Virus Research, Glasgow

**Correspondence to:**

Dr. Antonia Ho

MRC-University of Glasgow Centre for Virus Research

University of Glasgow

Glasgow, G61 1QH

**Email:** [Antonia.Ho@glasgow.ac.uk](mailto:Antonia.Ho@glasgow.ac.uk)

**Telephone:** 0141 330 5725

**Word count: 999**

**Keywords:** 4C Mortality Score, 4C Score, COVID-19, COVID-19 management guidance, inclusion of 4C score, documentation of 4C score

**ABSTRACT**

The 4C Mortality Score (4C Score) was designed to risk stratify hospitalised patients with COVID-19. We assessed inclusion of 4C Score in COVID-19 management guidance and its documentation in patients' case notes in January 2021 in UK hospitals. 4C Score was included within guidance by 50% of sites, though score documentation in case notes was highly variable. Higher documentation of 4C Score was associated with score integration within admissions proformas, inclusion of 4C Score variables or link to online calculator, and management decisions. Integration of 4C Score within clinical pathways may encourage more widespread use.

## **INTRODUCTION**

Early identification of patients with COVID-19 at risk of poor outcome could support early decision-making regarding treatment and escalation of care (1). The 4C Mortality score (4C Score) was developed to better inform clinical decision-making in patients hospitalised with COVID-19 (2), and has been validated in COVID-19 patients hospitalised between August 2020 and February 2021 (3). This study aimed to assess if, and how, the 4C Score was incorporated into the assessment of COVID-19 patients in UK hospitals in January 2021 during the peak of the second wave (4).

## **METHODS**

Between 3 June and 30 September 2021, investigators from all UK hospitals were invited to take part in a cross-sectional study on COVID-19-specific management guidance at their hospital site, and the role of the 4C Score within these documents. Additionally, they were invited to undertake a retrospective case note review of adult patients presenting with community-acquired COVID-19 during a 2-week period (11-24 January 2021) at the peak of the 'second wave' to ascertain the proportion of patients that had a 4C Score documented in their case notes. The protocol was developed in collaboration with the UK National Infection Trainee Collaborative for Audit and Research (NITCAR) following STROBE guidance (5). Site investigators were recruited through NITCAR and local trainee networks. Data were collected on a REDCap database (Research Electronic Data Capture, Vanderbilt University, US) (6). This study was classified as service evaluation by the West of Scotland Research Ethics Committee. Caldicott Guardian approval was obtained at each participating site. Details of the study protocol are included as an online Appendix.

## **RESULTS**

Forty-one hospitals in England and Scotland participated in the cross-sectional study of COVID-19 specific guidance, 32 of which took part in the retrospective cohort study (Supplementary Table S1). Characteristics of participating hospitals are summarised in Table 1.

**Table 1. Characteristics and guidance document use in participating hospitals**

	Hospital sites n (%)
<b>Location</b>	
England	22 (53.7)
Scotland	19 (46.3)
<b>Hospital size (number of beds)</b>	
<500	12 (29.2)
500-999	23 (56.1)
≥1000	6 (14.6)
<b>Intensive care unit on-site</b>	38 (92.7)
<b>Infectious diseases unit on-site</b>	23 (56.1)
<b>COVID-19 Guidance documents in use</b>	40 (97.6)
<b>Type of COVID-19 specific guidance documents</b>	
Guidance document	38 (95.0)
Admissions protocol	18 (45.0)
Admissions proforma	11 (27.5)
Poster	10 (25.0)
Sticker	2 (5.0)
Other*	6 (15.0)
<b>Number of guidance documents in use per hospital</b>	
0	1 (2.4)
1-2	28 (68.3)
3-5	12 (29.2)
<b>Prognostic scores included in guidance</b>	
4C mortality score	20 (50.0)
Clinical Frailty Scale (CFS)	18 (45.0)
NEWS 2**	17 (42.5)
CURB-65**	7 (17.5)
No prognostic score referenced	10 (25.0)
Other†	2 (5.0)

Abbreviations: NEWS2, National early warning score 2; CURB 65, Confusion, Urea, Respiration rate, Blood pressure.

Definitions: guidance document - providing an overview of management of patients with COVID-19; admissions protocol - document outlining local admissions pathways; admissions proforma - clinical document to be completed and included in patient case notes; Poster - in use in a clinical area to provide a visual prompt in the management of COVID-19; Sticker- in use in case notes to provide a visual prompt for clinicians in the management of COVID-19.

\*COVID-19-specific guidance documents: Antimicrobial Companion app (n=2); prescribing guidance for trial medications (n=2); emails and newsletters (n=1); daily ward round proforma (n=1)

†ASA (American Society of Anaesthesiologists) grade (n=1); Performance status (n=1).

All but one hospital (97.6%) reported the use of COVID-19-specific guidance. The most common forms of guidance were guidance documents (n=38; 95.0%) and admissions protocols (n=18; 45.0%) (Table 1; see Appendix for definitions). Twelve sites (30.0%) utilised three or more forms of COVID-19 specific guidance.

Thirty (75.0%) hospital sites referenced one or more prognostic scores in their COVID-19-specific guidance; 4C Score was the most commonly included (n=20; 50.0%), followed by Clinical Frailty Scale

(CFS; n=18) and NEWS2 (n=17) (Table 1). 4C Score was utilised alongside one or more other prognostic scores at 14 sites, most commonly with CFS (n=12). Thirteen (65.0%) sites that included 4C Score within their COVID-19 guidance used it to inform management decisions, including treatment escalation (n=8), discharge (n=4), admission destination (n=3), treatment limitation (n=3) as well as remdesivir prescription (n=3).

After assessment against eligibility criteria (see Appendix), 4,123 case records from 32 hospital sites were analysed (Figure 1). 4C Score was recorded in >10% patient records in eight (25.0%) sites, ≤10% records in 10 (31.3%), and 0% in 14 (43.8%) sites (Table 2).

**Table 2. Comparison of the use of COVID-19 management guidance in hospitals that recorded 4C Score in >10 or ≤10% of patient case notes**

	Hospital sites & percentage of reviewed cases with 4C Score documented		P-value†
	>10% case notes (n= 8 sites)	≤10% case notes (n= 24 sites)	
	n (%)	n (%)	
<b>Infectious diseases unit onsite</b>	6 (75.0)	11 (45.8)	0.15
<b>COVID-19 specific guidance in use</b>	8 (100.0)	23 (95.8)	1.00
<b>4C Score incorporated within guidance</b>	8 (100.0)	10 (41.7)	0.004
<b>Type of COVID-19 specific guidance documents</b>			
Guidance document	6 (75.0)	23 (95.8)	0.08
Admissions protocol	3 (37.5)	8 (33.3)	1.00
Admissions proforma	5 (62.5)	4 (16.7)	0.02
Poster	3 (37.5)	4 (16.7)	0.33
Sticker	0 (0)	2 (8.3)	1.00
<b>No. guidance documents in use</b>			
0	0 (0)	1 (4.1)	
1-2	4 (50.0)	18 (75.0)	
≥3	4 (50.0)	5 (23.8)	0.30
<b>Features of 4C use within guidance</b>			
4C Score variables detailed <u>or</u> link to online calculator	7 (87.5)	8 (33.3)	0.008
<b>4C Score used to inform management decision</b>	6 (75.0)	5 (20.8)	0.01
Patient discharge	2 (25.0)	1 (4.2)	
Patient admission	2 (25.0)	0 (0.0)	
Treatment escalation	5 (62.5)	2 (8.3)	
Treatment limitation	2 (25.0)	1 (4.2)	
Other*	1 (12.5)	3 (12.5)	

\*Other: Use of remdesivir (n=3), risk stratification (n=1); †Chi-squared or Fisher's Exact Test

Overall, it was documented in the records of 418 (10.1%) patients, with wide variation in frequency of 4C Score documentation between hospital sites (median 8.2%, range 1.0-74.1% among 18 sites with 4C Score documented in  $\geq 1$  case record). The score was mostly documented by junior doctors from foundation to specialty trainees (77.5%, n=324), and predominantly in medical assessment units or medical wards (93.8%), compared to the Emergency Department (3.8%).

The eight sites that recorded 4C Score for >10% of reviewed patients were more likely to include the score within the admissions proforma (62.5% (5/8) vs. 16.7% (4/24), p= 0.02), detail the variables or incorporate an electronic link to the online calculator (87.5% (7/8) vs. 33% (8/24), p=0.008), and utilise the score to inform management recommendations (75.0% (6/8) vs. 20.8% (5/24), p=0.01).

## DISCUSSION

No prior studies have explored the use of COVID-19-specific guidance or the application of 4C Score in the assessment of COVID-19 patients. In this nationwide study, most surveyed hospitals (97.6%) had COVID-19-specific guidance documents; around two-thirds employed multiple forms of guidance. The 4C Score was the most frequently referenced prognostic score, often in conjunction with CFS. The latter is the only clinical stratification tool referenced in the NICE COVID-19 management guidance (7). The 4C Score was chiefly used to inform treatment escalation, but was also used to inform decisions on patient destination (i.e. discharge or admission location) as well as treatment limitation.

Although half of participating sites incorporated the 4C Score within guidance, documentation of the score varied substantially between sites. Audits of documentation of established mortality risk scores such as CURB-65 (for patients with community-acquired pneumonia) have also demonstrated variable recording (8,9). Nevertheless, our findings suggest that documentation of the 4C Score may be encouraged by incorporation into clinicians' workflow through admissions proformas, detailing 4C Score variables or online calculator within guidance, in addition to linking the score to management recommendations.

This project demonstrates the capacity of national trainee networks to facilitate multi-centre studies. Collaboration with NITCAR and other trainee networks enabled the rapid recruitment of 41 UK sites and completion of >4,000 case note reviews within a short timeframe.

This study has several limitations. Although NITCAR is a national organisation, site participation was not representative, with a preponderance towards Scottish sites and no sites included from Wales or Northern Ireland. We assumed that guidance documents reported by sites between June and

September 2021 were in use in January 2021 (period selected for retrospective case note review); some of these documents may not have been in use at that time or may have since been updated. We equated documentation with use of the 4C Score; this would have omitted occasions when a score was calculated but undocumented in the case notes, or conversely, decisions that were made without the value being considered.

In summary, this national study highlights that 4C Score and other clinical stratification tools were widely adopted in COVID-19-specific management guidance. Ongoing validation of the 4C Score in the context of vaccinated patients and emerging SARS-CoV-2 variants is needed. However, ensuring that the score is accessible to clinicians through integration within clinical pathways could lead to more widespread deployment.



**Figure 1.** Flow chart of eligibility criteria for retrospective case note review

**Acknowledgements.** The authors would like to thank the following for their contribution to this study. NITCAR committee for support and assistance in facilitating this nationwide study. Vanesa Anton-Vazquez, Ali Alam, Matthew Kennedy, Imran Khan, Darver Malik, Andy Nicholson, Linda Provan, Matthew Stevens, Omolola Wilson for contributing hospital site data regarding COVID-19 specific guidance.

**Contributors.** AH is guarantor and corresponding author for this work and attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted. Contributor role taxonomy (CRediT): conceptualisation: AH, AEB, AM, JSP; data curation: AEB, JSP, AH; formal analysis AH; investigation: JSP, AA, SB, KB-E, SB-M, RB, TC, DC, AC, TC-B, JE, JG-B, BH, HH, AK, SL, CL, KM, CM, BM, CM, EM, JN, DP, LP, MS, DS, SS, JT, FT, SW, EW, RW; methodology: AH, AEB, AM, JSP; project administration: JSP, AEB, NITCAR, AH; writing – original draft: AEB, JSP, AH; writing – review and editing: AA, SB, KB-E, SB-M, RB, TC, DC, AC, TC-B, JE, JG-B, BH, HH, AK, SL, CL, KM, CM, BM, CM, EM, JN, DP, LP, MS, DS, SS, JT, FT, SW, EW, RW, AM, NITCAR.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-project sectors.

**Competing interests** AH has received grant funding from UKRI for work that included the development and validation of 4C Mortality Score. Presented data included in this manuscript at the St. Andrew’s Day Symposium: Updates on Acute Medicine, Royal College of Physicians Edinburgh.

**Patient consent for publication** Not required.

## References

1. Wynants L, Sotgiu G. Improving clinical management of COVID-19: the role of prediction models. *Lancet Respir Med* [Internet]. 2021;9(4):320–1. Available from: [http://dx.doi.org/10.1016/S2213-2600\(21\)00006-0](http://dx.doi.org/10.1016/S2213-2600(21)00006-0)
2. Knight SR, Ho A, Pius R, Buchan I, Carson G, Drake TM, et al. Risk stratification of patients admitted to hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: Development and validation of the 4C Mortality Score. *BMJ* [Internet]. 2020;370(September):1–13. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7116472/pdf/bmj.m3339.pdf>

3. Knight SR, Gupta RK, Ho A, Pius R, Buchan I, Carson G, et al. Prospective validation of the 4C prognostic models for adults hospitalised with COVID-19 using the ISARIC WHO Clinical Characterisation Protocol. 2021;1–10.
4. healthcare @ coronavirus.data.gov.uk [Internet]. Available from: <https://coronavirus.data.gov.uk/details/healthcare>
5. Von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: Guidelines for reporting observational studies. PLoS Med [Internet]. 2007;4(10):1623–7. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2020495/pdf/pmed.0040296.pdf>
6. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. J Biomed Inform [Internet]. 2009;42(2):377–81. Available from: <https://www.sciencedirect.com/science/article/pii/S1532046408001226>
7. NICE. COVID-19 rapid guideline: managing COVID-19. Natl Inst Heal Care Excell [Internet]. 2021;(September):1–78. Available from: <https://www.nice.org.uk/terms-and->
8. Foley SC, Kelly EM, O’Neill SJ. Audit of the management of patients admitted with community acquired pneumonia. Ir Med J. 2006 May;99(5):138–40.
9. Al-Abri SS, Al-Maashani S, Memish ZA, Beeching NJ. An audit of inpatient management of community-acquired pneumonia in Oman: A comparison with regional clinical guidelines. J Infect Public Health [Internet]. 2012;5(3):250–6. Available from: <http://dx.doi.org/10.1016/j.jiph.2012.03.002>

Patients aged  $\geq 18$  reviewed in ED, MAU or admitted to an acute hospital bed from 11/1/2021 to 24/1/2021 inclusively with a diagnosis of 'Covid-19', 'Covid-19 pneumonia', 'Covid-19 pneumonitis', 'SARS-CoV-2 infection' or 'SARS-CoV-2 PCR positive' on ED or hospital discharge documentation

6288

Excluded by local investigators

2094

Recorded in RedCap by local investigators

4194

Incomplete record submitted

4

Date of admission outwith study dates

7

First positive COVID-19 test result  $> 7$  days from hospital admission

19

First positive COVID-19 test result  $\geq 14$  days prior to admission

41

Included in analysis

4123

4C score recorded in case notes

418

4C not recorded in case notes

3705