

## How To Enhance Empathy Nursing Students In Education: Literature Review

Bhakti Permana<sup>1,2</sup>, Moses Glorino Rumambo Pandin<sup>3</sup>

<sup>1</sup> Doctoral Program of Nursing, Faculty of Nursing, Airlangga University

<sup>2</sup> STIKep PPNI Jawa Barat

<sup>3</sup> Departement of English Literature, Faculty of Humanity, Airlangga University

Corresponding author: Bhakti Permana

Email: [bhakti.permana-2021@fkp.unair.ac.id](mailto:bhakti.permana-2021@fkp.unair.ac.id),

### Abstract

**Introduction.** The tendency of reducing student empathy requires efforts to improve it through education or training. The purpose of this review is to identify education, learning, or teaching that is used to increase student empathy.

**Method.** The design used a Literature Systematic Review. Articles were conducted in three electronic databases guided by The PRISMA. Articles published in English and in 2019 to 2021.

**Results.** The result obtained 20 articles and will be reviewed. Eighteen articles show an effect or relationship between intervention and empathy. Training materials that can increase student empathy are the concept of empathy, communication, mindfulness, and transcultural nursing education. Education and training using patients/patient simulations to help improve student empathy, namely: expert patients, ECARE Program, polypharmacy effects; clinical Simulation of Inpatients, Lectures in class, Virtual Dementia Tour, living in poverty, and games for team interaction. Another method is through the KSS module, mannequin simulator experience, and peers.

**Conclusion.** Education and training on the concepts of empathy, communication, meditation, and cultural competence using Simulation and immersion methods with patients, being like patients, using mannequins, or interacting with vulnerable groups can increase student empathy. Faculty and lecturers can apply experiential learning methods with Simulation and immersion in learning or training courses.

Keywords: empathy, student, nursing, learning, education

### 1. Introduction

Nursing students are often involved in nursing services for patients when conducting the clinical nursing practice in hospital. Students may not be ready to care and have negative behaviors from patients like being afraid, incapable, helpless, anxious, sad, low self-esteem, unable to manage their emotions, and maintaining patient space (Kapucu & Bulut, 2018; Lin et al., 2017). Feeling under pressure, failing to use therapeutic communication skills, a lack of courage, eastern

cultural influences, and emotional management experience are all aspects that affect nursing students' views (Lin et al., 2017). Empathy problems, together with negative attitudes and stigma, are an issue for health care improvement. (McKenna et al., 2012).

Empathy is the process of placing oneself in another person's shoes, viewing events from a different perspective, understanding and feeling that others' emotions and ideas, and explaining this condition to him (Cunico et al., 2012; Paola Ferri et al., 2017). The dimensions of empathy are cognitive, emotional, and behavioral. The emotional dimension shows potential for empathy. The cognitive and behavioural dimensions are the ability to empathize.

Empathy in health services is an essential component of a quality therapeutic and nursing relationship (Hojat, 2016). Empathy for a health professional will reduce depression, anxiety, distress; and increase emotional well-being, satisfaction, and adherence to treatment regimens (Hojat & Hojat, 2016) and can be used to carry out patient-centered care and interactions (Omid et al., 2018). Therefore, high empathy is essential for the professional development of students in preparing for the medical profession (Sapiro et al., 2004).

Several studies have shown that students' empathy levels can decrease during nursing education (Ward et al., 2012), the average medical profession student is lower than non-medical students, and 18.2% of all medical program students (n = 412) showed the lowest level of empathy (Sobczak et al., 2021). It requires programs to implement educational interventions to increase students' empathy levels.

The importance of empathic skills and the tendency to decrease empathy for students requires a nursing education process to develop empathy skills. Training in communication skills, experiential learning, and medical interprofessional education interventions, and simulation-based education, are all effective techniques for increasing students' empathy levels (Batt-Rawden et al., 2013; Bearman et al., 2015). For this reason, it is necessary to explore the latest developments in empathy education interventions to increase nursing students' empathy.

## 2. Review Methods

### 2.1 Study Objectives

The goal of this review is to identify the education, learning, or teaching that is used to increase the empathy of nursing students.

### 2.2 Eligibility Criteria

The type of research used is experimental design, quasi-experimental and mixed methods with quasi-experiments published in English between January 2019 to 2021. The type of the participant is nursing students regardless of gender, age, or country. The type of intervention chosen is an educational intervention designed to increase the level of empathy of nursing students. The type of outcome measure is quantitative data that is objectively measured or self-reported about empathy.

### 2.3 Search Strategy

Search articles or journals using keywords. The keywords used are "Nursing student" OR "Nurse student" AND Education OR Training OR Learning AND Empathy. The literature search used three databases, namely Scopus, Scient Direct, and Sage journals. In addition to searching for articles using keywords, the characteristics of the journal are also determined, namely publications in 2019 and above, full text, and English.

#### 2.4 Study Selection

The identified articles are entered into the Mendeley database, and then duplicate titles are identified and deleted. Potential articles will be screened by title and abstract based on inclusion criteria.

#### 2.5 Critical Appraisal

Articles that have met the criteria will then be assessed for quality using the standard assessment format from the JBI Critical Appraisal checklist and MMAT. The selection of the assessment format is based on the design of each quasi-experiment article using the JBI Critical Appraisal Checklist for Quasi-Experimental Studies and Mixmethod using the Mixed Methods Appraisal Tool (MMAT). Critical appraisal assesses eligible studies if the research score is at least 60% of the cut-off point value. Screening results showed that 20 articles achieved a score higher than 60%.

#### 2.6 Data Collection Process

Articles that meet the criteria and study quality will be tabulated using a table of researchers and year of publication, objectives, design, sample, variables, instruments, analysis, intervention, and results (Table 1).

### 3. Results

#### 3.1 Study Selection

Searching for articles through the database found 739 articles. After going through the selection process, 20 articles were obtained that would be reviewed (figure 1).




Figure 1 searching literature

### 3.2 Characteristics of Included Studies

The selected articles were published in 2019-2021, as many as four in 2019, 11 in 2020, and 5 in 2021. The research was carried out in 12 countries, namely four articles in the USA, three articles each in China, and Turkey, two articles in Spain, and one each from Malaysia, Ghana, Palestine, UK, Italy, Brazil, Singapore, and South Korea. The designs used in the research articles were a quasi-experiment with 13 articles, randomized control trial with six articles, and a mixed-method (with quasi-experiment) with 1 article. Based on the distribution of respondents, they were divided into one group of 5 articles and two groups of 15 articles. The measurement of respondents has measured post-test as many as two articles and pre-post test as many as 18 articles.

### 3.3 Participant Characteristics

The sampling method used convenience sampling of as many as 13 articles and random sampling as five and RCT 2 articles. The smallest sample size is 34 respondents, and the most significant sample is 250 respondents.

### 3.4 Types of Outcome Measures

The instruments to measure the empathy of nursing students used were: Jefferson's Scale of Empathy with five articles, JSE HPS version with six articles, Kiersmae Chen Empathy Scale with two articles, Empathic Tendency Scale (ETS) with 1 article, Empathy Inventory with 1 article, Empathic skill scale-B form (ESS-B) with 1 article, Interpersonal Reactivity Index (IRI) with two articles, The Comprehensive State Empathy Scale (CSES) with 1 article, and Toronto Empathy Questionnaire (TEQ) with 1 article.

### 3.5 Empathy Outcomes

Research shows that education and training can increase student empathy in several ways.

#### Subject

Training materials that can increase student empathy are:

1. The concept of empathy through training for 8 hours where every week for 2 hours (Ahmad et al., 2020)
2. Interpersonal Communication Skills, given for 14 hours where every week for 2 hours (Bas-Sarmiento et al., 2019)
3. Mindfulness-based empathy training (MBET) teaches about Aging, communication, empathy, mindfulness, body screening meditation, acceptance and allow, awareness of the judiciary, and Effective/empathic/active listening. Training given for 16 sessions, every week carried out for two sessions (Can Gür & Yilmaz, 2020)
4. Empathy Educational Model (EEM) contains primary material of empathy, emotion and language skills, nonlinguistic skills, and practicing empathy, given for 12 hours every week 2-3 times (Yang et al., 2020)
5. Transcultural nursing education held with subject health, nursing, culture, transcultural nursing, cultural sensitivity, and nursing research, for seven weeks (Kaçan & Örsal, 2020)
6. A re-unification educational program containing a model of cultural competence, given for 1 hour 6 weeks (Kim et al., 2020)

#### Patient

Education and training using patients/patient simulations to help increase student empathy, namely:

1. Expert patients shared stories when illness, the experience of nursing care. This program was held for one day (7 hours) (P Ferri et al., 2019)
2. The Empathetic CARE and Response program (ECARE) use restrained like patients in the healthcare setting, 1-hour lecture (Goh et al., 2020)
3. Polypharmacy effects used seven medications with jelly beans, managed multidrug and multi-dose regime for seven days (Kaulback et al., 2021)
4. Clinical Simulation like Hospitalized Patient used disposable diapers in the elimination needs (Filho et al., 2020)
5. Dementia LiveVR is a simulation experience to immerse life with dementia for seven minutes (Kimzey et al., 2020)

6. Virtual Dementia Tour, participants wore three small balls, a pair of blurred goggles, and large gloves, with the thumb and forefinger of the dominant hand taped together, as well as the last three fingers of the non-dominant hand, and the VDT room was poorly light and decorated (Peng et al., 2020)
7. Games to promote team interactions in the clinical setting for 6-8 weeks every 2.5 hours (Zelenski et al., 2020)
8. The KSS module is done for nursing students who practice in the pediatric room. The components of KSS module are Knowledge, Simulation and Sharing. The internship includes lectures, simulation scenarios and a self-reflection. The module was delivered over five days during the nine-month internship (Ding et al., 2020).

Education and training exercises using 1) mannequins through simulators' have, integrating simulator experience with traditional didactic approach. The intervention is done in the laboratory for four weeks (Ayed et al., 2021); 2) Peer Tutoring, after the tutor describes the incident, the tutee must describe the issue (Gisbert & Rivas, 2021); and 3) Community; Participants are assigned to a poor family and must live in their order for a simulated month separated into four consecutive weeks (Phillips et al., 2020).

#### 4. Discussion

The results of this review illustrate that education and training are carried out to increase nursing students' empathy. Of the 20 articles, 18 articles showed that education or training could increase/there was a positive relationship with the level of empathy. The education and training are about the concept of empathy, Interpersonal Communication Skills, Mindfulness-based empathy training (MBET), transcultural nursing education, and cultural competence. The method uses patients/patient simulations, peers, and the community.

Simulation methods with actual patients or like patients by humans or mannequins are proven and long-standing methods that effectively increase student empathy. The results of another review show that the most effective method is immersion and experiential Simulation in vulnerable groups and providing opportunities for reflection (T Levett-Jones et al., 2019). Simulation may be an appropriate educational methodology for developing empathy and empathy behavior (Bearman et al., 2015). A hospital simulation can provide an experience of dealing with illness and hospital life, feeling dependent and the need for contact, and dealing with being a patient. Other research shows that can increase empathetic behavior by listening to the patient's voice as a consumer of health services (Heidke et al., 2018).

Simulation of vulnerable groups in the community can increase student empathy. Social simulations with the poverty conditions of (Menzel et al., 2014) and persons with disabilities (Tracy Levett-Jones et al., 2017) can be helpful and used in learning. Simulations with mannequins can increase student empathy following the thesis and goals that NLN has made (Haley et al., 2017).

Educational materials and lectures given to students to increase empathy are also reviewed, namely communication skills, interprofessional skills, and the concept of empathy (Batt-Rawden

et al., 2013), and cultural and linguistic differences through three-dimensional simulations (Everson et al., 2015).

Based on the material provided, the length of education or training carried out varied from 6 hours (Kim et al., 2020), 8 hours (Ahmad et al., 2020), 12 hours (Yang et al., 2020), and 14 hours (Bas-Sarmiento et al., 2019). Education and training time is carried out starting from 1 day (P Ferri et al., 2019), four weeks (Ayed et al., 2021), six weeks (Kim et al., 2020), seven weeks (Kaçan & Örsal, 2020), and eight weeks (Can Gür & Yilmaz, 2020; Zelenski et al., 2020). On average, each week face-to-face for 2 hours. The time determination is adjusted to the material discussed and according to the stages of the learning. It guides lecturers and faculties in planning education and training for nursing students.

### Implications for education

The experiential learning method with patient simulation becomes like a patient, mannequin, and immersion in the community and helps increase student empathy. The experiential learning method with Simulation and immersion can be used as a learning method in nursing lectures or training.

### 5. Conclusion

Nursing students' empathy can be increased by improving cognitive, affective, and psychomotor aspects. Education and training can improve cognitive aspects by providing knowledge about empathy, communication, meditation, and cultural competence. Simulation and immersion with patients, being like patients, using mannequins, or interacting with vulnerable groups improve students' affective and psychomotor aspects.

## References

- Ahmad, A., Wahab, H., & Seman, N. (2020). The effect of educational intervention on enhancing the empathy among nursing students. *Enfermeria Clinica*, 30, 100–103. <https://doi.org/10.1016/j.enfcli.2019.12.038>
- Ayed, A., Malak, M. Z., Al-amer, R. M., Batran, A., & Salameh, B. (2021). Effect of High Fidelity Simulation on Perceptions of Self-Awareness, Empathy, and Patient-Centered Care Among University Pediatric Nursing Classes. *Clinical Simulation in Nursing*, 56, 84–90. <https://doi.org/10.1016/j.ecns.2021.04.005>
- Bas-Sarmiento, P., Fernández-Gutiérrez, M., Díaz-Rodríguez, M., Carnicer-Fuentes, C., Castro-Yuste, C., García-Cabanillas, M. J., Gavira-Fernández, C., de los Ángeles Martelo-Baro, M., Paloma-Castro, O., del Carmen Paublete-Herrera, M., Rodríguez-Cornejo, M. J., & Moreno-Corral, L. (2019). Teaching empathy to nursing students: A randomised controlled trial. *Nurse Education Today*, 80, 40–51. <https://doi.org/https://doi.org/10.1016/j.nedt.2019.06.002>
- Batt-Rawden, S. A., Chisolm, M. S., Anton, B., & Flickinger, T. E. (2013). Teaching empathy to medical students: An updated, systematic review. *Academic Medicine*, 88(8), 1171–1177. <https://doi.org/10.1097/ACM.0b013e318299f3e3>
- Bearman, M., Palermo, C., Allen, L. M., & Williams, B. (2015). Learning empathy through simulation: A systematic literature review. *Simulation in Healthcare*, 10(5), 308–319. <https://doi.org/10.1097/SIH.0000000000000113>
- Can Gür, G., & Yilmaz, E. (2020). The effects of mindfulness-based empathy training on empathy and aged discrimination in nursing students: A randomised controlled trial. *Complementary Therapies in Clinical Practice*, 39, 101140. <https://doi.org/https://doi.org/10.1016/j.ctcp.2020.101140>
- Cunico, L., Sartori, R., Marognolli, O., & Meneghini, A. M. (2012). Developing empathy in nursing students: A cohort longitudinal study. *Journal of Clinical Nursing*, 21(13–14), 2016–2025. <https://doi.org/10.1111/j.1365-2702.2012.04105.x>
- Ding, X., Wang, L., Sun, J., Li, D., Zheng, B., He, S., Zhu, L., & Latour, J. M. (2020). Effectiveness of empathy clinical education for children's nursing students: A quasi-experimental study. *Nurse Education Today*, 85, 104260. <https://doi.org/https://doi.org/10.1016/j.nedt.2019.104260>
- Everson, N., Levett-Jones, T., Lapkin, S., Pitt, V., van der Riet, P., Rossiter, R., Jones, D., Gilligan, C., & Courtney-Pratt, H. (2015). Measuring the impact of a 3D simulation experience on nursing students' cultural empathy using a modified version of the Kiersma-Chen Empathy Scale. *Journal of Clinical Nursing*, 24(19–20), 2849–2858. <https://doi.org/10.1111/jocn.12893>
- Ferri, P., Rovesti, S., Padula, M. S., D'Amico, R., & Di Lorenzo, R. (2019). Effect of expert-patient teaching on empathy in nursing students: A randomized controlled trial. *Psychology Research and Behavior Management*, 12, 457–467. <https://doi.org/10.2147/PRBM.S208427>
- Ferri, Paola, Rovesti, S., Panzera, N., Marcheselli, L., Bari, A., & Di Lorenzo, R. (2017). Empathic



- attitudes among nursing students: A preliminary study. *Acta Biomedica*, 88(4), 22–30. <https://doi.org/10.23750/abm.v88i3-S.6610>
- Filho, C. K. C., Meska, M. H. G., Machado, G. C. C., Franzon, J. C., Mano, L. Y., & Mazzo, A. (2020). Empathy of nursing students in a simulated clinical activity [Empatia de estudantes de enfermagem em atividade clínica simulada] [Empatía de estudiantes de enfermería en actividad clínica simulada]. *Cogitare Enfermagem*, 25, 1–10. <https://doi.org/10.5380/ce.v25i0.68142>
- Gisbert, D. D., & Rivas, A. V. (2021). Implementing Peer Tutoring For The Development Of Empathy. *Investigacion y Educacion En Enfermeria*, 39(2). <https://doi.org/10.17533/udea.iee.v39n2e07>
- Goh, Y.-S., Seetoh, Y.-T. M., Chng, M.-L., Ong, S. L., Li, Z., Hu, Y., Ho, C.-M. R., & Ho, S. H. C. (2020). Using Empathetic CARE and REsponse (ECARE) in improving empathy and confidence among nursing and medical students when managing dangerous, aggressive and violent patients in the clinical setting. *Nurse Education Today*, 94, 104591. <https://doi.org/10.1016/j.nedt.2020.104591>
- Haley, B., Heo, S., Wright, P., Barone, C., Rettigantid, M. R., & Anders, M. (2017). Effects of Using an Advancing Care Excellence for Seniors Simulation Scenario on Nursing Student Empathy: A Randomized Controlled Trial. *Clinical Simulation in Nursing*, 13(10), 511–519. <https://doi.org/10.1016/j.ecns.2017.06.003>
- Heidke, P., Howie, V., & Ferdous, T. (2018). Use of healthcare consumer voices to increase empathy in nursing students. *Nurse Education in Practice*, 29, 30–34. <https://doi.org/10.1016/j.nepr.2017.11.007>
- Hojat, M. (2016). Empathy in health professions education and patient care. In *Empathy in Health Professions Education and Patient Care*. Springer International Publishing. <https://doi.org/10.1007/978-3-319-27625-0>
- Hojat, M., & Hojat, M. (2016). Empathy as Related to Personal Qualities, Career Choice, Acquisition of Knowledge, and Clinical Competence. In *Empathy in Health Professions Education and Patient Care* (pp. 151–167). Springer International Publishing. [https://doi.org/10.1007/978-3-319-27625-0\\_9](https://doi.org/10.1007/978-3-319-27625-0_9)
- Kaçan, C. Y., & Örsal, Ö. (2020). Effects of Transcultural Nursing Education on the Professional Values, Empathic Skills, Cultural Sensitivity and Intelligence of Students. *Journal of Community Health Nursing*, 37(2), 65–76. <https://doi.org/10.1080/07370016.2020.1736374>
- Kapucu, S., & Bulut, H. (2018). Nursing students' perspectives on assisting cancer patients. *Asia-Pacific Journal of Oncology Nursing*, 5(1), 99–106. [https://doi.org/10.4103/apjon.apjon\\_44\\_17](https://doi.org/10.4103/apjon.apjon_44_17)
- Kaulback, M., Barker, N., & Yocom, D. (2021). Experiencing Empathy Through A Polypharmacy Simulation Experience in Baccalaureate Nursing Education. *Clinical Simulation in Nursing*, 59, 94–97. <https://doi.org/10.1016/j.ecns.2021.06.004>
- Kim, M., Noh, G.-O., & Park, M. J. (2020). The Effect of Re-unification Education on Empathy and Multicultural Competence of Nursing-College Students. In *Medico-legal Update* (Vol.

- 20, Issue 3). <http://www.ijop.net/index.php/mlu/article/download/1449/1320>
- Kimzey, M., Patterson, J., & Mastel-Smith, B. (2020). Effects of Simulation on Nursing Students' Dementia Knowledge and Empathy: A Mixed Method Study. *Issues in Mental Health Nursing, 42*(3), 274–279. <https://doi.org/10.1080/01612840.2020.1797252>
- Levett-Jones, T, Cant, R., & Lapkin, S. (2019). A systematic review of the effectiveness of empathy education for undergraduate nursing students. *Nurse Education Today, 75*, 80–94. <https://doi.org/10.1016/j.nedt.2019.01.006>
- Levett-Jones, Tracy, Lapkin, S., Govind, N., Pich, J., Hoffman, K., Jeong, S. Y. S., Norton, C. A., Noble, D., Maclellan, L., Robinson-Reilly, M., & Everson, N. (2017). Measuring the impact of a 'point of view' disability simulation on nursing students' empathy using the Comprehensive State Empathy Scale. *Nurse Education Today, 59*(April), 75–81. <https://doi.org/10.1016/j.nedt.2017.09.007>
- Lin, M.-F., Hsu, W.-S., Huang, M.-C., Su, Y.-H., Crawford, P., & Tang, C.-C. (2017). "I couldn't even talk to the patient": Barriers to communicating with cancer patients as perceived by nursing students. *European Journal of Cancer Care, 26*(4), 1–9. <https://doi.org/10.1111/ecc.12648>
- McKenna, L., Brown, T., Williams, B., Nursing, R. L.-J. of P., & 2020, U. (2012). Empathic and listening styles of first year undergraduate nursing students: A cross-sectional study. *Elsevier*. <https://www.sciencedirect.com/science/article/pii/S8755722320301630>
- Menzel, N., Willson, L. H., & Doolen, J. (2014). Effectiveness of a poverty simulation in second life®: Changing nursing student attitudes toward poor people. *International Journal of Nursing Education Scholarship, 11*(1), 39–45. <https://doi.org/10.1515/ijnes-2013-0076>
- Omid, A., Haghani, F., & Adibi, P. (2018). Emotional Intelligence: An Old Issue and a New Look in Clinical Teaching. *Advanced Biomedical Research, 7*(1), 32–32. <https://doi.org/10.4103/2277-9175.225926>
- Peng, X., Wu, L., Xie, X., Dai, M., & Wang, D. (2020). Impact of Virtual Dementia Tour on empathy level of nursing students: A quasi-experimental study. *International Journal of Nursing Sciences, 7*(3), 258–261. <https://doi.org/https://doi.org/10.1016/j.ijnss.2020.06.010>
- Phillips, K. E., Roberto, A., Salmon, S., & Smalley, V. (2020). Nursing Student Interprofessional Simulation Increases Empathy and Improves Attitudes on Poverty. *Journal of Community Health Nursing, 37*(1), 19–25. <https://doi.org/10.1080/07370016.2020.1693095>
- Sobczak, K., Zdun-Ryżewska, A., & Rudnik, A. (2021). Intensity, dynamics and deficiencies of empathy in medical and non-medical students. *BMC Medical Education, 21*(1), 1–9. <https://doi.org/10.1186/s12909-021-02927-x>
- Ward, J., Cody, J., Schaal, M., & Hojat, M. (2012). The Empathy Enigma: An Empirical Study of Decline in Empathy Among Undergraduate Nursing Students. *Journal of Professional Nursing, 28*(1), 34–40. <https://doi.org/10.1016/j.profnurs.2011.10.007>
- Yang, C., Zhu, Y.-L., Xia, B.-Y., Li, Y.-W., & Zhang, J. (2020). The effect of structured empathy education on empathy competency of undergraduate nursing interns: A quasi-experimental study. *Nurse Education Today, 85*, 104296.

<https://doi.org/https://doi.org/10.1016/j.nedt.2019.104296>

Zelenski, A. B., Saldivar, N., Park, L. S., Schoenleber, V., Osman, F., & Kraemer, S. (2020). Interprofessional improv: Using theater techniques to teach health professions students empathy in teams. *Academic Medicine*, 95(8), 1210–1214. <https://doi.org/10.1097/ACM.00000000000003420>

Tabel 1 review the article

Author	Methods	Intervention	Result
Ahmad, Wahab & Seman	Design: quasi-experimental Sample: 64 nursing students Variable: students' empathy Instrumen: Jefferson's Scale of Empathy	Training educational intervention for eight hours didactic (2 h per week) about concept of empathy, and then presentation of short video and discussion.	The empathy increased during post-intervention
Alhassan	Design: design in a randomised controlled trial (RCT) Sample: 181 nursing students and 49 midwifery students Variable: students' empathy Instrumen: JSE HPS version	Training in communication skills was conceived and developed utilizing the 'Four Habits Model' and PCNF. Small group discussions, brainstorming, participant experience, group reports, roleplaying, questions and answers, videos, and summaries were all used as methods.	the scores of empathy were no difference between the group
Ayed et.al.	Design: pretest-posttest design Sample: 83 respondent Variable: self-awareness, empathy, and patient-centered care Instrumen: Self-Consciousness Scaled Revised, Kiersmae Chen Empathy Scale, Patient Practitioner Orientation Scale	For four weeks, the intervention included simulation scenarios and debriefing. Three simulation scenarios were downloaded from the National Library of Medicine's ACES website ( <a href="http://www.nlm.org/ACE">www.nlm.org/ACE</a> ). For twenty minutes, participants used a simulator in combination with a traditional didactic method. Based on students' verbal reflections in a separate room, a systematic debriefing procedure was applied.	The empathy difference between pre and post high fidelity simulation intervention
Bas-Sarmiento et.al.	Design: Multicentre randomised controlled trial. Sample: 116 nursing students Variable: students' empathy Instrumen: Jefferson Scale of Empathy (JSE)	Specific training in communication The university teaching personnel accomplished the training with specific communication training. The intervention included two 2-hour weekly meetings for a total of seven session (14 h).	The empathy were improved after the intervention.
Can Gür & Yilmaz	Design: randomized controlled trial (RCT) Sample: 123 nursing students Variable: empathy level Instrumen: JSENS, ADAS	Eight weeks of mindfulness-based empathy training (MBET) are held twice a week. The training program included a total of sixteen sessions. Pedagogical expression, role-playing, model learning, and homework were all incorporated in the training. Mindfulness breathing, body screening meditation, cognitive awareness, and dried grape meditation were all formal practices taught in class and at home.	The empathy level reported increase.
Yang et.al.	Design: quasi-experimental, with	Nine dimensions composed an	The empathy

Author	Methods	Intervention	Result
	<p>two-group comparison  Sample: 118 nursing students  Variable: empathy competence  Instrument: The Jefferson Scale of Empathy-Health Providers (JSE-HPs)  Analysis: t-test, Chi-squared test</p>	<p>Empathy Educational Model (EEM). Classroom-based instruction was provided during a two-week, 12-hour structured empathy education program (two 3-hour sessions per week). Didactics, video demonstrations, situational teaching, problem-based learning (PBL), and task-based learning were used as instructional methodologies.</p>	<p>competence was higher after intervention</p>
Dincer & Inangil	<p>Design: randomized controlled trial study  Sample: 70 nursing students  Variable: alexithymia, empathy, and attitude toward disabled persons  Intervention: the affective learning method  Instrument: TAS, ETS and ATDPS</p>	<p>The intervention group used a technique referred to as transformative learning. The transformative process is designed to help an individual transform his or her beliefs, assumptions, and experiences into new and more relevant perspectives. In education, transformative learning approaches should include four separate activities: investigative, collaborative, higher-order thinking, and interactive.</p>	<p>The empathy were no significant difference after the intervention</p>
Ferri et.al.	<p>Design: randomized controlled trial  Sample: 144 nursing students  Variable: empathy development  Instrument: Jefferson Scale of Empathy — Health Professions Student (JSE-HPS)</p>	<p>The educational intervention began with a 1.5-hour theoretical session on empathy, followed by a presentation by another teacher (1.5 hrs). Following that, the two expert patients served as trainers, sharing their disease stories, as well as their positive and negative experiences with nursing care. Finally, the nursing instructor led a debriefing session to allow students to reflect on their educational experience (1 hour).</p>	<p>The empathy were significant difference in the experimental group. Male students showed increased in empathy after training.</p>
Filho et.al.	<p>Design: quasi-experimental study  Sample: 107 nursing students  Variable: Empathy  Instrument: Empathy Inventory</p>	<p>Clinical of patient simulation made use of the scenario's hospital structure, and the mediators' learning objectives contributed to the defined level of fidelity. Five assessing nurses evaluated the design both visually and conceptually.</p>	<p>The cognitive components of empathy will promote after intervention of the simulated clinical activity</p>
Gisbert & Rivas	<p>Design: a mixed methods sequential explanatory design  Sample: 76 nursing students  Variable: Empathy  Intervention: six one-hour sessions of reciprocal peer tutoring</p>	<p>The program consists of six one-hour peer tutoring sessions held twice a week. Two distinct dossiers are distributed to the students (one for student A and one for student B). While the tutor describes the event, the tutee listens and, if necessary, asks</p>	<p>The empathy revealed improvements in intervention group.</p>

Author	Methods	Intervention	Result
	Instrumen: Jefferson Scale of Empathy	questions. The tutee is then required to explain the scenario as though it were their own.	
Goh et.al.	Design: quasi-experimental design Sample: 249 nursing and 50 medical students Variable: empathy and confidence Instrumen: JSE	The Empathetic CARE and REsponse program (ECARE) emphasized the importance of an empathic response to restraint use. A classroom-based session utilizing video-assisted learning and restrained patients is followed by a facilitator-led debriefing session to address any concerns and potential psychological impact on the students.	The level of empathy increases after attending ECARE.
Kaçan & Örsal	Design: quasi-experimental Sample: 125 respondent Variable: Professional Values, Empathic Skills, Cultural Sensitivity and Intelligence Instrumen: Nurses professional values scale revised form (NPVS-R), Empathic skill scale-B form (ESS-B), Intercultural sensitivity scale (ISS)	The TN course was held for seven weeks with the theoretical basis namely: “health and culture”, “nursing and culture”, “transcultural nursing and historical development”, “basic concepts in transcultural nursing”, “transcultural nursing models”, “ethics in transcultural nursing” and “culturally adequate and cultural sensitivity approach to nursing research”, “the transcultural approach to reproductive health, the transcultural approach in the prenatal and postnatal period, in child care and diseases, in chronic diseases and care, in psychiatric diseases and care and the cultural approach in pain”.	The experiment group demonstrated greater empathy than the control group.
Kaulback, Barker & Yocom	Design: a quasi-experimental, pretest-posttest design Sample: 34 nursing students Variable: Empathy Instrumen: Kiersma-Chen Empathy Scale (KCES)	The intervention was given a prescription that included seven medications with jelly beans and was instructed to take the medications exactly as prescribed.	The empathy demonstrated a higher post intervention
Kim, Noh, & Park	Design: an experimental study Sample: 65 students: 32 in the experimental group and 33 in the control group Variable: empathy and multicultural competence Instrumen: Cross-Cultural Competence Inventory (CCCI), Interpersonal Reactivity Index (IRI)	The intervention was a re-unification educational program based on a model of cultural competence. The subject session focused on the backgrounds other cultures, health, inequality, and education for North Koreans and defectors, cultural sensitivity, re-unification based on empathy and multicultural competence. There was one 60-minute session per week.	The Empathy increased in the experimental group
Kimzey,	Design: one group quasi-	The intervention was led by a	the empathy indicated

Author	Methods	Intervention	Result
Patterson & Mastel-Smith	experimental, Mixed methods research Sample: 55 students Variable: students' empathy and dementia knowledge Instrument: Dementia Knowledge Assessment Tool Version 2 (DKAT2), The Comprehensive State Empathy Scale (CSES)	qualified facilitator and focused on Dementia LiveVR, a seven-minute dementia simulation experience. The students wore glasses with customized lenses and wore headphones that covered their plastic gloves. Other pupils were instructed verbally to complete five straightforward tasks. The facilitator then led a systematic debriefing at the conclusion of the session.	in quantitative findings and qualitative findings showed six components of empathy
Peng et.al.	Design: quasi-experiment Sample: 45 nursing students Variable: students' empathy level Instrument: Jefferson Scale of Empathy-Health Professional Students (JSE-HPS)	Modified VDT used a pair of dried corn kernels and a needle to simulate foot pain in shoes. The participant wore a pair of blurred goggles and large gloves, thumb and forefinger were taped. Five tasks were assigned to the participants: putting yellow socks, folding towels, brushing teeth, writing a three-sentence memo or sketching a basic picture, and picking up three little balls.	Their empathy levels demonstrated improvements
Phillips et.al.	Design: quasy experiment Sample: 117 respondent Variable: attitudes, empathy, self reflection and insight Instrument: APPP scale, TEQ, self-reflection and insight scale (SRIS)	The simulation was six hours, including an orientation session and a three-hour simulation. Every session used a 15-minute debriefing weekly. The participant was assigned a role in the community that lives in poverty. Participants live in real time for 15 minutes and then participate in a community debriefing session for 30 minutes. The entire simulated event lasted approximately three and a half hours.	The empathy Increased after the intervention
Yu et.al.	Design: : A randomized trial with three phases. Sample: 238 nursing students Variable: empathy Intervention: The WITS Instrument: Jefferson Scale of Empathy (JSE) Health Professional version	The WITS intervention is an immersive digital story told by a real-life colon cancer patient. The story was captured in audio format and is comprised of nine digital segments (some still images). To avoid congestion in the suite, students began the intervention in small groups of 5–6 at 10-minute intervals.	The empathy were higher in the intervention group
Zelenski et.al.	Design: a convergent mixed-methods design Sample: 45 intervention group, and 41 a comparison group	Each 2.5-hour class session opened with a discussion of prescribed readings on relationship-centered care, spontaneity, curiosity, and	Empathy was greatly enhanced on five CARE items: ease, care, explain, help, and plan

Author	Methods	Intervention	Result
	Variable: interprofessional empathy Instrument: Interpersonal Reactivity Index [IRI], Consultative and Relational Empathy [CARE] measure and a facial expression recognition task	nonverbal communication. The instructor urged students to incorporate improv rules into their daily lives and led multiple improv games. At the conclusion of each lesson, students engaged in five minutes of freewriting about their recent experiences..	
Ding et.al.	Design: quasi -experimental design Sample: 250 nursing students Variable: empathy skills, communication skills and professional identity Instrument: Jefferson Scale of Empathy -Health Profession - Student, the Clinical Communication Competence Scale and the Professional Identity scale	The KSS module is intended for nursing students pursuing clinical experience in paediatrics. The components of KSS module are Knowledge, Simulation, and Sharing. Lectures, simulation scenarios, and self-reflection are all included in the internship. The module was presented over a five-day period over the internship's nine months.	the empathy had significantly higher scores after the internship