Natural history and epidemiology of cleft lip and palate: a registry-based study in Iran (2000-19)

Mahdi Biabani<sup>1</sup>, Saeed Dastgiri<sup>2</sup>, Elham Davtalab Esmaeili<sup>3</sup>

<sup>1</sup>School of Dentistry, Tabriz University of Medical Sciences, Tabriz, Iran.

<sup>2</sup> Tabriz Health Services Management Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

<sup>3</sup>Road Traffic Injury Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

#### **Corresponding author Address:**

Prof. Saeed Dastgiri

Tabriz Registry of Congenital Anomalies (TRoCA), School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran.

Post Code: 5166615739, Iran

Tel: 0098 914 415 7039

Fax: 0098 411 336 4673

Email: saeed.dastgiri@gmail.com

#### Abstract

The aim of this study was to provide the natural history and epidemiology of cleft lip and cleft palate in the northwest region of Iran between 2000 and 2019. Since 2000, infants born with birth defects have been registered in the Tabriz Registry of Congenital Anomalies (TRoCA). For this study, the information and data were collected using the TRoCA registry system. Prevalence of cleft lip and cleft palate was 1.48 (95% CI 1.34; 1.62) per 1000 live births over the past two decades in the region. The occurrence of cleft lip and cleft palate was more common in males than females. The fetal death ratio was 5 percent of live born children. The proportion of infants with cleft lip and cleft palate surviving to the second week was 54 percent. The results may have a role in planning and evaluating the strategies for primary prevention of cleft lip and cleft palate, particularly in high-risk populations.

Keywords: Cleft lip; Cleft palate; Epidemiology; Prevalence; Mortality; Survival

### Introduction

Cleft lip and palate is the most common major congenital anomalies affecting the orofacial area. It is the second most common defect in newborns in many places (1-4). Research reports indicate that both genetic and environmental factors contribute to the occurrence of these defects (5, 6).

Many studies reported different figures for the prevalence of cleft lip and cleft palate in different geographical areas of the world. Accordingly, Indian American communities showed the highest, and blacks the lowest occurrence rate of defect. This rate was reported higher in some of the Asian countries too (7). Prevalence rate of these anomalies was 1 (per 1000 births) in Iran. This is lower than other countries in the region (8).

Infants born with cleft lip and cleft palate have many problems including dental deformities and malocclusion as well as difficulty in chewing, swallowing, speaking and esthetic requiring rehabilitation and a team of specialists to repair (9). Assessing the prevalence of cleft lip and palate would be helpful in planning and providing health care services for these children, and the information obtained can also be used as a source for future researches (10). The main aim of this registry-based study was to provide the natural history and epidemiology of cleft lip and cleft palate in the northwest of Iran between 2000 and 2019.

### **Materials and Methods**

Since 2000, infants born with birth defects in the region have been registered in the Tabriz Registry of Congenital Anomalies (TRoCA). The registry continues to operate under the guidelines of the World Health Organization (WHO), and as a member of the International Clearinghouse for Birth Defects Surveillance and Research (ICBDSR).

For this study, the information and data for cleft lip and palate including gender, birth weight, maternal age at delivery, ABO and Rh blood groups, social status, and the presence of concomitant diseases and abnormalities were collected using the TRoCA registry system. The International Classification of Disease (ICD -10) was used for coding the congenital anomalies. For the purpose of data analysis, prevalence rates with 95 percent confidence intervals were calculated.

This study has received ethical approval from Tabriz University of Medical Science (IR.TBZMED.REC.1399.624).

### Results

A total of 291,569 live births were investigated by the TRoCA registry from which 431 cases were identified as having cleft lip and cleft palate presenting a total prevalence rate of 1.48 (95% CI 1.34; 1.62) per 1000 live births over the past two decades in the region. The prevalence rate of these anomalies during the first 10 years (2000-09) and the second 10 years (2010-19) estimated 1.05 (95% CI 0.91; 1.19) and 2.46 (95% CI 2.13; 2.78) per 1000 live births. The prevalence rate of cleft lip with/without cleft palate was 0.94 (95% CI 0.82; 1.05) and the prevalence rate of isolated cleft palate was 0.54 (95% CI 0.45; 0.62) per 1000 live births. Figure 1 and 2 shows the prevalence rate and the time trend of cleft lip with/without cleft palate and isolated cleft palate in the area during the study period.



Fig 1. The prevalence (95 percent CI) and the time trend of isolated cleft palate in 1000 live births in the northwest of Iran (2000-19)



Fig 2. The prevalence (95 percent CI) and the time trend of cleft lip with/without cleft palate in the northwest of Iran (2000-19)

The occurrence of cleft lip and cleft palate was more common in males than females while the prevalence of isolated cases was higher in females. The occurrence of these defects by newborn weight, maternal age at delivery, social status, and maternal blood type are shown in table 1.

Variable	ariable		Isolated cleft palate		Cleft lip with/without cleft palate	
		Frequency	(Percent)	Frequency	(Percent)	
Gender	Male	42	41.6	99	60	
	Female	59	58.4	66	40	
Consanguinity	No relation	52	67.5	77	65.8	
	First cousin	13	16.9	19	16.3	
	Second cousin	12	15.6	21	17.9	
	A+	15	30	26	29.5	
Blood group	A-	4	8	4	4.5	
	B+	12	24	18	20.4	
	B-	0	0	1	1.3	
	O+	10	20	26	29.5	
	0-	2	4	3	3.4	
	AB+	7	14	10	11.4	
	AB-	0	0	0	0	
Birth weight	Less than 1000gr	3	10	5	10.9	
	1001 to 2500 gr	27	90	39	84.8	
	More than 2500gr	0	0	2	4.3	
	Less than 18	0	0	2	2.1	
Maternal age	18 to 35	43	91.5	91	95.8	
-	More than 35	4	8.5	2	2.1	
Urban or rural	Urban area	74	73.3	112	71.8	
	village	27	26.7	44	28.2	
Concomitant	Yes	26	25	22	14	
anomalies	No	78	75	136	86	

 Table 1. Characteristics of the infant with cleft lip and cleft palate in the northwest of Iran

 (2000-19)

A total of 52 (67.5 %) and 77 (65.5 %) of parents of isolated cases with cleft palate and cleft lip with/without cleft palate had no consanguineous marriage, respectively. The A+ and O+ blood groups were more common types among cleft lip with/without cleft palate. Type A+ was reported in 15 (30 %) of isolated cleft palate. Birth weight of 1000-2500 grams observed in 27 (90 %) and 39 (84.8 %) of isolated cleft palate and cleft lip with/without cleft palate cases, respectively. A number of 43 (91.5 %) of isolated cleft palate and 91 (95.8 %) of cleft lip with/without cleft palate born from mother aged 18 to 35.

Fetal death ratio was 5 percent of live births. Mortality rate of these groups of anomalies was 5 percent among all birth defects. The proportion of infants with cleft palate and cleft lip surviving to the second week of life was 54 percent.

### Discussion

In this study, the total prevalence rate of cleft lip and cleft palate showed an increasing trend over the past two decades in the region. This rate is two times more than the rate reported from Iran by Khazaei et al (8). Although similar studies conducted in different region of Iran, reported prevalence rates of these anomalies lower than of the current investigation, Jalili et al in the study from the capital city, Tehran reported similar figure with us (11-15).

Saudi Arabia (0.3 per 1000 live births) and Jordan (1.4 per 1000 live births) have a lower prevalence rate than ours whereas Pakistani rate was higher with 1.9 per 1000 live births (16-18). Beside similar figures reported in many other studies, Greg et al and Fitzpatrick et al reported different rates (19-23). Environmental factors, ethnic variations, maternal nutrition, and consanguineous marriages might have a role in explaining these differences.

Most of the infants born with these anomalies were in the weight range of 1000-2500 grams. We had no control group to compare, however, Lei et al found in Taiwan a significant relationship between the birth of babies who weigh less than 1500 grams and the prevalence of facial clefts (24). The maternal age of infants with clefts was mostly between 18 and 38 years which might be due to the higher frequency of the most the deliveries at this age range of mothers. Bille et al reported that older age of parents might have a role as a risk factor for cleft lip and palate (25).

## Conclusion

With the control of infectious disease and malnutrition, congenital anomalies including cleft lip and cleft palate are now making a proportionally greater contribution to the ill health in childhood in many countries. Our finding may be useful in identifying clues to the etiology of cleft lip and cleft palate. Once identified, the influencing factors may be prevented by proper intervention in the early stage of occurrence. The results may also have a role in planning and evaluating the strategies for primary prevention of cleft lip and cleft palate, particularly in high-risk populations.

## Acknowledgments

We gratefully thank all the children and mothers in the region for their assistance in TRoCA programme over the past two decades. This study has been funded by the Tabriz University of Medical Sciences (IR.TBZMED.REC.1399.624). The authors report no conflict of interest.

# References

1. Saal HM. Cld'ssification dnd description of nonsyndromic clefts. Cleft lip and palate: from origin to treatment 2002: 47.

2. Derijcke A, Eerens A, Carels C. The incidence of oral clefts: A review. British Journal of Oral and Maxillofacial Surgery 1996;34: 488-94.

3. Jalili D, Fathi M, Jalili C. Frequency of cleft lip and palate among live births in Akbar Abadi Hospital. Acta Medica Iranica. 2012:704-6.

4. Mossey PA, Little J, Munger RG, Dixon MJ, Shaw WC. Cleft lip and palate. The Lancet 2009;374: 1773-85.

5. Murray JC. Face facts: Genes, environment, and clefts. American journal of human genetics 1995;57: 227.

6. Jones MC. Etiology of facial clefts: Prospective evaluation of 428 patients. The Cleft palate journal 1988;25: 16-20.

7. Panamonta V, Pradubwong S, Panamonta M, Chowchuen B. Global birth prevalence of orofacial clefts: A systematic review. J Med Assoc Thai 2015;98: S11-21.

8. Khazaei S, Shirani AM, Khazaei M, Najafi F. Incidence of cleft lip and palate in iran. Saudi Med J 2011;32: 390-93.

9. Donald M. Re and every, dr dentistry for the child and adolescent. St Louis: Mosby Year Book Inc 2016.

10. Hammond M, Stassen L. Do you care? A national register for cleft lip and palate patients. British journal of plastic surgery 1999;52: 12-17.

11. Yassaei S, Mehrgerdy Z, Zareshahi G. Prevalence of cleft lip and palate in births from 2003-2006 in iran. Community dental health 2010;27: 118-21.

12. Golalipour MJ, Mirfazeli A, Behnampour N. Birth prevalence of oral clefting in northern iran. The Cleft palate-craniofacial journal 2007;44: 378-80.

13. Zandi M, Heidari A. An epidemiologic study of orofacial clefts in hamedan city, iran: A 15-year study. The Cleft palate-craniofacial journal 2011;48: 483-89.

14. Rajabian MH, Aghaei S. Cleft lip and palate in southwestern iran: An epidemiologic study of live births. Annals of Saudi medicine 2005;25: 385-88.

15. Jalili D, Fathi M, Jalili C. Frequency of cleft lip and palate among live births in akbar abadi hospital. Acta Medica Iranica 2012: 704-06.

16. Kumar P, Hussain MT, Cardoso E, Hawary MB, Hassanain J. Facial clefts in saudi arabia: An epidemiologic analysis in 179 patients. Plastic and reconstructive surgery 1991;88: 955-58.

17. Al Omari F, Al-Omari I. Cleft lip and palate in jordan: Birth prevalence rate. The Cleft palate-craniofacial journal 2004;41: 609-12.

18. Elahi MM, Jackson IT, Elahi O, Khan AH, Mubarak F, Tariq GB, et al. Epidemiology of cleft lip and cleft palate in pakistan. Plastic and reconstructive surgery 2004;113: 1548-55.

19. Farhud D, Walizadeh G-R, Kamali MS. Congenital malformations and genetic diseases in iranian infants. Human genetics 1986;74: 382-85.

20. Kim S, Kim WJ, Oh C, Kim JC. Cleft lip and palate incidence among the live births in the republic of korea. Journal of Korean medical science 2002;17: 49.

21. Jensen BL, Kreiborg S, Dahl E, Fogh-Andersen P. Cleft lip and palate in denmark, 1976-1981: Epidemiology, variability, and early somatic development. Cleft Palate J 1988;25: 258-69.

22. Gregg T, Boyd D, Richardson A. The incidence of cleft lip and palate in northern ireland from 1980–1990. British Journal of Orthodontics 1994;21: 387-92.

23. FitzPatrick D, Raine P, Boorman J. Facial clefts in the west of scotland in the period 1980-1984: Epidemiology and genetic diagnoses. Journal of Medical Genetics 1994;31: 126-29.

24. Lei R-L, Chen H-S, Huang B-Y, Chen Y-C, Chen PK-T, Lee H-Y, et al. Population-based study of birth prevalence and factors associated with cleft lip and/or palate in taiwan 2002–2009. PLoS One 2013;8: e58690.

25. Bille C, Skytthe A, Vach W, Knudsen LB, Andersen A-MN, Murray JC, et al. Parent's age and the risk of oral clefts. Epidemiology (Cambridge, Mass) 2005;16: 311.