

Modern contraceptive Use and its associated factors among married women in Senegal: Based on the recent National Demographic and Health Survey Data

Bezawit Mulat ^{1*}, Menen Tsegaw ³, Kegniet Shitu ²

¹Department of Human Physiology, School of Medicine, College of Medicine and Health Sciences, University of Gondar, Gondar, Ethiopia

²Department of Health Education and Behavioral Sciences, Institute of Public Health, College of Medicine and Health Sciences, University of Gondar, Gondar, Ethiopia

³Department of Public Health, College of Medicine and Health Sciences, Ambo University, Ambo, Ethiopia

*Corresponding Author

Bezawit Mulat

E-mail: mulatbezawit@gmail.com

Cell Phone: +251963729635

19

20 **Abstract**

21 **Objective**

22 This study aimed to assess the prevalence and associated factors of modern contraceptive use
23 among married women in Senegal.

24 **Method**

25 The most recent Demographic and Health Survey (DHS), which was conducted in 2019 in
26 Senegal was used. For this study 5,659 women were included. Multivariable binary Logistic
27 regression analysis was used to identify factors associated with women's contraceptive use.

28 **Result**

29 According to the study, 25.5% (95%CI (24.4, 26.7)) of married women in Senegal uses modern
30 contraceptive methods. Being currently working (AOR=1.27, 95% CI (1.11, 1.46)), husband
31 educational level (primary educational level (AOR=1.32 ,95% CI (1.07,1.61), higher
32 educational levels (AOR= 1.83 , 95% CI (1.32, 2.54)), wealth index (middle (AOR=1.37,95%
33 CI (1.15,1.63) and rich (AOR=1.34,95% CI (1.11,1.62)), exposure to family planning messages
34 (AOR=1.29,95% CI (1.13,1.48)), having ANC visits(AOR=2.01, 95% CI(1.13,3.57)) and not
35 wanting a child (AOR=2.35, 95%CI (1.90,2.95)) were positively and significantly with women's
36 contraceptives use. On the other hand, number of children (having no child (AOR=0.03, 95% CI
37 (0.01,0.06)), 1-2 children (AOR=0.47,95% CI(0.38,0.59) and 3-4 children (AOR=0.79, 95%
38 CI(0.67,0.96)), women who thought ≥ 5 children are ideal (AOR=0.73,95% CI(0.61,0.88)) , and

had a desire for more children after 2 years (AOR=2.19, 95% CI(1.82,2.64) were negatively and significantly with women's contraceptives use.

Conclusion: contraceptive utilization was low among married women in Senegal. Women's contraceptive use was affected by family planning messages exposure, sociodemographic, and obstetric factors. Thus, a tailored behavior change intervention is required to increase contraceptive utilization among married women in Senegal.

Key words: Modern contraceptive use, Married women, Senegal

Strengths and limitations of this study

- The inclusion of nationally representative data and a large sample size, which strengthened the generalizability of the findings, was the study's key strength.
- Sample weighing was employed to compensate unequal sample allocations and non-responses.
- It may not be possible to establish causal links due to the cross-sectional nature of the data.
- It's possible that crucial factors that influence women's use of modern contraceptives might be neglected because the study relied on secondary data.

Introduction

Contraception today includes a variety of options: oral contraceptive pills, implants, injectable, contraceptive patch and vaginal ring, intrauterine device (IDU), female and male condoms, female and male sterilization, vaginal barrier methods (including the diaphragm, cervical cap and spermicidal agents), lactational amenorrhea method (LAM), emergency contraception pills, standard days method (SDM), basal body temperature (BBT) method, Two-day method and sympto-thermal method (1). Modern contraceptive use is characterized as a person's or couple's ability to have as many children as they want, when they want and has been found to lower mother and child morbidity and death (2,3).

The use of modern contraceptives is critical for achieving SDG 3's goal 3.7, which states that by 2030, everyone in the world would have access to sexual and reproductive health care (4). Furthermore, in 1987, the Safe Motherhood Initiative established family planning as one of four pillars, alongside prenatal care, safe delivery, and postnatal care, to reduce maternal mortality in developing countries, where 99 percent of all maternal deaths occur (5).

According to the World Health Organization (WHO) report, nearly 308 million unwanted pregnancies were averted in 2017 (6). The increased use of contraceptive techniques has resulted in gains in not only health-related outcomes such as reduced maternal and newborn mortality, but also in educational and economic outcomes, particularly for girls and women (3,7).

In Africa the use of modern contraceptive varies across different countries. For instance, it is 62% in Zimbabwe, 6% in Guinea, 56%, Malawi and 56 % Kenya (8). Different studies stated that age , educational attainment of a woman , educational attainment of husbands , residence, wealth index of women, number of living children, having antenatal care visits (ANC), desire to space and want no more child are factors which have significant association with modern contraceptive utilization (8–10). Because of the large unmet need for contraception in Sub-Saharan African countries, the number of unplanned pregnancies is higher, estimated to be over 14 million per year (11). According to reports, 31% of conceptions in Senegal are unintentional, with 24 percent of all unwanted pregnancies ending in induced abortions, 60% in unplanned births, and 16% in miscarriages (12) . All the aforementioned adverse outcomes would be prevented using contraceptives. However, there is a scarcity of data on the prevalence and associated aspects of modern contraception use in Senegal. As a result, the goal of this study is to determine the prevalence of modern contraceptive use and the factors that influence it among married women in Senegal.

Methods

Study setting, data source, and study design

Senegal is a West African country that borders the North Atlantic Ocean to the west. It is bordered on the north by Mauritania along the Senegal River, on the east by Mali, on the south by Guinea and Guinea-Bissau, and on the west by Mauritania along the Gambia River (13). Currently the population of the Senegal is estimated that 17,340,913 (14).

This analysis employed based on the data from the 2019 Continuous-Demographic and Health Survey (C-DHS) of Senegal .The C-DHS is a nationally representative data set developed by the

Senegalese Agency of Statistics and Demography (ANSD) and the survey lasts from April 2019 - December 2019. The research participants are chosen using a two-stage stratified sampling technique. This data set (IR file) is consist of information collected from all eligible women aged 15-49 years and the current study excludes unmarried women and employed with a total weighted sample of 5,659 married reproductive age women. An authorization letter for the use of this data was obtained from DHS program and the dataset was downloaded from DHS website www.measuredhs.com

Study variables

Dependent variable

The study's outcome variable was the use of modern contraceptive methods among married reproductive age women in Senegal. The variable was catigorized into two categories: 1 = “Use modern contraceptive” and 0 =”did not use modern contraceptive”.

Independent variables

In this study the independent variables included: sociodemographic factors (Age, religion, residence, educational status of women, educational status of husband, and current working (employment) status, wealth index, media exposure to family planning methods and obstetric factors: complications during previous pregnancies (terminated pregnancy), presence of antenatal care visits, number of living children, ideal number of children that the family want and desire for more children.

Data processing and Analysis

Individual records (IR) files were used to extract data, which was then coded and transformed using STATA version 14 statistical software. To account for differential chance of selection and non-response in the original survey, weighted samples were used for analysis. The presence of statistical significance was determined using a multivariable logistic regression analysis. It was fitted after the model's fitness was evaluated using the Hosmer and Lemeshow goodness of fit test. The variance inflation factor (VIF) was also used to analyze multicollinearity across the explanatory components, and it was found to be within acceptable range (1-5)(15). A p-value less than 0.05 is used to evaluate the presence of a meaningful effect or relationship of independent factors with the outcome variable.

Patient public involvement

The public was not involved in the design or conduct, choice of outcome measures and recruitment of the study.

Result

A total of 5,659 married women were included in the study. The mean age of the participants was 31 (\pm 8.5) years with an age range of 15-49 years. The majority (98.2%) of the mothers were Muslims and 3357 (59.3 %) were from a rural area. Concerning the reproductive history (98.6%) of women have ANC visits (table 1).

Table 1 Sociodemographic and reproductive characteristics of married women in Senegal (n = 5,659).

Variable	Category	Frequency	Percent
----------	----------	-----------	---------

Age (in years)	15-24	1,340	23.7
	25-34	2,216	39.2
	35-49	2,103	37.1
Residence	Rural	3,357	59.3
	Urban	2,302	40.7
Religion	Muslim	5,556	98.2
	Christian	103	1.8
Educational status of respondents	No formal education	3,443	60.8
	Primary	1,177	20.8
	Secondary	857	15.2
	Higher	182	3.2
Educational status of the husband/partner	No formal education	3,615	63.9
	Primary	640	11.3
	Secondary	635	11.2
	Higher	330	5.8
	Don't know	439	7.8
Wealth index	Poor	2,298	40.6
	Middle	1,103	19.5
	Rich	2,258	39.9
Current working status	Currently working	2,904	51.3
	Currently not working	2,755	48.7
Mass media exposure to family planning methods	Have exposure	3,375	59.6
	Haven't exposure	2,284	40.4
Ideal number of children	≤4	1,190	21.0
	≥5	3,364	59.5
	Non-numeric response	1,105	19.5
Desire for more children	Wants within 2 years	1,602	28.3
	Wants after 2 years	2,450	43.3
	Wants, unsure timing	325	5.7
	Undecided	84	1.5
Number of living children	Doesn't want	1,198	21.2
	No child	718	12.7
	1-2 children	1,922	34.0

	3-4 children	1,485	26.2
	5 or more children	1,534	27.1
Have ANC visit	Yes	5,578	98.6
	No	81	1.4
Ever had terminated pregnancy	Yes	1,200	20.2
	No	4,733	79.8
Decision maker for using modern contraceptive	Women	295	20.4
	Husband	240	16.6
	Joint decision	893	62.0
	Other	16	1.0

Modern contraceptive use among married women

In this study the prevalence of modern contraceptive use among married women in Senegal was 25.5% with (95% CI (24.4, 26.7)). The use of modern contraceptive method is higher among age group of 25-34 as compared to age group of women 15-24. Additionally, the utilization of modern contraceptives was higher (30.5%) among women who are currently working. Furthermore, women who are exposed to family planning methods through the media are more likely to use modern contraception (29.5%).

Factors associated with modern contraceptive use

Based on the output of multivariable binary logistic regression analysis, the following explanatory variables had a statistically significant association with modern contraceptive use : current working status of women (currently working (AOR=1.27, 95% CI (1.11, 1.46))), Husband education (husbands who attended primary educational level (AOR=1.32 ,95% CI (1.07,1.61), husbands who attended higher educational levels (AOR= 1.83 , 95% CI (1.32, 2.54), women whose wealth index in middle(AOR=1.37,95% CI (1.15,1.63) and rich (AOR=1.34,95% CI

(1.11,1.62), women who had exposed to family planning methods through different medias (AOR=1.29,95% CI (1.13,1.48)), women who had ANC visits(AOR=2.01, 95% CI(1.13,3.57)). Plus to the above factors, women who had no child (AOR=0.03, 95% CI (0.01,0.06)), 1-2 children (AOR=0.47,95% CI(0.38,0.59) and 3-4 children (AOR=0.79, 95% CI(0.67,0.96)), women who thought ≥ 5 children are ideal(AOR=0.73,95% CI(0.61,0.88)) , and women who had a desire for more children after 2 years (AOR=2.19, 95% CI(1.82,2.64) and women who doesn't want a child (AOR=2.35, 95%CI (1.90,2.95)) were significantly associated with womens modern contraceptive use.

The odds of using modern contraceptive was increased by 27% among women who were currently working than their counterparts. The odds of using modern contraceptives was increased by 32% among women whose husband had primary level of education. Moreover, having a husband with primary and higher education level increased the odds of using modern contraceptive by 32% and 16% respectively, when compared to women having an illiterate husband. The odds of using modern contraceptives was increased by 29% among women who had exposure to family planning methods through different Medias (Table 2).

Table 2 Factors associated with modern contraceptive use among married women in Senegal (n=5,659).

167
168
169
170
171
172
173
174

Variable	Modern contraceptive use		p-value	AOR
	Yes (n=1,445)	No(n=4,214)		
	Frequency (%)	Frequency (%)		
Age				
15-24	221(16.5)	1,119(83.5)		1
25-34	613(27.7)	1,603(72.3)	0.8	0.97(0.78,1.2)
35-49	611(29.1)	1,492(70.9)	0.14	0.82(0.64,1.1)
Residence				
Urban	745(32.4)	1,557(67.6)		1
Rural	700(20.9)	2,657(79.1)	0.06	0.85(0.72,1.00)
Religion				
Muslim	1,414(25.5)	4,142(74.5)		1
Christian	31 (29.7)	72 (70.3)	0.92	0.98 (0.62,1.55)
Womens' educational level				
No formal education	754(21.9)	2,690(78.1)	0.05	0.61(0.37,1.00)
Primary	374(31.8)	803(68.2)	0.58	0.87(0.53,1.42)
Secondary	240(28.0)	617(72.0)	0.54	0.86(0.53,1.40)
Higher	77(42.3)	105(57.7)		1
Current working status				
Currently working	841(30.5)	1,913(69.5)	0.00	1.27 (1.11,1.46)

Currently not working	604(20.8)	2,301(79.2)		1
Husband education				
No formal education	777(21.5)	2838(78.5)		1
Primary education	192(29.9)	448(70.1)	0.008	1.32 (1.07,1.61)
Secondary education	192(30.2)	443(69.8%)	0.19	1.16 (0.93,1.45)
Higher	151(45.9)	179(54.1)	0.00	1.83 (1.32,2.54)
Don't know	134(30.4)	306(69.6)	0.28	1.16 (0.89,1.51)
Wealth index				
Poor	434(18.9)	1863(81.1)		1
Medium	301(27.3)	802(72.7)	0.00	1.37 (1.15,1.63)
Rich	709(31.4)	1549(68.6)	0.00	1.34 (1.11,1.62)
Media exposure				
Exposed	995(29.5)	2,381(70.5)	0.00	1.29 (1.13,1.48)
Non exposed	450(19.7)	1,833(80.3)		1
Have ANC visits				
Yes	1436(25.8)	4,142(74.5)	0.01	2.01 (1.13,3.57)
No	9(10.7)	72(89.3)		1
Ever had terminated pregnancy				
Yes	312(26.3)	877(73.7)	0.41	0.94 (0.79,1.09)
No	1,132(25.3)	3,337(74.7)		1
Number of living children				
No children	8(1.1)	711(98.9)	0.00	0.03 (0.01,0.06)
1-2 children	474(24.7)	1,448(75.3)	0.00	0.47 (0.38,0.59)
3-4 children	470(31.7)	1,014(68.3)	0.02	0.79 (0.67,0.96)
≥5 children	493(32.1)	1,041(67.9)		1
Ideal number of children				
≤ 4 children	381(32.0)	809(68.0)		1
≥ 5 children	854(25.4)	2,509(74.6)	0.001	0.73(0.61,0.88)
Non-numeric response	210(19.0)	895(81.0)	0.00	0.52(0.42,0.66)
Desire for more children				

Want's within 2 years	195(12.2)	1,407(87.8)		1
Want's after 2 years	750(30.6)	1,700(69.4)	0.00	2.19(1.82,2.64)
Want's, unsure timing	37(11.4)	288(88.6)	0.49	1.13(0.79,1.60)
Undecided	14(17.4)	70(82.6)	0.40	1.27(0.72,2.24)
Doesn't want	447(37.4)	750(62.6)	0.00	2.35(1.90,2.95)

Discussion

This study assessed the prevalence and associated factors of modern contraceptive use among Senegalese women. The prevalence of modern contraceptive use was 25.5% with (95% CI (24.4, 26.7)). Womens' working status, husband education, wealth index, having media exposure, having ANC visit, number of living children, ideal number of children and desire for more children were factors that have statistically significant association with modern contraceptive use.

In the present study, the use of modern contraceptives among married Senegalese women was 25.5% (95% CI (24.4, 26.7)). The result is in harmony with the study conducted in Senegal (26.3%) (16). This alignment might be explained by similarities in sociodemographic characteristics of the population.

The result of the current study is higher than the studies conducted in Bale zone, south East Ethiopia (20.8%)(17) and Metekel zone North West Ethiopia (18.6%)(18). The discrepancy might be explained by socio-cultural difference between the countries that could affect their awareness on contraceptive use (19). On the other hand the result of the present study is lower than studies done in North Showa, Ethiopia (46.9%) (20) Zambia (43%)(21), and Egypt (46.5%) (22).

In the present study women who are currently enrolled in different jobs (currently working) had significant association with modern contraceptive use. This result is supported by study conducted in Ethiopia (19). This could be owing to the fact that women who work and have a private job have a higher income, greater access to the media, and better health care, all of which have a beneficial impact on modern contraception use (23–26).

In the current study, modern contraceptive use among married Senegalese women was significantly associated with husband's educational attainment, which is supported by a study conducted in Senegal (16,27,28). It's probable that educated husbands are aware of the benefits of modern contraception since they have a greater chance to read newspapers, watch television, and use social media. The other factor that had significant association with contraceptive use was wealth index of a woman. Women from households with a medium or high wealth index were more likely to use modern contraception than those from households with a low wealth index. The finding was in line with a study conducted in Ethiopia (29).

Modern contraceptive use was also significantly associated with womens' media exposure. This outcome is obedient with studies conducted in Senegal (16) and Ethiopia (9). This alignment could be explained by the fact that media can help communities raise awareness about health issues as well as address social and cultural issues (30,31). This critical role of the media can aid in lowering barriers to health-care access and use, including contraception.

In this study presence of ANC visits have significant association with modern contraceptive use. The result is supported by a multicenter study done in African countries (8). It's possible that the increased prevalence of contraceptive use among women who had ANC visits is due to the fact that antenatal care provides a unique opportunity to interact with, educate, and provide family

support to women of all socioeconomic backgrounds, which facilitates the uptake of modern contraceptives. The study also showed that women who hadn't child have lower tendency to use modern contraceptives than women who had three and more than three children. This result is supported by studies done in Senegal (16,32). This meant that the more children a woman has, the more likely she is to want to space or limit the number of children, and the more contraceptive she is likely to use.

Furthermore, the present study also revealed the presence of statistically significant association between modern contraceptive use, and womens' perception for ideal number of children they would have. The result is in harmony with a study done in Senegal (16). According to the study, women who wished five or more children had a lower likelihood of taking modern contraception than women who desired four or fewer children. Finally, the results of the current study also showed that womens' desire for more children also affects womens' modern contraceptive utilization. That is the odds of using modern contraceptives among women who wants child within 2 years lower than women who wants child after 2 years and totally doesn't want a child at all.

The study's main strength was the use of nationally representative data and a large sample size, which increased the generalizability of its conclusions. The DHS study, on the other hand, has flaws in that it relies on respondents' self-reporting and may be prone to recall bias because women were asked to recall past incidents. Furthermore, the design of this study limits its ability to demonstrate causation between the desired outcome and these critical independent factors.

Conclusion and recommendation

According to the findings, the usage of modern contraceptive methods among married women in Senegal was low. Women's contraceptive use was affected by family planning messages exposure, sociodemographic, and obstetric factors. Thus, a tailored behavior change intervention is required to increase contraceptive utilization among married women in Senegal and such interventions should take those factors into account.

Ethics approval and consent to participate

The study is based on DHS data which was collected by DHS program, the DHS program collect the data by gaining ethical approval from national IRB. Thus, we didn't received any ethical approval since the data used for the study is from a secondary source. However, we got the authentication letter from the DHS program to use the data set for this study. The study was conducted under the Declaration of Helsinki.

Consent for publication

Not applicable.

Availability of data and materials

All result-based data are available within the manuscript and anyone can access the data set online from www.measuredhs.com

Competing interests

The authors declare that they have no competing interests.

Funding

No, any funding was received from any organization.

Authors' contributions

All authors made substantial contributions to conception, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to the current journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Acknowledgments

We would like to acknowledge the MEASURE DHS program for permitting us to obtain and use DHS data sets of Senegal.

Reference

1. Contraceptive prevalence – use of modern methods (%)(Health Equity Monitor) [Internet]. Available from: <https://www.who.int/data/gho/indicator-metadata-registry/imr-details/3334>

- 276 2. Akintade OL, Pengpid S, Peltzer K. Awareness and use of and barriers to family planning
277 services among female university students in Lesotho. S Afr J Obstet Gynaecol.
278 2011;17(3):72–8.
- 279 3. UNDESA. World Family Planning 2017 Highlights. Econ Soc Aff [Internet]. 2017;1–43.
280 Available from:
281 [https://www.un.org/en/development/desa/population/publications/pdf/family/WFP2017_H](https://www.un.org/en/development/desa/population/publications/pdf/family/WFP2017_Highlights.pdf)
282 [ighlights.pdf](https://www.un.org/en/development/desa/population/publications/pdf/family/WFP2017_Highlights.pdf)
- 283 4. United Nations (UN). Sustainable Development Goals (SDG 5) | United Nations Western
284 Europe [Internet]. United Nations. 2020. Available from: [https://unric.org/en/sdg-](https://unric.org/en/sdg-12/%0Ahttps://unric.org/en/sdg-5/)
285 [12/%0Ahttps://unric.org/en/sdg-5/](https://unric.org/en/sdg-5/)
- 286 5. Ahmed S, Li Q, Liu L, Tsui AO. Maternal deaths averted by contraceptive use: An
287 analysis of 172 countries. Lancet [Internet]. 2012;380(9837):111–25. Available from:
288 [http://dx.doi.org/10.1016/S0140-6736\(12\)60478-4](http://dx.doi.org/10.1016/S0140-6736(12)60478-4)
- 289 6. WHO. WHO recommendations on self-care interventions. Human Reproduction
290 Programme. 2020.
- 291 7. Darroch JE. Adding it up: Investing in Contraception and Maternal and Newborn Health
292 for Adolescents in Ethiopia , 2018. Guttmacher Institute. 2018.
- 293 8. Apanga PA, Kumbeni MT, Ayamga EA, Ulanja MB, Akparibo R. Prevalence and factors
294 associated with modern contraceptive use among women of reproductive age in 20
295 African countries: A large population-based study. BMJ Open. 2020;10(9).
- 296 9. Asresie MB, Fekadu GA, Dagne GW, Gelaw YM. Modern Contraceptive Use and

Influencing Factors in Amhara Regional State: Further Analysis of Ethiopian Demographic Health Survey Data 2016. Adv Public Heal. 2020;2020(Mc).

10. Antarini A. Factors influencing the use of modern contraception among reproductive aged women in bangka belitung province, indonesia. Pan Afr Med J. 2021;39.

11. Kwabena E, Id A, Budu E, Sambah F, Baatiema L, Appiah F, et al. Prevalence and determinants of unintended pregnancy in sub-Saharan Africa: A multi- country analysis of demographic and health surveys. PLoS One. 2019;14(8):1–16.

12. Abortion in Senegal – Guttmacher Institute [Internet]. Available from: <https://www.guttmacher.org/fact-sheet/abortion-senegal>

13. Senegal - Republic of Senegal - Western Africa [Internet]. Available from: <https://www.nationsonline.org/oneworld/senegal.htm>

14. senegal-population @ worldpopulationreview.com [Internet]. Available from: <https://worldpopulationreview.com/countries/senegal-population>

15. Choueiry G. What is an Acceptable Value for VIF? (With References) – Quantifying Health [Internet]. 2020. Available from: <https://quantifyinghealth.com/vif-threshold/>

16. Zegeye B, Ahinkorah BO, Idriss-Wheeler D, Olorunsaiye CZ, Adjei NK, Yaya S. Modern contraceptive utilization and its associated factors among married women in Senegal: a multilevel analysis. BMC Public Health. 2021;21(1):1–13.

17. Belda SS, Haile MT, Melku AT, Tololu AK. Modern contraceptive utilization and associated factors among married pastoralist women in Bale eco-region, Bale Zone, South East Ethiopia. BMC Health Serv Res. 2017;17(1):1–12.

18. Adane AA, Bekele YA, Melese E, Worku GT, Netsere HB. Modern Contraceptive Utilization and Associated Factors among Married Gumuz Women in Metekel Zone North West Ethiopia. Biomed Res Int. 2020;2020.
19. Gebre MN, Edossa ZK. Modern contraceptive utilization and associated factors among reproductive-age women in Ethiopia: Evidence from 2016 Ethiopia demographic and health survey. BMC Womens Health. 2020;20(1):1–14.
20. A. M, D. W, A. F, B. M. Determinants of modern contraceptive utilization among married women of reproductive age group in North Shoa Zone, Amhara Region, Ethiopia. Reprod Health [Internet]. 2014;11(1):1–7. Available from: <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed12&NEWS=N&AN=2014111648>
21. Lasong J, Zhang Y, Gebremedhin SA, Opoku S, Abaidoo CS, Mkandawire T, et al. Determinants of modern contraceptive use among married women of reproductive age: A cross-sectional study in rural Zambia. BMJ Open. 2020;10(3):1–10.
22. Unit FH. Practice of Family Planning among Married Female Attendants to Shawa Family Health Unit, Dakahlia, Egypt. Egypt Fam Med J. 2020;4(1):24–41.
23. Tekelab T, Melka AS, Wirtu D. Predictors of modern contraceptive methods use among married women of reproductive age groups in Western Ethiopia: A community based cross-sectional study. BMC Womens Health [Internet]. 2015;15(1):1–8. Available from: <http://dx.doi.org/10.1186/s12905-015-0208-z>
24. Ofonime JE. Determinants of modern contraceptive uptake among Nigerian women:

Evidence from the national demographic and health survey. Afr J Reprod Health
[Internet]. 2017;21(3):89–95. Available from:

<https://www.scopus.com/inward/record.uri?eid=2-s2.0-85038842247&partnerID=40&md5=c36f097c56e5413d445dcaaf7d7a30a1>

25. Osmani AK, Reyer JA, Osmani AR, Hamajima N. Factors influencing contraceptive use among women in Afghanistan: Secondary analysis of Afghanistan Health Survey 2012. Nagoya J Med Sci. 2015;77(4):551–61.

26. Islam AZ, Mondal NI, Khatun ML. Prevalence and Determinants of Contraceptive use among Employed and Unemployed Women in Bangladesh. 2016;5(2):92–102.

27. Tessema ZT, Teshale AB, Tesema GA, Yeshaw Y, Worku MG. Pooled prevalence and determinants of modern contraceptive utilization in East Africa: A Multi-country Analysis of recent Demographic and Health Surveys. PLoS One [Internet]. 2021;16(3 March):1–16. Available from: <http://dx.doi.org/10.1371/journal.pone.0247992>

28. Gebremariam A. Modern Contraceptive Method Mix and Factors Affecting Utilization of Modern Contraceptives among Married Women in Adigrat Town, Tigray, Northern Ethiopia. Fam Med Med Sci Res. 2014;03(04).

29. Lakew Y, Reda AA, Tamene H, Benedict S, Deribe K. Geographical variation and factors influencing modern contraceptive use among married women in Ethiopia: evidence from a national population based survey. Reprod Health. 2013;1–10.

30. Ayanore MA, Pavlova M, Groot W. Unmet reproductive health needs among women in some West African countries: A systematic review of outcome measures and

determinants. Reprod Health [Internet]. 2016;13(1):1–10. Available from:
<http://dx.doi.org/10.1186/s12978-015-0104-x>

31. Ansary R, Anisujjaman M. Factors Determining Pattern of Unmet Need for Family
 Planning in. Int Res J Soc Sci. 2012;1(4):16–23.

32. Bulto GA, Zewdie TA, Beyen TK. Demand for long acting and permanent contraceptive
 methods and associated factors among married women of reproductive age group in Debre
 Markos Town, North West Ethiopia. BMC Womens Health. 2014;14(1).