

1 **Modern contraceptive Use and its associated factors among married women**
2 **in Senegal: Based on the recent National Demographic and Health Survey**
3 **Data**

4 Bezawit Mulat ^{1*}, Menen Tsegaw ³, Kegnie Shitu ²

5 ¹Department of Human Physiology, School of Medicine, College of Medicine and Health
6 Sciences, University of Gondar, Gondar, Ethiopia

7 ²Department of Health Education and Behavioral Sciences, Institute of Public Health, College of
8 Medicine and Health Sciences, University of Gondar, Gondar, Ethiopia

9 ³Department of Public Health, College of Medicine and Health Sciences, Ambo University,
10 Ambo, Ethiopia

11

12 ***Corresponding Author**

13 Bezawit Mulat

14 E-mail: mulatbezawit@gmail.com

15 Cell Phone: +251963729635

16

17

18

19

20 **Abstract**

21 **Objective**

22 This study aimed to assess the prevalence and associated factors of modern contraceptive use
23 among married women in Senegal.

24 **Method**

25 The most recent Demographic and Health Survey (DHS), which was conducted in 2019 in
26 Senegal was used. For this study 5,659 women were included. Multivariable binary Logistic
27 regression analysis was used to identify factors associated with women's contraceptive use.

28 **Result**

29 According to the study, 25.5% (95%CI (24.4, 26.7)) of married women in Senegal uses modern
30 contraceptive methods. Being currently working (AOR=1.27, 95% CI (1.11, 1.46)), husband
31 educational level (primary educational level (AOR=1.32 ,95% CI (1.07,1.61), higher
32 educational levels (AOR= 1.83 , 95% CI (1.32, 2.54)), wealth index (middle (AOR=1.37,95%
33 CI (1.15,1.63) and rich (AOR=1.34,95% CI (1.11,1.62)), exposure to family planning messages
34 (AOR=1.29,95% CI (1.13,1.48)), having ANC visits(AOR=2.01, 95% CI(1.13,3.57)) and not
35 wanting a child (AOR=2.35, 95%CI (1.90,2.95)) were positively and significantly with women's
36 contraceptives use. On the other hand, number of children (having no child (AOR=0.03, 95% CI
37 (0.01,0.06)), 1-2 children (AOR=0.47,95% CI(0.38,0.59) and 3-4 children (AOR=0.79, 95%
38 CI(0.67,0.96)), women who thought ≥ 5 children are ideal (AOR=0.73,95% CI(0.61,0.88)) , and

39 had a desire for more children after 2 years (AOR=2.19, 95% CI(1.82,2.64) were negatively and
40 significantly with women's contraceptives use.

41 **Conclusion:** contraceptive utilization was low among married women in Senegal. Women's
42 contraceptive use was affected by family planning messages exposure, sociodemographic, and
43 obstetric factors. Thus, a tailored behavior change intervention is required to increase
44 contraceptive utilization among married women in Senegal.

45 **Key words: Modern contraceptive use, Married women, Senegal**

46

Strengths and limitations of this study

- The inclusion of nationally representative data and a large sample size, which strengthened the generalizability of the findings, was the study's key strength.
- Sample weighing was employed to compensate unequal sample allocations and non-responses.
- It may not be possible to establish causal links due to the cross-sectional nature of the data.
- It's possible that crucial factors that influence women's use of modern contraceptives might be neglected because the study relied on secondary data.

47

48

49

50 **Introduction**

51 Contraception today includes a variety of options: oral contraceptive pills, implants, injectable,
52 contraceptive patch and vaginal ring, intrauterine device (IDU), female and male condoms,
53 female and male sterilization, vaginal barrier methods (including the diaphragm, cervical cap and
54 spermicidal agents), lactational amenorrhea method (LAM), emergency contraception pills,
55 standard days method (SDM), basal body temperature (BBT) method, Two-day method and
56 sympto-thermal method (1). Modern contraceptive use is characterized as a person's or couple's
57 ability to have as many children as they want, when they want and has been found to lower
58 mother and child morbidity and death (2,3).

59 The use of modern contraceptives is critical for achieving SDG 3's goal 3.7, which states that by
60 2030, everyone in the world would have access to sexual and reproductive health care (4).
61 Furthermore, in 1987, the Safe Motherhood Initiative established family planning as one of four
62 pillars, alongside prenatal care, safe delivery, and postnatal care, to reduce maternal mortality in
63 developing countries, where 99 percent of all maternal deaths occur (5).

64 According to the World Health Organization (WHO) report, nearly 308 million unwanted
65 pregnancies were averted in 2017 (6). The increased use of contraceptive techniques has resulted
66 in gains in not only health-related outcomes such as reduced maternal and newborn mortality,
67 but also in educational and economic outcomes, particularly for girls and women (3,7) .

68 In Africa the use of modern contraceptive varies across different countries. For instance, it is
69 62% in Zimbabwe, 6% in Guinea, 56%, Malawi and 56 % Kenya (8). Different studies stated
70 that age , educational attainment of a woman , educational attainment of husbands , residence,
71 wealth index of women, number of living children, having antenatal care visits (ANC), desire to
72 space and want no more child are factors which have significant association with modern
73 contraceptive utilization (8–10). Because of the large unmet need for contraception in Sub-
74 Saharan African countries, the number of unplanned pregnancies is higher, estimated to be over
75 14 million per year (11). According to reports, 31% of conceptions in Senegal are unintentional,
76 with 24 percent of all unwanted pregnancies ending in induced abortions, 60% in unplanned
77 births, and 16% in miscarriages (12) . All the aforementioned adverse outcomes would be
78 prevented using contraceptives. However, there is a scarcity of data on the prevalence and
79 associated aspects of modern contraception use in Senegal. As a result, the goal of this study is to
80 determine the prevalence of modern contraceptive use and the factors that influence it among
81 married women in Senegal.

82 **Methods**

83 **Study setting, data source, and study design**

84 Senegal is a West African country that borders the North Atlantic Ocean to the west. It is
85 bordered on the north by Mauritania along the Senegal River, on the east by Mali, on the south
86 by Guinea and Guinea-Bissau, and on the west by Mauritania along the Gambia River (13).
87 Currently the population of the Senegal is estimated that 17,340,913 (14).

88 This analysis employed based on the data from the 2019 Continuous-Demographic and Health
89 Survey (C-DHS) of Senegal .The C-DHS is a nationally representative data set developed by the

90 Senegalese Agency of Statistics and Demography (ANSD) and the survey lasts from April 2019
91 - December 2019. The research participants are chosen using a two-stage stratified sampling
92 technique. This data set (IR file) is consist of information collected from all eligible women aged
93 15-49 years and the current study excludes unmarried women and employed with a total
94 weighted sample of 5,659 married reproductive age women. An authorization letter for the use of
95 this data was obtained from DHS program and the dataset was downloaded from DHS website
96 www.measuredhs.com

97 **Study variables**

98 **Dependent variable**

99 The study's outcome variable was the use of modern contraceptive methods among married
100 reproductive age women in Senegal. The variable was catigorized into two categories: 1 = “Use
101 modern contraceptive” and 0 =”did not use modern contraceptive”.

102 **Independent variables**

103 In this study the independent variables included: sociodemographic factors (Age, religion,
104 residence, educational status of women, educational status of husband, and current working
105 (employment) status, wealth index, media exposure to family planning methods and obstetric
106 factors: complications during previous pregnancies (terminated pregnancy), presence of
107 antenatal care visits, number of living children, ideal number of children that the family want and
108 desire for more children.

109 **Data processing and Analysis**

110 Individual records (IR) files were used to extract data, which was then coded and transformed
111 using STATA version 14 statistical software. To account for differential chance of selection and
112 non-response in the original survey, weighted samples were used for analysis. The presence of
113 statistical significance was determined using a multivariable logistic regression analysis. It was
114 fitted after the model's fitness was evaluated using the Hosmer and Lemeshow goodness of fit
115 test. The variance inflation factor (VIF) was also used to analyze multicollinearity across the
116 explanatory components, and it was found to be within acceptable range (1-5)(15). A p-value
117 less than 0.05 is used to evaluate the presence of a meaningful effect or relationship of
118 independent factors with the outcome variable.

119 **Patient public involvement**

120 The public was not involved in the design or conduct, choice of outcome measures and
121 recruitment of the study.

122 **Result**

123 A total of 5,659 married women were included in the study. The mean age of the participants
124 was 31 (\pm 8.5) years with an age range of 15-49 years. The majority (98.2%) of the mothers were
125 Muslims and 3357 (59.3 %) were from a rural area. Concerning the reproductive history (98.6%)
126 of women have ANC visits (table 1).

127 Table 1 Sociodemographic and reproductive characteristics of married women in Senegal (n =
128 5,659).

Variable	Category	Frequency	Percent
----------	----------	-----------	---------

Age (in years)	15-24	1,340	23.7
	25-34	2,216	39.2
	35-49	2,103	37.1
Residence	Rural	3,357	59.3
	Urban	2,302	40.7
Religion	Muslim	5,556	98.2
	Christian	103	1.8
Educational status of respondents	No formal education	3,443	60.8
	Primary	1,177	20.8
	Secondary	857	15.2
	Higher	182	3.2
Educational status of the husband/partner	No formal education	3,615	63.9
	Primary	640	11.3
	Secondary	635	11.2
	Higher	330	5.8
	Don't know	439	7.8
Wealth index	Poor	2,298	40.6
	Middle	1,103	19.5
	Rich	2,258	39.9
Current working status	Currently working	2,904	51.3
	Currently not working	2,755	48.7
Mass media exposure to family planning methods	Have exposure	3,375	59.6
	Haven't exposure	2,284	40.4
Ideal number of children	≤4	1,190	21.0
	≥5	3,364	59.5
	Non-numeric response	1,105	19.5
Desire for more children	Wants within 2 years	1,602	28.3
	Wants after 2 years	2,450	43.3
	Wants, unsure timing	325	5.7
	Undecided	84	1.5
Number of living children	Doesn't want	1,198	21.2
	No child	718	12.7
	1-2 children	1,922	34.0

	3-4 children	1,485	26.2
Have ANC visit	5 or more children	1,534	27.1
	Yes	5,578	98.6
	No	81	1.4
Ever had terminated pregnancy	Yes	1,200	20.2
	No	4,733	79.8
Decision maker for using modern contraceptive	Women	295	20.4
	Husband	240	16.6
	Joint decision	893	62.0
	Other	16	1.0

129

130 **Modern contraceptive use among married women**

131 In this study the prevalence of modern contraceptive use among married women in Senegal was
 132 25.5% with (95% CI (24.4, 26.7)). The use of modern contraceptive method is higher among
 133 age group of 25-34 as compared to age group of women 15-24. Additionally, the utilization of
 134 modern contraceptives was higher (30.5%) among women who are currently working.
 135 Furthermore, women who are exposed to family planning methods through the media are more
 136 likely to use modern contraception (29.5%).

137 **Factors associated with modern contraceptive use**

138 Based on the output of multivariable binary logistic regression analysis, the following
 139 explanatory variables had a statistically significant association with modern contraceptive use :
 140 current working status of women (currently working (AOR=1.27, 95% CI (1.11, 1.46))), Husband
 141 education (husbands who attended primary educational level (AOR=1.32 ,95% CI (1.07,1.61),
 142 husbands who attended higher educational levels (AOR= 1.83 , 95% CI (1.32, 2.54), women
 143 whose wealth index in middle(AOR=1.37,95% CI (1.15,1.63) and rich (AOR=1.34,95% CI

144 (1.11,1.62), women who had exposed to family planning methods through different medias
145 (AOR=1.29,95% CI (1.13,1.48)), women who had ANC visits(AOR=2.01, 95% CI(1.13,3.57)).
146 Plus to the above factors, women who had no child (AOR=0.03, 95% CI (0.01,0.06)), 1-2
147 children (AOR=0.47,95% CI(0.38,0.59) and 3-4 children (AOR=0.79, 95% CI(0.67,0.96)),
148 women who thought ≥ 5 children are ideal(AOR=0.73,95% CI(0.61,0.88)) , and women who had
149 a desire for more children after 2 years (AOR=2.19, 95% CI(1.82,2.64) and women who doesn't
150 want a child (AOR=2.35, 95%CI (1.90,2.95)) were significantly associated with womens modern
151 contraceptive use.

152 The odds of using modern contraceptive was increased by 27% among women who were
153 currently working than their counterparts. The odds of using modern contraceptives was
154 increased by 32% among women whose husband had primary level of education. Moreover,
155 having a husband with primary and higher education level increased the odds of using modern
156 contraceptive by 32% and 16% respectively, when compared to women having an illiterate
157 husband. The odds of using modern contraceptives was increased by 29% among women who
158 had exposure to family planning methods through different Medias (Table 2).

159 Table 2 Factors associated with modern contraceptive use among married women in Senegal
160 (n=5,659).

161

162

163

164

165

166

167
168
169
170
171
172
173
174

Variable	Modern contraceptive use		p-value	AOR
	Yes (n=1,445)	No(n=4,214)		
	Frequency (%)	Frequency (%)		
Age				
15-24	221(16.5)	1,119(83.5)		1
25-34	613(27.7)	1,603(72.3)	0.8	0.97(0.78,1.2)
35-49	611(29.1)	1,492(70.9)	0.14	0.82(0.64,1.1)
Residence				
Urban	745(32.4)	1,557(67.6)		1
Rural	700(20.9)	2,657(79.1)	0.06	0.85(0.72,1.00)
Religion				
Muslim	1,414(25.5)	4,142(74.5)		1
Christian	31 (29.7)	72 (70.3)	0.92	0.98 (0.62,1.55)
Womens' educational level				
No formal education	754(21.9)	2,690(78.1)	0.05	0.61(0.37,1.00)
Primary	374(31.8)	803(68.2)	0.58	0.87(0.53,1.42)
Secondary	240(28.0)	617(72.0)	0.54	0.86(0.53,1.40)
Higher	77(42.3)	105(57.7)		1
Current working status				
Currently working	841(30.5)	1,913(69.5)	0.00	1.27 (1.11,1.46)

Currently not working	604(20.8)	2,301(79.2)		1
Husband education				
No formal education	777(21.5)	2838(78.5)		1
Primary education	192(29.9)	448(70.1)	0.008	1.32 (1.07,1.61)
Secondary education	192(30.2)	443(69.8%)	0.19	1.16 (0.93,1.45)
Higher	151(45.9)	179(54.1)	0.00	1.83 (1.32,2.54)
Don't know	134(30.4)	306(69.6)	0.28	1.16 (0.89,1.51)
Wealth index				
Poor	434(18.9)	1863(81.1)		1
Medium	301(27.3)	802(72.7)	0.00	1.37 (1.15,1.63)
Rich	709(31.4)	1549(68.6)	0.00	1.34 (1.11,1.62)
Media exposure				
Exposed	995(29.5)	2,381(70.5)	0.00	1.29 (1.13,1.48)
Non exposed	450(19.7)	1,833(80.3)		1
Have ANC visits				
Yes	1436(25.8)	4,142(74.5)	0.01	2.01 (1.13,3.57)
No	9(10.7)	72(89.3)		1
Ever had terminated pregnancy				
Yes	312(26.3)	877(73.7)	0.41	0.94 (0.79,1.09)
No	1,132(25.3)	3,337(74.7)		1
Number of living children				
No children	8(1.1)	711(98.9)	0.00	0.03 (0.01,0.06)
1-2 children	474(24.7)	1,448(75.3)	0.00	0.47 (0.38,0.59)
3-4 children	470(31.7)	1,014(68.3)	0.02	0.79 (0.67,0.96)
≥5 children	493(32.1)	1,041(67.9)		1
Ideal number of children				
≤ 4 children	381(32.0)	809(68.0)		1
≥ 5 children	854(25.4)	2,509(74.6)	0.001	0.73(0.61,0.88)
Non-numeric response	210(19.0)	895(81.0)	0.00	0.52(0.42,0.66)
Desire for more children				

Want's within 2 years	195(12.2)	1,407(87.8)		1
Want's after 2 years	750(30.6)	1,700(69.4)	0.00	2.19(1.82,2.64)
Want's, unsure timing	37(11.4)	288(88.6)	0.49	1.13(0.79,1.60)
Undecided	14(17.4)	70(82.6)	0.40	1.27(0.72,2.24)
Doesn't want	447(37.4)	750(62.6)	0.00	2.35(1.90,2.95)

175

176 **Discussion**

177 This study assessed the prevalence and associated factors of modern contraceptive use among
 178 Senegalese women. The prevalence of modern contraceptive use was 25.5% with (95% CI (24.4,
 179 26.7)). Womens' working status, husband education, wealth index, having media exposure,
 180 having ANC visit, number of living children, ideal number of children and desire for more
 181 children were factors that have statistically significant association with modern contraceptive
 182 use.

183 In the present study, the use of modern contraceptives among married Senegalese women was
 184 25.5% (95% CI (24.4, 26.7)). The result is in harmony with the study conducted in Senegal
 185 (26.3%) (16). This alignment might be explained by similarities in sociodemographic
 186 characteristics of the population.

187 The result of the current study is higher than the studies conducted in Bale zone, south East
 188 Ethiopia (20.8%)(17) and Metekel zone North West Ethiopia (18.6%)(18) . The discrepancy
 189 might be explained by socio-cultural difference between the countries that could affect their
 190 awareness on contraceptive use (19). On the other hand the result of the present study is lower
 191 than studies done in North Showa, Ethiopia (46.9%) (20) Zambia (43%)(21), and Egypt (
 192 46.5%) (22).

193 In the present study women who are currently enrolled in different jobs (currently working) had
194 significant association with modern contraceptive use. This result is supported by study
195 conducted in Ethiopia (19). This could be owing to the fact that women who work and have a
196 private job have a higher income, greater access to the media, and better health care, all of which
197 have a beneficial impact on modern contraception use (23–26).

198 In the current study, modern contraceptive use among married Senegalese women was
199 significantly associated with husband’s educational attainment, which is supported by a study
200 conducted in Senegal (16,27,28). It's probable that educated husbands are aware of the benefits
201 of modern contraception since they have a greater chance to read newspapers, watch television,
202 and use social media. The other factor that had significant association with contraceptive use was
203 wealth index of a woman. Women from households with a medium or high wealth index were
204 more likely to use modern contraception than those from households with a low wealth index.
205 The finding was in line with a study conducted in Ethiopia (29).

206 Modern contraceptive use was also significantly associated with womens’ media exposure. This
207 outcome is obedient with studies conducted in Senegal (16) and Ethiopia (9). This alignment
208 could be explained by the fact that media can help communities raise awareness about health
209 issues as well as address social and cultural issues (30,31). This critical role of the media can aid
210 in lowering barriers to health-care access and use, including contraception.

211 In this study presence of ANC visits have significant association with modern contraceptive use.
212 The result is supported by a multicenter study done in African countries (8). It's possible that the
213 increased prevalence of contraceptive use among women who had ANC visits is due to the fact
214 that antenatal care provides a unique opportunity to interact with, educate, and provide family

215 support to women of all socioeconomic backgrounds, which facilitates the uptake of modern
216 contraceptives. The study also showed that women who hadn't child have lower tendency to use
217 modern contraceptives than women who had three and more than three children. This result is
218 supported by studies done in Senegal (16,32). This meant that the more children a woman has,
219 the more likely she is to want to space or limit the number of children, and the more
220 contraceptive she is likely to use.

221 Furthermore, the present study also revealed the presence of statistically significant association
222 between modern contraceptive use, and womens' perception for ideal number of children they
223 would have. The result is in harmony with a study done in Senegal (16). According to the study,
224 women who wished five or more children had a lower likelihood of taking modern contraception
225 than women who desired four or fewer children. Finally, the results of the current study also
226 showed that womens' desire for more children also affects womens' modern contraceptive
227 utilization. That is the odds of using modern contraceptives among women who wants child
228 within 2 years lower than women who wants child after 2 years and totally doesn't want a child
229 at all.

230 The study's main strength was the use of nationally representative data and a large sample size,
231 which increased the generalizability of its conclusions. The DHS study, on the other hand, has
232 flaws in that it relies on respondents' self-reporting and may be prone to recall bias because
233 women were asked to recall past incidents. Furthermore, the design of this study limits its ability
234 to demonstrate causation between the desired outcome and these critical independent factors.

235 **Conclusion and recommendation**

236 According to the findings, the usage of modern contraceptive methods among married women in
237 Senegal was low. Women's contraceptive use was affected by family planning messages
238 exposure, sociodemographic, and obstetric factors. Thus, a tailored behavior change intervention
239 is required to increase contraceptive utilization among married women in Senegal and such
240 interventions should take those factors into account.

241 **Ethics approval and consent to participate**

242 The study is based on DHS data which was collected by DHS program, the DHS program collect
243 the data by gaining ethical approval from national IRB. Thus, we didn't received any ethical
244 approval since the data used for the study is from a secondary source. However, we got the
245 authentication letter from the DHS program to use the data set for this study. The study was
246 conducted under the Declaration of Helsinki.

247 **Consent for publication**

248 Not applicable.

249 **Availability of data and materials**

250 All result-based data are available within the manuscript and anyone can access the data set
251 online from www.measuredhs.com

252 **Competing interests**

253 The authors declare that they have no competing interests.

254 **Funding**

255 No, any funding was received from any organization.

256 **Authors' contributions**

257 All authors made substantial contributions to conception, acquisition of data, or analysis and
258 interpretation of data; took part in drafting the article or revising it critically for important
259 intellectual content; agreed to submit to the current journal; gave final approval of the version to
260 be published; and agree to be accountable for all aspects of the work.

261 **Acknowledgments**

262 We would like to acknowledge the MEASURE DHS program for permitting us to obtain and use
263 DHS data sets of Senegal.

264

265

266

267

268

269

270

271

272 **Reference**

273 1. Contraceptive prevalence – use of modern methods (%)(Health Equity Monitor)
274 [Internet]. Available from: [https://www.who.int/data/gho/indicator-metadata-registry/imr-](https://www.who.int/data/gho/indicator-metadata-registry/imr-details/3334)
275 [details/3334](https://www.who.int/data/gho/indicator-metadata-registry/imr-details/3334)

- 276 2. Akintade OL, Pengpid S, Peltzer K. Awareness and use of and barriers to family planning
277 services among female university students in Lesotho. *S Afr J Obstet Gynaecol.*
278 2011;17(3):72–8.
- 279 3. UNDESA. World Family Planning 2017 Highlights. *Econ Soc Aff [Internet].* 2017;1–43.
280 Available from:
281 [https://www.un.org/en/development/desa/population/publications/pdf/family/WFP2017_H](https://www.un.org/en/development/desa/population/publications/pdf/family/WFP2017_Highlights.pdf)
282 [ighlights.pdf](https://www.un.org/en/development/desa/population/publications/pdf/family/WFP2017_Highlights.pdf)
- 283 4. United Nations (UN). Sustainable Development Goals (SDG 5) | United Nations Western
284 Europe [Internet]. United Nations. 2020. Available from: [https://unric.org/en/sdg-](https://unric.org/en/sdg-12/%0Ahttps://unric.org/en/sdg-5/)
285 [12/%0Ahttps://unric.org/en/sdg-5/](https://unric.org/en/sdg-5/)
- 286 5. Ahmed S, Li Q, Liu L, Tsui AO. Maternal deaths averted by contraceptive use: An
287 analysis of 172 countries. *Lancet [Internet].* 2012;380(9837):111–25. Available from:
288 [http://dx.doi.org/10.1016/S0140-6736\(12\)60478-4](http://dx.doi.org/10.1016/S0140-6736(12)60478-4)
- 289 6. WHO. WHO recommendations on self-care interventions. *Human Reproduction*
290 *Programme.* 2020.
- 291 7. Darroch JE. Adding it up: Investing in Contraception and Maternal and Newborn Health
292 for Adolescents in Ethiopia , 2018. Guttmacher Institute. 2018.
- 293 8. Apanga PA, Kumbeni MT, Ayamga EA, Ulanja MB, Akparibo R. Prevalence and factors
294 associated with modern contraceptive use among women of reproductive age in 20
295 African countries: A large population-based study. *BMJ Open.* 2020;10(9).
- 296 9. Asresie MB, Fekadu GA, Dagnaw GW, Gelaw YM. Modern Contraceptive Use and

- 297 Influencing Factors in Amhara Regional State: Further Analysis of Ethiopian
298 Demographic Health Survey Data 2016. *Adv Public Heal.* 2020;2020(Mc).
- 299 10. Antarini A. Factors influencing the use of modern contraception among reproductive aged
300 women in bangka belitung province, indonesia. *Pan Afr Med J.* 2021;39.
- 301 11. Kwabena E, Id A, Budu E, Sambah F, Baatiema L, Appiah F, et al. Prevalence and
302 determinants of unintended pregnancy in sub-Saharan Africa: A multi- country analysis of
303 demographic and health surveys. *PLoS One.* 2019;14(8):1–16.
- 304 12. Abortion in Senegal _ Guttmacher Institute [Internet]. Available from:
305 <https://www.guttmacher.org/fact-sheet/abortion-senegal>
- 306 13. Senegal - Republic of Senegal - Western Africa [Internet]. Available from:
307 <https://www.nationsonline.org/oneworld/senegal.htm>
- 308 14. senegal-population @ worldpopulationreview.com [Internet]. Available from:
309 <https://worldpopulationreview.com/countries/senegal-population>
- 310 15. Choueiry G. What is an Acceptable Value for VIF? (With References) – Quantifying
311 Health [Internet]. 2020. Available from: <https://quantifyinghealth.com/vif-threshold/>
- 312 16. Zegeye B, Ahinkorah BO, Idriss-Wheeler D, Olorunsaiye CZ, Adjei NK, Yaya S. Modern
313 contraceptive utilization and its associated factors among married women in Senegal: a
314 multilevel analysis. *BMC Public Health.* 2021;21(1):1–13.
- 315 17. Belda SS, Haile MT, Melku AT, Tololu AK. Modern contraceptive utilization and
316 associated factors among married pastoralist women in Bale eco-region, Bale Zone, South
317 East Ethiopia. *BMC Health Serv Res.* 2017;17(1):1–12.

- 318 18. Adane AA, Bekele YA, Melese E, Worku GT, Netsere HB. Modern Contraceptive
319 Utilization and Associated Factors among Married Gumuz Women in Metekel Zone North
320 West Ethiopia. *Biomed Res Int.* 2020;2020.
- 321 19. Gebre MN, Edossa ZK. Modern contraceptive utilization and associated factors among
322 reproductive-age women in Ethiopia: Evidence from 2016 Ethiopia demographic and
323 health survey. *BMC Womens Health.* 2020;20(1):1–14.
- 324 20. A. M, D. W, A. F, B. M. Determinants of modern contraceptive utilization among married
325 women of reproductive age group in North Shoa Zone, Amhara Region, Ethiopia. *Reprod*
326 *Health* [Internet]. 2014;11(1):1–7. Available from:
327 [http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed12&NEWS=N&A](http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed12&NEWS=N&AN=2014111648)
328 [N=2014111648](http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed12&NEWS=N&AN=2014111648)
- 329 21. Lasong J, Zhang Y, Gebremedhin SA, Opoku S, Abaidoo CS, Mkandawire T, et al.
330 Determinants of modern contraceptive use among married women of reproductive age: A
331 cross-sectional study in rural Zambia. *BMJ Open.* 2020;10(3):1–10.
- 332 22. Unit FH. Practice of Family Planning among Married Female Attendants to Shawa Family
333 Health Unit, Dakahlia, Egypt. *Egypt Fam Med J.* 2020;4(1):24–41.
- 334 23. Tekelab T, Melka AS, Wirtu D. Predictors of modern contraceptive methods use among
335 married women of reproductive age groups in Western Ethiopia: A community based
336 cross-sectional study. *BMC Womens Health* [Internet]. 2015;15(1):1–8. Available from:
337 <http://dx.doi.org/10.1186/s12905-015-0208-z>
- 338 24. Ofonime JE. Determinants of modern contraceptive uptake among Nigerian women:

- 339 Evidence from the national demographic and health survey. *Afr J Reprod Health*
340 [Internet]. 2017;21(3):89–95. Available from:
341 [https://www.scopus.com/inward/record.uri?eid=2-s2.0-](https://www.scopus.com/inward/record.uri?eid=2-s2.0-85038842247&partnerID=40&md5=c36f097c56e5413d445dcaaf7d7a30a1)
342 [85038842247&partnerID=40&md5=c36f097c56e5413d445dcaaf7d7a30a1](https://www.scopus.com/inward/record.uri?eid=2-s2.0-85038842247&partnerID=40&md5=c36f097c56e5413d445dcaaf7d7a30a1)
- 343 25. Osmani AK, Reyer JA, Osmani AR, Hamajima N. Factors influencing contraceptive use
344 among women in Afghanistan: Secondary analysis of Afghanistan Health Survey 2012.
345 *Nagoya J Med Sci.* 2015;77(4):551–61.
- 346 26. Islam AZ, Mondal NI, Khatun ML. Prevalence and Determinants of Contraceptive use
347 among Employed and Unemployed Women in Bangladesh. 2016;5(2):92–102.
- 348 27. Tessema ZT, Teshale AB, Tesema GA, Yeshaw Y, Worku MG. Pooled prevalence and
349 determinants of modern contraceptive utilization in East Africa: A Multi-country Analysis
350 of recent Demographic and Health Surveys. *PLoS One* [Internet]. 2021;16(3 March):1–16.
351 Available from: <http://dx.doi.org/10.1371/journal.pone.0247992>
- 352 28. Gebremariam A. Modern Contraceptive Method Mix and Factors Affecting Utilization of
353 Modern Contraceptives among Married Women in Adigrat Town, Tigray, Northern
354 Ethiopia. *Fam Med Med Sci Res.* 2014;03(04).
- 355 29. Lakew Y, Reda AA, Tamene H, Benedict S, Deribe K. Geographical variation and factors
356 influencing modern contraceptive use among married women in Ethiopia: evidence from a
357 national population based survey. *Reprod Health.* 2013;1–10.
- 358 30. Ayanore MA, Pavlova M, Groot W. Unmet reproductive health needs among women in
359 some West African countries: A systematic review of outcome measures and

360 determinants. *Reprod Health* [Internet]. 2016;13(1):1–10. Available from:
361 <http://dx.doi.org/10.1186/s12978-015-0104-x>

362 31. Ansary R, Anisujjaman M. Factors Determining Pattern of Unmet Need for Family
363 Planning in. *Int Res J Soc Sci*. 2012;1(4):16–23.

364 32. Bulto GA, Zewdie TA, Beyen TK. Demand for long acting and permanent contraceptive
365 methods and associated factors among married women of reproductive age group in Debre
366 Markos Town, North West Ethiopia. *BMC Womens Health*. 2014;14(1).

367