

Correlates of functional physical capacity in physically active older adults: a conceptual-framework-based cross-sectional analysis of social determinants of health and clinical parameters

Lucinéia Orsolin Pfeifer¹, Lucas Helal^{2,3}, Nórton Luís Oliveira^{4,5}, Daniel Umpierre^{4,5,6}

1 Exercise Pathophysiology Research Laboratory, Graduate Program in Cardiology and Cardiovascular Sciences, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil.

2 Diretoria de Pesquisa, Hospital de Clínicas de Porto Alegre, Porto Alegre, RS, Brazil.

3 Graduate Program in Cardiology and Cardiovascular Sciences, School of Medicine, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil.

4 Exercise Pathophysiology Research Laboratory, Hospital de Clínicas de Porto Alegre, Clinical Research Center, Porto Alegre, RS, Brazil

5 National Institute of Science and Technology for Health Technology Assessment, Hospital de Clínicas de Porto Alegre, Clinical Research Center, Porto Alegre, RS, Brazil.

6 Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil.

Corresponding author:

Daniel Umpierre, PhD

Hospital de Clínicas de Porto Alegre

Rua Ramiro Barcelos, 2350 - Hospital de Clínicas de Porto Alegre, Clinical Research Center, 21301

Porto Alegre, RS, Brazil

Ph: +55 51 3359.6332 / Fax: +55 51 3334.6462

Email: danielumpierre@hcpa.edu.br

Abstract

This study aimed to explore social determinants of health and health/clinical determinants on two outcomes of functional physical capacity. Therefore, a population-based sample of 327 older adults (69±7 years; 83.5% women) underwent demographical and clinical questionnaires, risk factors assessments, six-minute walk testing (walking capacity), and handgrip strength testing. Based on multivariable linear regression models, age (-4.05m; -5.3 to -2.8), being men (71.40m; 50.5 to 92.3), body mass index (-3.88m; -5.6 to -2.1), and quality of life (18.48m; 6.3 to 30.6) remained as predictive variables for walking capacity ($R^2=30.8\%$). In the final model for handgrip strength, age (-0.6% kgf; 0.89 to 0.2) and male sex (65.2% kgf; 55.3 to 75.8) remained as predictive variables. Despite exploratory analyses including contextual factors as potential predictors of walking capacity and handgrip strength, only outcomes at the individual levels were associated, either positively or negatively, with the variations presented by this studied sample of older adults.

Keywords: Aged, Physical function, Sociodemographic factors.

INTRODUCTION

Functional decline in older adults may be affected by clinical status, physical fitness, and social determinants of health (Beaton et al., 2015). In this regard, sarcopenia, which is a progressive decline in skeletal muscle mass (Evans, 1995), has been consistently associated with a decline in functional capacity and disability in the elderly. Similarly, dynapenia, which relates to the loss in muscle strength independently of changes in muscle mass is also an independent predictor of overall mortality in this population (Clark & Manini, 2012). In terms of functional capacity outcomes, handgrip and lower limb muscle strength, and ambulatory gait speed are indexes of physical fitness and may predict adverse events in older adults, such as declines in cognition, mobility, functional status (Rijk et al., 2016), disability, (Donoghue et al., 2014; Shimada et al., 2015), hospitalization (Cesari et al., 2009) and all-cause mortality (De Buysers et al., 2013; Rantanen et al., 2003; Rijk et al., 2016; Studenski et al., 2011). Importantly, regular exercise may counteract these deleterious effects (Izquierdo et al., 2021), both by structured exercise training and physical activity advice, thus optimizing the maintenance of functional capacity during aging.

Similar to the many health-related events, the functional decline in older adults is potentially influenced by a combination of social determinants of health and health/clinical-related determinants. Age, sex, and body mass have a solid association with ambulatory functional outcomes (i.e., impaired walking test and handgrip strength) (ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories, 2002; Casanova et al., 2011; Corrêa et al., 2020; V. Z. Dourado, 2011; Gale et al., 2007; Lu et al., 2020; Salbach et al., 2015; Tveter et al., 2014) and are frequently used in trials as surrogate outcomes (V. Z. Dourado, 2011). Although some potential predictors more income, use of medications, diabetes, quality of life, and depression are, likewise, associated (Enright et al., 2003; Kuziemski et al., 2019; Liang et al., 2020; Ling et al., 2010; Love et al., 2020; Serra et al., 2015), although not explored as far as the other potential predictors aforementioned. The variability in results of

functional tests can be explained through a myriad of factors (standards of the tests, color, ethnicity, and geographic locations, even in the same country), which make the inter and intra-test comparability inappropriate to other populations.

We perform out the first exploratory study from a sample initially evaluated in a cross-sectional descriptive study carried out previously (F. M. Dourado et al., 2021). We aimed to estimate, in a representative sample of older adults engaged in a municipal program of physical activity, the extent of the contribution of social and clinical health determinants in functional physical capacity, here defined as the ambulatory walking capacity and the handgrip force. As this is an exploratory study, we do not present a directional hypothesis.

METHODS

Study setting

This is the first exploratory study within the descriptive cross-sectional study entitled: *"Health profile of elderly users of a town public program of physical activity: a cross-sectional study"*. In this study, we used questionnaires, clinical measurements, and functional tests to assess the health profile and social determinants of health of older adults spontaneously engaged in a public lifestyle program composed of several lifestyle items, including a physical activity program, in the city of Porto Alegre, Brazil. In the city public program, modalities such as gymnastics, resistance training, rhythms, and/or recreational sports were offered to participants. The majority of them took place twice a week lasting one hour of duration (F. M. Dourado et al., 2021). The study protocol is publicly available at <https://osf.io/q4r69/> and was approved by the Institutional Review Board of the Hospital de Clínicas de Porto Alegre (GPPG 2018-0100; CAAE 84093317600005327) and of the Municipal Secretariat of Health (SMS/POA). The procedures followed national guidelines for bioethics in research. All participants read and signed the informed consent before starting their enrollment in the study. The reporting of this manuscript is based on the STROBE Statement

recommendations (Strengthening the reporting of observational studies in epidemiology) in its extension for cross-sectional studies (von Elm et al., 2007). Study data are also available in a public repository for independent researchers (<https://zenodo.org/record/4341443#.X9u61thKhPY>; doi: 10.5281/zenodo.4341443).

Participants and eligibility criteria

The inclusion and exclusion criteria defined in the previous descriptive study were: being 60 years old or older and regularly attending the physical activity programs of the municipal community centers. Exclusion criteria were defined as the impossibility of moving to the assessment sites and/or any musculoskeletal injury which put the patient at risk in any setting. All participants included in the previous study were considered for inclusion in this study provided they had complete assessment data.

Sampling and sample size calculation

The sample size calculation was carried out for the previous descriptive study, using the prevalence of hypertension as the outcome of interest and considering the size of each of the centers in the city, totaling 443 participants. Detailed information is described in the previous study (F. M. Dourado et al., 2021).

Data collection

The recruitment and data collection were carried forward between April 2018 and February 2019. Data were collected at two moments. In visit 1, the research team went to 11 different sites attended by the participants. The data collection included self-administered questionnaires and the 6-minute walking test (6MWT). In visit 2, the participants went to the Hospital de Clínicas de Porto Alegre, Clinical Research Center, undergoing clinical measurements (e.g., blood pressure readings and anthropometric evaluation) and handgrip

testing. The procedures for data collection are more detailed elsewhere (F. M. Dourado et al., 2021).

The researchers received training in the standardization of procedures; some of the measurements were routinely performed by the research team for other studies (Umpierre et al., 2019), based on a manual of standard operating procedures. This was done to minimize measurement bias since different researchers were involved in the measurements and procedures in different centers visited.

Study Variables

Social determinants of health and clinical/health-related variables

A 62-item questionnaire was applied (visit 1), allowing the collection of demographics, clinical history, social determinants of health, risk factors, diet, physical activity behaviors, morbidity, and perception of health. Sex was categorized into men and women; the number of medications in four categories (none, up to 3 medications, 4 to 6 and 7 or more); diabetes as yes, no, or gestational; family income in four minimum wages (MW) categories (up to 2 MW, 2 to 4 MW, 4 to 10 MW, 10 to 20 MW).

The older adult's height (cm) and weight (kg) were transformed into body mass index (BMI) values by taking the weight in kilograms and dividing it by the square of height in meters (kg/m^2).

Quality of life was assessed using the Short Form 6 Dimension (SF-6D) self-administered questionnaire, which consists of six questions, within six domains, namely: functional capacity, physical and emotional aspects, social aspects, pain, mental health, and vitality. The questionnaire generates a score between 0.29 - 1.00, whether 1.00 means a full quality of life state.

Depressive symptoms were assessed by the Geriatric Depression Scale (GDS-15), which consists of 15 questions. In GDS-15, each positive answer associated with depression represents a point, generating scores from 0 to 15, where scores of 6 or higher were used as suggestive of depression (Krishnamoorthy et al., 2020). Using this cutoff point (score 6), the variable was categorized into presence or absence of depression.

Functional Capacity

6MWT

The 6MWT was used to indirectly assess the cardiorespiratory fitness (estimations) and walking capacity. We considered the distance covered during a 6-minute period, in a 30-meter delimited pathway. The participants were encouraged in a standardized manner, according to the established guidelines (ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories, 2002).

Handgrip Test

The handgrip strength test was used to measure the strength generated by the muscles of the hand and forearm, and assess the physical condition of the upper limb, an important proxy of health status. Three measurements were taken using a dynamometer (Jamar, model 2A, Asimow Engineering Co., Santa Monica, USA), at a standing position of participants, and the elbow at 90 degrees of flexion. We took a series of measurements for both hands and intervals of one minute were given between each attempt. For the purpose of this study, the highest value reached in kgf was considered the dominant hand.

Statistical methods

Continuous variables are described as mean and standard deviation whereas categorical variables are reported as prevalences and absolute frequencies. Most variables were maintained in the way they were collected, however, some variables were categorized to

facilitate data analysis and interpretation (i.e., number of medications, and family income).

We evaluated distributions visually and statistically - the first through a visual inspection of histograms and Kolmogorov-Smirnov tests which assume normality of the data as H_0 .

For inferential analysis, we used a multivariable linear regression model. Components of functional capacity were defined as dependent variables for this study in two separate models. To maintain similarities between analyses and preserve the natural units of the coefficients, we transformed the handgrip observations into a natural logarithm (log), which generated a Gaussian distribution and allowed us to carry on a linear regression model. Then, we regressed both log of handgrip strength and 6WMT by linear multivariable regression analysis using a Gaussian distribution for the outcome and an identity function as linkage, in a two-step manner (model 1 and model 2 - the last one adjusted to more variables based on our conceptual framework model - explained below).

In regression analyses, we diagnosed the basic assumptions of multivariable linear regressions (i.e., normality of the distribution of the outcome and its residual, multicollinearity, and constant variance of residuals), the adequacy of the chosen model predictors the model's effectiveness. We plotted adjusted outcomes against Pearson residuals and independent variables against residuals, as well as a tested correlation between them and the assumption of homoscedasticity. When exploring heterogeneity, we found evidence of heterogeneity of variances, mostly due to observations in sex, about the dependent variable handgrip. Based on this, we carried out a linear model for 6WMT and a linear model using Generalized Least Squares (GLS) for the log handgrip strength.

The variables included in the multivariable linear regression model were defined *a priori* based on a causal conceptual framework. Distal and proximal variables were inserted according to the knowledge base (e.g., obtained through published literature) for potential causality. We chose the conceptual framework method because stepwise models may have limitations if no causal relationship is considered. These models may have limited power to

select important predictors and correlates in small datasets and may result in a bias regarding the estimated regression coefficients (Steyerberg et al., 1999). Because thresholds of statistical significance are limited as a selection method to determine which variables should be added to a regression model (Bagherzadeh-Khiabani et al., 2016), our variable selection model for inclusion in the regression model is based on identified direct evidence of a possible association and physiological plausibility relationship of the influence of covariates. Considering that model 1 of the regression was composed of the proximal variables, which were loaded into model 2 (distal variables) only when they reached a correlation with the dependent variable ($P < 0.05$).

No data imputation was done. In this analysis model, only the complete data of all variables present in the regression model remained, that is, any participants that presented missing data were excluded. To claim statistical significance, we assumed a threshold of 0.05. Analyses were performed in the software SPSS (Statistical Package for the Social Sciences) version 18.0 and R (R Core Team 2020, version 4.0.3). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. R Packages used were tidyverse, nlme, emmeans.

RESULTS

In total, 327 participants were included in the current study. Participants had a mean age of 69 ± 7 years (min to max 60 up to 91 years old), and 83.5% of them were female (**Table 1**). The mean BMI was 27 ± 3 kg/m² and 47.4% of the participants were classified as "overweight". The prevalence of self-reported diabetes (any type) was 12.5%. Potential depression, as classified by the depression scores of the GDS-15 instrument, had a prevalence of 18%; and, importantly, QoL levels had a mean score of 0.87 ± 0.06 . Concerning family income, 73.2% of participants received up to 4 MW. 70 participants (21.4%) reported not using any type of continuous-use medication, although the vast majority took at least three medications continuously as prescribed. Commenting on our primary outcomes

(dependent variables), the mean walk capacity achieved was 498.4 ± 78 and the mean handgrip strength by the dominant hand was $27.1 \text{ kgf} \pm 8.2 \text{ kgf}$ (min to max from 302m up to 690m and 8 kgf to 60 kgf, respectively for both outcomes). **Table 2** shows the descriptive analysis of dependent variables and predictors.

We regressed the 6MWT in a multivariable linear regression of 312 participants (**Table 3**). Model 1, including predictors as (age, sex, BMI, number of medications, and quality of life (SF-6D)) explained 30% (adjusted R^2) of the variability in walking distance. For each unit of age ($P < 0.001$), BMI ($P < 0.001$), sex ($P < 0.001$), and QoL ($P = 0.001$), which remained independent correlates, modified the covered distance in 6 minutes about a decrease of -4.04m (-5.2m to -2.7m), -4.30m (-6.0m to -2.6m) and 72.84m (52.9m to 92.8m), respectively; and, the QoL increases, the 6MWT increases, in average, 20.58m (8.9m to 32.2m). In a second model, we maintained variables with statistical significance and added into the model 1 potential predictors/correlates as diagnosis of diabetes, family income, and potential depression. The addition of such variables roughly modified the variance (adjusted $R^2 = 30.8\%$ vs 30%). All previous independent variables remained correlated with 6MWT and the new inserted variables did not reach correlation to 6MWT.

To the handgrip strength, the results of the analysis of multivariable regression on generalized linear models with 325 participants are presented in **Table 4**. In the first model, age, sex, BMI, and the number of medications were included according to our conceptual framework. For each unit of age ($P = 0.009$), sex (male) ($P < 0.001$) and use of seven or more medications ($P = 0.02$), which remained independent correlates, modified handgrip performance in an average decrease of -0.5 kgf% (0.89 kgf to 0.09 kgf), and -13.1 kgf% (22.74 kgf to 2.17 kgf), for age and seven or more medications, respectively; and a 65.9 kgf (55.89 kgf to 76.47 kgf) increase for males, in average. In model 2, we maintained variables with statistical significance from model 1 and added diagnosis of diabetes. Only the independent variables age ($P = 0.005$) and sex ($P < 0.001$) remained correlated with the

handgrip test, the number of medications (seven or more) lost correlation ($P=0.07$) and the newly inserted variable did not reach a correlation with the test. However, age and sex (men) were practically not impacted either after adjustment for the additional variables, neither their precision estimators -0.5 kgf (0.89 kgf to 0.09 kgf) vs -0.6 kgf (0.89 kgf to 0.2 kgf); 65.9 kgf (55.89 kgf to 76.47 kgf) vs 65.2 kgf (55.3 kgf to 75.8 kgf). Compared to model 1, model 2 provides minimally better results (Pseudo R^2 : 0.29 vs 0.28).

DISCUSSION

In the present exploratory study, we assessed whether predictors would contribute to predicting the 6MWT and handgrip test based on a conceptual framework construct. The main findings of our study are that social determinants of health and clinical parameters can have both direct and inverse associations with functional outcomes. Importantly, the addition of such well-identified explanatory variables (model 2) did not substantially modify the magnitude of effect, precision, and mean explanatory capacity of the outcome variance. Therefore, we infer that these variables did not present themselves as confounders nor modifiers of effect. While for 6MWT, age, sex (women), BMI, and QoL had a relevant mean clinical impact as predictors, only sex (men) impacted importantly handgrip strength, regardless of any statistical significance.

Several variables can impact the 6MWT performance, among them, height, age, sex, body weight are variables that can be highlighted because they are part of the reference equations for the distance covered in the test (ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories, 2002; Britto et al., 2013). Age, sex, and BMI are well-established predictors of 6MWT performance (ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories, 2002; Casanova et al., 2011; V. Z. Dourado, 2011; Salbach et al., 2015; Tveter et al., 2014) and the same was found in our study. Walking capacity is a proxy of QoL in a causal relationship (Lord & Menz, 2002; Serra et al., 2015). In his regard, we observed that for each additional score in the SF-6D questionnaire, around

21m was added to the walking capacity on average. Although clinically important, the cross-sectional nature and causal principles (such as Hill's criteria for causality) (Hill, 1965) do limit a claim and put this independent relationship at risk of reverse causality. Lastly, the use of medications, diabetes diagnosis, depression, and family income did not explain any variability in the mean 6MWT values of the sample. While expected, this may have been an approximation of the QoL scores of our sample, which were roughly impaired at baseline, and consequently did not change the model.

Regarding the analyses using the handgrip strength as an outcome, sex and age were the predictors linked to the test performance, consistent with previous studies (Corrêa et al., 2020; Lu et al., 2020; Tveter et al., 2014). Diabetes did not predict changes in this outcome, although this disease leads to peripheral neuropathy and overall inflammation (Matsuda et al., 2004; Tsalamandris et al., 2019). To note, only 13% of the sample self-reported a diagnosis of diabetes and, additionally, we did not collect data regarding comorbidity related to diabetes (e.g., peripheral neuropathy). In both walking and handgrip strength tests, one possible contributing factor for diabetes not being a significant predictor may be due to the physical-active status of study participants. Some classes of medications may also impair strength production, such as antihypertensives and lipid-lowering drugs (Ashfield et al., 2010; Love et al., 2020; Onder et al., 2002), which were prevalent in our sample (F. M. Dourado et al., 2021). Nonetheless, BMI may be collinear to diabetes and reflect an inherent effect of disease, mirroring the effect of the disease in the model. After all, even for the variables which appeared to be correlated, the contribution to force magnitude was clinically marginal, except for sex, since being male explained to perform an average of 66.0 kgf more.

Our results should be interpreted in light of limitations. First, the mean values for our predicted outcomes (walking and handgrip) were similar to those of healthy and physically active ones, which may be a consequence of the engagement of our sample in a lifestyle program. Also, although using the conceptual framework model to choose explanatory

variables with a solid rationale, some of them may present reverse causality in this study setting (cross-sectional study), regardless of our efforts to annulate this type of bias; also, we considered only one conceptual framework model for each multivariable analysis. The addition of more models and the comparability could allow us to better explain the phenomenon, which was diagnosed, at least statistically, by the presented R^2 (around 30% of the mean variability of outcomes). For the handgrip strength variable, the original distribution was a Poisson-like distribution. As we decided to maintain betas to see contributions in real units, the variable was log-transformed. However, a sensitivity analysis using a robust variance Poisson regression with exponentialized betas (prevalence ratios - PRs) is a recommended future direction from our side.

On the other hand, we emphasize the exploratory analysis of correlates of these two outcomes, by its clinical relevance, particularly in this setting - in the older and majority in women. There is a pathophysiological rationale to assume a decline in the functional capacity during the aging process, here assumed as a combination of 6WMT performance and handgrip test performance in the older adults, due to impairments in various physiological systems of the body. To quantify the contribution of such parameters, as attenuators of contributors to the decline, is of importance, and we did this in a population-based study. Caution should be taken into account for extrapolation in terms of sex, as a matter of importance. Then, our results should be considered for older people who are not healthy, but under treatment (pharmacological and non-pharmacological). Further exploration of the diagnosis of diabetes, number of medications, depression, and family income are also welcome.

CONCLUSION

We conclude that in an older population engaged in a municipal program physical activity program, the walking capacity and handgrip strength were directly and inversely associated with social determinants of health and clinical parameters (sex, age, BMI, and QoL). The

results may inform of programs or studies with comparable subjects cohorts and provide evidence for model adjustment in observational intervention studies targeting functional capacity in older subjects.

Acknowledgments

We would like to thank Rogério Boff Borges, a statistician at the Biostatistics Unit of the Hospital de Clínicas of Porto Alegre, for his help and guidance in carrying out the statistical analysis for this article.

Funding source

This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001; National Institute of Science and Technology for Health Technology Assessment (IATS) – FAPERGS/Brasil; National Council on Technology and Scientific Development (CNPq).

REFERENCES

- Ashfield, T. A., Syddall, H. E., Martin, H. J., Dennison, E. M., Cooper, C., & Aihie Sayer, A. (2010). Grip strength and cardiovascular drug use in older people: findings from the Hertfordshire Cohort Study. *Age and Ageing*, *39*(2), 185–191. <https://doi.org/10.1093/ageing/afp203>
- ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories. (2002). ATS statement: guidelines for the six-minute walk test. *American Journal of Respiratory and Critical Care Medicine*, *166*(1), 111–117. <https://doi.org/10.1164/ajrccm.166.1.at1102>
- Bagherzadeh-Khiabani, F., Ramezankhani, A., Azizi, F., Hadaegh, F., Steyerberg, E. W., & Khalili, D. (2016). A tutorial on variable selection for clinical prediction models: feature selection methods in data mining could improve the results. *Journal of Clinical Epidemiology*, *71*, 76–85. <https://doi.org/10.1016/j.jclinepi.2015.10.002>
- Beaton, K., McEvoy, C., & Grimmer, K. (2015). Identifying indicators of early functional decline in community-dwelling older people: a review. *Geriatrics & Gerontology International*, *15*(2), 133–140. <https://doi.org/10.1111/ggi.12379>
- Britto, R. R., Probst, V. S., de Andrade, A. F. D., Samora, G. A. R., Hernandez, N. A., Marinho, P. E. M., Karsten, M., Pitta, F., & Parreira, V. F. (2013). Reference equations for the six-minute walk distance based on a Brazilian multicenter study. *Brazilian Journal of Physical Therapy*, *17*(6), 556–563. <https://doi.org/10.1590/S1413-35552012005000122>
- Casanova, C., Celli, B. R., Barria, P., Casas, A., Cote, C., de Torres, J. P., Jardim, J., Lopez, M. V., Marin, J. M., Montes de Oca, M., Pinto-Plata, V., Aguirre-Jaime, A., & Six Minute Walk Distance Project (ALAT). (2011). The 6-min walk distance in healthy subjects: reference standards from seven countries. *The European Respiratory Journal: Official Journal of the European Society for Clinical Respiratory Physiology*, *37*(1), 150–156.

<https://doi.org/10.1183/09031936.00194909>

- Cesari, M., Kritchevsky, S. B., Newman, A. B., Simonsick, E. M., Harris, T. B., Penninx, B. W., Brach, J. S., Tykavsky, F. A., Satterfield, S., Bauer, D. C., Rubin, S. M., Visser, M., Pahor, M., & Health, Aging, and Body Composition Study. (2009). Added value of physical performance measures in predicting adverse health-related events: results from the Health, Aging And Body Composition Study. *Journal of the American Geriatrics Society*, 57(2), 251–259. <https://doi.org/10.1111/j.1532-5415.2008.02126.x>
- Clark, B. C., & Manini, T. M. (2012). What is dynapenia? *Nutrition*, 28(5), 495–503. <https://doi.org/10.1016/j.nut.2011.12.002>
- Corrêa, T. G. C., Donato, S. V. S., Lima, K. C. A., Pereira, R. V., Uygur, M., & de Freitas, P. B. (2020). Age- and Sex-Related Differences in the Maximum Muscle Performance and Rate of Force Development Scaling Factor of Precision Grip Muscles. *Motor Control*, 24(2), 274–290. <https://doi.org/10.1123/mc.2019-0021>
- De Buyser, S. L., Petrovic, M., Taes, Y. E., Toye, K. R. C., Kaufman, J.-M., & Goemaere, S. (2013). Physical function measurements predict mortality in ambulatory older men. *European Journal of Clinical Investigation*, 43(4), 379–386. <https://doi.org/10.1111/eci.12056>
- Donoghue, O. A., Savva, G. M., Cronin, H., Kenny, R. A., & Horgan, N. F. (2014). Using timed up and go and usual gait speed to predict incident disability in daily activities among community-dwelling adults aged 65 and older. *Archives of Physical Medicine and Rehabilitation*, 95(10), 1954–1961. <https://doi.org/10.1016/j.apmr.2014.06.008>
- Dourado, F. M., Oliveira, N. L., Helal, L., Ricardo, L. I. C., Santos, L. dos, De Nardi, A. T., Botton, C. E., Pfeifer, L. O., Vasques, L. M., Santos, L. P., & Others. (2021). Health profile of elderly users of a town public program of physical activity: a cross-sectional study. In *Health profile of elderly users of a town public program of physical activity: a cross-sectional study*. <https://pesquisa.bvsalud.org/portal/resource/pt/pps-1678>
- Dourado, V. Z. (2011). Reference equations for the 6-minute walk test in healthy individuals. *Arquivos Brasileiros de Cardiologia*, 96(6), e128–e138.

<https://www.scielo.br/scielo.php?pid=S0066->

[782X2011000600022&script=sci_arttext&tIng=pt](https://www.scielo.br/scielo.php?pid=S0066-782X2011000600022&script=sci_arttext&tIng=pt)

Enright, P. L., McBurnie, M. A., Bittner, V., Tracy, R. P., McNamara, R., Arnold, A., Newman, A. B., & Cardiovascular Health Study. (2003). The 6-min walk test: a quick measure of functional status in elderly adults. *Chest*, *123*(2), 387–398.

<https://doi.org/10.1378/chest.123.2.387>

Evans, W. J. (1995). What is sarcopenia? *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, *50 Spec No*, 5–8.

https://doi.org/10.1093/gerona/50a.special_issue.5

Gale, C. R., Martyn, C. N., Cooper, C., & Sayer, A. A. (2007). Grip strength, body composition, and mortality. *International Journal of Epidemiology*, *36*(1), 228–235.

<https://doi.org/10.1093/ije/dyl224>

Hill, A. B. (1965). THE ENVIRONMENT AND DISEASE: ASSOCIATION OR CAUSATION? *Proceedings of the Royal Society of Medicine*, *58*, 295–300.

<https://www.ncbi.nlm.nih.gov/pubmed/14283879>

Izquierdo, M., Merchant, R. A., Morley, J. E., Anker, S. D., Aprahamian, I., Arai, H., Aubertin-Leheudre, M., Bernabei, R., Cadore, E. L., Cesari, M., Chen, L.-K., de Souto Barreto, P., Duque, G., Ferrucci, L., Fielding, R. A., García-Hermoso, A., Gutiérrez-Robledo, L. M., Harridge, S. D. R., Kirk, B., ... Singh, M. F. (2021). International Exercise

Recommendations in Older Adults (ICFSR): Expert Consensus Guidelines. *The Journal of Nutrition, Health & Aging*. <https://doi.org/10.1007/s12603-021-1665-8>

Krishnamoorthy, Y., Rajaa, S., & Rehman, T. (2020). Diagnostic accuracy of various forms of geriatric depression scale for screening of depression among older adults: Systematic review and meta-analysis. *Archives of Gerontology and Geriatrics*, *87*, 104002.

<https://doi.org/10.1016/j.archger.2019.104002>

Kuziemski, K., Słomiński, W., & Jassem, E. (2019). Impact of diabetes mellitus on functional exercise capacity and pulmonary functions in patients with diabetes and healthy persons. *BMC Endocrine Disorders*, *19*(1), 2. <https://doi.org/10.1186/s12902-018-0328->

1

- Liang, X., Jiang, C. Q., Zhang, W. S., Zhu, F., Jin, Y. L., Cheng, K. K., Lam, T. H., & Xu, L. (2020). Glycaemia and hand grip strength in aging people: Guangzhou biobank cohort study. *BMC Geriatrics*, *20*(1), 399. <https://doi.org/10.1186/s12877-020-01808-0>
- Ling, C. H. Y., Taekema, D., de Craen, A. J. M., Gussekloo, J., Westendorp, R. G. J., & Maier, A. B. (2010). Handgrip strength and mortality in the oldest old population: the Leiden 85-plus study. *CMAJ: Canadian Medical Association Journal = Journal de l'Association Medicale Canadienne*, *182*(5), 429–435. <https://doi.org/10.1503/cmaj.091278>
- Lord, S. R., & Menz, H. B. (2002). Physiologic, psychologic, and health predictors of 6-minute walk performance in older people. *Archives of Physical Medicine and Rehabilitation*, *83*(7), 907–911. <https://doi.org/10.1053/apmr.2002.33227>
- Love, B. L., Mitchell, E., & Norris, L. A. B. (2020). Association Between Reduced Handgrip Strength and Commonly Prescribed Medications. *medRxiv*. <https://www.medrxiv.org/content/10.1101/2020.01.15.20017087v1.abstract>
- Lu, X., Chu, H., Wang, L., Yang, R., Li, Y., Sun, W., Yan, C., Liu, Y., Guo, Z., & Cheng, X. (2020). Age- and sex-related differences in muscle strength and physical performance in older Chinese. *Aging Clinical and Experimental Research*, *32*(5), 877–883. <https://doi.org/10.1007/s40520-019-01263-x>
- Matsuda, M., Kawasaki, F., Inoue, H., Kanda, Y., Yamada, K., Harada, Y., Saito, M., Eto, M., Matsuki, M., & Kaku, K. (2004). Possible contribution of adipocytokines on diabetic neuropathy. *Diabetes Research and Clinical Practice*, *66 Suppl 1*, S121–S123. <https://doi.org/10.1016/j.diabres.2004.05.010>
- Onder, G., Penninx, B. W. J. H., Balkrishnan, R., Fried, L. P., Chaves, P. H. M., Williamson, J., Carter, C., Di Bari, M., Guralnik, J. M., & Pahor, M. (2002). Relation between use of angiotensin-converting enzyme inhibitors and muscle strength and physical function in older women: an observational study. *The Lancet*, *359*(9310), 926–930. [https://doi.org/10.1016/s0140-6736\(02\)08024-8](https://doi.org/10.1016/s0140-6736(02)08024-8)

- Rantanen, T., Volpato, S., Ferrucci, L., Heikkinen, E., Fried, L. P., & Guralnik, J. M. (2003). Handgrip strength and cause-specific and total mortality in older disabled women: exploring the mechanism. *Journal of the American Geriatrics Society*, 51(5), 636–641. <https://doi.org/10.1034/j.1600-0579.2003.00207.x>
- Rijk, J. M., Roos, P. R., Deckx, L., van den Akker, M., & Buntinx, F. (2016). Prognostic value of handgrip strength in people aged 60 years and older: A systematic review and meta-analysis. *Geriatrics & Gerontology International*, 16(1), 5–20. <https://doi.org/10.1111/ggi.12508>
- Salbach, N. M., O'Brien, K. K., Brooks, D., Irvin, E., Martino, R., Takhar, P., Chan, S., & Howe, J.-A. (2015). Reference values for standardized tests of walking speed and distance: a systematic review. *Gait & Posture*, 41(2), 341–360. <https://doi.org/10.1016/j.gaitpost.2014.10.002>
- Serra, A. J., de Carvalho, P. de T. C., Lanza, F., de Amorim Flandes, C., Silva, S. C., Suzuki, F. S., Bocalini, D. S., Andrade, E., Casarin, C., & Silva, J. A., Jr. (2015). Correlation of six-minute walking performance with quality of life is domain- and gender-specific in healthy older adults. *PLoS One*, 10(2), e0117359. <https://doi.org/10.1371/journal.pone.0117359>
- Shimada, H., Makizako, H., Doi, T., Tsutsumimoto, K., & Suzuki, T. (2015). Incidence of Disability in Frail Older Persons With or Without Slow Walking Speed. *Journal of the American Medical Directors Association*, 16(8), 690–696. <https://doi.org/10.1016/j.jamda.2015.03.019>
- Steyerberg, E. W., Eijkemans, M. J., & Habbema, J. D. (1999). Stepwise selection in small data sets: a simulation study of bias in logistic regression analysis. *Journal of Clinical Epidemiology*, 52(10), 935–942. [https://doi.org/10.1016/s0895-4356\(99\)00103-1](https://doi.org/10.1016/s0895-4356(99)00103-1)
- Studenski, S., Perera, S., Patel, K., Rosano, C., Faulkner, K., Inzitari, M., Brach, J., Chandler, J., Cawthon, P., Connor, E. B., Nevitt, M., Visser, M., Kritchevsky, S., Badinelli, S., Harris, T., Newman, A. B., Cauley, J., Ferrucci, L., & Guralnik, J. (2011). Gait speed and survival in older adults. *JAMA: The Journal of the American Medical*

Association, 305(1), 50–58. <https://doi.org/10.1001/jama.2010.1923>

Tsalamandris, S., Antonopoulos, A. S., Oikonomou, E., Papamikroulis, G.-A., Vogiatzi, G., Papaioannou, S., Deftereos, S., & Tousoulis, D. (2019). The Role of Inflammation in Diabetes: Current Concepts and Future Perspectives. *European Cardiology*, 14(1), 50–59. <https://doi.org/10.15420/ecr.2018.33.1>

Tveter, A. T., Dagfinrud, H., Moseng, T., & Holm, I. (2014). Health-related physical fitness measures: reference values and reference equations for use in clinical practice. *Archives of Physical Medicine and Rehabilitation*, 95(7), 1366–1373. <https://doi.org/10.1016/j.apmr.2014.02.016>

Umpierre, D., Santos, L. P., Botton, C. E., Wilhelm, E. N., Helal, L., Schaun, G. Z., Ferreira, G. D., De Nardi, A. T., Pfeifer, L. O., da Silveira, A. D., Polanczyk, C. A., Mendes, G. F., Tanaka, H., Alves, L., Galliano, L., Pescatello, L. S., Brizio, M. L., Bock, P. M., Campelo, P., ... The HAEL Study Group. (2019). The “Hypertension Approaches in the Elderly: a Lifestyle study” multicenter, randomized trial (HAEL Study): rationale and methodological protocol. *BMC Public Health*, 19(1), 657. <https://doi.org/10.1186/s12889-019-6970-3>

von Elm, E., Altman, D. G., Egger, M., Pocock, S. J., Gøtzsche, P. C., Vandenbroucke, J. P., & STROBE Initiative. (2007). The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Bulletin of the World Health Organization*, 85(11), 867–872. <https://doi.org/10.2471/blt.07.045120>

Tables

Table 1. Baseline characteristics of all participants.

Variables	Count (%) / Mean (SD)
Sex (n=351)	
Men	63 (18)
Women	288 (82)
Age (n=351)	
60 – 69 years	209 (59.5)
70 – 79 years	118 (33.5)
≥ 80 years	24 (7)
Body mass index (kg/m²) (n=334)	
Underweight ≤ 22	47 (14)
Normal weight > 22 and > 27	129 (39)
Overweight ≥ 27	158 (47)
Waist Circumference (cm)	
Women (n=281)	90 ± 10.7
Men (n=54)	99 ± 10.2
Education level (n=351)	
Incomplete elementary	62 (18)
Complete elementar	31 (9)
Incomplete high school	22 (6)
Complete high school	114 (32)
Incomplete undergraduate	20 (6)
Complete undergraduate	59 (17)
Graduate	43 (12)
Marital status (n=348)	
Never married	53 (15)
Married	141 (40.5)
Widower	81 (23.5)
Divorced	73 (21)
Ethnicity/Color (n=349)	
White	279 (80)
Black	36 (10.5)
Brown	26 (7.5)
Indigenous	8 (2)
Family income (n=347)	
Up to 2 MW	121 (35)
From 2 to 4 MW	135 (39)
From 4 to 10 MW	71 (20.5)
From 10 to 20 MW	20 (5.5)

Self-reported hypertension (n=347)	
No	161 (47)
Yes	171 (49)
Only in pregnancy	15 (4)
Office blood pressure (n=337)	
No self-reported hypertension (n=161)	
Systolic BP (mmHg)	120±18
Diastolic BP (mmHg)	72±10
With self-reported hypertension (n=165)	
Systolic BP (mmHg)	124±16
Diastolic BP (mmHg)	72± 9
Self-reported cardiovascular disease (n=349)	
No	299 (86)
Yes	50 (14)
Stable Angina	13 (26)
Acute myocardial infarction	10 (20)
Heart failure	9 (18)
Other	1 (2)
Self-reported diabetes (any type) (n=348)	
No	296 (85)
Yes	46 (13)
Only in pregnancy	6 (2)
Glycated Hemoglobin (%) (n=336)	
No self-reported diabetes (n=285)	5.5% ± 0.4
With self-reported diabetes (n=42)	6.9% ± 1.4
Self-reported hypercholesterolemia (n=349)	
No	160 (46)
Yes	189 (54)
Self-reported Arthritis/Rheumatism (n=341)	
No	240 (70)
Yes	101 (30)
Self-reported respiratory disease (n=350)	
No	313 (89.5)
Yes	37 (10.5)
Self-reported depression (n=349)	
No	284 (81)
Yes	65 (19)
Depression measured by the GDS-15 (n=352)	
No	283 (80.5)
Yes	69 (19.5)
General health status self-assessment (n=349)	
Very good	95 (27)
Good	181 (52)
Regular	63 (18)
Bad	7 (2)
Too bad	3 (1)

QoL measured by SF-6D (n=339)

0.87 ± 0.06

Note: SD: Standard Deviation; kg/m²: Kilogram Per Square Meter; cm: centimeter; MW: Minimum Wage; BP: Blood Pressure; QoL: Quality of Life; SF-6D: Short Form 6 Dimension; GDS-15: Geriatric Depression Scale.

Table 2. Dependent variable with the respective categorical variables used as predictors in each analysis.

Outcomes	Prevalence (%)	Mean 6WMT	CI 95% 6WMT
6MWT(m) (n=312)		500.0	(491.1 - 508.4)
Sex			
Female (n=262)	84.0	488.4	(479.7 - 497.1)
Male (n=50)	16.0	559.1	(536.1 - 582.1)
Number of medications			
None (n=69)	22.0	515.6	(497.0 - 534.1)
Up to 3 (n=149)	48.0	501.5	(489.7 - 513.4)
From 4 to 6 (n=79)	25.2	483.1	(466.5 - 499.8)
7+ (n=15)	4.8	463.9	(430.3 - 497.6)
Diabetes diagnosis			
No (n=266)	85.3	503.4	(494.0 - 512.9)
Yes (n=40)	12.8	476.0	(452.4 - 499.6)
Gestational (n=6)	1.9	495.3	(460.3 - 530.4)
Family income			
Up to 2 MW (n=109)	35.0	479.6	(466.6 - 492.6)
2 to 4 MW (n=118)	37.8	499.1	(484.2 - 513.9)
4 to 10 MW (n=65)	20.8	525.4	(508.3 - 542.5)
10 to 20 MW (n=20)	6.4	530.7	(489.5 - 571.8)
Depression - GDS-15			
No (n=257)	82.4	504.9	(495.6 - 514.2)
Yes (n=55)	17.6	475.8	(453.7 - 497.9)

		Mean Handgrip	95%CI Handgrip
Handgrip (kgf) (n=325)		27.0	(26.2 - 28)
Sex			
Women (n=271)	83.4	24.5	(24.0 - 25.1)
Men (n=54)	16.6	40.0	(37.8 - 42.3)
Diabetes diagnosis			
No (n=279)	85.9	27.2	(26.2 - 28.2)
Yes (n=40)	12.2	26.5	(24.0 - 29.0)
Gestational (n=6)	1.9	25.2	(22.4 - 28.0)
Number of medications			
None (n=69)	21.3	27.8	(25.8- 29.8)
Up to 3 (n=156)	48.0	26.6	(25.4 - 27.9)
From 4 to 6 (n=83)	25.5	27.8	(25.9 - 29.7)
7+ (n=17)	5.2	24.9	(21.3 - 28.5)

Note: 6MWT: 6-minute walk test; kgf: kilograms/force SE: Standard Error; CI: Confidence Interval; MW: Minimum Wage; GDS-15: Geriatric Depression Scale.

Table 3. Multivariable linear regression between 6MWT test, social determinants of health and clinical parameters.

Model 1: adjusted analysis for all predictors below			
Predictors (N=312)	β	CI 95%	P-value
Age	-4.04	-5.2 to -2.7	<0.001
Sex (male)	72.84	52.9 to 92.8	<0.001
BMI	-4.30	-6.0 to -2.6	<0.001
Number of medications (up to 3)	-0.56	-19.2 to 18.1	0.95
Number of medications (from 4 to 6)	-3.34	-24.7 to 18.0	0.76
Number of medications (7+)	-17.51	-55.0 to 20.0	0.36
QoL (SF-6D)	20.58	8.9 to 32.2	0.001
R ² = 0.32; Adjusted R ² = 0.30			
Model 2: adjusted analysis for significant variables in model 1 plus diabetes diagnosis, family income and depression			
Age	-4.05	-5.3 to -2.8	<0.001
Male sex	71.40	50.5 to 92.3	<0.001
BMI	-3.88	-5.6 to -2.1	<0.001
QoL (SF-6D)	18.48	6.3 to 30.6	0.003
Diabetes diagnosis (yes)	-14.05	-70.6 to 42.5	0.63
Diabetes diagnosis (no)	4.40	-48.6 to 57.4	0.87
Family income (up to 2 MW)	-13.53	-45.8 to 18.7	0.41
Family income (2 to 4 MW)	-17.19	-48.6 to 14.2	0.28
Family income (4 to 10 MW)	-4.22	-37.5 to 29.0	0.80
Depression - GDS-15 (no)	16.56	-4.0 to 37.1	0.11
R ² = 0.33; Adjusted R ² = 0.308			

Note: CI: Confidence Interval; BMI: Body Mass Index; QoL: Quality of Life; SF-6D: Short Form 6 Dimension; MW: Minimum Wage; GDS-15: Geriatric Depression Scale. The reference category for sex: female; number of medications: no medications; diabetes diagnosis: gestational diabetes; family income: 10 to 20 minimum wages and for depression: yes.

Table 4. Multivariable regression on generalized least squares, between handgrip test, social determinants of health and clinical parameters.

Model 1: adjusted analysis for all predictors below			
Predictors (n=325)	Variance	CI 95%	P-value
Age	-0.50	0.89 to 0.09	0.009
Sex (male)	65.86	55.89 to 76.5	<0.001
BMI	0.40	-0.09 to 1.0	0.13
Number of medications (up to 3)	-2.57	8.51 to -3.7	0.41
Number of medications (from 4 to 6)	-0.20	7.04 to -7.1	0.96
Number of medications (7+)	-13.06	22.74 to 2.2	0.02
Pseudo R²: 0.28			
Model 2: adjusted analysis for significant variables in model 1 plus diabetes diagnosis			
Age	-0.60	0.89 to 0.2	0.005
Male sex	65.2	55.3 to 75.8	<0.001
Number of medications (up to 3)	-1.88	7.8 to -4.6	0.56
Number of medications (from 4 to 6)	-1.51	-5.7 to 9.3	0.69
Number of medications (7+)	-10.77	21.3 to -1.2	0.07
Diabetes diagnosis (yes)	-2.76	10.1 to -5.0	0.47
Diabetes diagnosis (no)	2.94	-14.3 to 23.7	0.75
Pseudo R²: 0.29			

Note: 95%CI: Confidence Interval; BMI: Body Mass Index; R²: R Square. Estimation method: generalized least squares method. The reference category for sex: is female; number of medications: no medications and diabetes diagnosis: gestational diabetes.