

1 **Individual and community-level determinants of intention to use**
2 **contraceptive among married women in Ethiopia: A multi-level analysis of**
3 **National Survey**

4 Kegnie Shitu¹, Adugnaw Zeleke Alem², Tesfa Sewunet Alemneh², Bewuketu Terefe^{3*}

5
6 ¹Department of Health Education and Behavioral Science, Institute of Public Health, College of
7 Medicine and Health Sciences, University of Gondar, Gondar, Ethiopia

8 ²Department of Epidemiology and Biostatistics, Institute of Public Health, College of Medicine
9 and Health Sciences, University of Gondar, Gondar, Ethiopia

10 ^{3*}Department of Community Health Nursing, School of Nursing, College of Medicine and
11 Health Sciences, University of Gondar, Gondar, Ethiopia

12 **Abstract**

13 **Background:** Contraceptives are the most effective strategies to prevent unwanted pregnancies
14 and their consequences. Realizing intention to use contraceptives is a crucial stage to draft and
15 implement a successful family planning program.

16 **Objective:** This study aimed to identify individual and community level of factors affecting
17 women's intention to use contraceptives in Ethiopia.

18 **Method:** This study was based on a large national survey, Ethiopian Demographic and Health
19 Survey. A total weighted sample of 6,555 married/in union reproductive-age women were
20 included. Because of the hierarchical nature of the DHS data, a multilevel logistic regression
21 model was used to study individual and community-level factors that may influence intention to

22 contraceptive use. A 95% confidence interval and a p value of less than 0.05 were used to
23 declare statistical significance.

24 **Result:** The overall intention to use contraceptives was 48.63% (95% CI 47.42, 49.84).
25 Participants' age range of 25-34 years (AOR = 0.42, 95 CI% (0.22, 0.79)) and 35-49 years (AOR
26 = 0.12, 95% CI: (0.05, 0.28)), husband with primary education (AOR = 1.60, 95% CI: (1.02,
27 2.50)), heard of contraceptives from their community (AOR = 1.91. 95% CI: (1.29, 2.83)), ever
28 used contraceptives (AOR = 4.48, 95% CI: (2.91, 6.88)) and having six or more children (AOR =
29 0.46, 95% CI: (0.23, 0.9)) were individual factors significantly associated with interceptive
30 intention. From community level factors, high community family planning utilization rate
31 (AOR = 2.29, 95% CI: (1.36,3.86)) was associated with intention to use contraceptive.

32 **Conclusion:** More than half of married women were not intended to use contraceptives.
33 Intention to use contraceptive was affected by individual and community level attributes. Thus,
34 public health interventions particularly that could increase information dissemination regarding
35 contraceptives among the communities and enhance community level contraceptive utilization
36 rate are required at the national level to improve contraceptive utilization.

37 **Keywords:** Intention to contraception, married women, Multilevel analysis, Contraceptive,
38 Ethiopia

39 **Introduction**

40 A couple's purposeful effort to restrict or space the number of children they have through the use
41 of contraceptive techniques is referred to as family planning. Contraceptive methods are divided
42 into two categories: modern and old. Male and female sterilization, the intrauterine contraceptive
43 device (IUD), implants, injectables, the pill, male and female condoms, emergency

44 contraception, the standard 9 days method (SDM), and the lactational amenorrhea method are all
45 examples of modern approaches (LAM). Traditional approaches include things like rhythm,
46 withdrawal, and other folk methods(1).

47 Reports pointed out that, in 2019 there were about 1.1 billion reproductive age women who have
48 a need to use contraceptive methods, however, around 270 million of them had faced unmet need
49 of contraceptive methods(2, 3). Sustainable development goals (SDG) and united nations world
50 family planning report indicated that the total of rate satisfaction of mothers in family planning is
51 stopped at around 78%, nevertheless due to several hindering factors the rate satisfaction in
52 Africa regions has shown lowest rate (56%)(4). If all low and middle countries can satisfy
53 women in contraceptive services, they will have a chance of reduced unintended pregnancies,
54 unplanned births, and induced abortions approximately by three-quarters from 39 million to 22
55 million, from 30 million to 7 million and from 48 million to 12 million every single year
56 respectively with 76,000 fewer maternal deaths(5).

57 In the present scientific fact, the global community has reached on consensus agreements
58 regarding family planning by reducing the number of women's exposure to pregnancies, by
59 limited more than 4 births (6, 7), reducing unsafe abortion from unintended pregnancies (8, 9),
60 declining newborn and infant mortality rates (5), preventing the transmission of HIV/AIDS from
61 the mother to the child by 93% (10), empowering mothers and girls by staying them in
62 school(11), by improving adolescents' reproductive health, social and economic wellbeing(12).

63 The task of family planning is exponentially difficult and daunting in sub Saharan Africa (SSA),
64 hence According to the world bank project report the number of populations SSA has shown an
65 increasement of about ten times between 1990 and 2050(13), this in turn shows population

66 growth is fast in Africa(14). SSA has shown a few declines from 6.3 to 4.6 births per women
67 from 1990 to 2019(14).

68 Like the rest of the world community, Ethiopia is struggling with family planning issues and has
69 tried to increase contraception coverage by 69% by 2020, however has failed and it needs other
70 fertility and family planning implications (15). Studies on contraception prevalence in Ethiopia
71 show that it is low and far apart in terms of prevalence from one to other studies. For
72 illustrations, a study conducted in Benishangul Gumuz revealed that the prevalence of
73 contraceptive was 18.6%(16), 38% in southern Ethiopia Wolayita Sodo(17), 19.9% in eastern
74 Ethiopia (18). Other two important studies were also conducted based on demographic and
75 health survey data on contraceptive prevalence to use and discontinuation discovered that
76 20.42% and 32.2% respectively(19, 20). Children born from mothers who use modern
77 contraceptive methods have high rate of survival implications than their counterparts(21). Such
78 studies also identified some crucial factors which have positive or negative statistical
79 implications on contraceptive method use. This includes age, educational level, knowledge,
80 attitude (16, 17), residence, region, occupational status, number of children, women self-
81 decision, television watching (19, 20), counseling on contraceptive methods, (16, 20), myths and
82 misconceptions(17). There were also some other factors associated with intention to use
83 contraceptive methods. Fear of side effects, male partner opposition, inter spousal discussion,
84 discussion with health care providers, perceived cultural acceptability, status of gravidity, marital
85 status and post-natal care utilization were among the variables associated with the outcome
86 variable(22-27).

87 This study addresses two major research objectives. First, studies done on contraceptive methods
88 in Ethiopia are so limited in study area and study population that is difficult to determine for the

89 general population of the country. Second, these studies did not include events that may have a
90 potential effect on intention to use contraception methods at community level factors. Therefore,
91 this study aimed to include both individual and community level factors to determine the
92 intention to use contraceptive methods among married reproductive women. Identifying
93 important factors affecting intention to use contraceptive methods at the community and
94 individual level, this study contributes to Ethiopia, would have a paramount importance to
95 increase birth control by controlling birth rates. Policymakers and other stakeholders working on
96 family planning, women and children's health will access an up-to-date nationwide result from
97 the findings of this study to develop, improve and implement their plans accordingly.

98 **Methods**

99 **Data source, population, and sampling procedure**

100 The present study was based on the most recent Ethiopian Demographic and Health Survey
101 (EDHS) data. The survey was conducted from January 18, 2016, to June 27, 2016. For this
102 survey, a complete list of 84, 915 Enumeration Areas (EAs) from the Ethiopian Population and
103 Housing Census was used as a sampling frame. A stratified two-stage cluster sampling technique
104 was employed. In the first stage, 645 EAs were selected. In the second stage, an average of 28
105 households was selected per cluster/EA. The data is freely available in public and we accessed
106 the dataset used for the present study after we registered and received an authentication letter
107 from the Demographic and Health Survey (DHS) program at The DHS Program - Ethiopia:
108 Standard DHS, 2016 Dataset. For this study, a total weighted sample of 6, 555 married/in union
109 women who are not using contraceptive were included. Detailed information about the sampling
110 strategy, questioner, or other important information is available in EDHS of 2016(28).

111 **Variables of the Study**

112 The outcome variable of the present study was the intention to use contraceptives among
113 contraceptive non-user women. The variable was dichotomized into **1** = “Intended” and **0** = “Not
114 intended”. The independent variables were further classified into individual level (level 1)
115 variables and community level (level 2) variables. Individual-level variables included age,
116 religion, educational status, husband’s/partner’s age and education, family wealth index, current
117 working status, family planning message exposure, knowledge of family planning methods, ever
118 use of contraceptive, fertility preference, desire to have children, and a number of children.
119 Whereas, community variables involved variables directly taken with no aggregation (residence
120 and contextual region), and variables obtained by aggregating individual variables into their
121 respected community (community poverty, community female education, community family
122 planning message exposure, community antenatal care service utilization rate, community family
123 planning utilization rate, community women’s education, and community health facility
124 distance). The aggregates were computed using the mean values of the proportions of women in
125 each category of a given variable. Since the aggregate values of each variable don’t follow a
126 normal distribution curve, we categorized the aggregate values of a cluster into groups based on
127 median values.

128 **Operational definitions**

129 **Household wealth quintile:** The wealth index classifications were done in quintiles: poorest;
130 poor; middle; rich; richest. These were computed using principal component analyses (PCA).
131 This variable was further categorized into three categories (Poor, Medium, and Rich) by merging
132 the lower two (poorest and poor) quintiles and the upper two (richest and rich) quintiles.

133 **Family planning message exposure:** This variable was computed from the frequency of
134 exposure to family planning messages from television, radio, magazine or mobile messages The
135 variable was categorized into two parts: Not exposed when the participant is exposed to none of
136 the four channels at least once a week; and exposed if the participant was exposed to family
137 planning messages at least once a week from one of the four channels.

138 **Contextual region:** For this survey regions were categorized into three categories (agrarian,
139 pastoral, and metropolitan) that may have a strong relationship to health information seeking and
140 intention to use contraceptives. The Tigray, Amhara, Oromia, SNNP, Gambella, and
141 Beneshangul Gumuz were recorded as agrarian. The Somali and Afar regions were merged to
142 form the pastoralist region and the city administrations- Addis Ababa, Dire Dawa, and Harar
143 were combined as metropolitan (1).

144 **Community female education:** This is the aggregate value of the educational levels of women
145 based on the average of proportions of educational levels in the community. It was defined as
146 low if the proportion of women with secondary education & above in the community was 0 –
147 12.4 % and high if the proportion was 12.5 –100 %.

148 **Community media exposure:** This variable was derived from the individual responses for
149 exposure to radio or television. It was defined as low if the proportion of women exposed to
150 media in the community was 0–18.7 % and high if the proportion was 18.8–100 %.

151 **Community ANC utilization rate:** This variable is also derived from the individual values for
152 ANC utilization. It was defined as low if the proportion of women who attended at least one
153 ANC visit in the community was 0 – 81.3 % and high if the proportion was between 81.4 –100
154 %.

155 **Community poverty:** With the same procedure, this variable is also derived from an individual
156 household's wealth index. It was defined as high if the proportion of women from the two lowest
157 wealth quintiles in a given community was 25.9–100 % and low if the proportion was 0–26 %.

158 **Community distance to the health facility:** The variable was aggregated from individual
159 perceived distance to a health facility is a big problem. It was as categorized as low if the
160 proportion of women who perceived health facility distance as a big problem in the community
161 was 0–42.2% and categorized as high if the proportion was between 42.2% and 100%.

162 **Data processing and Analysis**

163 Data were extracted from individual records (IR) files and further coding and transformations
164 were done using statistical software, STATA version 14. The weighted samples were used for
165 analysis to adjust for unequal probability of selection and non-response in the original survey.
166 Since the EDHS employed multi-stage stratified cluster sampling techniques, the data have a
167 hierarchical structure. In such kinds of situations, single-level logistic regression is not
168 recommended because Traditional multiple regression techniques treat the units of analysis as
169 independent observations. One consequence of failing to recognize hierarchical structures is that
170 standard errors of regression coefficients will be underestimated, leading to an overstatement of
171 statistical significance. In this point of view, to draw valid inference and conclusion an advanced
172 statistical model which takes the hierarchy of the data into account is required. Therefore, a
173 multivariable multilevel binary logistic regression model was used to estimate the fixed and
174 random effects of the factors associated with intention to use contraceptives. Four models were
175 constructed. The first model also called an empty model which was fitted without any
176 explanatory variables. This model was specified to decompose the amount of variance that
177 existed between communities. The null model is important for understanding the community

178 variations, and we used it as a reference to estimate how much community factors were able to
179 explain the observed variations in the intention to use contraceptives. Moreover, this model was
180 used to justify the use of a multilevel statistical framework as it is a litmus paper on whether
181 multilevel or the traditional logistic regression should be used. It was assessed using the Log
182 Likelihood Ratio test (LLR), Median Odds Ratio (MOR), Intra-class Correlation Coefficient
183 (ICC), and Proportional Change of Variance (PCV). The second model contained only
184 individual-level factors. The third contained only community-level factors. Whereas, the final
185 (fourth) model containing both individual and community-level factors. Moreover, the model
186 comparison was done using model deviance, a model with the lowest deviance selected for
187 reporting and interpretation results.

188 **Result**

189 **Individual level characteristics**

190 A total of 6,555 weighted sample of married/ in union and contraceptive non-users were
191 involved in the present study. The median age of the women was 30 years old with an
192 interquartile range (IQR) of 13 years. The median (IQR) husband's/partner's age was 38 (16)
193 years. More than half of (56.6%) of the participants were affiliated to Christianity. Majority
194 (65.68%) and more than half (50.51%) of the participants and their husbands/partners did not
195 attend formal education respectively. About 2999 (46%) of the participant were form a house
196 hold with poor wealth status. Concerning exposure to family planning messages, only 1572
197 (24%) of the participants were exposed to family planning message from at least one of the four
198 main (television, radio, magazine and mobile message) channels.

199 Almost all (98%) of the participants knew at least one of the family planning methods. However,
 200 only 2257 (34.4%) of the participants had ever used at least once a time. Decision for not using
 201 contraceptive was mainly made jointly by the respondents and their partner whereas 569 (10.43)
 202 participants reported that their partners were the main decision makers for not using
 203 contraceptives. Regarding to fertility, about 2840 (43.3%) participants hadn't desire to have
 204 children in the future. In more than on third (35.3%) the cases both the respondents and their
 205 partners want the same number of children to have in the future (Table 1).

206 Table 1: Sociodemographic and other individual characteristics of married/in union women in
 207 Ethiopia (n=6555)

Variables	Intended to use FP		Total, n (%)
	No, n (%)	Yes, n (%)	
Age			
15-24	436 (30.15)	1,011 (69.85)	1447 (22.07)
25-34	1,166 (43.17)	1,535 (56.83)	2701 (41.21)
35-49	1,765 (73.35)	642 (26.65)	2407 (36.72)
Husband's/partner's age			
15-24	89(28.14)	228 (71.86)	317 (4.84)
25-34	698 (33.91)	1361 (66.09)	2059 (31.42)
35-44	1011 (49.06)	1050 (50.94)	2061 (31.45)
45-54	894(69.73)	388 (30.27)	1282(19.56)
>=55	674 (80.83)	160 (19.17)	834 (12.73)
Religion			
Cristian	1,634 (45)	1,991 (55)	3625 (56.67)
Muslim	1639 (59.14)	1133 (40.86)	2772(43.33)
Highest educational level			
No-formal education	2510 (58.29)	1795 (41.71)	4305 (65.68)
Primary	700 (40.1)	1048 (59.9)	1748 (26.67)
Secondary	82 (26.24)	230 (73.76)	312 (4.76)
Higher	75 (39.55)	115 (60.45)	190 (2.90)
Husband's/partner's education			
No formal education	1999 (60.88)	1285 (39.12)	3284 (50.51)
Primary education	972(41.77)	1355 (58.23)	2327 (35.79)

Secondary education		205 (37.7)	338 (62.3)	543 (8.35)
Higher education		164 (47.11)	184 (52.89)	348 (5.35)
Currently working				0
No		2,471 (51.99)	2,282 (48.01)	4753 (72.51)
Yes		897 (49.76)	905 (50.24)	1802 (27.49)
Household wealth Index				
Poor		1,628 (54.28)	1,371 (45.72)	2999 (45.75)
Medium		630 (48.76)	662 (51.24)	1292 (19.71)
Rich		1,110 (49.02)	1,154 (50.98)	2264 (34.54)
Exposure to family planning (FP) messages				0
No		2,740 (55)	2,243 (45.02)	4983 (76.02)
Yes		628(39.93)	944 (60.07)	1572 (23.98)
Source of FP message exposure				
Television	No	3051 (52)	2765 (47.54)	5816 (88.73)
	Yes	316 (42.84)	421 (57.16)	739 (11.27)
Radio	No	2874(54)	2448 (46)	5322 (81.19)
	Yes	494 (40.05)	739 (59.95)	1233 (18.81)
Magazine/newspaper	No	3311 (52)	3085 (48.23)	6396 (97.57)
	Yes	56 (35.59)	103 (64.41)	159 (2.43)
Mobile phone	No	3304 (51.33)	3132 (48.67)	6436 (98.18)
	Yes	64 (53.57)	55 (46.43)	119 (1.82)
Told of family planning at health facility				
No		887 (48)	948 (51.66)	1835 (62.46)
Yes		396 (35.92)	707 (64.08)	1103 (37.54)
Told of family planning by fieldworker				
No		396 (54.8)	327 (45.17)	723 (40.89)
Yes		430 (41.15)	615 (58.85)	1045 (59.11)
Heard of family planning from the community				
No		2,149 (54. 43)	1,799 (45.57)	3948 (60.23)
Yes		1,219 (46.75)	1,388 (53.25)	2607 (39.77)
Know a modern contraceptive				
Don't know		123 (90.0)	14 (10.38)	138 (2.11)
Know		3,244 (50.56)	3,173 (49.44)	6417 (97.89)
Ever use of contraceptive				
No		2708 (63.0)	1590 (37.0)	4298 (65.57)
Yes		660 (29.24)	1597 (70.76)	2257 (34.43)
Who decide for not using contraceptive				
Mainly respondent		917(55.86)	725 (44.14)	1642 (30.09)
mainly husband/partner		338 (59.39)	231 (40.61)	569 (10.43)
joint decision		1716 (54.45)	1436 (45.55)	3152 (57.76)

Other	73 (77.47)	21 (22.53)	94 (1.72)
Fertility preference			
Have another	1691 (46.52)	2024 (54.48)	3715 (56.67)
Undecided	233 (60.43)	152 (39.57)	385 (5.87)
No more	1288(56.1)	1007 (43.9)	2295 (35.01)
Declared infecund	156 (98.03)	4 (1.97)	160 (2.44)
Desire for more children			
No	1676 (59.04)	1164 (40.96)	2840 (43.33)
Yes	1691 (45.52)	2024 (54.48)	3715 (56.67)
Husband's children desire			
Both wants the same	1,094(47.26)	1,221 (52.74)	2315 (35.32)
Husband wants more	1,019 (55.18)	828 (44.82)	1847 (28.18)
Husband wants fewer	233 (51.7)	218 (48.3)	451 (6.88)
Don't know	1,021 (52.59)	921 (47.41)	1942 (29.63)
No of Children			
0-2	906 (40.61)	1,325 (59.39)	2231 (34.04)
3-5	1,280 (49.57)	1,301 (50.43)	2581 (39.37)
6 and above	1,182 (67.83)	561 (32.17)	1743 (26.59)

208 **Community level factor**

209 Majority (87.84%) of the participants were from rural residency. Whereas, 4178 (63.7%), 4734
 210 (72.2%), 2803 (42.76), 4808 (73.35) and 3748 (57.18) of the participants were from a
 211 community with high poverty, low community female's education, low exposure to family
 212 planning, low community ANC utilization rate and low women's empowerment respectively.
 213 Moreover, in the chi2 analysis, residence, community poverty, region, community level female's
 214 education, community ANC utilization rate, community exposure to family planning messages,
 215 community family planning utilization rate and community health facility distance problem were
 216 significantly associated with intention to use family planning at p-value <0.05 (Table 2).

217 Table 2: Community level variables descriptive result by informed choice among contraceptive
 218 user women in Ethiopia (n= 6,555)


Variables	Intention to use FP, n (%)	Total, n (%)	p-value
-----------	----------------------------	--------------	---------

	No	Yes		
Residence				
Urban	394 (49.50)	403 (50.50)	797 (12.16)	
Rural	2,973 (51.63)	2,785 (48.37)	5758 (87.84)	0.003
Contextual Region				
Agrarian	2,836 (48.41)	3,023 (51.59)	5859 (89.38)	
Pastoral	425 (84.35)	79 (15.65)	504 (7.69)	
Metropolitan	106 (55.50)	86 (44.50)	192 (2.93)	<0.001
Community poverty				
Low	1,202 (50.59)	1,175 (49.41)	2377 (36.26)	
High	2,165 (51.82)	2,013 (48.18)	4178 (63.74)	<0.001
Community female education				
Low	2468 (52.14)	2266 (47.86)	4734 (72.22)	
High	900 (49.39)	921 (50.61)	1821 (27.78)	<0.001
Community exposure to FP messages				
Low	1,498 (53.45)	1,305 (46.55)	2803 (42.76)	
High	1,869 (49.83)	1,882 (50.17)	3751 (57.22)	<0.001
Community ANC utilization rate				
Low	2558 (53.21)	2250 (46.79)	4808 (73.35)	
High	809 (46.33)	938 (53.67)	1747 (26.65)	<0.001
Community FP utilization rate				
Low	1,401 (64.27)	780 (35.73)	2181 (33.27)	
High	1,966 (44.95)	2,408 (55.05)	4374 (66.73)	<0.001
Community health facility distance problem				
Low	996 (48.86)	1,043 (51.14)	2039 (31.11)	
High	2,371 (52.51)	2,145 (47.49)	4516 (68.89)	<0.001
Community women's empowerment				
Low	1,979 (52.82)	1,769 (47.18)	3748 (57.18)	
High	1,388 (49.45)	1,419 (50.55)	2807 (42.82)	0.287

219 *Not: ANC= Antenatal care, FP= Family planning*

220 **Intention to use contraceptive**

221 Only 3368 (48.6%, 95% CI 47.42, 49.84) of married reproductive age women were intended to
 222 use contraceptive in the future (Figure 1).



223

224 **Figure 1:** Intention to use contraceptives among married reproductive age women in Ethiopia

225 (n= 6, 555)

226 **Factors associated with intention to use family planning**

227 In the multilevel multivariable analysis, age, education, heard of family planning from the
228 community, ever use of contraceptives, number of children, and community level family
229 planning utilization rate were significant factors associated with participant's intention to use
230 contraceptives in the future.

231 Participants within the age range of 25-34 years (AOR = 0.42, 95 CI% (0.22, 0.79)) and 35-49
232 years (AOR = 0.12, 95% CI: (0.05, 0.28)) were less likely to be intended to use contraceptives in
233 the future compared to participants with the age range of 15-24 years. Women's husband/partner
234 with primary education were 1.6 times more likely to be intended to use contraceptives
235 compared to husbands with no formal education (AOR = 1.60, 95% CI: (1.02, 2.50)).
236 Participants heard of contraceptives from their community (AOR = 1.91, 95% CI: (1.29, 2.83))
237 and ever used contraceptives (AOR = 4.48, 95% CI: (2.91, 6.88)) were almost two and 4.5 times

238 more likely to be intended to use contraceptives in the future respectively. In addition to this,
 239 women who had six or more children (AOR = 0.46, 95% CI: (0.23, 0.9)) were less likely to be
 240 intended to use contraceptives. Regarding community level variables, Women from a community
 241 with high rate of family planning utilization were 2.3 times more likely to be intended to use
 242 contraceptives (AOR = 2.29, 95% CI: (1.36,3.86)).

243 Table 3: Individual and community level facto associated with intention to use family planning
 244 among married/ in union contraceptive non-user women in Ethiopia (n=6,555).

Independent variables	Null model	Model I	Model III	Model IV
		AOR [95% CI]	AOR [95% CI]	AOR [95% CI]
Age of the respondent				
15-24	-	1	-	1
25-34	-	0.39 (0.21, 0.73) *	-	0.42 (0.22, 0.79) *
35-49	-	0.10 (0.04, 0.24) *	-	0.12 (0.05, 0.28) *
Husband/partner's age				
15-24	-	1	-	1
25-34	-	1.05 (0.42, 2.61)	-	0.90 (0.36, 2.27)
35-44	-	1.45 (0.54, 3.89)	-	1.27 (0.47, 3.47)
45-54	-	1.12 (0.38, 3.32)	-	1.03 (0.34, 3.08)
>=55	-	0.39 (0.12, 1.27)	-	0.40 (0.12, 1.33)
Religion				
Christian	-	1	-	1
Muslim	-	0.55 (0.36, 0.85) *	-	0.76 (0.48, 1.22)
Education				
No formal education	-	1	-	1
primary	-	1.01 (0.64, 1.62)	-	1.11 (0.69, 1.77)
secondary	-	1.23 (0.56, 2.71)	-	1.86 (0.83, 4.20)

			2.72)	
higher	-	0.64 (0.23, 1.80)	-	0.96 (0.34, 2.71)
Husband/partner's education				
No formal education	-	1	-	1
Primary education	-	1.62 (1.04, 2.54) *	-	1.60 (1.02, 2.50) *
Secondary education	-	0.84 (0.43, 1.65)	-	1.06 (0.54, 2.10)
Higher education	-	1.29 (0.58, 2.90)	-	1.58 (0.70, 3.56)
Exposure to FP messages				
Not exposed	-	1	-	1
Exposed	-	1.13 (0.73, 1.75)	-	1.17 (0.74, 1.84)
Told of FP at health facility				
No	-	1	-	1
yes	-	1.45 (0.93, 2.26)	-	1.35 (0.87, 2.10)
Told of FP by field worker				
No	-	1	-	1
Yes	-	1.03 (0.65, 1.62)	-	1.01 (0.64, 1.58)
Heard of FP from the community				
No	-	1	-	1
Yes	-	2.13 (1.44, 3.15) *	-	1.91 (1.29, 2.83) **
Knowledge of FP				
Don't know	-	1	-	1
Know	-	8.62 (0.97, 76.33)	-	7.51 (0.85, 66.53)
Ever use of FP				
No	-	1	-	1
Yes	-	4.89 (3.18, 7.52) *	-	4.48 (2.91, 6.88) **
Decision maker for not using FP				
Mainly the respondent	-	1	-	1
Mainly husband, partner	-	1.02 (0.49, 2.10)	-	1.07 (0.52, 2.23)
Joint decision	-	1.23 (0.82, 1.86)	-	1.24 (0.83, 1.88)
Other	-	1.20 (0.20, 2.72)	-	1.29 (0.21, 7.90)

			7.23)	
Fertility preference				
Have more/undecided	-	1	-	1
No more/declared infecund	-	1.17 (0.49, 2.81)	-	1.03 (0.44, 2.40)
Desire to have children				
No	-	1	-	1
Yes	-	1.04 (0.44, 2.42)	-	1.03 (0.45, 2.34)
Household wealth index				
Poor	-	1	-	1
Medium	-	1.20 (0.70, 2.06)	-	1.00 (0.58, 1.73)
Rich	-	0.86 (0.52, 1.42)	-	0.92 (0.51, 1.65)
Husbands desire for children				
Husband wants the same	-	1	-	1
husband wants more	-	0.70 (0.44, 1.13)	-	0.68 (0.42, 1.09)
husband wants fewer	-	1.18 (0.59, 2.39)	-	1.16 (0.58, 2.33)
don't know	-	0.91 (0.57, 1.44)	-	0.94 (0.59, 1.48)
No. of children				
0-2	-	1	-	1
3-5	-	0.81 (0.48, 1.35)	-	0.74 (0.44, 1.24)
6 and above	-	0.51 (0.26, 1.00)	-	0.46 (0.23, 0.91) *
Residence				
Rural	-	-	1	1
Urban	-	-	0.92 (0.67, 1.26)	0.68 (0.32, 1.45)
Region				
Pastoral	-	-	1	1
Agrarian	-	-	2.23 (1.80, 2.76)	1.52 (0.91, 2.56)
			*	
Metropolitan	-	-	1.28 (0.94, 1.75)	0.86 (0.41, 1.79)
Community poverty				
High	-	-	1	1
Low Poverty	-	-	0.89 (0.70, 1.13)	1.24 (0.68, 2.23)
Community female education				
Low	-	-	1	1

High	-	-	1.10 (0.88, 1.38)	0.61 (0.35, 1.10)
Community FP message exposure				
Low	-	-	1	1
High	-	-	1.31(1.06,1.62)*	0.94 (0.56, 1.58)
Community ANC utilization rate				
Low	-	-	1	1
High	-	-	1.22(0.98, 1.52)	1.07 (0.66, 1.76)
Community FP utilization				
Low	-	-	1	1
High	-	-	3.21 (2.61,3.95)	2.29 (1.36,3.86) *
Community HF distance problem				
High	-	-		1
Low	-	-	0.84 (0.68, 1.04)	0.71 (0.44, 1.13)
Community women's empowerment				
Low	-	-	1	1
High	-	-	0.95 (0.79, 1.14)	0.90 (0.59, 1.39)
Random parameters and model comparison				
Community level variance	1.36	0.74	0.55	0.50
ICC (%)	25.8	15.9	12.2	11.31
MOR (95% CI)	3.03(2.74, 3.37)	2.26 (1.62, 3.99)	2.02 (1.86, 2.21)	1.96(1.40, 3.79)
PCV (%)	Reference	45.6	59.9	63.25
DIC (-2LLR)	-4154.98	931.7	7962.8	900.0

245 Note: * significant at p-value <0.05, ** Significant at p-value <0.001, ICC= Intra cluster
 246 correlation, MOR= Median Odds Ratio, PCV= Proportional Change of Variance, DIC=
 247 Deviance information criterion, LLR=Log Likelihood Ratio

248 Discussion

249 The aim of this study was to assess future intention to use contraceptives and associated factors
 250 among married/ in union women who were not current contraceptive user. The proportion of
 251 participants who were intended to use contraceptive in the future was 48.63 (47.42-49.84). This
 252 figure is less observed than studies conducted in Indonesia 63%(29), in Ghana 70%(30, 31), in

253 Uganda(32), in Malawi (33), in Ethiopia 84.3% (34) and it is almost in agreement with studies
254 conducted in Ghana 49.3%(35), however this figure is higher than studies investigated in
255 Ethiopia (17, 36). The possible justification why the observed findings of this study is less than
256 the above mentioned one might be due to countries profiles of family planning experience,
257 participants attitude, knowledge and education backgrounds towards family planning intention.
258 On the other hand, this study founds significantly higher prevalence of intention to use
259 contraceptives, might be regarding to minimizing confounding factors both on the individuals
260 and community level factors that could have positive or negative implication on the intention to
261 use contraceptives. Majority of those studies concluded the intention to use contraceptive based
262 on individual factors with a very limited sample size and setting of the study. This might have
263 increased their prevalence findings.

264 This study discovered that age was one of the main predictor's variables. Participants whose age
265 group is from 25 to 34 years and 35 to 49 years old have less likely intention to use
266 contraceptives in their future family than the younger. This finding is in agreement with many
267 other studies conducted at various study setting with different times. A study conducted in
268 Cameron (37), in Zambia(38), in Uganda (32), in Malawi (33), and in Ethiopia (39). This is
269 related to the fact that older women may be in less need of contraceptive methods(40). This also
270 can be explained by as mothers gets older and older, their ability to give birth may decrease and
271 they may experience health problems, sometimes due to pregnancy or related reasons. On the
272 other hand, they may think they are infertile due to their age and also may have a negative view
273 of modern contraceptives.

274 The study also depicted that those women's husband who have formal education has more likely
275 intention to use contraceptive in the near future than their counterparts having no formal

276 education. This investigation almost similar with other studies done in Ethiopia (23, 41). In
277 Pakistan (42), in Cameroon (43), in Uttar Pradesh (44) and in Nigeria (45). Male partners,
278 especially in rural Ethiopia, have a tradition of domination family matters, sometimes by their
279 own interests. Education is the major key to solve this challenges of self-determination in family
280 issues (46, 47). Education can enhance humans romantic relationship, sense of humor and ability
281 to make informed decision making cultures with their female partners (48, 49). The results of
282 this study also illustrate this fact.

283 The other interesting variable was those participants who had heard from the community about
284 contraceptive or family planning have more tendency of intention to use contraceptives. This
285 finding is supported by several studies conducted in low and middle income countries' (50), in
286 Ethiopia(51), in SSA (52), in Rwanda(53), and in Nigeria (54). This shows that a favorable
287 community level information of contraceptives were significant predictors of intention to use
288 modern contraceptive methods and should take into consideration during the development of
289 Ethiopian family planning programs. The increase in the number of people who have heard of
290 the positive benefits of contraceptives in the community is one indication that mothers are ready
291 to do whatever is best for the community.

292 Individuals who have ever used family planning at least once in their life time has more
293 favorable interest to use the contraceptive methods once again in their future life. These
294 participants who have ever used contraceptives at least once may increase their awareness and
295 knowledge about the service over time. Their awareness and knowledge in turn might increase
296 the intention to use contraceptives through the exposure and counseling received from health
297 care providers. This idea is supported by studies conducted in Ethiopia (34, 39), in Yemen(55),
298 and in Malaysia (56)

299 Mothers having six and more than six children have shown less positive intention to apply family
300 planning in the near future than their counterparts. This finding is in disagreement studies
301 conducted in Ethiopia (39, 57). The reason why mothers with six or more than six number of
302 children do not use contraceptives is because they might have a low and negative attitude of
303 contraception or have much pressure and future need of children on themselves, their partners or
304 the community to have more children or think they will not have children later due to their aging
305 or uneducated. They may be also using contraceptive method is offensive and forbidden in their
306 religion and cultural practices (58). Those participants might have also faced geographic
307 challenges and unavailability of resources (59). Some wealthier women also mentioned that
308 using contraceptive methods have health concern issues and infrequent sex (50).

309 Participants living in a community which family planning coverage is more utilized have a
310 greater positive tendency to apply family planning for future upcoming tears than those living in
311 less utilized communities. This finding is supported by studies conducted in Pakistan, Rwanda
312 and Taiwan with a conclusion of accessing of family planning services did not depend only
313 mothers own individual level factors but also by the factors of her household and community
314 level factors (60-62). Other study in Ethiopia also again showed that, Ethiopia will more
315 privilege through community based outreach, and interpersonal communication that could
316 effectively modified the knowledge and behavior of women of intention to use contraceptives
317 (63). Certain community members are almost similar in the knowledge, attitudes and way of
318 life. Thus, it is important for mothers to decide whether or not to use contraceptives. For
319 example, when people around her who have a positive view of modern contraception and have a
320 better quality and a healthier baby, mother will develop more courage and self-confidence the
321 intention to use modern contraceptives. Perhaps she could explain to others in her community

322 that using contraception would be better for her, her baby and her family. Or, if she has a
323 misconception view of birth control, she is more likely to benefit from it if the majority of the
324 community benefits. She can easily find the information he needs through her friends and
325 community.

326 **Conclusion**

327 The findings of this study revealed that intention to use contraceptives in the near future among
328 married women current contraceptive nonusers was very low. Age of women, educational status
329 of male partners, ever use of contraceptives, heard of contraceptives information from the
330 community, having six and a greater number of children and higher community utilization family
331 planning utilization rate were factors that statistically significant variables associated with
332 intention to use contraceptives. Thus, public health interventions particularly that could increase
333 information dissemination regarding contraceptives among the communities and enhance
334 community level contraceptive utilization rate are urgently required at the national level to
335 address potential hindering factors and to improve the rate of contraceptive utilization in
336 Ethiopia.

337 **Strength and limitation of the study**

338 The strength of this study includes the use of a large sample size. As the sample size increases,
339 the sample gets closer to the actual population, which decreases the potential for deviations from
340 the actual population. In addition to this, an advanced statistical model that can take the nature of
341 the data into account was employed. However, the present study should be interpreted with
342 several limitation that includes; because of the cross-sectional nature of the study, making casual

343 inferences about the observed associations might not be possible. Moreover, social desirability
344 bias might be introduced as far as the data were entirely based on self-reports.

345 **Declarations**

346 **Acknowledgments**

347 Authors acknowledged the DHS program office to take the ethical consideration of this study

348 **Statement of authors contribution**

349 KS: Conceptualization, data curation, formal analysis, investigation, methodology, resources,
350 software, validation, visualization, writing – original draft, writing – review & editing. BT, AZA,
351 TSA: Data curation, formal analysis, investigation, methodology, resources, software, validation,
352 visualization, writing – original draft, writing – review & editing. All the authors read and
353 approved the final manuscript.

354 **Funding statement**

355 This research did not receive any specific grant from funding agencies in the public, commercial,
356 or not-for-profit sectors.

357 **Competing interest statement**

358 The authors declare no conflict of interest.

359 **Availability of data**

360 Data for this study were obtained from the DHS Program.

361 **Consents for publication**

362 Not applicable

363 **Ethical approval and consent to participate**

364 Ethical approval and permission letter was requested online to DHS program at
365 www.dhsprogram.com to access the data for this study and DHS program was granted the
366 permission through email.

367 **References**

- 368 1. EPHIEE I. Ethiopia mini demographic and health survey 2019: key indicators. USA:
369 EPHI and ICF; 2019.
- 370 2. Kantorová V, Wheldon MC, Ueffing P, Dasgupta AN. Estimating progress towards
371 meeting women's contraceptive needs in 185 countries: A Bayesian hierarchical modelling
372 study. PLoS medicine. 2020;17(2):e1003026.
- 373 3. United Nations Department of Economic and Social Affairs Population Division NY.
374 Family Planning and the 2030 Agenda for Sustainable Development • Data Booklet. 2019.
- 375 4. United Nations DoEaSA, Population Division World Family Planning -Highlights
376 (ST/ESA/SER.A/414). 2017.
- 377 5. Darroch JE, Sully E, Biddlecom A. Adding it up: investing in contraception and maternal
378 and newborn health, 2017—supplementary tables. New York, NY: The Guttmacher Institute.
379 2017.
- 380 6. Stover J, Ross J. How increased contraceptive use has reduced maternal mortality.
381 Maternal and child health journal. 2010;14(5):687-95.

- 382 7. Cleland J, Conde-Agudelo AP. H., Ross, J., & Tsui, A. Contraception and health,
383 Lancet.380(9837):149-56.
- 384 8. Bearak J, Popinchalk A, Alkema L, Sedgh G. Global, regional, and subregional trends in
385 unintended pregnancy and its outcomes from 1990 to 2014: estimates from a Bayesian
386 hierarchical model. The Lancet Global Health. 2018;6(4):e380-e9.
- 387 9. Sedgh G, Bearak J, Singh S, Bankole A, Popinchalk A, Ganatra B, et al. Abortion
388 incidence between 1990 and 2014: global, regional, and subregional levels and trends. The
389 Lancet. 2016;388(10041):258-67.
- 390 10. Singh S, Darroch JE, Ashford LS. Adding it up: the costs and benefits of investing in
391 sexual and reproductive health 2014. 2014.
- 392 11. United Nations Educational SaCOU. Developing an education sector response to early
393 and unintended pregnancy. Paris (France): UNESCO; 2014. 2014.
- 394 12. Darroch JE, Woog V, Bankole A, Ashford LS. Adding it up: costs and benefits of
395 meeting the contraceptive needs of adolescents. 2016.
- 396 13. Catalog WBD. Population Estimates And Projections. 2020.
- 397 14. Affairs UNDoEaS. . World Fertility and Family Planning 2020.
- 398 15. Asefa G, Birhanu Z. FERTILITY AND FAMILY PLANNING IMPLICATIONS OF
399 ETHIOPIA'S FP2020 TARGET. 2014.
- 400 16. Adane AA, Bekele YA, Melese E, Worku GT, Netsere HB. Modern contraceptive
401 utilization and associated factors among married Gumuz women in Metekel Zone North West
402 Ethiopia. BioMed Research International. 2020;2020.

- 403 17. Meskele M, Mekonnen W. Factors affecting women's intention to use long acting and
404 permanent contraceptive methods in Wolaita Zone, Southern Ethiopia: A cross-sectional study.
405 BMC women's health. 2014;14(1):1-9.
- 406 18. Degefa Hidru H, Dingeta T, Menigiste B, Etsay B, Gebremedhin H, Berwo M, et al.
407 Modern Contraceptive Utilization and Its Associated Factors among Indigenous and
408 Nonindigenous Married Women of Reproductive Age Group in Jiggiga Town, Eastern Ethiopia,
409 2018. BioMed Research International. 2020;2020.
- 410 19. Gebre MN, Edossa ZK. Modern contraceptive utilization and associated factors among
411 reproductive-age women in Ethiopia: evidence from 2016 Ethiopia demographic and health
412 survey. BMC women's health. 2020;20(1):1-14.
- 413 20. Mekonnen BD, Wubneh CA. Prevalence and associated factors of contraceptive
414 discontinuation among reproductive-age women in Ethiopia: using 2016 Nationwide Survey
415 Data. Reproductive Health. 2020;17(1):1-10.
- 416 21. Bitew F, Nyarko SH. Modern contraceptive use and intention to use: implication for
417 under-five mortality in Ethiopia. Heliyon. 2019;5(8):e02295.
- 418 22. Belda SS, Haile MT, Melku AT, Tololu AK. Modern contraceptive utilization and
419 associated factors among married pastoralist women in Bale eco-region, Bale Zone, South East
420 Ethiopia. BMC health services research. 2017;17(1):1-12.
- 421 23. Debebe S, Limenih MA, Biadgo B. Modern contraceptive methods utilization and
422 associated factors among reproductive aged women in rural Dembia District, northwest Ethiopia:
423 Community based cross-sectional study. International Journal of Reproductive BioMedicine.
424 2017;15(6):367.

- 425 24. Geremew AB, Gelagay AA. Modern contraceptive use and associated factors among
426 married women in Finote Selam town Northwest Ethiopia: a community based cross-sectional
427 study. *Women's midlife health*. 2018;4(1):1-8.
- 428 25. Mohammed A, Woldeyohannes D, Feleke A, Megabiaw B. Determinants of modern
429 contraceptive utilization among married women of reproductive age group in North Shoa Zone,
430 Amhara Region, Ethiopia. *Reproductive health*. 2014;11(1):1-7.
- 431 26. Abraha TH, Teferra AS, Gelagay AA. Postpartum modern contraceptive use in northern
432 Ethiopia: prevalence and associated factors. *Epidemiology and health*. 2017;39.
- 433 27. Tekelab T, Melka AS, Wirtu D. Predictors of modern contraceptive methods use among
434 married women of reproductive age groups in Western Ethiopia: a community based cross-
435 sectional study. *BMC women's health*. 2015;15(1):1-8.
- 436 28. Csa I. Central statistical agency (CSA)[Ethiopia] and ICF. Ethiopia demographic and
437 health survey, Addis Ababa, Ethiopia and Calverton, Maryland, USA. 2016.
- 438 29. Utomo B, Suahya PK, Romadlona NA, Robertson AS, Aryanty RI, Magnani RJ. The
439 impact of family planning on maternal mortality in Indonesia: what future contribution can be
440 expected? *Population Health Metrics*. 2021;19(1):1-13.
- 441 30. Eliason S, Baiden F, Quansah-Asare G, Graham-Hayfron Y, Bonsu D, Phillips J, et al.
442 Factors influencing the intention of women in rural Ghana to adopt postpartum family planning.
443 *Reproductive health*. 2013;10(1):1-8.
- 444 31. Der AD, Anaman-Torgbor JA, Charles-Unadike VO, Tarkang EE. Predictors of intention
445 to use modern contraceptives among female senior secondary school students in the Kpando
446 Municipality, Ghana. *African Health Sciences*. 2021;21(3):1375-84.

- 447 32. Lutalo T, Gray R, Santelli J, Guwatudde D, Brahmbhatt H, Mathur S, et al. Unfulfilled
448 need for contraception among women with unmet need but with the intention to use
449 contraception in Rakai, Uganda: a longitudinal study. *BMC women's health*. 2018;18(1):1-7.
- 450 33. Forty J, Rakgoasi SD, Keetile M. Patterns and determinants of modern contraceptive use
451 and intention to use contraceptives among Malawian women of reproductive ages (15–49 years).
452 *Contraception and reproductive medicine*. 2021;6(1):1-12.
- 453 34. Abraha TH, Belay HS, Welay GM. Intentions on contraception use and its associated
454 factors among postpartum women in Aksum town, Tigray region, northern Ethiopia: a
455 community-based cross-sectional study. *Reproductive health*. 2018;15(1):1-8.
- 456 35. Bawah AA, Asuming P, Achana SF, Kanmiki EW, Awoonor-Williams JK, Phillips JF.
457 Contraceptive use intentions and unmet need for family planning among reproductive-aged
458 women in the Upper East Region of Ghana. *Reproductive health*. 2019;16(1):1-9.
- 459 36. Tiruneh FN, Chuang K-Y, Ntenda PA, Chuang Y-C. Factors associated with
460 contraceptive use and intention to use contraceptives among married women in Ethiopia. *Women
461 & health*. 2016;56(1):1-22.
- 462 37. Rai RK. Future intention of contraceptive use among Comorian women, 2012. *Journal of
463 Public Health*. 2015;23(5):289-96.
- 464 38. Lasong J, Zhang Y, Gebremedhin SA, Opoku S, Abaidoo CS, Mkandawire T, et al.
465 Determinants of modern contraceptive use among married women of reproductive age: a cross-
466 sectional study in rural Zambia. *BMJ open*. 2020;10(3):e030980.
- 467 39. Oumer M, Manaye A, Mengistu Z. Modern contraceptive Method utilization and
468 associated factors among women of reproductive age in Gondar City, Northwest Ethiopia. *Open
469 Access Journal of Contraception*. 2020;11:53.

- 470 40. Endriyas M, Eshete A, Mekonnen E, Misganaw T, Shiferaw M, Ayele S. Contraceptive
471 utilization and associated factors among women of reproductive age group in Southern Nations
472 Nationalities and Peoples' Region, Ethiopia: cross-sectional survey, mixed-methods.
473 Contraception and reproductive medicine. 2017;2(1):1-9.
- 474 41. Wondim G, Degu G, Teka Y, Diress G. Male Involvement in Family Planning Utilization
475 and Associated Factors in Womberma District, Northern Ethiopia: Community-Based Cross-
476 Sectional Study. Open Access Journal of Contraception. 2020;11:197.
- 477 42. Asif MF, Pervaiz Z, Afridi JR, Abid G, Lassi ZS. Role of husband's attitude towards the
478 usage of contraceptives for unmet need of family planning among married women of
479 reproductive age in Pakistan. BMC Women's Health. 2021;21(1):1-7.
- 480 43. Egbe T, Ketchen S, Egbe E, Ekane G, Belley-Priso E. Risk factors and barriers to male
481 involvement in the choice of family planning methods in the Buea Health District, south west
482 region, Cameroon: a cross-sectional study in a semi-urban area. Women Health Open J.
483 2016;1(3):82-90.
- 484 44. Singh A, Singh K, Verma P. Knowledge, attitude and practice GAP in family planning
485 usage: an analysis of selected cities of Uttar Pradesh. Contraception and reproductive medicine.
486 2016;1(1):1-10.
- 487 45. Ijadunola MY, Abiona TC, Ijadunola KT, Afolabi OT, Esimai OA, OlaOlorun FM. Male
488 involvement in family planning decision making in Ile-Ife, Osun State, Nigeria. African journal
489 of reproductive health. 2010;14(4).
- 490 46. Adera A, Belete T, Gebru A, Hagos A, Gebregziabher W. Assessment of the role of men
491 in family planning utilization at Edaga-Hamuse town, Tigray, North Ethiopia. Am J Nurs Sci.
492 2015;4(4):174.

- 493 47. Demissie DB, Kurke A, Awel A, Oljira K. Male involvement in family planning and
494 associated factors among Marriedin Malegedo town west Shoa zone, Oromia, Ethiopia. *planning*.
495 2016;15.
- 496 48. Weisfeld GE, Nowak NT, Lucas T, Weisfeld CC, Imamoğlu EO, Butovskaya M, et al.
497 Do women seek humorousness in men because it signals intelligence? A cross-cultural test.
498 2011.
- 499 49. Hall JA. Humor in romantic relationships: A meta-analysis. *Personal Relationships*.
500 2017;24(2):306-22.
- 501 50. Moreira LR, Ewerling F, Barros AJ, Silveira MF. Reasons for nonuse of contraceptive
502 methods by women with demand for contraception not satisfied: an assessment of low and
503 middle-income countries using demographic and health surveys. *Reproductive health*.
504 2019;16(1):1-15.
- 505 51. Abate MG, Tareke AA. Individual and community level associates of contraceptive use
506 in Ethiopia: a multilevel mixed effects analysis. *Archives of Public Health*. 2019;77(1):1-12.
- 507 52. Zimmerman LA, Bell SO, Li Q, Morzenti A, Anglewicz P, Group PPI, et al. Individual,
508 community and service environment factors associated with modern contraceptive use in five
509 Sub-Saharan African countries: A multilevel, multinomial analysis using geographically linked
510 data from PMA2020. *PloS one*. 2019;14(6):e0218157.
- 511 53. Habyarimana F, Ramroop S. The analysis of socio-economic and demographic factors
512 associated with contraceptive use among married women of reproductive age in Rwanda. *The*
513 *Open Public Health Journal*. 2018;11(1).
- 514 54. Ejembi CL, Dahiru T, Aliyu AA. Contextual factors influencing modern contraceptive
515 use in Nigeria. *DHS Working Papers*. 2015(120).

- 516 55. Masood MS, Alsonini NA. Knowledge and attitude about reproductive health and family
517 planning among young adults in Yemen. *International Journal of Population Research*.
518 2017;2017.
- 519 56. Elkalmi RM, Khan MU, Ahmad A, Srikanth AB, Abdurhaman NS, Jamshed SQ, et al.
520 Knowledge, awareness, and perception of contraception among senior pharmacy students in
521 Malaysia: A pilot study. *Journal of research in pharmacy practice*. 2015;4(2):94.
- 522 57. Alemayehu GA, Fekadu A, Yitayal M, Kebede Y, Abebe SM, Ayele TA, et al.
523 Prevalence and determinants of contraceptive utilization among married women at Dabat Health
524 and Demographic Surveillance System site, northwest Ethiopia. *BMC women's health*.
525 2018;18(1):1-7.
- 526 58. Agha S. Intentions to use contraceptives in Pakistan: implications for behavior change
527 campaigns. *BMC public health*. 2010;10(1):1-13.
- 528 59. UNFPA. 5 Upsetting Reasons Women Aren't Using Family Planning Around the World
529 Today. 2022.
- 530 60. Stephenson R, Hennink M. Barriers to family planning service use among the urban poor
531 in Pakistan. *Asia-Pacific Population Journal*. 2004;19(2):5-26.
- 532 61. Farmer DB, Berman L, Ryan G, Habumugisha L, Basinga P, Nutt C, et al. Motivations
533 and constraints to family planning: a qualitative study in Rwanda's Southern Kayonza District.
534 *Global Health: Science and Practice*. 2015;3(2):242-54.
- 535 62. Wang RH, Cheng CP, Chou FH. A causal model of contraceptive intention and its gender
536 comparison among Taiwanese sexually inexperienced adolescents. *Journal of clinical Nursing*.
537 2008;17(7):930-9.

538 63. Bang K-S, Chae S-M, Lee I, Yu J, Kim J. Effects of a community outreach program for
539 maternal health and family planning in Tigray, Ethiopia. Asian nursing research.
540 2018;12(3):223-30.

541