

1 Comparing Sexual Network Mean Active Degree Measurement 2 Metrics among Men who have Sex with Men

3 4 **Authors**

5 Christina Chandra MPH¹, Martina Morris PhD², Connor Van Meter MPH¹, Steven M. Goodreau
6 PhD³, Travis Sanchez DVM¹, Patrick Janulis PhD⁴, Michelle Birkett PhD⁴, Samuel M. Jenness
7 PhD¹

8 ¹ Department of Epidemiology, Rollin School of Public Health, Emory University, Atlanta, GA

9 ² Department of Sociology, University of Washington, Seattle, WA

10 ³ Department of Anthropology, University of Washington, Seattle, WA

11 ⁴ Department of Medical Social Sciences, Feinberg School of Medicine, Northwestern University,
12 Chicago, IL

13

14 **Correspondence**

15 *Christina Chandra, Emory University, 1520 Clifton Road, Atlanta, GA 30323. Email:*

16 christina.lynne.chandra@emory.edu.

17

18 **Funding**

19 This work was supported by National Institutes of Health grants R21 MH112449 and R01
20 AI138783, and a grant from the MAC AIDS Fund.

21

22 **Conflicts of Interest**

23 The authors declare no conflicts of interest.

24

25 **Word Counts**

26 Abstract: 249 words

27 Text: 3393 words

28

29 **Keywords**

30 Sexual networks, HIV, sexually transmitted infection, men who have sex with men, partnerships

31

32 **Summary**

33 Survey designs can lead to potential bias, such as underestimation, in the measurement of
34 mean active degree in sexual networks of men who have sex with men.

35 **ABSTRACT**

36 **Background** Mean active degree is an important proxy measure of cross-sectional network
37 connectivity commonly used in HIV/STI epidemiology research. No current studies have
38 compared measurement methods of mean degree using cross-sectional surveys for men
39 who have sex with men (MSM) in the United States.

40 **Methods** We compared mean degree estimates based on reported ongoing main and casual
41 sexual partnerships (*current method*) against dates of first and last sex (*retrospective*
42 *method*) from 0–12 months prior to survey date in ARTnet, a cross-sectional survey of
43 MSM in the U.S. (2017–2019). ARTnet collected data on the number of sexual partners
44 in the past year but limited reporting on details used for calculating mean degree to the 5
45 most recent partners. We used linear regression to understand the impact of truncated
46 partnership data on mean degree estimation.

47 **Results** Retrospective method mean degree systematically decreased as the month at which it
48 was calculated increased from 0–12 months prior to survey date. Among participants with
49 >5 partners in the past year compared to those with ≤5, the average change in main
50 degree between 12 and 0 months prior to survey date was -0.05 (95% CI: -0.08, -0.03)
51 after adjusting for race/ethnicity, age, and education. The adjusted average change in
52 casual degree was -0.40 (95% CI: -0.45, -0.35).

53 **Conclusions** The retrospective method underestimates mean degree for MSM in surveys with
54 truncated partnership data, especially for casual partnerships. The current method is less
55 prone to bias from partner truncation when the target population experiences higher
56 cumulative partners per year.

57 INTRODUCTION

58 In the United States, men who have sex with men (MSM) experience a disproportionate burden of
59 HIV and bacterial sexually transmitted infections (STIs). In 2018, MSM accounted for 66% of all new
60 HIV diagnoses, 54% of primary and secondary syphilis cases, and 43% of gonorrhea cases.^{1,2}
61 Young, non-white MSM are particularly affected by HIV/STIs despite evidence that individual risk
62 behaviors are not different between Black and white MSM.³ Differences in sexual network
63 connectivity, the mechanistic pathway for HIV/STI transmission, may explain these disparities
64 among MSM in the U.S.^{4,5} Networks are also essential for deploying prevention tools, such as HIV
65 PrEP or STI partner services.^{6,7}

66 The potential effectiveness of a network-informed HIV/STI public health response depends
67 on good empirical metrics: accurate and unbiased estimates of observable behaviors that determine
68 the unobservable network connectivity. One common metric is active degree: the count of current,
69 ongoing partners at a point in time, which may be summarized as the mean active degree across
70 nodes (persons) at the population level. It is mathematically established and intuitive that network
71 connectivity rises as mean degree increases, though the relation is non-linear.⁷ Active degree also
72 forms the definitional basis of partnership concurrency (active degree of two or more). Modeling has
73 demonstrated that higher mean active degree and higher prevalence of concurrency create network
74 conditions that lead to more rapid and pervasive HIV/STI spread within networks.^{5,8} Even small
75 changes in mean active degree can have a substantial impact on network connectivity, and
76 subsequently epidemic persistence or elimination, due to the non-linear threshold effects.⁹ For this
77 reason, accurate measurement of mean active degree is needed to assess epidemic potential.

78 UNAIDS focused on this problem in a reference group meeting in 2009, leading to published
79 recommendations for measuring active degree and calculating point prevalence of concurrency as
80 an indicator for monitoring national HIV epidemics.¹⁰ Approaches they considered were: 1) active
81 degree as measured on the day of survey (“current method”); 2) using reported dates of first and last
82 sex with each partner during the last year (“retrospective method”). UNAIDS recommended the
83 latter. **Figure 1** provides a schematic of how the current and retrospective methods are used to
84 calculate mean degree with hypothetical data.

85 The goal of the current method is to measure the person’s ongoing sexual partnerships on the
86 day of the survey. This can be done on a partner-by-partner basis after enumerating relevant
87 partners (e.g., “Do you expect to have sex with this person again?”), or with a single summary

88 question (e.g., “With how many people do you currently have an active sexual relationship?”). Active
89 degree is measured as the sum of ongoing partnerships reported. This approach does not depend
90 on recall of events or dates and can be asked in a single question if time and respondent burden are
91 concerns. The drawback is that respondents must accurately predict whether a partnership will
92 continue. While that may be reasonable for populations characterized by having few long-term stable
93 partnerships, it may fail in populations with frequent short-term casual partnerships.¹¹

94 With the retrospective method, partners are enumerated for some period (e.g., all in the last
95 year) and the partnership’s date of first and last sex are recorded. The partnership intervals are then
96 evaluated for active overlap at specified time points prior to the survey date. The UNAIDS group
97 recommended using 6 months prior to survey date to get a point estimate of the active degree. This
98 method does not require the respondent to predict whether current partnerships will continue, but it
99 requires accurate recollection of the dates of first/last sex. For surveys that ask for partnership dates
100 for a limited number of partners (e.g., 5 most recent partners), participants with more sex partners
101 may reach this limit before the specified number of months prior to survey date. For example, if a
102 participant’s 5 most recent partnerships occurred within 3 months of the survey date, any additional
103 partnerships at 6 months prior to the survey date would not be counted. The higher the cumulative
104 number of partners over time, the more potential there is for downward bias in active degree as you
105 go back in time.

106 While both methods have been used in surveys of MSM in the U.S.,^{12–15} there have been no
107 comparative studies of these methods for MSM. Compared to heterosexual men and women, active
108 degree is often higher among MSM, and more heterogenous, reflecting different partnership types of
109 varying durations and typologies.¹⁶ Furthermore, the biases from truncated partnership data in cross-
110 sectional studies (also known as fixed choice design and right-censoring of degree) has been well-
111 documented in the field of network science.^{17–19} It would be useful to understand how these
112 measurement approaches perform for MSM with this survey design.

113 In this study, we compared estimates using the current and the retrospective methods among
114 U.S. MSM in a survey that allows for estimation with both methods and collects truncated
115 partnership data. We aimed to understand if partnership data truncation was the main source of bias
116 for the differences between the current and retrospective method, even after controlling for
117 demographic variables that may be correlated with both truncation of partnership data and other
118 mechanisms for bias in mean degree estimates.

119

120 **METHODS**

121 *Study Design.* We used data from ARTnet, a cross-sectional web-based study of MSM in the U.S.
122 conducted between 2017 and 2019.²⁰ ARTnet recruited participants through the American Men’s
123 Internet Survey (AMIS), an ongoing study about MSM sexual health.²¹ ARTnet eligibility criteria
124 included age between 15 and 65 years, cisgender male identity, male sex at birth, and having ever
125 had sex with a male partner. ARTnet recruited AMIS participants and collected data in two waves:
126 July 2017–February 2018 and September 2018–January 2019. Participants were deduplicated
127 within and across waves. For participants with more than one survey record, we retained the most
128 recent survey record from either wave. The Emory University Institutional Review Board approved
129 this study.

130 *Measures.* ARTnet collected information on egocentric sexual network data, recent sexual behavior,
131 and utilization of HIV and STI prevention services. Participants were asked to provide information on
132 up to 5 of their most recent male sexual partners within the last 12 months. Participants reported on
133 type of partnership (main, casual, or one-time), dates of first and last sex by month and year, and
134 sexual activity (e.g., condomless anal sex). Participants also answered the question, “Is this
135 relationship with this partner active and ongoing?” If participants answered “Yes” but reported sex
136 only once with the partner, then the partnership was considered a one-time partnership and not
137 factored into mean degree calculations. We ran a sensitivity analysis to understand how an
138 alternative definition where these partnerships were considered ongoing, casual partnerships and
139 how imputing “Don’t know” answers to “Yes” impacted the mean degree of casual partnerships.

140 Our main exposure was the truncation of partnership data from limiting partnership information
141 to the 5 most recent partners. ARTnet also asked participants for the total number of male sexual
142 partners within the last 12 months, although details such as dates of first/last sex were not collected
143 on all partners. The main exposure of partnership truncation was operationalized by a binary
144 variable that split participants by whether they had 5 or fewer partners, including main, casual, or
145 one-time, or more than 5 partners in the past 12 months. Those with more than 5 male partners in
146 the past year are considered to have provided truncated partnership data. Our outcome of interest
147 was a summary measure of active degree (henceforth, degree): the mean degree across all
148 participants at a specified time point. This measure did not include one-time partnerships, as they

149 are not ongoing. We calculated mean degree using the current method and compared it to mean
150 degree calculated using the retrospective method at specified months prior to the survey date,
151 ranging from 0 (current study month) to 12 months. We refer to the months prior to the survey date
152 as monthly “offsets” (e.g., 6 months prior to the survey date is a 6-month offset). Since participants
153 reported only month and year for dates of first/last sex in ARTnet, we randomly imputed days within
154 these months. We conducted a sensitivity analysis to understand how date imputation impacted
155 mean degree estimates from the retrospective method.

156 *Statistical Analysis.* For descriptive analyses, we plotted mean degree by both current and
157 retrospective methods by the primary exposure variable of having 5 or fewer or more than 5
158 partners. Our main outcome was the difference in degree between 12- and 0-month offsets. This
159 difference provides a measure of stability of the retrospective method across offset months.
160 Bivariable linear regression was used to estimate that 12-month difference and variations in that
161 difference by the primary exposure variable and selected demographic variables (e.g., race/ethnicity,
162 age, geography, education, income) that may also be correlated with bias in mean degree estimates.
163 In multiple linear regression, we investigated whether variations in the 12- and 0-month degree
164 difference were caused by the truncation of partnership data, after controlling for race, age, and
165 education, which we hypothesized as associated with the primary exposure and the outcome. All
166 analyses were conducted in R 4.2.0.²² Analysis scripts are provided in a GitHub repository
167 (<https://github.com/EpiModel/Mean-Degree-Analysis>).

168

169 **RESULTS**

170 Of the 4,904 MSM who completed the ARTnet study, most were non-Hispanic white, less than 35
171 years old, completed college or above, and had an annual household income of at least \$40,000
172 (**Table 1**). A total of 16,198 partnerships were reported by the participants. Of these, 7,602 (46.9%)
173 were one-time partnerships, 5,978 (36.9%) were casual partnerships, and 2,618 (16.2%) were main
174 partnerships. On the day of survey, 5,875 (68.3%) of main and casual partnerships were reported as
175 ongoing. Overall, 1,962 (40.0%) of participants had more than 5 partners (main, casual, and one-
176 time) in the past 12 months.

177 Mean degree estimates varied by partnership type for both measurement approaches. Using
178 the current method, total mean degree was 1.19 across partnership types: 0.45 for main and 0.74 for

179 casual partnerships (**Figure 2**). Mean degree estimates ranged from 1.01 at the 12-month offset to
180 1.23 at the 1-month offset across all partnerships. Mean degree ranged from 0.38 (12-month offset)
181 to 0.45 (0-month offset) for main partnerships and from 0.64 (12-month offset) to 0.78 (1-month
182 offset) for casual partnerships.

183 The mean degrees estimated by the retrospective method decreased as the month offsets
184 increased from 1 to 12 months (**Figure 2**). Mean degree estimated at the 0-month offset was the
185 same as the mean degree estimated from the current method across all partnership types, which is
186 expected given that the 0-month offset date is the same as the survey date for each participant.
187 Mean degree of main partnerships from the current method (0.45) was most similar to estimates
188 between 1- and 2-month offsets (0.45–0.44), while for casual partnerships, the current estimate
189 (0.75) was most similar to mean degree estimated at 5- and 6-month offsets (0.76, 0.73). Overall
190 mean degree estimated with the current method (1.19) was most similar to mean degree at the 4-
191 month offset (1.19) across all partnerships.

192 When stratifying by having 5 or fewer or more than 5 partners in the past 12 months, the
193 decreasing trend of mean degree was largely explained by participants who had more than 5
194 partners (**Figure 3**). Participants with more than 5 partners in the past year did not have the
195 opportunity to share information on more than 5 of their partners, and therefore, the partnerships
196 reported may be biased to the most recent partnerships. For main partnerships, there continued to
197 be a decreasing trend of mean degree observed between the 1- and 12-month offsets (0.48 to 0.42)
198 for participants with 5 or fewer partnerships in the past year; however, this did not impact the stability
199 of the overall mean degree when restricted to participants with 5 or fewer partnerships. Mean degree
200 remained stable from 0.83 at 1-month offset to 0.80 at 12-month offset for participants with 5 or
201 fewer partners in the past year, but mean degree decreased from 1.83 at 1-month offset to 1.34 at
202 12-month offset for those with more than 5 partners. A similar pattern was found for casual mean
203 degree. Among participants with 5 or fewer partners, casual mean degree was relatively stable from
204 0.35 at 1-month offset to 0.37 at 12-month offset. For participants with more than 5 partners, casual
205 mean degree decreased from 1.44 at 1-month offset to 1.04 at 12-month offset. **Supplemental**
206 **Table 1** describes mean degree estimates using both methods by participant and partnership
207 characteristics for 0-, 3-, 6-, and 12-month offsets.

208 Bivariable and multivariable linear regression further confirmed that the primary exposure of
209 partnership truncation was the principal cause of the decreasing trend of retrospective mean degree

210 estimates, particularly for casual partnerships. Race/ethnicity, age, and education were found to be
211 potential confounders of the relationship between truncation of partnership data and the downward
212 bias of mean degree in the retrospective method (**Table 2; Supplemental Table 2; Supplemental**
213 **Figures 1–5**). The negative association between average change in degree and number of male
214 partners in the past year remained after adjusting for demographics. The average change in main
215 degree between 12- and 0-month offsets among participants with more than 5 partners in the past
216 year compared to those with 5 or fewer partners was -0.04 (95% CI: -0.07, -0.02) in unadjusted
217 analysis and -0.05 (95% CI: -0.08, -0.03) in adjusted analysis (**Table 2**). For casual partnerships
218 among participants with more than 5 partners compared to 5 or fewer partners, the unadjusted
219 average change in degree was -0.41 (95% CI: -0.46, -0.36) and adjusted average change in degree
220 was -0.40 (95% CI: -0.45, -0.35).

221 In a sensitivity analysis on the influence of date imputation on retrospective mean degree,
222 random date imputation had little effect on mean degree (**Supplemental Table 5**). When
223 considering the minimum mean degree (sexual partnerships were not active during the entire month
224 reported) and maximum mean degree (sexual partnerships were active during the entire month
225 reported), there was an average percent change of 2–3%.

226 Twenty percent of one-time partnerships were considered active and ongoing on the day of the
227 survey, but these partnerships were considered one-time partnerships and not included in mean
228 degree estimation because participants had sex with these partners once. In a sensitivity analysis
229 we redefined these partnerships as active, casual partnerships and considered a scenario where a
230 proportion of “Don’t know” responses were “Yes.” These alternative definitions suggest our original
231 definition may be underestimating the casual mean degree using the current method and the
232 retrospective method at the 0-month offset (**Supplemental Table 6**). There was no effect on the
233 retrospective mean degree beyond the 0-month offset.

234

235 **DISCUSSION**

236 In this study, we compared the current and the retrospective methods of measuring mean degree in
237 a sample of U.S. MSM. The two methods produced similar estimates of mean degree for participants
238 with fewer past-year partners. With the retrospective method, mean degree estimates decreased as
239 the retrospective measure of degree was calculated at more months prior to the survey date. This

240 was due to truncated partnership data from the survey design: information used to calculate mean
241 degree such as dates of first/last sex were collected on up to 5 of the most recent partners. Our
242 results suggest that this downward bias leads to underestimation of degree, especially of casual
243 partnerships, using the retrospective method when surveys have truncated partnership data.
244 Underestimation of degree may impact projected transmission of HIV in simulated networks.^{23,24} The
245 current method of measurement is less prone to bias from truncated partnership data but may have
246 its own flaws.

247 With 2 in 5 ARTnet participants reporting more than 5 main, casual, and one-time male
248 sexual partners in the past year, the truncation of partnership data had a major impact on mean
249 degree estimation with the retrospective method. This was particularly true for casual partnerships,
250 which were more prevalent compared to main partnerships in our sample. An exploration of a few
251 cases with large differences in degree between 12- and 0-month offsets may offer some explanation
252 for the downward bias (**Supplemental Figures 6–7**). For example, in **Supplemental Figure 7**, study
253 participant 4 reported 15 male sexual partners in the past 12 months but was only able to report on
254 the 5 most recent partners, which were all considered ongoing on the day of survey but not active at
255 12 months prior to the survey date. This person's degree would be 0 at the 12-month offset but 5 at
256 the 0-month offset, resulting in a difference of -5. Since participants were asked to report on their
257 most recent partnerships, participants with truncated partnership data were likely to only report on
258 partnerships closer to the survey date before exhausting their responses, leading to underestimation
259 of degree as month offsets increased. Furthermore, reporting on more one-time partnerships, which
260 don't contribute to mean degree, may censor main and casual partnerships in the past year and
261 underestimate mean degree.

262 A direct comparison of the current method to the UNAIDS-recommended retrospective
263 method at 6 months using ARTnet data shows that the percent of participants reporting more than 1
264 partner at both time points was similar and total, main, and casual mean degree was slightly smaller
265 when estimated at the 6-month offset (**Supplemental Table 7**). The distribution of degree for both
266 time points were also similar (**Supplemental Figure 8**). The retrospective method may still be a
267 suitable alternative to the current method if there was no truncation of partnership data in the
268 behavioral survey or if information on the active status of current partnerships was not collected and
269 if mean degree is estimated within 6 months of the survey date.

270 The current or retrospective measurement methods have been applied to the estimation of
271 partnership concurrency (degree of two or more), aligning generally with our current study. One
272 study of young adults in the U.S. directly comparing reported partnership concurrency within the last
273 6 years through direct questioning and overlapping partnership dates found both methods reported a
274 similar prevalence of concurrency.²⁵ However, the agreement about concurrent partnerships
275 between the two methods was only modest, with the authors suggesting that measuring concurrency
276 using overlapping partnership dates was more likely to be complicated by missing or uninterpretable
277 data. Another study of heterosexual partnerships in Malawi found that measuring concurrency using
278 partnership dates underestimated self-reported concurrency due to difficulty of recalling partnership
279 start and end dates and general underreporting of partnerships.²⁶ The current method may be
280 statistically preferable because both mean active degree and partnership age can be jointly
281 estimated with typical distributional assumptions.^{20,27} On the other hand, study participants may not
282 be able to accurately predict whether current partnerships will continue, leading to overestimation of
283 mean active degree using the current method.¹¹ The current method is not perfect, and bias
284 adjustment or improvement of the measurement method is needed. Our analysis was unable to
285 identify current partners that were truly active and ongoing because we did not collect follow-up data.
286 Future studies may consider the use of longitudinal data where behavioral surveys administered at
287 multiple time points collect data on a limited number of existing and new partners. This study design
288 could evaluate the severity of the bias from truncation partnership data compared to bias from
289 participants' failure to predict ongoing partnerships.

290 *Limitations.* There are some limitations to this study. First, only month and year of dates of first/last
291 sex were reported for each partnership. Partnerships with start or end months that were the same as
292 the month of the retrospective date from the date of survey could be considered ongoing or not
293 ongoing depending on the exact day of first/last sex. To address this, we randomly imputed the day
294 for the start and end dates to identify partnerships overlapping with the retrospective date. A
295 sensitivity analysis found that the average percent change of mean degree for more extreme
296 assumptions was minimal. Second, this study was limited to cross-sectional data, which may be
297 useful because many studies estimate mean degree from cross-sectional data, but longitudinal
298 studies may be able to describe unique biases from cross-sectional data such as the prediction of
299 ongoing partnerships.¹¹ Third, ARTnet coding decisions around one-time partnerships and “Do not
300 know” responses for the question of whether a partnership is active and ongoing may lead to

301 underestimation of mean degree using the current method or retrospective 0-month offset, which
302 was confirmed through a sensitivity analysis. Given prior research that suggests MSM may be
303 overconfident in predicting which sexual partnerships are ongoing,¹¹ we used the original method as
304 a conservative measure.

305 *Conclusion.* Sexual mean active degree is an important measure in HIV and STI epidemiology. It
306 should therefore be measured rigorously and estimated consistently across studies. Future network-
307 based studies should justify their methods for mean active degree measurement with special
308 consideration of how data is collected. Our analysis suggests that cross-sectional data with
309 truncated partnership data may underestimate mean degree using the retrospective method,
310 favoring the current method. However, improvements can further be made to the current method to
311 address bias in predicting persistence of partnerships.¹¹ The accuracy of the active degree estimates
312 will have important impacts on the delivery of HIV/STI prevention interventions that target networks.

313 REFERENCES

- 314 1. Centers for Disease Control and Prevention. HIV Surveillance Report, 2018 [Internet]. 2018 Nov. Report No.:
315 29. Available from: <http://www.cdc.gov/hiv/library/reports/hiv-surveillance.html>.
- 316 2. Centers for Disease Control and Prevention. Sexually Transmitted Disease Surveillance 2018 [Internet]. 2019.
317 Available from: <https://www.cdc.gov/std/stats>
- 318 3. Millett GA, Flores SA, Peterson JL, et al. Explaining disparities in HIV infection among black and white men
319 who have sex with men: a meta-analysis of HIV risk behaviors. *AIDS*. 2007 Oct 1;21(15):2083–91.
- 320 4. Mustanski B, Morgan E, D'Aquila R, et al. Individual and Network Factors Associated With Racial Disparities in
321 HIV Among Young Men Who Have Sex With Men. *J Acquir Immune Defic Syndr*. 2019 Jan;80(1):24–30.
- 322 5. Morris M, Kurth AE, Hamilton DT, et al. Concurrent partnerships and HIV prevalence disparities by race: linking
323 science and public health practice. *Am J Public Health*. 2009;99(6):1023–31.
- 324 6. Teixeira da Silva D, Bouris A, Ramachandran A, et al. Embedding a Linkage to Pre-Exposure Prophylaxis
325 (PrEP) Care Intervention in Social Network Strategy and Partner Notification Services: Results from a Pilot
326 Randomized Controlled Trial. *J Acquir Immune Defic Syndr*. 2021;86(2):191–9.
- 327 7. Erdos P, Rényi A. On the evolution of random graphs. *Publ Math Inst Hung Acad Sci*. 1960;5(1):17–60.
- 328 8. Eaton JW, Hallett TB, Garnett GP. Concurrent sexual partnerships and primary HIV infection: a critical
329 interaction. *AIDS Behav*. 2011;15(4):687–92.
- 330 9. Carnegie NB, Morris M. Size matters: concurrency and the epidemic potential of HIV in small networks. *PLoS*
331 *ONE*. 2012;7(8):e43048.
- 332 10. Eaton J, Case, Kelsey, secretariat of the UNAIDS Reference Group on Estimates, Modelling, and Projections.
333 Consultation on Concurrent Sexual Partnerships [Internet]. UNAIDS; 2009 Jun p. 19. Available from:
334 [http://www.epidem.org/sites/default/files/content/resources/attachments/Concurrency%20meeting%20recomm](http://www.epidem.org/sites/default/files/content/resources/attachments/Concurrency%20meeting%20recommendations_Final.pdf)
335 [endations_Final.pdf](http://www.epidem.org/sites/default/files/content/resources/attachments/Concurrency%20meeting%20recommendations_Final.pdf)
- 336 11. Uong S, Rosenberg ES, Goodreau SM, et al. Assessment of Bias in Estimates of Sexual Network Degree
337 using Prospective Cohort Data. *Epidemiology*. 2020 Mar;31(2):229–37.
- 338 12. Jenness SM, Johnson JA, Hoover KW, et al. Modeling an integrated HIV prevention and care continuum to
339 achieve the ending the HIV epidemic goals. *AIDS*. 2020;34(14):2103–13.
- 340 13. Maloney KM, Driggers RA, Sarkar S, et al. Projected Impact of Concurrently Available Long-Acting Injectable
341 and Daily-Oral HIV Pre-Exposure Prophylaxis. *J Infect Dis*. 2021;223(1):72–82.
- 342 14. Jenness SM, Le Guillou A, Chandra C, et al. Projected HIV and Bacterial Sexually Transmitted Infection
343 Incidence Following COVID-19-Related Sexual Distancing and Clinical Service Interruption. *J Infect Dis*. 2021
344 Mar 29;223(6):1019–28.
- 345 15. Janulis P, Phillips G, Birkett M, et al. Sexual Networks of Racially Diverse Young MSM Differ in Racial
346 Homophily But Not Concurrency. *J Acquir Immune Defic Syndr*. 2018 Apr;77(5):459–66.
- 347 16. Glick SN, Morris M, Foxman B, et al. A Comparison of Sexual Behavior Patterns Among Men Who Have Sex
348 With Men and Heterosexual Men and Women. *J Acquir Immune Defic Syndr*. 2012;60(1):83–90.
- 349 17. Holland PW, Leinhardt S. The structural implications of measurement error in sociometry. *J Math Sociol*. 1973
350 Jul 1;3(1):85–111.
- 351 18. Kossinets G. Effects of missing data in social networks. *Soc Netw*. 2006 Jul 1;28(3):247–68.

- 352 19. Gommans R, Cillessen AHN. Nominating under constraints: A systematic comparison of unlimited and limited
353 peer nomination methodologies in elementary school. *Int J Behav Dev.* 2015 Jan 1;39(1):77–86.
- 354 20. Weiss KM, Goodreau SM, Morris M, et al. Egocentric sexual networks of men who have sex with men in the
355 United States: Results from the ARTnet study. *Epidemics.* 2020 Jan 24;30:100386.
- 356 21. Sanchez TH, Sineath RC, Kahle EM, et al. The Annual American Men’s Internet Survey of Behaviors of Men
357 Who Have Sex With Men in the United States: Protocol and Key Indicators Report 2013. *JMIR Public Health
358 Surveill.* 2015;1(1):e3.
- 359 22. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria: R
360 Foundation for Statistical Computing; 2019. Available from: <https://www.R-project.org/>
- 361 23. Harling G, Onnela J-P. Impact of degree truncation on the spread of a contagious process on networks. *Netw
362 Sci.* 2018 Mar;6(1):34–53.
- 363 24. Anderson EJ, Weiss KM, Morris MM, et al. HIV and Sexually Transmitted Infection Epidemic Potential of
364 Networks of Men Who Have Sex With Men in Two Cities. *Epidemiology.* 2021 Sep 1;32(5):681–9.
- 365 25. Nelson SJ, Manhart LE, Gorbach PM, et al. Measuring sex partner concurrency: it’s what’s missing that counts.
366 *Sex Transm Dis.* 2007;34(10):801–7.
- 367 26. Glynn JR, Dube A, Kayuni N, et al. Measuring concurrency: an empirical study of different methods in a large
368 population-based survey and evaluation of the UNAIDS guidelines. *AIDS.* 2012 May 15;26(8):977–85.
- 369 27. Krivitsky P, Morris M. Inference for Social Network Models from Egocentrically-Sampled Data, with Application
370 to Understanding Persistent Racial Disparities in HIV Prevalence in the US. *Ann Appl Stat.* 2017;11(1):427–55.
- 371

TABLES

Table 1. Individual- and Sexual Partnership-level Characteristics of Participants in ARTnet Study (2017–2019) of Men who have Sex with Men in the U.S.

Individual	N	%
Total Sample	4904	100.0
Age (Mean and SD)	36.5	14.2
Age Category		
15–24	1324	27.0
25–34	1268	25.9
35–44	694	14.2
45–54	833	17.0
55–65	785	16.0
Race/Ethnicity		
Black (Non-Hispanic)	266	5.4
Hispanic	676	13.8
Other (Non-Hispanic)	439	9.0
White (Non-Hispanic)	3523	71.8
Residence in Census Region		
Northeast	882	18.0
Midwest	994	20.3
South	1782	36.3
West	1246	25.4
Residence in Census Division		
New England	250	5.1
Middle Atlantic	632	12.9
East North Central	698	14.2
West North Central	296	6.0
South Atlantic	1062	21.7
East South Central	222	4.5
West South Central	498	10.2
Mountain	404	8.2
Pacific	842	17.2
Education		
High school or below	616	12.6
Some college	1519	31.1
College and above	2742	56.2
Annual Household Income		
\$0 to \$19,999	605	13.7
\$20,000 to \$39,999	867	19.6
\$40,000 to \$74,999	1258	28.5
\$75,000 or more	1686	38.2
Number of Male Partners in the Past 12 Months		
≤ 5 Partners	2939	59.9
> 5	1962	40.0
Sexual Partnerships^a	N	%
Total Sample	16198	100.0
Sexual Partnership Type		
Main	2618	16.2
Casual	5978	36.9
One-time	7602	46.9
Self-Reported Ongoing on Day of Survey		
Yes	7289	45.0
No	8909	55.0

^a The ARTnet survey allows participants to report details on up to 5 of the most recent sexual male partners within the past 12 months of taking the survey.

Table 2. Bivariable linear regression results of average change in degree between 12 months and 0 months prior to survey date using the retrospective method for main and casual sexual partnerships of male participants (N = 4,904) in the ARTnet study (2017–2019)

	Main Partnerships			Casual Partnerships		
	Estimate	95% CI	F-Test P-value	Estimate	95% CI	F-Test P-value
Number of Male Partners^a			<0.01			<0.01
Intercept (≤ 5 Partners)	-0.06	-0.07, -0.04		0.06	0.03, 0.09	
> 5	-0.04	-0.07, -0.02		-0.41	-0.46, -0.36	
Race/Ethnicity			0.05			0.16
Intercept (Black)	-0.02	-0.07, 0.02		-0.01	-0.11, 0.10	
Hispanic	-0.08	-0.13, -0.02		-0.07	-0.20, 0.04	
Other	-0.04	-0.10, 0.01		-0.09	-0.22, 0.04	
White	-0.05	-0.10, -0.01		-0.11	-0.23, 0.01	
Age (continuous)	0.003	0.002, 0.004	<0.01	-0.003	-0.005, -0.001	<0.01
Age Category			<0.01			<0.01
Intercept (15-24 years)	-0.15	-0.17, -0.13		-0.03	-0.08, 0.01	
25-34 years	0.09	0.06, 0.12		-0.07	-0.14, -0.01	
35-44 years	0.09	0.05, 0.12		-0.10	-0.18, -0.02	
45-54 years	0.11	0.08, 0.15		-0.15	-0.23, -0.07	
55-65 years	0.13	0.09, 0.16		-0.11	-0.19, -0.03	
Census Region			0.28			0.15
Intercept (Northeast)	-0.06	-0.08, -0.03		-0.14	-0.20, -0.08	
Midwest	-0.03	-0.07, 0.00		0.07	-0.01, 0.15	
South	-0.02	-0.05, 0.01		0.05	-0.02, 0.12	
West	-0.02	-0.05, 0.02		0.00	-0.08, 0.07	
Highest Level of Education			<0.01			<0.01
Intercept (High school or below)	-0.12	-0.15, -0.09		-0.06	-0.13, 0.01	
Some college	0.05	0.01, 0.08		0.03	-0.05, 0.11	
College and above	0.06	0.03, 0.09		-0.11	-0.18, -0.03	
Annual Household Income			0.44			<0.01
Intercept (\$0 to \$19,999)	-0.08	-0.11, -0.05		-0.01	-0.08, 0.05	
\$20,000 to \$39,999	0.00	-0.04, 0.04		-0.08	-0.17, 0.01	
\$40,000 to \$74,999	0.00	-0.04, 0.04		-0.07	-0.15, 0.02	
\$75,000 or more	0.02	-0.02, 0.05		-0.16	-0.24, -0.08	

^a The ARTnet survey allows participants to report details such as dates of first and last sex on up to 5 of the most recent sexual male partners within the past 12 months of taking the survey. Participants are also asked about the total number of male sexual partners they have had within the past 12 months but do not report details used to calculate mean degree on all of these partners.

Figure 1. Schematic representation of how mean degree is calculated using hypothetical sexual partnership data with the retrospective method when measuring mean degree at three months prior to survey date and the current method where mean degree is measured on survey date. Mean degree can be calculated for main and casual sexual partnerships, and one-time partnerships are not included in the calculation by definition as they are not ongoing sexual partnerships.

Study ID	Partner ID	Partner Type	Months Prior to Survey Date							Day of Survey	Retrospective Method Measured at 3 Months Prior to Survey Date		Current Method Measured on Day of Survey			
			6	5	4	3	2	1	0		Partnership Ongoing?	Mean Degree	Partnership Ongoing?	Mean Degree		
1	1	Casual									No	1		No	1	
	2	Main									Yes	1		Yes	1	
2	1	Casual									No			No		
	2	One-Time									No			No		
	3	Main									Yes	2	1	No	1	1
	4	Casual									Yes			Yes		
3	1	Casual									No	0		Yes	1	

	Mean Degree		
	Main	Casual	Total
Retrospective Method Measured at 3 Months Prior to Survey Date	2/3	1/3	1
Current Method Measured on Day of Survey	1/3	2/3	1

Figure 2. Comparison of mean degree calculated using the retrospective method from 0 to 12 months prior to survey date and mean degree calculated by the current method on the day of survey among all, main, and casual male sexual partnerships of 4,904 ARTnet participants.




Figure 3. Comparison of mean degree calculated using the retrospective method from 0 to 12 months prior to survey date and mean degree calculated by the current method on the day of survey among all, main, and casual male sexual partnerships of 4,904 ARTnet participants (2017–2019) stratified by having 5 or fewer partners or more than 5 partners. ARTnet participants were only able to report on dates of first and last sex and whether partnerships were ongoing on a maximum of 5 partners, restricting calculation of mean degree to the 5 most recent partners.

