

1 Targeted transcutaneous cervical spinal cord stimulation promotes 2 upper limb recovery in spinal cord and peripheral nerve injury

3 Santosh Chandrasekaran¹, Nikunj A Bhagat^{1,9}, Richard Ramdeo¹, Sadegh Ebrahimi¹, Pawan D Sharma²,
4 Adam Stein⁶, Doug G Griffin⁷, Susan J Harkema^{2,3,4,5} and Chad E Bouton^{1,8}

5 ¹Neural Bypass and Brain Computer Interface Laboratory, Institute of Bioelectronic Medicine, Feinstein Institutes
6 for Medical Research, Northwell Health, Manhasset, New York

7 ²Kentucky Spinal Cord Injury Research Center, University of Louisville, Louisville, Kentucky

8 ³Department of Bioengineering, University of Louisville, Louisville, Kentucky

9 ⁴Frazier Rehabilitation Institute, University of Louisville Health, Louisville, Kentucky

10 ⁵Department of Neurological Surgery, University of Louisville, Louisville, Kentucky

11 ⁶Department of Physical Medicine and Rehabilitation, Donald and Barbara Zucker School of Medicine at Hofstra,
12 Northwell Health, Manhasset, New York

13 ⁷Northwell Health STARS Rehabilitation, East Meadow, New York

14 ⁸Donald and Barbara Zucker School of Medicine at Hofstra/Northwell, Manhasset, New York

15 ⁹Department of Physical Medicine & Rehabilitation, University of Texas Health Science Center, Houston, Texas

16 **Corresponding authors:** Santosh Chandrasekaran, schandraseka@northwell.edu and Chad E Bouton,
17 cbouton@northwell.edu

18 **Abstract**

19 Long-term recovery of limb function is a significant unmet need in people with paralysis.
20 Targeted neuromodulation through epidural stimulation of the spinal cord, when paired with
21 intense physical therapy, has shown promising results towards restoring volitional control of
22 the upper limb in people with tetraplegia. Non-invasive neuromodulation of the cervical spinal
23 cord using transcutaneous spinal cord stimulation (tSCS) has shown similar improvements in
24 upper-limb motor control rehabilitation. However, the potential rehabilitative effects of
25 activating specific cervical spinal segments in a targeted fashion using tSCS has largely remained
26 unexplored. We show in two individuals with SCI that tSCS of the cervical spinal cord resulted in
27 up to 1136% increase in exerted force, with minimal physical therapy. We also show, for the
28 first time, the effectiveness of targeted tSCS in restoring strength (407% increase) and dexterity
29 to the digits in an individual with paralysis of the hand due to a peripheral nerve injury.
30 Furthermore, we believe this is the first study to document a 3-point improvement in sensation
31 in people with SCI or peripheral injury after receiving tSCS. Lastly, participant gains persisted
32 after a one-month period void of stimulation, suggesting tSCS may lead to lasting benefits. Non-
33 invasive targeted spinal cord stimulation shows tremendous promise as a safe and effective
34 therapeutic approach with broad applications.

35 **Introduction**

36 Spinal cord and peripheral nerve injury can result in paralysis due to the disruption in
37 the transmission of neural signals. More than half of spinal cord injuries (SCIs) occur at the
38 cervical level (NSCISC and University of Alabama at Birmingham, 2021), and regaining voluntary
39 control of the hand and arm is the highest priority in such cases (Anderson, 2004). However,
40 the chances of regaining hand and arm function are exceedingly low beyond 12-18 months post
41 injury (Fawcett et al., 2006).

42 Electrical stimulation of the lumbar spinal cord using epidurally placed electrodes has
43 recently shown great promise in evoking voluntary movements of the lower limb during
44 stimulation after SCI, in rats (van den Brand et al., 2012), non-human primates (Capogrosso et
45 al., 2016) and humans (Harkema et al., 2011; Angeli et al., 2014, 2018; Gill et al., 2018; Wagner
46 et al., 2018). Additionally, when paired with intense motor training, lumbar epidural stimulation
47 has also demonstrated persistent increase in voluntary control of the lower limbs even in the
48 absence of stimulation (Rejc et al., 2017; Angeli et al., 2018; Wagner et al., 2018; Seáñez and
49 Capogrosso, 2021; Rowald et al., 2022). Epidural stimulation has been shown to primarily
50 engage the large-to-medium size sensory afferent fibers present in the dorsal column and roots
51 of the spinal cord (Rattay et al., 2000). Local spinal circuits constitute of these afferent fibers
52 forming synaptic connections with spinal interneurons and motoneurons. Pharmacological and
53 computational experiments (Capogrosso et al., 2013) suggest that activation of the dorsal fibers
54 increases the excitability of the local spinal circuitry including the efferent fibers. Expanding this

55 idea further, recent studies have demonstrated that epidural stimulation targeting cervical
56 spinal cord could activate (Greiner et al., 2020) and restore upper limb movement (Lu et al.,
57 2016). Though extremely promising, the invasive nature of epidural stimulation is a major
58 hurdle for clinical translation and long-term rehabilitation (James et al., 2018; Taccola et al.,
59 2020).

60 Meanwhile, non-invasive, persistent stimulation of the spinal cord using transcutaneous
61 electrodes has been demonstrated to evoke voluntary movements in both upper (Freyvert et
62 al., 2018; Gad et al., 2018; Inanici et al., 2021) and lower limbs (Sayenko et al., 2015, 2019).
63 Moreover, recent evidence suggests that transcutaneous spinal cord stimulation (tSCS), like
64 epidural stimulation, exerts its neuromodulatory effect on motoneurons trans-synaptically via
65 activation of large-to-medium size sensory afferent fibers (Ladenbauer et al., 2010; Hofstoetter
66 et al., 2018) and can also improve descending supraspinal drive (Guiho et al., 2021). Only
67 recently, has it been shown that pairing tSCS at the cervical levels with intense motor training
68 can result in sustained improvements in hand and arm function (Benavides et al., 2020; Zhang
69 et al., 2020; Inanici et al., 2021). Taken together, tSCS can be a promising rehabilitative tool for
70 people with SCI.

71 The commercially available electrodes employed for tSCS, however, are relatively large
72 resulting in a wider distribution of the applied current. Additionally, being placed over the skin,
73 the electrodes are also distant from the dorsal fibers of the spinal cord with layers of
74 intervening connective and bony tissue including the dorsal aspects of the cervical vertebrae.
75 These factors may limit the effectiveness and specificity of tSCS for the recruitment of target
76 motoneuron pools which could be a major drawback since epidural stimulation studies have
77 shown the importance of targeted stimulation (Wagner et al., 2018) in restoring voluntary
78 muscle control.

79 We performed targeted transcutaneous stimulation of the cervical spinal cord paired
80 with minimal physical therapy in two individuals classified as having a motor complete SCI and
81 one individual with a peripheral nerve injury. Stimulation was targeted to specific cervical levels
82 using a custom electronically configurable electrode array, with each electrode having a small
83 form factor (10 mm x 10 mm). Modulating the spatial configuration of the stimulation
84 electronically allowed us to extensively map the recruitment profile of the upper-limb muscles
85 based on the location of stimulation. This enabled us to choose the precise location of
86 stimulation to achieve maximal recruitment of the muscle group of interest. Even though
87 participants received stimulation and motor training only once per week, we observed a rapid
88 increase in both volitionally controlled muscle activity and effective force within a period of 5-6
89 weeks. However, the observed gains were restricted to muscles that generated at least a

90 measurable amount of force at the beginning of the intervention. In the subject with peripheral
91 nerve injury, we observed a rapid increase in functionality of the affected digits.

92 Taken together, this study demonstrates the advantages of using a highly configurable
93 electrode array in conjunction with minimal physical therapy in restoring volitional control of
94 upper-limb movement using tSCS. Also, this is the first study to demonstrate the benefits of
95 tSCS for improving hand function in a case of peripheral injury.

96 **Methods**

97 **Participants**

98 We performed transcutaneous stimulation of the cervical spinal cord in two individuals
99 with tetraplegia resulting from C5 level motor complete spinal cord injuries and one individual
100 with paralysis of the left hand due to peripheral nerve injury (median nerve at the wrist and
101 suspected brachial plexus injury), but with an otherwise intact spinal cord. The details of the
102 participants are summarized in Table 1. All procedures were approved by the Northwell Health
103 Institutional Review Board. The study has been registered with ClinicalTrials.gov
104 (NCT04755699).

Table 1 Study participants information. Demographic and injury-related information for each participant.

Participant	Age Range	Gender	Injury type	Time since injury (yrs.)
CTS01	late 40s	F	Median nerve, Brachial plexus	4
CTS02	early 20s	M	SCI: C5, ASIA A	2
CTS03	mid 30s	M	SCI: C5, ASIA B	7

105 **Experimental sessions**

106 Participants visited the lab for their intervention sessions once per week. Their heart
107 rate and blood pressure were measured at the beginning and end of each session. Baseline
108 GRASSP assessments were carried out in two initial sessions, scheduled a month apart, to
109 establish an initial baseline. For finer measurements of voluntary muscle control, we measured
110 both the evoked force and electromyographic (EMG) activity in the muscles of the hand and
111 arm. For efficiency, we restricted our measurements to only one side for all participants. For
112 participants with SCI, we chose the left side. For the participant with the peripheral nerve
113 injury, we chose the side that was affected by the resultant paralysis, i.e., the left side. Bipolar
114 surface EMG electrodes were used to record muscle activity from the following muscles of the
115 left arm and hand: biceps brachii (BIC), triceps brachii (TRI), flexor digitorum profundus (FDP),
116 extensor digitorum communis (EDC) and abductor policis brevis (ABP). Force generated by the
117 isometric flexion of the muscles of interest and EMG activity were simultaneously recorded at
118 the start of each session.

119 For evaluating the forces and EMG activity generated by the tricep muscle, as in the case
120 of participants CTS02 and CTS03, the participants' arms were extended out in front of them,
121 with the hand in a neutral position and a load cell placed under the ulnar head prominence at
122 the wrist (Figure 1C). The participants were instructed pushed down on to the load cell in this
123 position. The elbow was resting on the table and prevented from lifting during the task. For
124 evaluation of the D1 interphalangeal (D1 IP) joint and lateral pinch, as in the case of participant
125 CTS01, a custom rig was used as shown in Figure 1D-E. The participant was asked to push down
126 on the sensor using their thumb. For all force measuring tasks, the experimenter cued the
127 participants to push against the force sensor for 3-5 secs and then to relax for 3-5 secs. The task
128 was performed in 2 sets of 5 trials each with a 60-90 secs of rest period between the sets.
129 Verbal encouragement was provided to encourage the participants to generate maximal
130 maintainable force.

131 While administering tSCS as a therapeutic intervention, the participant performed the
132 same tasks as described above. This served the dual purpose of providing physical therapy
133 while receiving tSCS as well as measuring the evoked volitional EMG activity and force.

134 **Transcutaneous spinal cord stimulation (tSCS)**

135 Transcutaneous spinal cord stimulation was provided using a custom-built stimulator
136 and electrode array. The flexible PCB electrode array consisted of electroless nickel immersion
137 gold (ENIG) or immersion silver-plated square contacts (10 mm x 10 mm) arranged in an 8 x 5
138 pattern with a 1 mm inter-electrode separation. Electrical stimulation to any combination of
139 the 40 electrodes within the array, was controlled using solid-state relays mounted above each
140 of the electrodes and a custom MATLAB-based GUI. Miniature green LEDs mounted on the
141 dorsal surface of the electrode array, visually indicated the active electrodes. To target specific
142 cervical segments (Figure 1A), an electrode configuration of 1 x 3 (Figure 1B) was used wherein
143 3 adjacent contacts within a single row, spanning the midline, were used. The electrode array
144 was affixed to the back of the neck using a rectangular piece of proprietary hydrogel. To ensure
145 consistency in placement of the array between sessions, we used the inion of the external
146 occipital protuberance as a landmark. Distances measured from the inion were used to place
147 the electrode array and identify the location of stimulation. Two 5 x 10 cm rectangular self-
148 adhesive hydrogel electrodes (Axelgaard Manufacturing Co., Ltd., USA) placed along the midline
149 over the lumbar spinal cord served as return electrodes (anodes).

150 Stimulation consisted of a 10 kHz multiphasic sinusoidal waveform with pulse duration of 1 ms
151 for generating recruitment profiles. The waveform length was reduced to 0.5 ms during therapy
152 to reduce neck muscle activation and increase participant comfort. Similar pulse duration has




Figure 1. Experiment setup. A) Schematic showing the location of a 1 x 3 activated electrode configuration superimposed over the human spinal cord showing the dorsal column and roots. B) The custom electronically configurable electrode array placed over the cervical spinal cord of a study participant with a 1 x 3 configuration of activated electrodes (green LEDs). C) Setup for measuring tricep force. The ulnar protrusion of the wrist is placed over the 25 lb load cell during the task (red dashed square). D-E) Using the custom-built rig with a 10 lb load cell (gray puck) for measuring the force using D1 interphalangeal joint and lateral pinch, respectively in case of participant CTS01.

153 been used in previously published work (Alam et al., 2020; Benavides et. al, 2020). The
154 stimulation frequency was chosen to be 3 Hz for characterizing recruitment curves and 50 Hz
155 during therapy.

156 While administering tSCS as a therapeutic intervention, the stimulation amplitude was
157 chosen to be 140-160 mA. This amplitude was chosen based on patient comfort but at the
158 same time was an amplitude that did not interfere with the tasks being performed by the
159 participant. Stimulation was delivered for 45-60 minutes and was paired with the participant
160 performing the force measurement tasks described above.

161 After 16 weeks of receiving of tSCS, we administered a 3-week period when the
162 participant received no stimulation at all. They continued to perform the tasks that constituted
163 their physical therapy as described above. Weekly administration of tSCS was resumed after
164 this 'No Stim' period.

165 **EMG signal processing**

166 We simultaneously recorded bipolar EMG from the following muscles of the left arm
167 and hand: biceps brachii (BIC), triceps brachii (TRI), flexor digitorum profundus (FDP), extensor
168 digitorum communis (EDC) and abductor policis brevis (ABP). We used pre-gelled Ag/AgCl
169 electrodes, a differential amplifier (AD6221) with a bandpass filter (20-498Hz), and a signal
170 digitizer (PicoScope® Model 4824A). The sampling rate used was 10 MHz while characterizing
171 the recruitment profile of the upper limb muscles and 10-20 kHz during task performance. For
172 characterizing the area-under-the curve (AUC), the EMG signal was filtered using a 60Hz IIR
173 comb filter and Butterworth bandpass filter between 10-1000 Hz using a digital filter in
174 MATLAB.

175 **Recruitment profile**

176 Stimulation consisted of a 1ms-long multiphasic pulse of the 10 kHz sinusoidal
177 waveform at a stimulation frequency of 3 Hz delivered using an electronically selected
178 configuration of three contacts within a single row spanning the midline. Stimulation
179 amplitudes tested ranged from 100 mA to up to 225 mA in intervals of approximately 25 mA.
180 This was repeated for each of the eight rows of electrodes on the array. The Picoscope 6
181 acquisition software (Pico Technology, Cambridgeshire, UK) was used to trigger acquisition of a
182 100 ms-long EMG signal following each stimulation pulse. For each stimulation amplitude and
183 at each electrode row, we recorded such an EMG signal from 20-30 repetitions of the
184 stimulation pulse. All the data was imported into MATLAB for further analysis. From each of the
185 EMG signals recorded, we isolated a snippet starting from 5 ms and ending at 55ms after the
186 stimulation artifact. We measured the peak-to-peak amplitude (P2P) for this snippet. It was
187 included for analysis only if the maximum amplitude of the snippet was greater than 5 times
188 the standard deviation of baseline signal of that recording channel. For each muscle, the P2P
189 amplitudes were normalized to the maximal P2P amplitude recorded across all amplitudes and
190 electrode rows.

191 We also measured the stimulation current of maximal activation for each muscle. For
192 this, at each cervical level, we increased the amplitude until the reflexive activation of the
193 muscle was visually determined to be consistent for every stimulation pulse and did not change
194 for any small increase in amplitude. This stimulation amplitude was noted as the Maximal
195 Activation Current for that muscle at the cervical level. This procedure was repeated for each
196 muscle.

197 **Functional Outcome**

198 In addition to the areas recommended in the GRASSP assessment for sensibility
199 evaluation, we tested few additional relevant spots on the dorsal and volar aspect of the hand
200 to obtain a comprehensive profiling of the sensory changes secondary to the intervention
201 sessions. To provide encouragement and also observe the true capability of the participant, we
202 initially let the GRASSP motor assessment tasks run for more than the stipulated time of 75
203 secs.

204 **Statistical Analysis**

205 For statistical analysis of the evoked force, we compared the forces recorded in three stages of
206 the study (start, no-stim and end of study). For the early stage, we chose the first three
207 consecutive sessions after start of weekly tSCS sessions. For the no-stim stage, we chose three
208 sessions during the period when tSCS was not being delivered. For the end of study stage, we
209 chose the last three consecutive sessions. For the force measurements of the tricep muscle in
210 case of the participants with SCI, we performed a repeated-measures Analysis of Variance
211 (ANOVA) between the average values of force obtained in the three epochs. For the force
212 measurements involving the D1 IP muscle in case of the participant with the peripheral nerve
213 injury, we performed a t-test. We employed the Bonferroni correction for all multiple
214 comparisons.

215 **Results**

216 Our results demonstrate that transcutaneous spinal cord stimulation can be used to activate
217 specific motoneuron pools in the cervical spinal cord. Moreover, targeted transcutaneous spinal
218 cord stimulation paired with minimal physical therapy resulted in a substantial and sustained
219 increase in muscle activity and strength in specific upper-limb muscles in two patients with
220 motor complete cervical SCI and one participant with a peripheral injury. All three participants
221 received stimulation for up to 1hr, once per week. Blood pressure and heart rate monitoring at
222 the start and end of each session showed no adverse effects of the stimulation.

223 **Recruitment of upper limb muscles through targeted tSCS**

224 We characterized the recruitment profile of the different motor pools innervating the
225 upper limb muscles. To determine the effect of the location of stimulation along the rostro-
226 caudal axis on the recruitment of the upper limb muscles, we sequentially delivered stimulation
227 to different electrode triplets in the array (Figure 2A-C); also see Supplementary Figure 1 for an
228 example stimulation-evoked EMG activity. Simultaneously, we recorded EMG activity from 5
229 muscles of the left arm and hand. Stimulation amplitude was increased and the corresponding
230 increase in EMG amplitude was used as a measure of recruitment. We hypothesized that the
231 recruitment pattern through transcutaneous stimulation would reflect the rostrocaudal
232 segment-wise distribution of the upper limb motor nuclei in the cervical spinal cord. The

233 recruitment profiles showed a distinct and consistent shape in all participants. Almost all the
234 muscles exhibited the best recruitment when stimulation was delivered around the C5-C6 level.
235 Stimulating using rostral electrodes primarily activated the biceps (Figure 2D, G and E, H),
236 except in CTS03. Activation of the triceps invariably was strongest when stimulation was
237 localized around the C5-C6 level. Stimulation using electrodes positioned more caudally
238 resulted in decreased recruitment across all muscles. C8-T1 level stimulation showed
239 recruitment of FDP and APB in participant CTS03 (Figure 2C, F and I). In participant CTS01, APB
240 showed only weak recruitment irrespective of the location of stimulation, possibly due to the
241 suspected brachial plexus injury. The fact that two participants had a spinal cord injury and
242 metallic implants in the cervical vertebrae did not result in an obvious difference in the
243 recruitment profile of the upper-limb motor pools is noteworthy. Additionally, we also
244 determined the current required for maximal activation (Max. Activation Current) for each of
245 the upper limb muscles. The thresholds of activation showed similar results as the recruitment
246 profiles with biceps and/or triceps muscles having a low threshold of activation during rostral
247 stimulation (Figure 2G, H and I).

248




Figure 2. Muscle recruitment profile during cervical tCS. A-C) X-ray images in the sagittal plane with radio-opaque markers on the neck (white dots) for the three participants. The topmost marker identifies the inion of the external occipital protuberance. The second and third markers identify points 7 cm and 9.2 cm respectively, from the inion signifying the first and third rows of a putative electrode array whose first row of electrodes was aligned at 7cm from the inion. The last marker identifies the location of the last row of the putative electrode array at 15.7 cm from the inion. The cervical labels mark the exit point of the respective dorsal roots. **D-F)** Mean activation of the 5 muscles across all stimulation amplitudes mediated by tCS through each of the 8 electrode rows. **G-I)** Stimulation amplitude that resulted in maximal activation of each of the 5 muscles.

249 Increased muscle activity and force generated with tSCS

250 Previous spinal cord stimulation studies have generally shown improvements in muscles
251 that have at least some residual activity. Given this, we focused on muscles with at least some
252 residual activity in our participants. Since both SCI participants exhibited no movement in their




Figure 3. Increase in muscle activity and force after tSCS. A) The force generated by the left D1 IP joint in case of participant CTS01. * indicates a significant difference (Two-sample t-test with Bonferroni correction for multiple comparisons) **B)** The AUC of the EMG activity recorded from the left APB muscle during the task of force generation using the D1 IP joint. **C)** The force generated by the left tricep muscle in case of participant CTS02 (blue) and CTS03 (orange). * indicates significant difference (Repeated measures ANOVA and Bonferroni correction for multiple comparison). The blue (CTS02) and orange (CTS03) bars indicate the specific data points from each participant that were included for the statistical analysis. **D)** The AUC of the EMG activity recorded from the left tricep muscle during the task of force generation using the left tricep muscle.

253 fingers, we focused on the volitional control of their tricep muscle. Meanwhile, as the
254 participant with the brachial plexus injury (CTS01) had residual movement in the fingers of the
255 left hand, we focused on the volitional control of the D1 interphalangeal (D1 IP) joint and lateral
256 pinch. Here we report the forces and EMG recorded from these specific muscles before any
257 tSCS was delivered during the session. We chose the site of stimulation based on the choice of
258 muscle and its activation profile. For the participant with the brachial plexus injury, we focused
259 on thumb muscles, and delivered tSCS targeted at the C5 level (about 8 cm from theinion of
260 the external occipital protuberance) based on the activation profile for the APB muscle (see
261 Figure 2A, D and G). For participants CTS02 and CTS03, we delivered tSCS targeted at the C6
262 level (about 10 cm from theinion of the external occipital protuberance).

263 We observed an average increase of 407% (t-statistic = -5.26, $p < 0.05$) in the force
264 generated by the flexion of D1 IP joint (Figure 3A) in the 'No Stim' period at the end of 16
265 sessions compared to the forces at the start of the study. This reduced a little to 287% increase
266 by the end of the study (30 sessions). This was accompanied by an increase in EMG activity of
267 the left APB muscle (Figure 3B). We did not observe any major increase in the lateral pinch
268 force during this period. We observed an average increase of force generated by the left tricep
269 muscle of up to 893% (at the end of 17 sessions) and 825% (at the end of 16 sessions) for CTS02
270 and CTS03, respectively, in the 'No Stim' period compared to the start of the study. This
271 increased to 1136% (29 sessions) and 1035% (31 sessions), respectively ($F = 13.99$, $p < 0.01$),
272 (Figure 3C, Supp. Fig. 1). We also observed a corresponding increase in EMG activity for the
273 tricep muscle (Figure 3D).

274 **GRASSP assessments**

275 To evaluate the clinical significance of the progress showed by each participant, we also
276 performed standard clinical assessments, namely the GRASSP test. The GRASSP strength test
277 showed only minor increases in the movements assessed, especially in the participants with SCI
278 including the tricep muscle which was specifically targeted in this study (Figure 4A). GRASSP
279 sensibility test showed decreases in sensation threshold at many locations in the hand in all
280 participants (Figure 4B). For the participant with the brachial plexus injury, most of the
281 improvement was observed in the thumb as well as the ulnar side of the hand. GRASSP strength

282 showed increases in D1 IP and D5 abduction movements (Figure 4A). GRASSP sensibility
283 showed up to a 3-point increase in sensation at the pinky tip (Figure 4B).




Figure 4. GRASSP assessments. A) GRASSP strength change from baseline **B)** GRASSP sensibility change from baseline. Circles show 1- to 3-point increase in sensory perception as defined by the GRASSP sensibility scale. **C)** GRASSP prehension change in nut threading task for CTS01.

284

285 The nut threading task included in the GRASSP prehension tests showed an
286 improvement in performing the right grasp (Figure 5A) and a decrease in the time taken to
287 perform the task (Figure 5B). Participant CTS01 also showed improvement in range of
288 movement of thumb abduction, flexion, and extension as well as index abduction (Supp. Fig. 2).
289 Participants also shared their anecdotal descriptions of the effect of receiving tSCS. These
290 described increased control in moving their arms and being able to give “stronger hugs.”




Figure 5. GRASSP Prehension Test – Nut threading. A) Snapshots showing improvement in grasping behavior from before tSCS therapy (first row) and week 42 of therapy (second row). **B)** Scatter plot shows the improvement in the time required to perform the task. 'x' denotes that the participant never managed to complete the task.

291 Discussion

292 In this study, we observed a substantial increase in volitionally generated force and EMG
293 activity in specific muscles after paralysis from tSCS in two individuals with SCI and one with a
294 peripheral injury (median nerve and suspected brachial plexus injury). All three participants
295 showed these changes within a brief period of receiving tSCS. Interestingly, the improvements
296 were restricted to the muscles that showed at least some measurable amount of force at the
297 beginning of therapy. For instance, neither of the participants with SCI registered any force
298 during finger flexion. We did not observe any improvement in the forces and EMG activity
299 generated during finger flexion in either participant with SCI even when the activity was
300 performed over many weeks while receiving tSCS. This could be due to non-optimal tSCS
301 patterns or that the severity of the injuries was too severe to benefit from this type of
302 therapeutic intervention.

303 Our results are comparable to those demonstrated by other studies involving cervical tSCS in
304 people with SCI (Inanici et al., 2021). However, our design involved participants receiving spinal
305 cord stimulation and minimal physical therapy for 1-2 hrs only once every week. Importantly,
306 our stimulation was targeted to achieve maximal recruitment of the muscle group of interest.
307 This suggests that targeted tSCS could improve the efficacy of spinal cord activation and achieve
308 restoration of volitional control even with minimal physical therapy.

309 The electrode configuration we used in this study was a 1 row x 3 columns configuration
310 spanning the spinal cord midline. The effectiveness of such an electrode configuration in
311 recruitment of specific motoneuron pools has been demonstrated in able-bodied individuals
312 (Gerasimenko et al., 2015; Krenn et al., 2015). It is highly likely that this configuration resulted
313 in the stimulation of the dorsal column fibers thereby activating motor nuclei in off-target
314 cervical levels and thus reducing specificity. Activation of the dorsal roots via lateralized
315 stimulation could result in greater selectivity in the motoneuron pools being activated.
316 However, an earlier study with lateralized stimulation showed an increase in side-specific
317 activation and not across spinal levels (Calvert et al., 2019). Previous studies have explored the

318 idea of targeting stimulation above and below the injury level with the aim of enhancing the
319 activity of the descending inputs as well as local circuitry below the site of injury (Zhang et al.,
320 2020; Inanici et al., 2021). With a high cervical location of SCI (C5) for the participants in this
321 study, most of the stimulation was restricted to being targeted at or below the injury level
322 (Freyvert et al., 2018). In fact, stimulation targeted at higher cervical levels was perceived as
323 uncomfortable by the participants. This demonstrates that tSCS can be tailored to suit patient
324 comfort, target only those cervical levels innervating the muscle of interest, and still result in
325 significant motor improvements.

326 Furthermore, we used a custom electrode array that could be electronically configured. This
327 allowed efficient mapping and can support dynamic spatial pattern switching. In future studies,
328 this feature can be combined with brain-computer interfaces (BCIs) to switch spatial patterns
329 based on movement-related information decoded from intracortical activity (Bouton et al.,
330 2016; Sharma et al., 2016; Friedenberget al., 2017; Chandrasekaran et al., 2021b). Such
331 automated configuration of targeted tSCS based on user movement intentions could improve
332 the usability of this technology by people with SCI while performing activities of daily living.

333 It was interesting to observe improvement in somatosensory perception in regions innervated
334 by spinal roots from below the injury level in both participants with SCI. To our knowledge, this
335 is the first study to document improvements in sensation in people with SCI or peripheral injury
336 after receiving tSCS. Stimulating dorsal roots of the spinal cord has been demonstrated to relay
337 somatotopically relevant sensory information (Chandrasekaran et al., 2020). Meanwhile,
338 restoring somatosensation through intracortical stimulation has been repeatedly demonstrated
339 in humans as well (Flesher et al., 2016; Chandrasekaran et al., 2021a; Fifer et al., 2021). It may
340 be worthwhile to explore the benefits of pairing intracortical stimulation and tSCS for the long-
341 term rehabilitation of somatosensation in SCI.

342 We believe this is the first study that performed tSCS in an individual with peripheral injury
343 (Lopez and Sdrulla, 2021) that resulted in the rehabilitation of volitional control of hand
344 movement. The rapid and significant improvement observed in the hand function in this case
345 signifies major implications for a broader range of conditions affecting the central and
346 peripheral nervous systems, including traumatic injuries and stroke.

347 **Acknowledgements**

348 We would like to thank our participants for their extraordinary commitment to this study, their
349 patience with the experiments, and the deep insights provided by them, the clinicians, and
350 researchers at Feinstein Institutes for Medical Research. We would like to extend our deepest
351 gratitude to Dan Sczubba, MD, David Bonda, MD and Karen Black, MD at Northwell Health for
352 useful discussions and assistance related to the imaging sessions.

353 **Funding**

354 This study was funded through support provided by Feinstein Institutes for Medical Research at
355 Northwell Health.

356 **Author Contributions**

357 SC, NAB and CEB designed the study. SC, NAB, RR, SE and CEB performed the experiments and
358 SC analyzed the data. NAB designed and tested the electrode array, with guidance from CEB.
359 DGG performed an independent assessment of the motor and sensory capacity of the
360 participants. PDS and SJH provided critical feedback and input. All authors contributed towards
361 interpreting the results of the experiments. SC and CEB finished the initial draft of the
362 manuscript and all authors provided critical review, edits, and approval of the final manuscript.

363 **Competing Interest Statement**

364 CEB has financial interests in Sanguistat, LLC and Neuvotion, Inc. and multiple patents in the
365 neurostimulation field. SE has financial interests in Neuvotion, Inc. The remaining authors have
366 no conflicting financial interests.

367 **References**

- 368 Anderson KD (2004) Targeting Recovery: Priorities of the Spinal Cord-Injured Population. *Journal*
369 *of Neurotrauma* 21:1371–1383.
- 370 Angeli CA, Boakye M, Morton RA, Vogt J, Benton K, Chen Y, Ferreira CK, Harkema SJ (2018)
371 Recovery of Over-Ground Walking after Chronic Motor Complete Spinal Cord Injury. *New*
372 *England Journal of Medicine* 379:1244–1250.
- 373 Angeli CA, Edgerton VR, Gerasimenko YP, Harkema SJ (2014) Altering spinal cord excitability
374 enables voluntary movements after chronic complete paralysis in humans. *Brain* 137:1394–
375 1409 Available at: [https://academic.oup.com/brain/article-](https://academic.oup.com/brain/article-lookup/doi/10.1093/brain/awu038)
376 [lookup/doi/10.1093/brain/awu038](https://academic.oup.com/brain/article-lookup/doi/10.1093/brain/awu038) [Accessed June 25, 2019].
- 377 Benavides FD, Jo HJ, Lundell H, Reggie V, Perez MA (2020) Cortical and Subcortical Effects of
378 Transcutaneous Spinal Cord Stimulation in Humans with Tetraplegia.
- 379 Bouton CE, Shaikhouni A, Annetta N v., Bockbrader MA, FriedenberG DA, Nielson DM, Sharma G,
380 Sederberg PB, Glenn BC, Mysiw WJ, Morgan AG, Deogaonkar M, Rezai AR (2016) Restoring
381 cortical control of functional movement in a human with quadriplegia. *Nature* 533:247–250.
- 382 Calvert JS, Manson GA, Grahn PJ, Sayenko DG (2019) Preferential activation of spinal
383 sensorimotor networks via lateralized transcutaneous spinal stimulation in neurologically
384 intact humans. *Journal of Neurophysiology* 122:2111–2118 Available at: www.jn.org
385 [Accessed February 14, 2022].
- 386 Capogrosso M et al. (2016) A brain–spine interface alleviating gait deficits after spinal cord injury
387 in primates. *Nature* 539:284–288 Available at:
388 <http://www.nature.com/articles/nature20118> [Accessed June 26, 2019].
- 389 Capogrosso M, Wenger N, Raspopovic S, Musienko P, Beauparlant J, Luciani LB, Courtine G,
390 Micera S (2013) A computational model for epidural electrical stimulation of spinal
391 sensorimotor circuits. *J Neurosci* 33:19326–19340.
- 392 Chandrasekaran S, Bickel S, Herrero JL, Kim J won, Markowitz N, Espinal E, Bhagat NA, Ramdeo R,
393 Xu J, Glasser MF, Bouton CE, Mehta AD (2021a) Evoking highly focal percepts in the
394 fingertips through targeted stimulation of sulcal regions of the brain for sensory restoration.
395 *Brain Stimulation* 14:1184–1196.
- 396 Chandrasekaran S, Fifer M, Bickel S, Osborn L, Herrero J, Christie B, Xu J, Murphy RKJ, Singh S,
397 Glasser MF, Collinger JL, Gaunt R, Mehta AD, Schwartz A, Bouton CE (2021b) Historical
398 perspectives, challenges, and future directions of implantable brain-computer interfaces for
399 sensorimotor applications. *Bioelectronic Medicine* 2021 7:1 7:1–11 Available at:
400 <https://link.springer.com/articles/10.1186/s42234-021-00076-6> [Accessed May 17, 2022].

- 401 Chandrasekaran S, Nanivadekar AC, McKernan G, Helm ER, Boninger ML, Collinger JL, Gaunt RA,
402 Fisher LE (2020) Sensory restoration by epidural stimulation of the lateral spinal cord in
403 upper-limb amputees. *Elife* 9:1–26.
- 404 Fawcett JW et al. (2006) Guidelines for the conduct of clinical trials for spinal cord injury as
405 developed by the ICCP panel: spontaneous recovery after spinal cord injury and statistical
406 power needed for therapeutic clinical trials. *Spinal Cord* 2007 45:3 45:190–205 Available at:
407 <https://www.nature.com/articles/3102007> [Accessed December 5, 2021].
- 408 Fifer MS, McMullen DP, Osborn LE, Thomas TM, Christie BP, Nickl RW, Candrea DN, Pohlmeier
409 EA, Thompson MC, Anaya MA, Schellekens W, Ramsey NF, Bensmaia SJ, Anderson WS,
410 Wester BA, Crone NE, Celnik PA, Cantarero GL, Tenore F v (2021) Intracortical
411 Somatosensory Stimulation to Elicit Fingertip Sensations in an Individual With Spinal Cord
412 Injury. *Neurology*:10.1212/WNL.00000000000013173.
- 413 Flesher SN, Collinger JL, Foldes ST, Weiss JM, Downey JE, Tyler-Kabara EC, Bensmaia SJ, Schwartz
414 AB, Boninger ML, Gaunt RA (2016) Intracortical microstimulation of human somatosensory
415 cortex. *Science Translational Medicine* 8.
- 416 Freyvert Y, Yong NA, Morikawa E, Zdunowski S, Sarino ME, Gerasimenko Y, Edgerton VR, Lu DC
417 (2018) Engaging cervical spinal circuitry with non-invasive spinal stimulation and buspirone
418 to restore hand function in chronic motor complete patients. *Scientific Reports* 8:15546
419 Available at: <https://www.nature.com/articles/s41598-018-33123-5>.
- 420 FriedenberG DA, Schwemmer MA, Landgraf AJ, Annetta N v., Bockbrader MA, Bouton CE, Zhang
421 M, Rezai AR, Mysiw WJ, Bresler HS, Sharma G (2017) Neuroprosthetic-enabled control of
422 graded arm muscle contraction in a paralyzed human. *Scientific Reports* 2017 7:1 7:1–10
423 Available at: <https://www.nature.com/articles/s41598-017-08120-9> [Accessed May 17,
424 2022].
- 425 Gad P, Lee S, Terrafranca N, Zhong H, Turner A, Gerasimenko Y, Edgerton VR (2018) Non-invasive
426 activation of cervical spinal networks after severe paralysis. *Journal of Neurotrauma*
427 35:2145–2158 Available at:
428 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6119225/pdf/neu.2017.5461.pdf>.
- 429 Gerasimenko Y, Gorodnichev R, Moshonkina T, Sayenko D, Gad P, Reggie Edgerton V (2015)
430 Transcutaneous electrical spinal-cord stimulation in humans. *Annals of Physical and*
431 *Rehabilitation Medicine* 58:225–231.
- 432 Gill ML, Grahn PJ, Calvert JS, Linde MB, Lavrov IA, Strommen JA, Beck LA, Sayenko DG, van
433 Straaten MG, Drubach DI, Veith DD, Thoreson AR, Lopez C, Gerasimenko YP, Edgerton VR,
434 Lee KH, Zhao KD (2018) Neuromodulation of lumbosacral spinal networks enables
435 independent stepping after complete paraplegia. *Nature Medicine* 2018 24:11 24:1677–

- 436 1682 Available at: <https://www.nature.com/articles/s41591-018-0175-7> [Accessed
437 February 7, 2022].
- 438 Greiner N, Barra B, Schiavone G, James N, Fallegger F, Borgognon S, Lacour S, Bloch J, Courtine G,
439 Capogrosso M (2020) Recruitment of Upper-Limb Motoneurons with Epidural Electrical
440 Stimulation of the Primate Cervical Spinal Cord. *bioRxiv:2020.02.17.952796* Available at:
441 <https://www.biorxiv.org/content/10.1101/2020.02.17.952796v1>.
- 442 Guiho T, Baker SN, Jackson A (2021) Epidural and transcutaneous spinal cord stimulation
443 facilitates descending inputs to upper-limb motoneurons in monkeys. *Journal of Neural
444 Engineering* 18:046011.
- 445 Harkema S, Gerasimenko Y, Hodes J, Burdick J, Angeli C, Chen Y, Ferreira C, Willhite A, Rejc E,
446 Grossman RG, Edgerton VR (2011) Effect of epidural stimulation of the lumbosacral spinal
447 cord on voluntary movement, standing, and assisted stepping after motor complete
448 paraplegia: A case study. *The Lancet* 377:1938–1947 Available at:
449 [http://dx.doi.org/10.1016/S0140-6736\(11\)60547-3](http://dx.doi.org/10.1016/S0140-6736(11)60547-3) [Accessed June 25, 2019].
- 450 Hofstoetter US, Freundl B, Binder H, Minassian K (2018) Common neural structures activated by
451 epidural and transcutaneous lumbar spinal cord stimulation: Elicitation of posterior root-
452 muscle reflexes. *PLoS One* 13 Available at: <https://pubmed.ncbi.nlm.nih.gov/29381748/>
453 [Accessed December 21, 2021].
- 454 Inanici F, Brighton LN, Samejima S, Hofstetter CP, Moritz CT (2021) Transcutaneous Spinal Cord
455 Stimulation Restores Hand and Arm Function after Spinal Cord Injury. *IEEE Transactions on
456 Neural Systems and Rehabilitation Engineering* 29:310–319.
- 457 James ND, McMahon SB, Field-Fote EC, Bradbury EJ (2018) Neuromodulation in the restoration
458 of function after spinal cord injury. *Lancet Neurol* 17:905–917.
- 459 Krenn M, Hofstoetter US, Danner SM, Minassian K, Mayr W (2015) Multi-Electrode Array for
460 Transcutaneous Lumbar Posterior Root Stimulation. *Artificial Organs* 39:834–840 Available
461 at: <https://onlinelibrary.wiley.com/doi/full/10.1111/aor.12616> [Accessed February 14,
462 2022].
- 463 Ladenbauer J, Minassian K, Hofstoetter US, Dimitrijevic MR, Rattay F (2010) Stimulation of the
464 human lumbar spinal cord with implanted and surface electrodes: A computer simulation
465 study. *IEEE Transactions on Neural Systems and Rehabilitation Engineering* 18:637–645.
- 466 Lopez L, Sdrulla AD (2021) Success with dorsal root entry zone lesioning after a failed trial of
467 spinal cord stimulation in a patient with pain due to brachial plexus avulsion. *PAIN Reports*
468 6:e973.
- 469 Lu DC, Edgerton VR, Modaber M, Auyong N, Morikawa E, Zdunowski S, Sarino ME, Sarrafzadeh
470 M, Nuwer MR, Roy RR, Gerasimenko Y (2016) Engaging Cervical Spinal Cord Networks to

- 471 Reenable Volitional Control of Hand Function in Tetraplegic Patients. *Neurorehabilitation*
472 and *Neural Repair* 30:951–962 Available at:
473 <https://journals.sagepub.com/doi/full/10.1177/1545968316644344> [Accessed December 6,
474 2021].
- 475 NSCISC, University of Alabama at Birmingham (2021) National Spinal Cord Injury Statistical
476 Center, Facts and Figures at a Glance. Available at: [www.msctc.org/sci/model-system-](http://www.msctc.org/sci/model-system-centers)
477 [centers](http://www.msctc.org/sci/model-system-centers). [Accessed December 15, 2021].
- 478 Rattay F, Minassian K, Dimitrijevic MR (2000) Epidural electrical stimulation of posterior
479 structures of the human lumbosacral cord: 2. quantitative analysis by computer modeling.
480 *Spinal Cord* 38:473–489.
- 481 Rejc E, Angeli CA, Atkinson D, Harkema SJ (2017) Motor recovery after activity-based training
482 with spinal cord epidural stimulation in a chronic motor complete paraplegic. *Scientific*
483 *Reports* 7:1–12.
- 484 Rowald A et al. (2022) Activity-dependent spinal cord neuromodulation rapidly restores trunk
485 and leg motor functions after complete paralysis. *Nature Medicine* 2022 28:2 28:260–271
486 Available at: <https://www.nature.com/articles/s41591-021-01663-5> [Accessed March 22,
487 2022].
- 488 Sayenko DG, Atkinson DA, Dy CJ, Gurley KM, Smith VL, Angeli C, Harkema SJ, Edgerton VR,
489 Gerasimenko YP (2015) Spinal segment-specific transcutaneous stimulation differentially
490 shapes activation pattern among motor pools in humans. *Journal of Applied Physiology*
491 118:1364–1374.
- 492 Sayenko DG, Rath M, Ferguson AR, Burdick JW, Havton LA, Edgerton VR, Gerasimenko YP (2019)
493 Self-Assisted Standing Enabled by Non-Invasive Spinal Stimulation after Spinal Cord Injury.
494 *Journal of Neurotrauma* 36:1435–1450.
- 495 Seáñez I, Capogrosso M (2021) Motor improvements enabled by spinal cord stimulation
496 combined with physical training after spinal cord injury: review of experimental evidence in
497 animals and humans. *Bioelectronic Medicine* 2021 7:1 7:1–13 Available at:
498 <https://bioelecmed.biomedcentral.com/articles/10.1186/s42234-021-00077-5> [Accessed
499 February 14, 2022].
- 500 Sharma G, FriedenberG DA, Annetta N, Glenn B, Bockbrader M, Majstorovic C, Domas S, Mysiw
501 WJ, Rezai A, Bouton C (2016) Using an artificial neural bypass to restore cortical control of
502 rhythmic movements in a human with quadriplegia. *Scientific Reports* 6.
- 503 Taccola G, Barber S, Horner PJ, Bazo HAC, Sayenko D (2020) Complications of epidural spinal
504 stimulation: lessons from the past and alternatives for the future. *Spinal Cord* 58:1049–
505 1059.

506 van den Brand R, Heutschi J, Barraud Q, DiGiovanna J, Bartholdi K, Huerlimann M, Friedli L,
507 Vollenweider I, Moraud EM, Duis S, Dominici N, Micera S, Musienko P, Courtine G (2012)
508 Restoring voluntary control of locomotion after paralyzing spinal cord injury. *Science* (1979)
509 336:1182–1185 Available at: <https://www.science.org/doi/abs/10.1126/science.1217416>
510 [Accessed December 6, 2021].


511 Wagner FB et al. (2018) Targeted neurotechnology restores walking in humans with spinal cord
512 injury. *Nature* 2018 563:7729 563:65–71 Available at:
513 <https://www.nature.com/articles/s41586-018-0649-2> [Accessed December 5, 2021].

514 Zhang F, Momeni K, Ramanujam A, Ravi M, Carnahan J, Kirshblum S, Forrest GF (2020) Cervical
515 Spinal Cord Transcutaneous Stimulation Improves Upper Extremity and Hand Function in
516 People with Complete Tetraplegia: A Case Study. *IEEE Transactions on Neural Systems and*
517 *Rehabilitation Engineering* 28:3167–3174.

518

519


520 Supplementary Figures



Supplementary Figure 1. Example trace of muscle recruitment profile of the 5 muscles upon tSCS stimulation. This was for participant CTS02 when stimulated at 10.2 cm at 150 mA. Colored traces indicate the reflex activity evoked in each of the 5 muscles as shown in the legend. Dashed black line shows average of all the trials.

521


522



Supplementary Figure 2. Example traces of cue, force measured during tricep extension and EMG recorded from triceps brachii for participant CTS02 from A) week 11 and B) week 32.

523

524



Supplementary Figure 3. Improvement in range of movement (ROM) in D1 abduction, D1 flexion, D1 extension and D2 abduction for participant CTS01. A) shows superimposed images from week 5 and week 31; B) Change in ROM for the 4 different movements.