

Single Injection Nerve Blocks Plus IV Lidocaine Infusions as an Alternative to Continuous Nerve Blocks for Perioperative Pain Management: A retrospective review.

Dmitriy Gromov,¹ Jeremy Kearns^{1,2}, Jacques E Chelly³

Authors Affiliations:

¹Regional Anesthesiology and Acute Pain Medicine Fellow, Department of Anesthesiology and Perioperative Medicine, University of Pittsburgh Medical Center, Pittsburgh, PA, ² current affiliation: Virtua Health System; ³ Professor, Department of Anesthesiology and Perioperative Medicine and Orthopedic Surgery, Director of Research, Co-Director of the Center for Innovation in Pain Care, Director of the Center of Complementary and Alternative Medicine, Director of the Center for Innovation in Pain Care, Founder and Former Chief, and member of the Acute Interventional Perioperative Pain Service, UPMC Presbyterian- Shadyside Hospital, Former Fellowship director, Regional Anesthesiology and Acute Pain Medicine Fellowship, Department of Anesthesiology and Perioperative Medicine, University of Pittsburgh and University of Pittsburgh Medical Center, Pittsburgh, PA

Correspondence: Chelly, Jacques E

Posner Pain Center, 532 S. Aiken Ave, Suite 407, Pittsburgh, PA, 15232

Tel: 1 412 623 6904; Fax: 1 412 623 4188

Email: chelje@anes.upmc.edu

Abstract:

Purpose: Continuous nerve blocks (CNBs) and intravenous lidocaine infusions (IV Lido) represent an effective approach to perioperative pain management. We hypothesized that a single injection nerve block (SNB) plus intravenous lidocaine infusion (IV Lido) would be as effective as CNBs. Furthermore, since recently, the use of facial plane blocks are increasingly advocated, we compared CNBs vs SNBs plus IV Lido in patients undergoing erector spinae plane (ESP) and quadratus lumborum (QL) blocks for video assisted thoracic surgery (VATS) and abdominal/retroperitoneal surgeries, respectively.

Patients and Method: Using our IRB approved registry (PRO10120146), we retrospectively reviewed the electronic record of 105 patients, including 51 patients who underwent VATS and received either single injection erector plane block (SESPB) plus IV lido or continuous erector spinae plane block (CESPB), and 54 patients who underwent major abdominal surgery and received either single injection quadratus lumborum block (SQLB) plus IV Lido or continuous quadratus lumborum block (CQLB). Demographics, verbal pain scores (0-10), and opioid consumption (morphine intravenous equivalent; MIVE), all in the context of the same multimodal approach (acetaminophen, ketamine, dexmedetomidine, and ketorolac) were collected at 24, 48, and 72 hours after surgery. Alpha was set to 0.05.

Results: SNBs plus IV Lido were as effective as CNBs regarding pain control and total MIVE at 24, 48, or 72 hours after surgery. Subgroup analysis indicated similar findings were observed in patients who underwent VATS or major abdominal surgery.

Conclusions: This analysis suggests that SNBs plus IV Lido are as effective as CNBs for perioperative pain management when ESP or QL blocks are used for VATS or major abdominal surgery, respectively.

Keywords: Regional anesthesia, quadratus lumborum block, erector spinae plane block, acute pain

Introduction

A continuous nerve block (CNB) is an established approach to provide effective perioperative pain management. One of the recognized benefits of CNBs is that, compared to a single injection nerve block (SNB), CNBs can significantly prolong the duration of analgesia.^{1,2} Despite the prolonged duration of analgesia, there are disadvantages and complications associated with CNBs. Disadvantages include frequent dislodgment of the perineural catheter, leaking of the local anesthetic solution, exposure of the perineural catheters related to the patients' movements. Complications include major hematoma, infection, and failure to provide effective analgesia because of the initial misplacement or secondary dislodgment of the catheter³ and limiting patients' mobility. Furthermore, limitations exist to the use of CNB in patients receiving anticoagulants. Last, in 2009, Centers for Medicare and Medicaid Services (CMS) unbundled the payment for a CNB from the payment for the follow up visit, thus reducing reimbursement. This led to a significant decrease in the number of CNBs being performed, especially in a busy practice, as placement of a CNB takes significantly more time than placement of a SNB and no additional payment was provided.⁴ Developing an alternative approach to CNBs to allow provision of prolonged perioperative analgesia without CNB limitations is of major value.

For several years, our standard protocols to manage perioperative pain for a wedge resection and laparoscopic abdominal surgery included the use of an SNB combined with a continuous infusion of lidocaine at a rate of 50 mg/hr for 48 hrs or a total dose of 2,400 mg. This

technique has been used in our institution in more than 1,000 patients and has been proven to be effective and safe in the context of a multimodal approach to perioperative pain management. In addition, lidocaine infusions have been used safely in abdominal, thoracic, and urologic procedures. A number of authors have confirmed the safety of the postoperative infusion of lidocaine in doses ranging from 3,200mg to 6,000mg over a 48 hour period.⁵⁻¹⁹

In the past few years, the use of interfascial plane blocks has been more and more advocated,²⁰ including erector spinae plane blocks (ESPB) for thoracic surgery,²¹⁻²⁵ pectoralis and serratus blocks for breast surgery²⁶⁻²⁹ and quadratus lumborum (QL) blocks for abdominal surgery,³⁰

Intravenous lidocaine (IV lido) infusions have been demonstrated to be an effective technique for postoperative analgesia in several different types of surgery, including abdominal,³¹⁻³⁴ thoracic,^{35,36} and spine.^{37,38} We hypothesized that SNBs followed by IV lido would be as effective as CNBs. More specifically, single injection erector spinae plane block (SESPB) plus IV Lido would be as effective as continuous erector spinae plane block (CESPB) for video assisted thoracic surgery (VATS), and single injection quadratus lumborum block (SQLB) plus IV Lido would be as effective as continuous quadratus lumborum block (CQLB) for abdominal/retroperitoneal surgery.

Material and methods

Methods

Using our IRB approved registry (PRO10120146), we retrospectively reviewed the electronic records of 105 patients who received regional anesthesia and multimodal pain control from an acute pain service and underwent either unilateral video assisted thoracic surgery

(VATS) or major abdominal surgery (exploratory laparotomy, gastrectomy, hyperthermic intraperitoneal chemotherapy (HIPEC), and cysto-prostatectomy surgery). Each patient included in this analysis underwent surgery using the same enhanced recovery after surgery (ERAS) protocol. For the general anesthetic, the ERAS protocol includes an induction with IV propofol and rocuronium, followed by maintenance anesthesia consisting of 1) propofol, dexmedetomidine 0.2-0.5 mcg/kg/hr, and ketamine 0.2-1 mg/kg/hr IV infusions, 2) intraoperative magnesium 2-4g IV, and 3) rocuronium boluses for muscle relaxation to maintain a train of four (TOF) 2/4 twitches. Opioid medications were strictly avoided intraoperatively. The perioperative pain protocol included acetaminophen 1g PO preoperatively, postoperatively acetaminophen 1g IV or PO every 6 hours according to each patient's ability to tolerate oral medication, ketamine 5-10 mg/hr IV infusion, and ketorolac 15 mg IV every 8 hours for 72 hours starting in the PACU. In addition, patients who underwent VATS received a dexmedetomidine 0.2 mcg/kg/hr IV infusion for the first 24 hours of recovery in the ICU. All patients were eligible to receive postoperative opioid medications on an as needed basis (oxycodone PO, hydromorphone IV). No patients included in this study received opioid based patient-controlled analgesia (PCA).

Prior to surgery, an interfascial plane block was performed while the patient was in the pre-operative area. Each block was performed following 1) the placement of monitors to monitor blood pressure, pulse oximetry, and EKG, 2) the proper positioning of the patient, either the sitting position in the case of a unilateral ESP block for VATS or the lateral decubitus in the case of a bilateral QL2 block for abdominal surgery or a single QL2 block performed for nephrectomy, 3) verifying the patients' consent, allergies, and the laterality of the procedure, and 4) the administration of midazolam IV 1-2 mg and/or Fentanyl IV 50-100 mcg for sedation.

After proper disinfection of the skin, a curvilinear 5 MHz ultrasound probe encased in a sterile sheath and connected to a Sonosite X-Porte® (Fujifilm Sonosite, San Diego, CA) was used to determine the proper position of the nerve block needle. The landmarks for the ESP block included the transverse process of T5 vertebrae and the erector spinae plane situated immediately above the transverse process. Next, in the case of a single injection block, a 90mm 22 Gauge Touhy needle (B-Braun, Bethlehem, PA) was properly positioned under ultrasound guidance using an in-plane approach. For the QL2 block, the probe was first placed anteriorly to identify the external oblique, internal oblique, transversalis abdominis. Then, moved posteriorly to identify the transversalis fascia, and quadratus lumborum muscles. The placement of the needle in the erector spinal plan (ESP) or below the transversalis fascia (QL2) was followed by the injection of 20ml of 0.5% ropivacaine in an incremental fashion with negative aspiration between each incremental dose. In the case of a continuous block, a 20 gauge catheter was placed 5 cm beyond the tip of the 100 mm 18 gauge Touhy needle (B-Braun, Bethlehem, PA). The procedure was repeated on the contralateral side in the case of a single injection or continuous QL2 block performed for major abdominal surgery. In the recovery room, an infusion of 0.25% lidocaine at a rate of 17.5-25 mg/hr for a unilateral infusion and 35-50 mg/hr for bilateral infusion) with additional as needed boluses of 3 ml of 0.25% lidocaine (7.5 mg) every hour as needed, was begun for each continuous peripheral nerve block (2 infusions for patients who received bilateral continuous nerve blocks). These infusions continued for 3 days after surgery. A continuous infusion of IV lido 50 mg/hr was begun in the recovery room for patients who received a single injection nerve block. These infusions were continued for 2 days after surgery.

Statistics

The primary endpoint was opioid consumption at 24 hours postoperatively (reported as morphine IV equivalents or MIE). Opioid consumption included as needed oxycodone PO and Hydromorphone IV. No patients included in this study received opioid based patient-controlled analgesia (PCA). Secondary endpoints included OPS (oral pain scores obtained from the patients using a scale of 1 (no pain) to 10 (worst possible pain) at 24, 48, and 72 hours and opioid consumption at 48 and 72 hours.

Demographic data were reported as mean \pm SD, whereas opioid consumption and pain scores were reported as median (95% interval). A t-test was performed to compare demographics, and a Kruskal-Wallis test was performed to compare opioid and non-opioid consumption, pain scores, and total non-opioid consumption at each point of time. Alpha was set up at 0.05.

Results

The overall database was comprised of 51 patients who underwent video assisted thoracic surgery (VATS) with either continuous erector spinae plane block (CESPB; n=34) or single injection erector spinae block (SESPB) plus IV lidocaine infusion (SESPB; n=17), and 54 patients who underwent major abdominal surgery (exploratory laparotomy, HIPEC, gastrectomy, cysto-prostatectomy, or partial nephrectomy) and received either continuous quadratus lumborum block (CQLB; n=31) or single injection quadratus lumborum block (SQLB; n=23) plus IV lidocaine infusion. Continuous nerve block infusions CESPB & CQLB were administered for 72 hours postoperatively. The study covered a period of 4 months from July 1 to November 1, 2019 and was conducted at The University of Pittsburgh Shadyside Hospital.

Demographics

Overall group and sub-group comparison did not show any statistical difference with respect to age, weight, or gender (Table 1).

Overall continuous nerve blocks (CNB) vs single nerve blocks (SNB) plus IV Lido

1. Pain scores

There were no statistical differences among groups. OPS scores were as follows (Table 2):

- a) 24 hours postoperatively, CNB (5, 0.59) vs SNB + IV Lido (5, 0.75)
- b) 48 hours, CNB (4, 0.61) vs SNB + IV Lido (4, 0.94)
- c) 72 hours, CNB (4, 0.64) vs SNB + IV Lido (4, 0.94)

2. Opioid consumption

There were no statistical differences among groups. MIE were as follows (Table 2):

- a) 24 hours postoperatively, CNB (6.4, 3.29) vs SNB + IV Lido (8, 2.29)
- b) 48 hours, CNB (2.4, 5.92) vs SNB + IV Lido (4, 2.00)
- c) 72 hours, CNB (2, 2.82) vs SNB + IV Lido (2.8, 3.12)

3. Non-opioids analgesics

There were no statistical differences among groups. Total non-narcotic analgesic usage was as follows (Table 3):

- a) Acetaminophen (0-72 hours): acetaminophen was given at scheduled dosing periods for a maximum of 72 hours, CNB (3000, 136.57) vs SNB + IV Lido (3000, 195.29)
- b) Dexmedetomidine (0-24 hours): dexmedetomidine was infused for a maximum of 24 hours postoperatively while the patient was in the ICU following VATS, CNB (181.90, 43.59) vs SNB + IV Lido (99.99, 57.92).

c) Ketamine (0-72 hours): ketamine was infused postoperatively for a maximum of 72 hours postoperatively, CNB (87.82, 8.51) vs SNB + IV Lido (78.53, 19.41)

d) Ketorolac (0-72 hours): ketorolac was given at scheduled dosing periods for a maximum of 72 hours, CNB (30.00, 4.39) vs SNB + IV Lido (30.00, 6.07)

Thoracic Surgery – CESPb vs SESPb + IV Lido

1. Pain scores

There were no statistical differences among groups. VAS scores were as follows (Table 2):

- a. 24 hours postoperatively, CNB (5, 0.85) vs SNB + IV Lido (4, 1.01)
- b. 48 hours, CNB (4.5, 0.84) vs SNB + IV Lido (4, 2.05)
- c. 72 hours, CNB (5, 1.12) vs SNB + IV Lido (4, 2.31)

2. Opioid consumption

There were no statistical differences among groups. MIE were as follows (Table 2):

- a. 24 hours postoperatively, CNB (4.4, 5.47) vs SNB + IV Lido (7.6, 2.52)
- b. 48 hours, CNB (1, 2.16) vs SNB + IV Lido (4, 2.46)
- c. 72 hours, CNB (2, 2.46) vs SNB + IV Lido (4.2, 7.74)

3. Non-opioids analgesics

There were no statistical differences among groups. Total non-narcotic analgesic usage was as follows (Table 3):

- a. Acetaminophen (0-72 hours): acetaminophen was given at scheduled dosing periods for max 72 hours, CNB (3000, 197.82) vs SNB + IV Lido (3000, 322.27)
- b. Dexmedetomidine (0-24 hours): dexmedetomidine was infused for a maximum of 24 hours postoperatively while the patient was in the ICU following VATS, CNB (184.49, 44.46) vs SNB + IV Lido (99.99, 57.92)

- c. Ketamine (0-72 hours): ketamine was infused postoperatively for a maximum of 72 hours postoperatively, CNB (84.86, 10.13) vs SNB + IV Lido (78.53, 13.64)
- d. Ketorolac (0-72 hours): ketorolac was given at scheduled dosing periods for a maximum of 72 hours, CNB (30, 5.60) vs SNB + IV Lido (30, 7.49)

Abdominal Surgery – CQLB vs SQLB + IV Lido

1. Pain scores

There were no statistical differences among groups. OPS were as follows (Table 2):

- a. 24 hours postoperatively, CNB (5, 0.85) vs SNB + IV Lido (5, 1.13)
- b. 48 hours, CNB (4, 0.93) vs SNB + IV Lido (4, 1.07)
- c. 72 hours, CNB (4, 0.80) vs SNB + IV Lido (4, 1.09)

2. Opioid consumption

There were no statistical differences among groups. MIE were as follows (Table 2):

- a. 24 hours postoperatively, CNB (9.2, 3.72) vs SNB + IV Lido (8.8, 3.63)
- b. 48 hours, CNB (7.6, 11.72) vs SNB + IV Lido (2.4, 2.96)
- c. 72 hours, CNB (4, 4.77) vs SNB + IV Lido (2, 3.57)

3. Non-opioids analgesics

There were no statistical differences among groups. Total non-narcotic analgesic usage was as follows (Table 3):

- a. Acetaminophen (0-72 hours): acetaminophen was given at scheduled dosing periods for a maximum of 72 hours, CNB (3000, 191.50) vs SNB + IV Lido (3000, 250.66)
- b. Dexmedetomidine (0-24 hours): dexmedetomidine was infused for a maximum of 24 hours postoperatively while the patient was in the ICU following VATS, CNB (91.01, 135.12) vs SNB + IV Lido (0)

- c. Ketamine (0-72 hours): ketamine was infused postoperatively for a maximum of 72 hours postoperatively, CNB (104.90, 12.48) vs SNB + IV Lido (81.39, 81.39)
- d. Ketorolac (0-72 hours): ketorolac was given at scheduled dosing periods for a maximum of 72 hours, CNB (15, 4.99) vs SNB + IV Lido (30, 10.86)

Discussion

The use of intravenous lidocaine infusions for analgesia has been advocated since 1951.³⁹ Although, it is possible to find studies published on the topic in the 80s, the recent development of laparoscopic surgery has led to a renewed interest in the technique. The average dose of lidocaine reported in the case of an infusion of 24hrs and 48hrs is 2,894 mg and 4,678 mg, respectively. Cassuto et al, 1985⁴⁰ advocated the use of 3,460 mg for 24 hrs in an article titled “Inhibition of postoperative pain by continuous low dose intravenous infusion of lidocaine”. Our protocol is based on the use of 2,400 mg, which is less than most authors reported for both 24 and 48 hrs of infusions. The chosen dose is based on our previous experience with the use of the same protocol in patients undergoing procedures such as lung wedge and abdominal surgery.

Our data support that, in the context of a multimodal approach to perioperative pain following either VATS or major open abdominal surgery, the use of a single injection nerve block followed by a continuous infusion of IV lidocaine at 50 mg/hr provides similar perioperative analgesia than a continuous ESP or QL plane block in patients undergoing VATS and major abdominal surgery, respectively.

Recent reviews provide evidence supporting the use of continuous intravenous infusions of lidocaine for postoperative analgesia following several different types of surgery, including

abdominal, VATS, and spine.³¹⁻⁴⁰ Our study demonstrated that a continuous intravenous infusion of lidocaine started in the recovery room after a single injection nerve block performed prior to surgery is as effective as a continuous nerve block. Both techniques include a preoperative injection of local anesthetic into a fascial plane that is followed by a continuous local anesthetic infusion started in the recovery room, administered either intravenously or through the nerve block catheter.

Although only a limited number of studies have compared fascial plane nerve blocks to more established approaches, such as paravertebral blocks and/or lumbar plexus blocks, the use of ultrasound-guided fascial plane blocks, including ESP and QL blocks, have gained popularity. Fascial plane blocks offer the advantage of reduced risks for complications, such as pneumothorax and bleeding. The associated risk of major bleeding is reduced, as local anesthetic is injected into a fascial plane, far away from the major vessels that travel near large nerves. The risk of pneumothorax is reduced, as the ESP is found superficial to the transverse process, which is easily identified and located centimeters superficial to the pleura. When comparing single injection nerve blocks to continuous nerve blocks, there is no risk of intravascular catheter migration in a single injection nerve block, as there is no catheter. This makes our model not only relevant but practical as well.

Lidocaine's half-life is estimated to be between 90-120 min. In the absence of an initial bolus, reaching steady state requires between 7.5 and 12 hours of infusion. The administration of an initial bolus, although of theoretical value to reach steady state more rapidly, would not be of value in the context of surgery, as it would require cardiac monitoring, which is not always available prior to surgery. Under these conditions, the performance of a single injection nerve

block allows for the provision of an analgesic bridge prior to a lidocaine infusion reaching a steady state concentration.

Evidence supports the concept that preemptive analgesia can significantly reduce postoperative pain and opioid consumption when compared to beginning pain management either intra-operatively or postoperatively. This is best achieved by performing a regional nerve block prior to surgery. Unfortunately, the skills and resources required to perform a nerve block are a finite entity. A single injection nerve block requires much less time and resources when compared to a continuous technique. SNBs have several advantages when compared to CNBs: their performance requires less time, there is a lower risk of bleeding or infections, there is no risk of perineural catheter dislodgment or migration, and they require a lower overall cost.

Although the main mechanism of action of lidocaine is blocking voltage-gated sodium channels (VGSC/NaVs), it has been demonstrated that lidocaine can also reduce the peak currents of sodium channels and accelerate the deactivation process to reduce the excitability of sensory fibers, leading to a reduction of pain perception.²⁴ At the level of the central nervous system, lidocaine has been reported to control the presynaptic release of glutamate from presynaptic terminals of the spinal substantia gelatinosa neuron. It has also been reported to interfere with the G-couple proteins of the GABA, substance P, and neurokinin-1 receptor.³⁹

Our findings need to be confirmed using a prospective randomized design, which is currently in progress, but our data are especially relevant in modern healthcare systems where increasing emphasis is placed upon quality, efficiency, and appropriate utilization of resources.

The limitation of this study includes the fact it is a retrospective analysis. However, in each case, the surgery was performed under the same ERAS protocol. Additionally, patients

were not randomized to each treatment arm, which can only be achieved using a prospective, randomized design.

Conclusion

This retrospective analysis suggests that a single injection nerve block followed by a continuous IV lidocaine infusion may be as effective as a continuous nerve block in patients undergoing major abdominal surgery and VATS. A prospective randomized study is required to confirm these findings.

Acknowledgments

None.

Disclosure

The author reports no conflicts of interest in this work.

References

1. Chelly, JE, Ghisi, D, Fanelli A. Continuous Peripheral Nerve Blocks in Acute Pain Management. *Br J Anaeth*, 2010; 105(S1):i86-i96
2. Bingham AE, Fu R, Horn JL, Abrahams MS. Continuous peripheral nerve block compared with single-injection peripheral nerve block: a systematic review and meta-analysis of randomized controlled trials. *Reg Anesth Pain Med*, 2012;37(6):583-94.
3. Joshi G, Gandhi K, Shah N, Gadsden J, Corman SL. Peripheral nerve blocks in the management of postoperative pain: challenges and opportunities. *J Clin Anesth*, 2016;35:524-29.
4. Chelly JE, Horne JL, Hudson ME, Williams JP. Factors impacting the on time transfer to the operating room in patients undergoing peripheral nerve blocks in the preoperative area. *J Clin Anesth* 2010, 22: 115-21
5. Dale GJ, Phillips S, Falk GL. The analgesic efficacy of intravenous lidocaine infusion after laparoscopic fundoplication: a prospective, randomized, double-blind, placebo-controlled trial. *Local and Regional Anesthesia* 2016; 9: 87-93.
6. Kim HO, Lee SR, Choi WJ, Kim H. Early oral feeding following laparoscopic colorectal cancer surgery. *ANZ Journal of Surgery* 2014; 84: 539-44
7. Cassuto J, Wallin G, Högström S, Faxén A, Rimbäck G. Inhibition of postoperative pain by continuous low-dose intravenous infusion of lidocaine. *Anesth Analg* 1985;64: 971-4.
8. Grady MV, Mascha E, Sessler DI, Kurz A. The effect of perioperative intravenous lidocaine and ketamine on recovery after abdominal hysterectomy. *Anesth Analg* 2012;115:1078-84.

9. Kaba A, Laurent SR, Detroz BJ, Sessler DI, Durieux ME, Lamy ML, Intravenous lidocaine infusion facilitates acute rehabilitation after laparoscopic colectomy. *Anesthesiology* 2007;106:11–8.
10. Kim HJ, Kim WH, Kim G, Kim E, Park MH, Shin BS, et al. A comparison among infusion of lidocaine and dexmedetomidine alone and in combination in subjects undergoing coronary artery bypass graft: a randomized trial. *Contemporary Clinical Trials* 2014;39: 303–9.
11. Rimbäck G, Cassuto J, Tollesson PO. Treatment of postoperative paralytic ileus by intravenous lidocaine infusion. *Anesthesia and Analgesia* 1990;70(4):414–9
12. Striebel HW, Klettke U. Is intravenous lidocaine infusion suitable for postoperative pain management?. *Schmerz* 1992;6 :245–50
13. Tikuisis R, Miliauskas P, Samalavicius NE, Zurauskas A, Samalavicius R, Zabulis V. Intravenous lidocaine for post-operative pain relief after hand-assisted laparoscopic colon surgery: A randomized, placebo-controlled clinical trial. *Techniques in Coloproctology* 2014;18: 373–80
14. Maquoi I, Joris JL, Dresse C, Vandenbosch S, Venneman I, Brichant JF. Transversus abdominis plane block or intravenous lignocaine in open prostate surgery: a randomized controlled trial. *Acta Anaesth Scand* 2016 ;60: 1453–60.
15. Wallin G, Cassuto J, Hogstrom S, Linden I, Faxen A, Rimback G, et al. Effects of lidocaine infusion on the sympathetic response to abdominal surgery. *Anesth Analg* 1987;66:1008–13

16. Wuethrich PY, Romero J, Burkhard FC, Curatolo M. No benefit from perioperative intravenous lidocaine in laparoscopic renal surgery: a randomised, placebo-controlled study. *Europ J Anaesth* 2012 ; 29:537-43.
17. Insler SR, O'Connor M, Samonte AF, Bazaral MG. Lidocaine and the inhibition of postoperative pain in coronary artery bypass patients. *J Card Vasc Anesth* 1995;9(5):541-6
18. Mathew JP, Mackensen GB, Phillips B, Bute B, Grocott HP, Glower DD, Laskowitz DT, et al. Randomized, double-blind, placebo controlled study of neuroprotection with lidocaine in cardiac surgery. *Stroke* 2009;40(3):880-7.
19. Wongyingsinn M, Baldini G, Charlebois P, Liberman S, Stein B, Carli F. Intravenous lidocaine versus thoracic epidural analgesia: a randomized controlled trial in patients undergoing laparoscopic colorectal surgery using an enhanced recovery program. *Reg Anes Pain Med* 2011;36):241-8.
20. Jakobsson J, Wickerts L, Forsberg S, Ledin G Transversus abdominal plane (TAP) block for postoperative pain management: a review. *F1000Res*, 2015 Nov 26;4:F1000 Faculty Rev-1359. doi: 10.12688/f1000research.7015.1.
21. Mallan D, Sharan S, Saxena S, Singh TK, Faisal . Anesthetic techniques: focus on transversus abdominis plane (TAP) blocks. *Local Reg Anesth*, 2019 Sep 5;12:81-88. doi: 200.2147/LRA.S138537.
22. Jones JH, Aldwinckle R Interfascial Plane Blocks and Laparoscopic Abdominal Surgery: A Narrative Review. *Local Reg Anesth*, 2020 Oct 23;13:159-169. doi: 12.2147/LRA.S272694.
23. Helander EM, Webb MP, Kendrick J, Montet T, Kaye AJ, Cornett EM, Kaye AD

- PECS, serratus plane, erector spinae, and paravertebral blocks: A comprehensive review.
Best Pract Res Clin Anaesth, 2019 ;33(4):573-581. doi: 10.1016/j.bpa.2019.07.003.
24. George RM, Wilson SH. Serratus plane blocks: not quite plane and simple.
Reg Anesth Pain Med, 2019 2018-100338. doi: 10.1136/rapm-2018-100338
25. Bang S, Chung K, Chung J et al. The erector spinae plane block for effective analgesia after lung lobectomy: Three cases report. Medicine (Baltimore), 2019 Jul;98(29):e16262.
26. Thomas M, Philip FA, Mathew AP, Jagathnath Krishna KM. Intraoperative pectoral nerve block (Pec) for breast cancer surgery: A randomized controlled trial. J Anaesth Clin Pharm. 2018 ; 34:318-23.
27. Franco CD, Inozemtsev K . Refining a great idea: the consolidation of PECS I, PECS II and serratus blocks into a single thoracic fascial plane block, the SAP block. Reg Anesth Pain Med, 2019 Sep 26:rapm-2019-100745. doi: 10.1136/rapm-2019-100745
28. Homsy C, Lindsey JT. Ann . Regional Anesthetic Blocks in Plastic Surgery Using Portable Ultrasound: A Simplified Approach. Plast Surg, 2019 ;82(6S Suppl 5):S374-S379. doi: 10.1097/SAP.0000000000001805
29. Tsui BCH, Fonseca A, Munshey F. The erector spinae plane (ESP) block: A pooled review of 242 cases. J Clin Anesth, 2019 ;53:29-34.
30. Dewinter G, Coppens S, Van de Velde M, D'Hoore A, Wolthuis A, Cuypers E, Rex S. Quadratus Lumborum Block Versus Perioperative Intravenous Lidocaine for Postoperative Pain Control in Patients Undergoing Laparoscopic Colorectal Surgery: A Prospective, Randomized, Double-blind Controlled Clinical Trial. Ann Surg, 2018; 268:769-75
31. Mccarthy GC, Megalla SA, Habib AS. Impact of intravenous lidocaine infusion on

- postoperative analgesia and recovery from surgery: a systematic review of randomized controlled trials. *Drugs*, 2010;70:1149-63.
32. Kranke P, Jokinen J, Pace NL, Schnabel A, Hollmann MW, Hahnenkamp K, Eberhart LH, Poepping DM, Weibel S. Continuous intravenous perioperative lidocaine infusion for postoperative pain and recovery. *Cochrane Database Syst Rev*, 2015 Jul 16;(7):CD009642. doi: 10.1002/14651858.CD009642.pub2
33. Dunn LK, Durieux ME. Perioperative Use of Intravenous Lidocaine. *Anesthesiology*, 2017;126:729-37.
34. Beaussier M, Delbos A, Maurice-szamburski A, Ecoffey C, Mercadal L. Perioperative Use of Intravenous Lidocaine. *Drugs*, 2018;78:1229-46
35. Weibel S, Jelting Y, Pace NL et al. Continuous intravenous perioperative lidocaine infusion for postoperative pain and recovery in adults. *Cochrane Database Syst Rev*, 2018 ;6: CD0009642. doi: 10.1002/14651858.CD009642.pub3
36. Slovack M, Taylor B, Bryce R et al. Does intravenous lidocaine infusion during video-assisted thoracoscopic surgery reduce postoperative analgesia? A randomized controlled study. *Can J Anaesth*, 2015 Jun;62: 676-7.
37. Ibrahim A, Aly M, Farrag W . Effect of intravenous lidocaine infusion on long-term postoperative pain after spinal fusion surgery. *Medicine (Baltimore)*, 2018 Mar; 97(13): e0229. Published online 2018 Mar 30. doi: 10.1097/MD.00000000000010229
38. Kurabe M, Furue H, Kohno T. Intravenous administration of lidocaine directly acts on spinal dorsal horn and produces analgesic effect: An in vivo patch-clamp analysis. *Sci Rep*, 2016; 6: 26253. Published online 2016 May 18. doi: 10.1038/srep26253

39. Gilbert CRA, Hanson IR, Brown AB, Hingson RA. Intravenous use of xylocaine. *Anesth Analg* 1951;30:301-13.
40. Cassuto J, Wallin G, Högström S, Faxén A, Rimbäck G. Inhibition of postoperative pain by continuous low-dose intravenous infusion of lidocaine. *Anesthesia and Analgesia* 1985;64: 971-4.
41. Xi Y Xinchuan W, Yi M, Qian L, Jin L. A review of the mechanism of the central analgesic effect of lidocaine. A review of the mechanism of the central analgesic effect of lidocaine. *Medicine (Baltimore)*, 2020; 99: e19898. Published online 2020 Apr 24. doi: 10.1097/MD.00000000000019898

Table 1 Demographic characteristics of patients (mean \pm SD)

	Age (mean \pm SD)	Weight (kg) (mean \pm SD)	Gender M/F %
CESFB, SESFB, CQLB, SQLB	64.15 \pm 11.49	83.85 \pm 21.33	48.57 / 51.43
CESPB & CQLB	65.00 \pm 9.76	85.73 \pm 21.13	47.69 / 52.31
SESPB & SQLB + IV Lido	62.78 \pm 13.87	80.78 \pm 21.58	50.00 / 50.00
CESPB (Thoracic)	65.71 \pm 9.08	82.91 \pm 21.54	41.18 / 58.82
SESPB + IV Lido (Thoracic)	59.35 \pm 15.39	78.53 \pm 22.43	35.29 / 64.71
CQLB (Abdominal)	64.23 \pm 10.55	88.84 \pm 20.56	54.84 / 45.16
SQLB + IV Lido (Abdominal)	65.30 \pm 12.37	82.43 \pm 21.27	60.87 / 39.13

Abbreviations: CESPB, continuous Erector Spinae block; SESPB, single shot Erector Spinae block; CQLB, continuous Quadratus Lumborum block; SQLB, single shot continuous Quadratus Lumborum block; IV Lido, Intravenous Lidocaine infusion; CNB, continuous nerve block; SNB, single shot nerve block. Overall Data represented by CESPB, SESPB, CQLB, SQLB. CNBs represented by CESPB & CQLB. SNBs represented by SESPB + IV Lido. Thoracic represents VATS surgery. Abdominal represents major abdominal surgery.

Table 2 Comparison of pain scores and opiate consumption (MIVE) between groups (median, CI)

	Pain Scores 24 hrs	Pain Scores 48 hrs	Pain Scores 72 hrs	MIVE 0-24 hrs	MIVE 24-48 hrs	MIVE 48-72 hrs
CESPB, SESP, CQLB, SQLB						
Count	105	96	73	105	96	73
Median (CI 95%)	5 (0.45)	4 (0.51)	4 (0.52)	7.2 (2.19)	3.6 (3.75)	2 (2.09)
CESPB & CQLB						
Count	65	60	47	65	60	47
Median (CI 95%)	5 (0.59)	4 (0.61)	4 (0.64)	6.4 (3.29)	2.4 (5.92)	2 (2.82)
SESPB & SQLB + IV Lido						
Count	40	36	26	40	36	26
Median (CI 95%)	5 (0.75)	4 (0.94)	4 (0.94)	8 (2.29)	4 (2.00)	2.8 (3.12)
CESPB (Thoracic)						
Count	34	30	21	34	30	21
Median (CI 95%)	5 (0.85)	4.5 (0.84)	5 (1.21)	4.4 (5.47)	1 (2.16)	2 (2.46)
SESPB + IV Lido (Thoracic)						
Count	17	13	8	17	13	8
Median (CI 95%)	4 (1.01)	4 (2.05)	4 (2.31)	7.6 (2.52)	4 (2.46)	4.2 (7.74)
CQLB (Abdominal)						
Count	31	30	26	31	30	26
Median (CI 95%)	5 (0.85)	4 (0.93)	4 (0.80)	9.2 (3.72)	7.6 (11.72)	4 (4.77)
SQLB + IV Lido (Abdominal)						
Count	23	23	18	23	23	18
Median (CI 95%)	5 (1.13)	4 (1.07)	4 (1.09)	8.8 (3.63)	2.4 (2.96)	2 (3.57)

Abbreviations: CESPB, continuous Erector Spinae block; SESP, single shot Erector Spinae block; CQLB, continuous Quadratus Lumborum block; SQLB, single shot continuous Quadratus Lumborum block; IV Lido, Intravenous Lidocaine infusion; CNB, continuous nerve block; SNB, single shot nerve block. Overall Data represented by CESPB, SESP, CQLB, SQLB. CNBs represented by CESPB & CQLB. SNBs represented by SESP + IV Lido. Thoracic represents VATS surgery. Abdominal represents major abdominal surgery.

Table 3 Comparison of non-opiate analgesics usage between groups (median, CI)

	Acetaminophen (mg) 0-72 hrs	Dexmedetomidine (mcg) 0-72 hrs	Ketamine IV (mg) 0-72 hrs	Ketorolac IV (mg) 0-72 hrs
CESPB, SESP, CQLB, SQLB				
Count	105	18	105	105
Median (CI 95%)	3000 (111.90)	137.41 (36.29)	85.21 (8.26)	30.00 (3.56)
CESPB & CQLB				
Count	65	15	65	40
Median (CI 95%)	3000 (136.57)	181.90 (43.59)	87.82 (8.51)	30.00 (4.39)
SESPB & SQLB + IV Lido				
Count	40	3	40	40
Median (CI 95%)	3000 (195.29)	99.99 (57.92)	78.53 (19.41)	30.00 (6.07)
CESPB (Thoracic)				
Count	34	10	34	34
Median (CI 95%)	3000 (197.82)	184.49 (44.46)	84.86 (10.13)	30.00 (5.60)
SESPB + IV Lido (Thoracic)				
Count	17	3	17	17
Median (CI 95%)	3000 (322.27)	99.99 (57.92)	78.53 (13.64)	30.00 (7.49)
CQLB (Abdominal)				
Count	31	5	31	22
Median (CI 95%)	3000 (191.50)	91.01 (135.12)	104.90 (12.48)	15.00 (4.99)
SQLB + IV Lido (Abdominal)				
Count	23	0	23	19
Median (CI 95%)	3000 (250.66)		81.39 (81.39)	30.00 (10.86)

Abbreviations: CESPB, continuous Erector Spinae block; SESP, single shot Erector Spinae block; CQLB, continuous Quadratus Lumborum block; SQLB, single shot continuous Quadratus Lumborum block; IV Lido, Intravenous Lidocaine infusion; CNB, continuous nerve block; SNB, single shot nerve block. Overall Data represented by CESPB, SESP, CQLB, SQLB. CNBs represented by CESPB & CQLB. SNBs represented by SESP + IV Lido. Thoracic represents VATS surgery. Abdominal represents major abdominal surgery.

