

1 **COVID-19 mortality in the United States: It's been two**
2 **Americas from the start**

3

4 Michael A. Stoto^{1*}¶, Samantha Schlageter^{1¶}, John Kraemer¹

5 ¹ Department of Health Systems Administration, Georgetown University, Washington D.C.,
6 United States of America

7

8 * Corresponding Author

9 E-mail: stotom@georgetown.edu (MAS)

10 ¶These authors contributed equally to this work

11

12 **Abstract**

13 During the summer of 2021, a narrative of "two Americas" emerged: one with high demand for
14 the COVID-19 vaccine and the second with widespread vaccine hesitancy and opposition to
15 masks and vaccines.

16 We analyzed "excess mortality" rates (the difference between total deaths and what would
17 have been expected based on earlier time periods) prepared by the CDC for the United States
18 from January 3, 2020 to September 26, 2021.

19 Between Jan. 3, 2020 and Sept. 26, 2021, there were 895,693 excess deaths associated with
20 COVID-19, 26% more than reported as such. The proportion of deaths estimated by the excess
21 mortality method that were reported as COVID-19 was highest in Northeast (92%) and lowest in
22 the West (72%) and South (76%).

23 Of the estimated deaths, 43% occurred between Oct. 4, 2020 and Feb. 27, 2021. Before May
24 31, 2020, approximately 56% of deaths were in the Northeast, where 17% of the population
25 resides. Subsequently, 48% of deaths were in the South, which makes up 38% of the
26 population. Since May 31, 2020, the South experienced COVID-19 mortality 26% higher than
27 the national rate, whereas the Northeast's rate was 42% lower.

28 If each region had the same mortality rate as the Northeast, more than 316,234 COVID-19
29 deaths between May 31, 2020 and Sept. 26, 2021 were "avoidable." More than half (63%) of
30 the avoidable deaths occurred between May 31, 2020 and February, 2021, and more than half
31 (60%) were in the South.

32 Regional differences in COVID-19 mortality have been strong throughout the pandemic. The
33 South has had higher mortality rates than the rest of the U.S. since May 31, 2020, and
34 experienced 62% of the avoidable deaths. A comprehensive COVID-19 policy, including
35 population-based restrictions as well as vaccines, is needed to control the pandemic.

36 Introduction

37 During the summer of 2021, as vaccine uptake slowed, a narrative of "two Americas"
38 emerged: one with a high demand for the COVID-19 vaccine and the second with high vaccine
39 hesitancy, and later widespread opposition to mask and vaccine mandates. The second
40 America was mostly concentrated in Southern states and rural areas, especially in counties that
41 voted for Donald Trump. Through the summer, the numbers of cases, hospitalizations, and
42 deaths increased dramatically in the second America [1,2]. This narrative shapes not only our
43 understanding of what happened, but also what should, or could, be done to control the
44 ongoing pandemic.

45 However, our analysis of how COVID-19 mortality evolved over time shows that stark
46 regional differences existed from the start of the pandemic, both in cases and deaths as well as
47 testing and vaccine uptake. To see this, we analyzed "excess mortality," the difference
48 between total deaths and what would have been expected based on earlier time periods.
49 Reported cases, hospitalizations, and deaths are known to substantially underestimate actual
50 infections and deaths by a variable fraction depending on testing availability, patient and
51 physician awareness and attitudes, hospital resources, and other factors [3,4]. Since COVID-19
52 awareness and concern, as well as test availability and use, vary markedly throughout the U.S.,
53 excess mortality estimates can help avoid potential bias due to systematic differences in
54 reporting between areas with high and low prevalence areas.

55

56 Methods

57 In order to identify when regional differences in the pandemic emerged, and their
58 temporal association with vaccine uptake, we analyzed the evolution of COVID-19 mortality
59 patterns. Our analysis is based on state-level weekly excess mortality calculations published by
60 CDC [5]. Estimates from other sources might be slightly different, but the differences are not so
61 great as to affect the overall conclusions. In order to avoid reporting lags that potentially vary
62 by region (especially during the summer of 2021 when some states experienced far greater
63 caseloads than others), we concluded the analysis on September 26, 2021.

64 We grouped the states into the four standard Census regions. Other *ad hoc* groupings
65 could exaggerate or minimize differences. The on-line appendix includes figures comparing the
66 per-capita weekly rates of excess mortality to demonstrate the similarity of patterns within
67 each region. This is especially true in the Northeast and South, but there is more internal
68 variation in the West, where California is somewhat of an outlier. Based on a preliminary

69 analysis of the data, we designated five periods with substantially different patterns of excess
70 mortality.

71 Vaccine coverage rates were calculated by Eva Rest of Georgetown University based on
72 a data set compiled by Tiu and colleagues [6].

73

74 **Results**

75 Between Jan. 3, 2020 and Sept. 26, 2021, the excess mortality associated with COVID-19
76 in the U.S. totaled 895,693, a per capita rate of 270 per 100,000 population. During the same
77 period, 710,999 COVID-19 deaths were reported, amounting to 79% of the estimated excess
78 mortality. In other words, during this period the U.S. experienced 26% more COVID-19 deaths
79 than reported as such.

80 As can be seen in Figure 1, over time there are clear differences among the regions,
81 especially between the Northeast and South. Figure 2 shows 43% of the excess deaths
82 occurred between Oct. 4, 2020 and Feb. 27, 2021. Before May 31, 2020 (period 1),
83 approximately 56% of deaths were in the Northeast; subsequently, 48% were in the South.

84

85 **Figure 1. Excess mortality by week and region, U.S., January 3, 2020 – September 26, 2021.**
86 **Vertical lines correspond to May 30, 2020, Oct. 3, 2020, Feb. 27, 2021, and June 25, 2021.**
87 **Source: authors' calculations based on CDC data**

88

89 **Figure 2. Excess mortality by period and region, U.S., January 3, 2020 – September 26, 2021.**
90 **Periods defined as in Table 1. Source: authors' calculations based on CDC data**

91

92 Controlling for the different sizes of the regions and lengths of the periods, Table 1
93 displays excess mortality by region per 100,000 population per day. Before May 31, 2020, the
94 daily mortality rate in the Northeast (0.881 per 100,000) was 3.3 times the national rate while
95 the rate for the South (0.132 per 100,000) was 48% of the national rate. Subsequently, the
96 South experienced COVID-19 mortality 26% higher than the national rate, whereas the
97 Northeast's rate was 42% lower.

98

99 **Table 1. Excess and avoidable mortality by region and time period, U.S., January 3, 2020 –**
 100 **September 26, 2021**

	Period 1 (1/3– 5/30/20)	Period 2 (5/31– 10/3/20)	Period 3 (10/4/20– 2/27/21)	Period 4 (2/28– 6/26/21)	Period 5 (6/27– 9/26/21)	Entire period
Excess mortality by region and period						
Midwest	22,325	23,169	84,049	12,142	23,345	165,030
Northeast	75,143	7,986	51,200	7,613	10,572	152,514
South	24,667	85,031	158,350	30,270	93,712	392,030
West	12,177	36,869	92,891	10,824	33,358	186,119
U.S.	134,312	153,055	386,490	60,849	160,987	895,693
Excess mortality per capita by region and period (per 100,000 population) per day						
Midwest	0.219	0.227	0.823	0.119	0.229	0.381
Northeast	0.881	0.111	0.609	0.112	0.202	0.422
South	0.132	0.539	0.859	0.203	0.816	0.494
West	0.105	0.375	0.810	0.117	0.466	0.377
U.S.	0.274	0.368	0.799	0.156	0.534	0.430
Excess mortality as a proportion of reported COVID-19 deaths						
Midwest	85.1%	61.0%	95.0%	84.7%	61.7%	83.4%
Northeast	82.7%	98.8%	108.7%	121.3%	55.6%	92.3%
South	78.8%	67.1%	84.6%	56.0%	76.2%	76.2%
West	74.9%	57.0%	84.7%	54.1%	56.7%	71.8%
U.S.	81.7%	65.4%	90.1%	69.5%	68.7%	79.4%
Avoidable excess mortality by region and period						
Midwest	0	13,606	22,738	3,026	10,685	50,055
Northeast	0	0	0	0	0	0
South	0	67,528	46,131	13,584	70,541	197,783
West	0	25,975	23,046	439	18,936	68,395
U.S.	0	107,108	91,915	17,048	100,162	316,234
Proportion fully vaccinated by region at the end of the period (% of the entire population)						
Midwest			7.3%	44.4%	52.9%	
Northeast			6.8%	51.9%	60.3%	
South			7.5%	42.9%	49.8%	
West			7.3%	39.8%	48.9%	
U.S.			7.5%	47.7%	56.5%	

101

102 Table 1 also shows that the ratio of estimated to reported COVID-19 deaths varies over
103 time and among the regions. The proportion is lowest in the West (71.8%) and South (76.2%).
104 The proportions were similar in all regions (approximately 82%) in Period 1, but vary markedly
105 afterwards, dropping to less than 60% in the West in some periods. The overall proportion is
106 highest in Northeast (92%), and is greater than 100% in the third and fourth period (109% and
107 121% respectively. This is probably because some people who died of another primary cause
108 had a positive COVID-19 test and were included in the reported counts [7], as called for by the
109 National Center for Health Statistics [8].

110 Because the rates of excess mortality vary so markedly, one can calculate how many
111 deaths could have been avoided if each region had the same rates as the lowest region in each
112 time period. We assume that no deaths were avoidable in the first period, before much was
113 known about treating or preventing COVID-19. Subsequently, we counted as “avoidable” the
114 difference between the rates seen and those in the Northeast, which were the lowest in each
115 period. According to this calculation, 316,234 COVID-19 deaths between May 31, 2020 and
116 Sept. 26, 2021 were "avoidable." Figure 3 shows that more than half of the avoidable deaths
117 (62%) were in the South. More than half (63%) of the avoidable deaths occurred between May
118 31, 2020 and February, 2021, and an additional 36% between June 27 and Sept. 26, 2021.

119

120 **Figure 3. Avoidable mortality by period and region, U.S., January 3, 2020 – September 26,**
121 **2021. Periods defined as in Table 1. Source: authors’ calculations based on CDC data**

122

123 Discussion

124 The theme of "two Americas" arose in the summer of 2021, regarding at first vaccine
125 refusal, and later opposition to vaccine and mask mandates and more generally, covid-
126 denialism [9]. However, our analysis of excess mortality demonstrates that large disparities
127 have existed since the beginning of the pandemic in the U.S. The starkest contrast is between
128 the Northeast and South. As first noted by Woolf and colleagues [10] for 2020, the first wave of
129 the pandemic was highly concentrated in the Northeast, and particularly in the New York
130 metropolitan area. Since May 31, 2020, however, approximately 48% all excess deaths were in
131 the South, which makes up 38% of the population. The disparity was most apparent in the
132 summer of 2020 (May 31 – Oct. 3), when the daily excess mortality rates were 0.539 per
133 100,000 in the South, 0.369 per 100,000 nationally, and 0.111 per 100,000 in the Northeast.
134 Similarly, of the 316,234 avoidable COVID-19 deaths between May 31, 2020 and Sept. 26, 2021,
135 the majority (62%) were in the South.

136 As a sensitivity analysis, we estimated the numbers of deaths that could have been
137 avoided in Period 1 as one-half of the difference between the actual rates and those in the
138 West (which had the lowest excess mortality rate in that period). This increases the number of
139 avoidable deaths to 402,700 an increase of 86,466. With this assumption the proportion of
140 avoidable deaths in the South was still high, 56%.

141 The regional level of analysis is simple enough for disparities to be clearly apparent, but
142 masks more extreme disparities at the state and local level. A *Washington Post* analysis, for
143 instance, demonstrates a strong relationship between the proportion who voted for Donald
144 Trump in 2020 and COVID-19 mortality at the county-level [11]. A second analysis shows a
145 strong correlation at the state level between vaccine uptake and the Trump vote [11]. Thus, the
146 disparities in this analysis are likely underestimates of actual differences at finer levels of
147 geography.

148 Strong resistance to vaccine and mask mandates emerged during the summer of 2021,
149 especially in the South and in states with Republican governors [12,13]. Consistent with this,
150 excess mortality in the summer of 2021 was correlated with vaccine uptake. As seen in Table 1,
151 the Northeast had the lowest COVID-19 mortality rates and the highest vaccination coverage:
152 52% of the population was fully vaccinated on June 27, 2021 and 60% on September 26, 2021.
153 The South, at the other extreme, had the highest COVID-19 mortality rates and lowest vaccine
154 coverage (40% and 49% on the same dates).

155 Nationally, vaccines have already saved many lives [14] and boosters have the potential
156 to save many more [15]. However, our analysis suggests that major differences in COVID-19
157 mortality emerged in the summer of 2020, well before vaccines became available. Indeed, 63%
158 of the avoidable deaths occurred by the end of February, 2021, when the vaccine rollout was
159 just beginning. These avoidable deaths occurred before the Delta variant became dominant in
160 the United States, so Delta is not part of the explanation. Similarly, because the South has had
161 dramatically higher COVID-19 mortality than the Northeast (since June, 2020) during all seasons
162 of the year, weather is not a likely explanation. Northeasters may have carried some natural
163 immunity into the summer of 2020 [16], but by July, 2020, Anand and colleagues estimate
164 substantially higher seroprevalence rates in the South (37.9%) than in the Northeast (17.5%), so
165 natural immunity cannot explain the large differences starting in the summer of 2020 [17].
166 Excess COVID-19 mortality in the South and other areas of the country, therefore, is likely to be
167 due at least in part to higher transmission resulting from differences in mask use, school
168 attendance, social distancing, and other behaviors.

169 Compared to estimates based on epidemiologic models, the excess mortality methods
170 and assumptions are simple and straightforward. While they cannot explain *why* some regions
171 had different mortality rates, excess mortality estimates can accurately document when and
172 where COVID-19 occurred. Although causal patterns are complex and difficult to ascertain,

173 differential implementation of and adherence to stay-at-home orders, mask use, and other
174 non-pharmaceutical interventions seem to be at least a partial explanation for the regional
175 differences in COVID-19 mortality.

176 As we write this in January, 2022, the nation is concerned with a resurgence of cases
177 driven primarily by the Omicron variant. While improving vaccination coverage is clearly
178 important, our analysis demonstrates the potential impact of population-based restrictions as
179 well as vaccines, indicating that a comprehensive COVID-19 policy is needed to control the
180 pandemic.

181 This analysis demonstrates the benefits of the excess mortality approach [18]. This
182 method provides objective estimates of COVID-19's impact on mortality that do not rely on test
183 availability, clinical decisions, and reporting processes that can lead to under - and over -
184 counting. The ratio of estimated to reported COVID-19 deaths ranged from 121% (Northeast in
185 period 4) to 54% in the West and 56% in the South in the same period. An analysis of reported
186 COVID -19 deaths, therefore would have shown both fewer avoidable deaths and regional
187 disparities. The presence of reporting delays, which can be greater in some areas than others,
188 means that excess mortality estimates and differentials are not reliable for a period of weeks,
189 so excess mortality estimates are limited as real-time surveillance tools. And the reliability of
190 these methods will decline as we go further beyond the base period. However, over the period
191 studies, they can be quite informative.

192 Although the causes are not fully understood, COVID-19 clearly has played out
193 differently across the country over the nearly two years since it emerged. The analysis
194 therefore illustrates how the importance of going beyond cumulative case counts for the U.S. as
195 a whole that dominate the news cycle. Beyond the regional differences that are the focus of
196 this paper, there are substantial differences between rural and urban areas with states and
197 among socio-demographic groups. Because they are less sensitive to differences in reporting
198 patterns than case counts, excess mortality methods can be especially useful in understanding
199 these patterns [19].

200

201 **References**

- 202 1. Baumgaertner E. Here's how the most- and least-vaccinated states fared against the Delta
203 variant. Los Angeles Times [Internet]. 2021 Sep 20 [cited 2022 Jan 9]; Available from:
204 [https://www.latimes.com/world-nation/story/2021-09-20/heres-how-the-most-and-least-](https://www.latimes.com/world-nation/story/2021-09-20/heres-how-the-most-and-least-vaccinated-states-fared-against-the-delta-variant)
205 [vaccinated-states-fared-against-the-delta-variant](https://www.latimes.com/world-nation/story/2021-09-20/heres-how-the-most-and-least-vaccinated-states-fared-against-the-delta-variant)


- 206 2. Leonhardt D. U.S. Covid Deaths Get Even Redder. The New York Times [Internet]. 2021
207 Nov 8 [cited 2022 Jan 9]; Available from:
208 <https://www.nytimes.com/2021/11/08/briefing/covid-death-toll-red-america.html>
- 209 3. National Academies of Sciences, Engineering, and Medicine. Evaluating Data Types: A
210 Guide for Decision Makers using Data to Understand the Extent and Spread of COVID-19
211 [Internet]. Washington, DC: The National Academies Press; 2020 [cited 2022 Jan 9]. 15 p.
212 Available from: [https://www.nap.edu/catalog/25826/evaluating-data-types-a-guide-for-](https://www.nap.edu/catalog/25826/evaluating-data-types-a-guide-for-decision-makers-using-datai)
213 [decision-makers-using-datai](https://www.nap.edu/catalog/25826/evaluating-data-types-a-guide-for-decision-makers-using-datai)
- 214 4. Fricker Jr RD. Covid-19: One year on.... Significance. 2021 Feb 3;18(1):12–5.
- 215 5. Centers for Disease Control and Prevention. Excess Deaths Associated with COVID-19
216 [Internet]. 2022 [cited 2022 Jan 9]. Available from:
217 https://www.cdc.gov/nchs/nvss/vsrr/covid19/excess_deaths.htm
- 218 6. Tiu A, Susswein Z, Merritt A, Bansal S. Characterizing the spatiotemporal heterogeneity of
219 the COVID-19 vaccination landscape. MedRxiv: 2021.10.04.21263345 [Preprint]. 2021
220 [cited 2022 Jan 9]. Available from:
221 <https://www.medrxiv.org/content/10.1101/2021.10.04.21263345v2>
- 222 7. Ackley CA, Lundberg DJ, Elo IT, Preston SH, Stokes AC. County-Level Estimates of Excess
223 Mortality Associated with COVID-19 in the United States. MedRxiv [Preprint]. 2021
224 medRxiv 2021.04.23.21255564 [posted 2021 Apr 25; revised 2021 May 5; cited 2022 Jan
225 9]. Available from: <https://www.medrxiv.org/content/10.1101/2021.04.23.21255564v3>
- 226 8. National Vital Statistics System. Guidance for Certifying Deaths Due to Coronavirus Disease
227 2019 (COVID–19). Vital Stat Report Guide [Internet]. 2020 Apr;3. Available from:
228 <https://www.cdc.gov/nchs/data/nvss/vsrg/vsrg03-508.pdf>
- 229 9. Rubin J. We’re becoming two Americas: One healthy, one deliberately at risk. Washington
230 Post [Internet]. 2021 Jul 19 [cited 2022 Jan 9]; Available from:
231 [https://www.washingtonpost.com/opinions/2021/07/19/two-americans-one-healthy-](https://www.washingtonpost.com/opinions/2021/07/19/two-americans-one-healthy-one-deliberately-risk/)
232 [one-deliberately-risk/](https://www.washingtonpost.com/opinions/2021/07/19/two-americans-one-healthy-one-deliberately-risk/)
- 233 10. Woolf SH, Chapman DA, Sabo RT, Zimmerman EB. Excess Deaths From COVID-19 and
234 Other Causes in the US, March 1, 2020, to January 2, 2021. JAMA. 2021 Apr 2;
- 235 11. Bump P. What if, actually, vaccine advocates want people *not* to die? Washington Post
236 [Internet]. 2021 Sep 20 [cited 2022 Jan 9]; Available from:
237 [https://www.washingtonpost.com/politics/2021/09/20/what-if-actually-vaccine-](https://www.washingtonpost.com/politics/2021/09/20/what-if-actually-vaccine-advocates-want-people-not-die/)
238 [advocates-want-people-not-die/](https://www.washingtonpost.com/politics/2021/09/20/what-if-actually-vaccine-advocates-want-people-not-die/)
- 239 12. Krugman P. How Covid Became a Red-State Crisis. The New York Times [Internet]. 2021 Jul
240 29 [cited 2022 Jan 9]; Available from:
241 <https://www.nytimes.com/2021/07/29/opinion/covid-vaccinations-republicans.html>

- 242 13. Jarvie J. One nation, divisible — over vaccines. Los Angeles Times [Internet]. 2021 [cited
243 2022 Jan 9]; Available from:
244 [https://enewspaper.latimes.com/infinity/article_share.aspx?guid=34b29cc1-f8bf-4da9-
245 82b8-282102286176](https://enewspaper.latimes.com/infinity/article_share.aspx?guid=34b29cc1-f8bf-4da9-82b8-282102286176)
- 246 14. Schneider E, Shah A, Sah P, Moghadas S, Vilches T, Galvani A. The U.S. COVID-19
247 Vaccination Program at One Year: How Many Deaths and Hospitalizations Were Averted?.
248 The Commonwealth Fund [Internet]. 2021 Dec [cited 2022 Jan 9]. Available from:
249 [https://www.commonwealthfund.org/publications/issue-briefs/2021/dec/us-covid-19-
250 vaccination-program-one-year-how-many-deaths-and](https://www.commonwealthfund.org/publications/issue-briefs/2021/dec/us-covid-19-vaccination-program-one-year-how-many-deaths-and)
- 251 15. Schneider E, Shah A, Sah P, Moghadas S, Vilches T, Galvani A. Responding to Omicron:
252 Aggressively Increasing Booster Vaccinations Now Could Prevent Many Hospitalizations
253 and Deaths. Commonwealth Fund [Internet]. 2022 Jan [cited 2022 Jan 9]. Available from:
254 <https://www.commonwealthfund.org/blog/2022/responding-omicron>
- 255 16. Havers FP, Reed C, Lim T, Montgomery JM, Klena JD, Hall AJ, et al. Seroprevalence of
256 Antibodies to SARS-CoV-2 in 10 Sites in the United States, March 23-May 12, 2020. JAMA
257 Intern Med. 2020 Jul 21; 180(12):1576-1586.
- 258 17. Anand S, Montez-Rath M, Han J, Bozeman J, Kerschmann R, Beyer P, et al. Prevalence of
259 SARS-CoV-2 antibodies in a large nationwide sample of patients on dialysis in the USA: a
260 cross-sectional study. Lancet Lond Engl. 2020 Oct 24;396(10259):1335–44.
- 261 18. Ritchie H, Mathieu E, Rodés-Guirao L, Appel C, Giattino C, Ortiz-Ospina E, et al.
262 Coronavirus Pandemic (COVID-19). Our World Data [Internet]. 2020 Mar 5 [cited 2022 Jan
263 9]; Available from: <https://ourworldindata.org/coronavirus>
- 264 19. Stoto MA, Rothwell C, Lichtveld M, Wynia MK. A National Framework to Improve
265 Mortality, Morbidity, and Disparities Data for COVID-19 and Other Large-Scale Disasters.
266 Am J Public Health. 2021 Jul 1;111(S2):S93–100.


267

268

269



270



271

272




Figure 3




Figure 2




Figure 1