

1 **Reasons underlying the intention to vaccinate children aged 5-11 against COVID-19: A**
2 **cross-sectional study of parents in Israel, November 2021**

3
4 **Authors:**

5 Nicole G. Morozov¹, Amiel A. Dror^{2,3*}, Amani Daoud^{2,3}; Netanel Eisenbach^{2,3}; Edward
6 Kaykov^{3,4}; Masad Barhoum⁵; Tsvi Sheleg⁵; Eyal Sela^{2,3}; Michael Edelstein³

7
8 *Morozov and Dror are equal first authors

9
10 ¹Tel Aviv University, Tel Aviv, Israel

11 ²Department of Otolaryngology, Head and Neck Surgery, Galilee Medical Center, Nahariya,
12 Israel

13 ³Azrieli Faculty of Medicine, Bar-Ilan University, Safed, Israel

14 ⁴Geriatric Medicine Department, Galilee Medical Center, Nahariya, Israel

15 ⁵Galilee Medical Center, Nahariya, Israel

16
17 *Corresponding author: Amiel A. Dror, Department of Otolaryngology, Head and Neck Surgery,
18 Galilee Medical Center, Nahariya, Israel; Azrieli Faculty of Medicine, Bar-Ilan University,
19 Safed, Israel Tel: +972544530534; email: amieldror@gmail.com

20

21

22

23

24 **Abstract**

25 Vaccination is a key tool to mitigate impacts of the COVID-19 pandemic. In Israel, COVID-19
26 vaccines became available to adults in December 2020 and to 5–11-year-old children in
27 November 2021. Ahead of the vaccine roll-out in children, we aimed to determine whether
28 surveyed parents intended to vaccinate their children and describe reasons for their intentions.
29 We collected information on parental socio-demographic characteristics, COVID-19 vaccine
30 history, intention to vaccinate their children against COVID-19, and reasons for parental
31 decisions using an anonymous online survey. We identified associations between parental
32 characteristics and plans to vaccinate children using a logistic regression model and described
33 reasons for intentions to vaccinate or not. Parental non-vaccination and having experienced
34 major vaccination side effects were strongly associated with non-intention to vaccinate their
35 children (OR 0.09 and 0.18 respectively, $p < 0.001$). Parents who were younger, lived in the socio-
36 economically deprived periphery, and belonged to the Arab population had lower intentions to
37 vaccinate their children. Reasons for non-intention to vaccinate included concerns about vaccine
38 safety and efficacy (53%, 95%CI 50-56) and the belief that COVID-19 is a mild disease (73%,
39 95%CI 73-79), while a frequent motive for vaccination was the return to normal social and
40 educational life (89%, 95%CI 87-91). Understanding rationales for COVID-19 vaccine rejection
41 or acceptance, as well as parental demographic data, can pave the way for intentional educational
42 campaigns to encourage not only vaccination against COVID-19, but also regular childhood
43 vaccine programming.

44

45 **Keywords:** COVID-19; vaccine hesitancy; childhood vaccines; parental vaccine hesitancy;
46 COVID-19 and children

47

48 **Highlights**

- 49 • Parental intention to vaccinate children aged 5-11 is much lower than vaccine coverage in
50 parental age groups
- 51 • Being unvaccinated and having experienced side effects following vaccination were the
52 greatest negative predictors in parents of intention to vaccinate their children
- 53 • Parents were more likely to accept a COVID-19 vaccine for their children to allow them
54 to return to daily social life and to ensure economic security in the family
- 55 • Parents were more likely to reject a COVID-19 vaccination for health reasons such as
56 safety concerns or due the belief that COVID-19 was a mild disease in children

57

58 **Introduction**

59

60 The rapid, global spread of COVID-19, an infection caused by the SARS-CoV-2 virus, spurred
61 the expeditious development of several vaccines and the eventual implementation of vaccine
62 programs at record speeds. In Israel, COVID-19 vaccines became available in December 2020
63 for adults and eligibility gradually extended to younger ages. Teenagers aged 12-15 became
64 eligible in June 2021 and children aged 5-11 in November 2021. Most high-income countries
65 now offer teenage vaccination, but vaccination of younger children is less common despite
66 evidence of its safety and effectiveness¹. Outside of Israel, the United States' (USA) Centers for
67 Disease Control and Prevention (CDC) recommends vaccination in the 5-11 age group while the
68 European Centre for Disease prevention and Control has also recommended considering
69 universal vaccination in this demographic^{2,3}.

70
71 The short period between the pandemic's onset and vaccine availability has added additional
72 challenges for vaccine acceptance among the public ⁴. Almost two years after the onset of the
73 pandemic and one year after the first administration of COVID-19 vaccines in adults, the USA's
74 Food and Drug Administration (FDA) approved the Pfizer/BioNTech mRNA vaccine for 5- to
75 11-year-old children ^{1, 5}. Studies conducted in the UK, USA, and Canada suggest that many
76 parents remain concerned about the vaccine's side effects and safety, leading parents to be
77 hesitant to administer the COVID-19 vaccine to their children ⁶⁻⁸. While new research has
78 examined the reasons why parents would be motivated or hesitant to vaccinate their children in
79 Israel, ours is one of the first to examine attitudes of parents to children aged 5-11.

80
81 Demographic criteria which have been associated with higher rates of vaccine refusal for
82 children include younger parental age, non-white or minority ethnicity, lower education level,
83 history of vaccination non-compliance to recommended childhood vaccinations, and lower
84 income in countries such as England, Canada, and Brazil ^{7, 9-11}. Parents concerned about future
85 finances were more likely to accept a vaccination for their offspring ¹¹. Larson et al. utilized a
86 four-question survey across emergency departments in 2016 to uncover the differing beliefs
87 towards vaccine safety across 67 countries, finding that vaccine safety skepticism relied on both
88 the specific vaccine type and the country itself ¹². Compared with most high-income countries,
89 Israel typically achieves high coverage for routine childhood vaccines. The 2020 national
90 coverage for the first MMR dose among 24-month-olds was 99% in Israel vs 91% in the United
91 States and in the UK ¹³. Of note, Arab Israelis typically achieve higher coverage for routine
92 childhood vaccines compared with the general population ¹⁴. It is, therefore, important to

93 understand and document attitudes to COVID-19 vaccinations due to their implications for
94 planning public health interventions.

95
96 We aimed to investigate the intentions of surveyed parents in Israel to vaccinate their 5-11-year-
97 old children against COVID-19 and the underlying reasons for their intentions. We sought to
98 determine what common reasons parents listed for accepting or rejecting vaccinating their
99 children. Our survey was initiated two weeks before children in this age group became eligible
100 for COVID-19 vaccination.

101

102 **Methods**

103

104 **Study design**

105

106 This study was designed as a cross sectional survey to understand the willingness of surveyed
107 parents of children 5-11 years old to vaccinate their children against COVID-19 and the reasons
108 for their acceptance or refusal of the vaccine using an online questionnaire in Hebrew and
109 Arabic. The survey was distributed through social media channels (Facebook and Twitter).
110 Participants voluntarily participated in the survey. Eligibility criteria for inclusion in this study
111 included whether the participant is a parent and if the parent has a child between the ages of five
112 and 11. Appendix 1 provides an example in English of the study with the goal of the survey to
113 determine eligibility for inclusion into our study, to obtain demographic data of the participant,
114 and to establish the participant's intention to vaccinate their 5–11-year-old children. Informed
115 consent was requested on the introductory page of the online questionnaire, prior to survey

116 enrollment. The anonymous web-based survey followed the American Association for Public
117 Opinion Research (AAPOR) reporting guidelines¹⁵. The survey was anonymous and no
118 identifiable information such as name, date of birth, or identification was collected. Participants
119 were permitted to terminate participation at will. Approval of this study was granted by the
120 Research Ethics Committee of the Galilee Medical Center. The questionnaire was designed and
121 distributed using Qualtrics (Qualtrics, Provo, UT, United States of America).

122
123 The survey was distributed between November 1-14, 2021, spanning over 14 days and
124 terminating one day before the planned opening of COVID-19 vaccinations to 5-11-year-old
125 children in Israel. The outreach to popular social media channels in Israel was conducted through
126 Facebook, Instagram, and Twitter advertisement platforms. To encourage participation from a
127 wide range of individuals representing Israeli society, we used advertisement sets in both Arabic
128 and Hebrew and tailored to males and females. We used the social media channel's advertising
129 algorithm to target parents of young children. Only respondents who were older than 18 years,
130 were parents of children aged 5-11, lived in Israel, and who completed the survey were included
131 in this study. The survey was distributed evenly in all geographic regions of Israel. The
132 advertisement invitation to participate in our survey included a direct callout with the phrase
133 "Can you help us to understand parent intention to vaccinate children for COVID-19" followed
134 by "Voluntary survey to study parents' intention to vaccinate children aged 5-11 for COVID-
135 19". The impression segmentation and click conversion rate are described in Figure 1.

136

137 **Survey Instrument**

138

139 The survey had two sections: The first section included nine questions about socio-demographic
140 characteristics of the participating parents including parental age, gender, area of residence,
141 household composition, number of children, ethnicity, parental education, parental COVID-19
142 vaccination status, and self-reported side effects from the vaccine (major, mild, or no symptoms),
143 and one question on intention to vaccinate their child(ren) against COVID-19.

144
145 In the second section, responders who indicated they were not intending to vaccinate their
146 children against COVID-19 were asked to state their reasons from a list of predefined
147 explanations, and those who indicated intention to vaccinate were also asked to affirm why from
148 a list of potential reasons. A free text answer was also provided.

149
150 The full questionnaire is available in Appendix 1. The estimated answering time for completing
151 the survey was under two minutes.

152
153 **Statistical analysis**

154
155 We calculated the proportion of parents intending to vaccinate their children together with 95%
156 confidence intervals, overall and in each stratum. We calculated that in order to detect a 5%
157 difference in response between vaccination intenders and non-intenders with an alpha of 0.05
158 and a power of 80%, 1531 participants were required. We compared parents who intended to
159 vaccinate their children to those who did not in terms of all socio-demographic factors recorded
160 in the questionnaire using a univariate logistic regression model. To determine the extent to
161 which each factor was an independent predictor, we conducted a multivariate logistic regression

162 analysis adjusting for all other factors found to be statistically significant (at the $\alpha=0.05$
163 level) in the univariate analysis. We also calculated the proportion of parents giving each reason
164 for intending or not intending to vaccinate together with 95% confidence intervals.

165

166 Data were prepared using Microsoft Excel and analyzed using STATA 15.

167

168 **Results**

169

170 We received 1837 responses to our survey, published online between November 1-14, 2021. Of
171 those, 1514 (82.4%) completed the survey in its entirety. The distribution of responders
172 according to age, sex, ethnicity, education level, and other demographic criteria is described in
173 Table 1. In our study parental self-reported ethnicity was 70.7% Jewish, 25.5% Arab, and 3.8%
174 “other”. In the general population these proportions, as of December 2021 were, 73.9% , 21.1%
175 is Arab, and 5.0% respectively.¹⁶ Overall intention to vaccinate was 44% (95% CI 41.3-45.9).

176

177 Female parents and parents above the age of 35 were more likely to vaccinate than male parents
178 and parents aged 35 and younger (47% vs. 40%, $p<0.002$; 45% vs. 39%, $p<0.02$, Table 1).

179 Compared with non-Arab Israeli parents, Israeli Arab parents reported a lower intention to
180 vaccinate their children (34% vs. 47%, $p<0.001$, Table 1). Other parental factors significantly
181 associated with a lower intention to vaccinate included having a single child, living in the
182 periphery (Northern or Southern, socioeconomically depressed regions of the country), having
183 received fewer than three doses of a COVID-19 vaccine and having experienced major side
184 effects of COVID-19 vaccination. After adjusting for other factors, being male, living in the

185 periphery, belonging to the Arab population, reception of fewer than three doses of the COVID-
186 19 vaccine and having experienced major side effects following COVID-19 vaccination all
187 remained significantly associated with a lower intention to vaccinate children (Table 1). Being
188 completely unvaccinated and having suffered major side effects were associated with the largest
189 reduction in intention to vaccinate children (-91% and -81% respectively, $p < 0.001$).

190
191 Of 801 parents intending to vaccinate their children, 93% were COVID-19 vaccinated at least
192 once. Intention to vaccinate their children increased with the number of doses the parents had
193 received; while only 13% of unvaccinated parents intended to vaccinate their children against
194 COVID-19, 32% of parents vaccinated with one dose, 48% of parents vaccinated with two doses,
195 and 62% of parents vaccinated with three doses were willing to vaccinate their children
196 ($p < 0.001$, Table 1).

197
198 The most often cited reason for choosing to vaccinate their children was to allow their children to
199 return to everyday social, educational, and financial life, followed by protecting other family
200 members from COVID-19 infection (89 and 77.9% respectively, Table 2). The protection of the
201 children against viral COVID-19 infection followed by vaccine safety were the least often
202 mentioned explanations for parents who intended to vaccinate their children (37.9%).

203
204 The most frequently mentioned explanation for not vaccinating their children is the belief that
205 COVID-19 leads to mild or asymptomatic disease in children (76.1%, Table 2). Concerns of the
206 parents regarding vaccine safety were ordered second in their consideration not to vaccinate their
207 children (53.1%, Table 2).

208

209 **Discussion**

210

211 Routine childhood vaccination offers the prospect of protection against infectious diseases. In
212 most settings, there is no ongoing endemic transmission of poliovirus, measles, or diphtheria
213 infections due to effective global vaccination campaigns¹⁷⁻¹⁹. However, COVID-19 is unique in
214 several ways: first, the pandemic is continuing to develop and therefore vaccine protection is
215 against an immediate and tangible threat; second, the volume and availability of information
216 around COVID-19 vaccines, offering arguments both for and against vaccination with very few
217 verification pathways for parents to assess the accuracy of the information they are accessing, is
218 an unprecedented phenomenon; third, the speed at which COVID-19 vaccines were developed have
219 raised some concerns about safety among the general public; fourth, the shifting understanding
220 that the purpose of the COVID-19 vaccines, including mRNA vaccines, is not to prevent
221 infection but rather to decrease the likelihood of severe adverse outcomes^{20,21}.

222

223 Existing research has established several factors which may influence vaccine acceptance,
224 including the type of vaccine, home country of the individual, religiosity, age, and other socio-
225 demographic criteria^{7,9-11}. These links were studied also in association with the global outbreak
226 of the H1N1 influenza pandemic of 2009-2010. According to a World Health Organization
227 systematic review report in 2016 of attitudes towards vaccination during this particular influenza
228 pandemic, parents of children aged 6-59 months were less likely to accept a vaccine for their
229 children if the parents were younger and had a lower educational achievement²². The most cited
230 reasons for hesitancy, during the 2009-10 influenza pandemic, included a decreased perception

231 of benefit from the influenza vaccine, decreased perception of danger from the virus, and
232 difficulties with ease of access to vaccination²². Similarly, we found that the most commonly
233 held reason by surveyed parents to not vaccinate their children is the belief that the disease is
234 mild or asymptomatic in children.

235
236 The COVID-19 pandemic has seen the digital spread of information, misinformation, and
237 disinformation on social media platforms on an unprecedented scale²³. Mistrust in government
238 and international health organizations has been suggested as a potential contributor to hesitancy
239 to follow government and international health guidelines, as fewer than 30% of individuals opted
240 to search for their own government social media publications for vaccine information and
241 guidance²⁴. Demographic data from social media sites shows that the largest user age groups of
242 social media sites such as Facebook, Instagram, and YouTube are ages up to 35 years old²⁵. Our
243 results indicate that parents younger than 35, the greatest social media users, were less likely to
244 vaccinate their children.

245
246 Of additional importance in the discussion of COVID-19 vaccine hesitancy is whether an
247 individual has experienced adverse effects attributed to the vaccine. Parental experience was
248 clearly a key factor in the decision, with unvaccinated, surveyed parents and those who
249 experienced perceived severe side effects very unlikely to vaccinate their children. Vaccine
250 hesitant individuals who experienced an adverse effect attributed to a vaccine did receive the
251 vaccine, indicating a likely initial willingness to be vaccinated but subsequent hesitancy based on
252 experience. Acknowledging these experiences while correcting potential factual misconceptions
253 regarding vaccine safety may be a technique to assuage concerned parents, as suggested by a

254 clinical report from The American Academy of Pediatrics ²⁶. Healthcare providers may be better
255 prepared to repair their trust in vaccines in the future, thus paving the way for parents to have
256 greater trust in vaccines and subsequent willingness to vaccinate their children.

257
258 We found that the intention to vaccinate children aged 5-11 was low, overall. Less than half of
259 surveyed parents intended to vaccinate their 5-11 children, and as of February 2022 just over
260 20% had vaccinated their offspring ²⁷. Even among parents who had received three doses of the
261 vaccine, intention to vaccinate was only 62%. This is consistent with experience from other
262 countries where vaccinated parents did not intend to vaccination their children ²⁸. Data from a
263 separate study, available online but not yet in a peer-reviewed publication, conducted at the same
264 time showed that, similar to our results, only 37% of parents intended to vaccinate their children
265 ²⁹. An Israeli study conducted prior to the initiation of ours found that Israeli parents of children
266 of any age, with a bachelor's degree, were found to have higher rates of vaccination ³⁰. In
267 contrast, we found that higher educational attainment did not reflect higher vaccination levels;
268 the differences in these findings may be related to the age of parents. While the parents in our
269 study had at least one child in the 5-11-year-old range, the parents in Heller et al.'s investigation
270 were not necessarily parents to children in that age bracket.

271
272 Our study has highlighted that the main reasons for intending to vaccinate children were not
273 biomedical. Key factors within Israeli society associated with an increased intention to vaccinate
274 children against COVID-19 include the parent's desire for the child to return to normal social
275 and educational life along with secondary motives such as the parent's desire to maintain job
276 security. Our findings suggest that, for COVID-19, the traditional perception of vaccination

277 benefits such as protection against severe illness has been superseded by indirect benefits such as
278 returning to regular societal life and education institutions, as well as assuring financial resilience
279 for the family. While this finding is not surprising considering the severe disruption to normal
280 life caused by the COVID-19 pandemic, it is not clear whether this perception of the societal
281 utility of vaccines will continue to impact the perception of other vaccines beyond the pandemic.

282
283 The main drivers against intention to vaccinate are more biomedical, encompassing beliefs that
284 COVID-19 causes a mild disease course in children and concerns about vaccine safety. Heller et
285 al. found vaccine hesitancy to be greater among respondents who did not believe COVID-19
286 vaccines are safe, with those who believe it is safe having higher rates of vaccination ³⁰.
287 Interestingly, the results of their study, which was able to demonstrate longitudinal attitudes
288 towards COVID-19 and vaccination, indicated a decreasing fear of COVID-19 and a greater
289 distrust in vaccines ³⁰. Despite that finding, Heller et al. discovered that a greater proportion of
290 respondents who had initially been worried about vaccine safety were vaccinated over the
291 study's time period, a finding which may suggest a change in policy and incentives or education
292 about vaccines ³⁰. These findings have implications for health promotion and communication
293 campaigns; ensuring parents' confidence in the safety of the vaccine and highlighting the
294 sociocultural benefits of vaccination may be key in guiding communities to higher levels of
295 vaccine uptake.

296
297 Groups who may be targets for intersectional outreach include minorities, parents with a lower
298 educational level, and families in lower financial brackets as these parents have been highlighted
299 as more likely to be vaccine hesitant ^{10, 31}. Our research found that Arab parents were less likely

300 to vaccinate their children as compared to other parents, consistent with research finding a higher
301 rate of COVID-19 vaccine hesitancy among minority communities in other countries, as well as
302 in Israel's Arab and Ultra-Orthodox communities ^{7, 32, 33}.

303
304 Strategies to reduce vaccine hesitancy have been discussed in other literature. Financial or other
305 incentivizing techniques for parents have been shown to increase vaccine uptake among their
306 children in certain communities such as in Nicaragua and India ³⁴. A major motivator for vaccine
307 uptake in our study is a desire to return to normal social and educational life in Israel, which is an
308 incentive in and of itself. Use of technology and social marketing has also been shown to
309 increase vaccine acceptance for HPV and MMR vaccines ³⁴. Digital health education
310 interventions specific to parental concerns, such as about vaccine technologies, may be key to
311 targeting certain parental groups.

312
313 The key limitation of our study is inherent to its design and concerns the self -selected nature of
314 the sample.. Parents who do not use social media channels, and therefore would not have been
315 exposed to this study, may not have been appropriately represented. Although use of social
316 media use is widespread in parental age groups in Israel, certain groups, for example the Ultra-
317 Orthodox, may be underrepresent due to lower smartphone and Internet penetration ³⁵. Because
318 the survey is entirely anonymous, comparisons between the sample and the general population
319 are difficult. Nevertheless, the ethnicity distribution of respondents was comparable to the
320 general population. Another potential selection bias is that individuals who have stronger
321 opinions either for or against vaccinating their children for COVID-19 may be more likely to
322 complete a questionnaire about the topic. It is also theoretically possible for individuals to have

323 completed the study more than once given that identifying participant information was not
324 collected. It must be noted that the age, sex, ethnicity, and geographical distribution of
325 participants suggest that all segments of the population were represented.

326
327 It is not clear to what extent our findings are generalizable to other countries. Globally, there are
328 wide discrepancies in vaccine uptake from one country to another with some countries enjoying
329 remarkably low levels of vaccine hesitancy among parents ^{9, 11}. Understanding what factors
330 enable childhood COVID-19 vaccination in countries with high parental intent and transferring
331 this knowledge to other contexts could help countries struggling with COVID-19 childhood
332 vaccination programs. Because a growing number of individuals, including parents, are seeking
333 health and vaccine information online or through peers, the availability of trustworthy sources
334 and early educational campaigns cannot be overstated ³⁶.

335

336 **Conclusion**

337
338 The availability of COVID-19 vaccines and licensing of these vaccinations for use in children
339 aged five to 11 is not a guarantee that parents will inoculate their children. Our study showed that
340 intention to vaccinate 5- to 11-year-old children was low even among fully vaccinated parents,
341 with socio-cultural factors, as well as vaccination experience in parents, being associated with
342 intention to vaccinate their children. Our study showed that indirect benefits of vaccination such
343 as returning to regular societal activities, including educational institutions and personal
344 economic stability, were bigger motivators than direct protection against the virus, while the
345 most common reasons for intending not to vaccinate children against COVID-19 were classical

346 biomedical concerns such disease severity and vaccine safety. These findings are important for
347 health authorities to shape educational and communication campaigns: focusing on the message
348 that vaccines protect against COVID-19 may not be as effective as messages stating that
349 vaccines will allow children to go to school and adults to work. Different messaging campaigns
350 should be implemented and evaluated to test this possibility.

351

352 **Conflicts of interest:** The authors declare no conflicts of interest.

353

354 **Attestation:** All authors attest they meet the ICMJE criteria for authorship

355

356 **Data availability statement:** Data is available upon request from the corresponding author.

357

358 Citations

- 359 1. Walter EB, Talaat KR, Sabharwal C, Gurtman A, Lockhart S, Paulsen GC, et al.
360 Evaluation of the BNT162b2 Covid-19 Vaccine in Children 5 to 11 Years of Age. *New England*
361 *Journal of Medicine* 2021.
- 362 2. Control ECfDPa. Interim public health considerations for COVID-19 vaccination of
363 children aged 5-11 years. Stockholm: European Centre for Disease Prevention and Control,
364 2021.
- 365 3. <https://www.cdc.gov/vaccines/covid-19/planning/children.html> CC-VfC-
366 YOICfDCaPcDAf.
- 367 4. Soares P, Rocha JV, Moniz M, Gama A, Laires PA, Pedro AR, et al. Factors Associated
368 with COVID-19 Vaccine Hesitancy. 2021; 9:300.
- 369 5. Woodworth KR, Moulia D, Collins JP, Hadler SC, Jones JM, Reddy SC, et al. The
370 Advisory Committee on Immunization Practices' Interim Recommendation for Use of Pfizer-
371 BioNTech COVID-19 Vaccine in Children Aged 5-11 Years - United States, November 2021.
372 *MMWR Morb Mortal Wkly Rep* 2021; 70:1579-83.
- 373 6. Ruggiero KM, Wong J, Sweeney CF, Avola A, Auger A, Macaluso M, et al. Parents'
374 Intentions to Vaccinate Their Children Against COVID-19. *Journal of Pediatric Health Care*
375 2021; 35:509-17.
- 376 7. Bell S, Clarke R, Mounier-Jack S, Walker JL, Paterson P. Parents' and guardians' views
377 on the acceptability of a future COVID-19 vaccine: A multi-methods study in England. *Vaccine*
378 2020; 38:7789-98.

- 379 8. Humble RM, Sell H, Dubé E, MacDonald NE, Robinson J, Driedger SM, et al. Canadian
380 parents' perceptions of COVID-19 vaccination and intention to vaccinate their children: Results
381 from a cross-sectional national survey. *Vaccine* 2021; 39:7669-76.
- 382 9. Bagateli LE, Saeki EY, Fadda M, Agostoni C, Marchisio P, Milani GP. COVID-19
383 Vaccine Hesitancy among Parents of Children and Adolescents Living in Brazil. *Vaccines* 2021;
384 9.
- 385 10. McKinnon B, Quach C, Dubé È, Tuong Nguyen C, Zinszer K. Social inequalities in
386 COVID-19 vaccine acceptance and uptake for children and adolescents in Montreal, Canada.
387 *Vaccine* 2021; 39:7140-5.
- 388 11. Goldman RD, Krupik D, Ali S, Mater A, Hall JE, Bone JN, et al. Caregiver Willingness
389 to Vaccinate Their Children against COVID-19 after Adult Vaccine Approval. *International*
390 *Journal of Environmental Research and Public Health* 2021; 18.
- 391 12. Larson HJ, de Figueiredo A, Xiaohong Z, Schulz WS, Verger P, Johnston IG, et al. The
392 State of Vaccine Confidence 2016: Global Insights Through a 67-Country Survey.
393 *EBioMedicine* 2016; 12:295-301.
- 394 13. Organization WH. Measles vaccination coverage World Health Organization
- 395 14. Stein-Zamir C, Israeli A. Timeliness and completeness of routine childhood vaccinations
396 in young children residing in a district with recurrent vaccine-preventable disease outbreaks,
397 Jerusalem, Israel. *Euro surveillance : bulletin European sur les maladies transmissibles =*
398 *European communicable disease bulletin* 2019; 24.
- 399 15. Transparency Initiative Disclosure Elements. American Association for Public Opinion
400 Research, 2021.
- 401 16. Statistics CBo. Population of Israel on the Eve of 2022. In: Center SI, ed., 2021.
- 402 17. Obregón R, Chitnis K, Morry C, Feek W, Bates J, Galway M, et al. Achieving polio
403 eradication: a review of health communication evidence and lessons learned in India and
404 Pakistan. *Bulletin of the World Health Organization* 2009; 87:624-30.
- 405 18. Clarke KEN, MacNeil A, Hadler S, Scott C, Tiwari TSP, Cherian T. Global
406 Epidemiology of Diphtheria, 2000-2017(1). *Emerging infectious diseases* 2019; 25:1834-42.
- 407 19. Holzmann H, Hengel H, Tenbusch M, Doerr HW. Eradication of measles: remaining
408 challenges. *Medical microbiology and immunology* 2016; 205:201-8.
- 409 20. The Lancet Infectious D. The COVID-19 infodemic. *The Lancet Infectious diseases*
410 2020; 20:875.
- 411 21. Salerno L, Craxì L, Amodio E, Lo Coco G. Factors Affecting Hesitancy to mRNA and
412 Viral Vector COVID-19 Vaccines among College Students in Italy. *Vaccines* 2021; 9.
- 413 22. Organization WH. Barriers of influenza vaccination intention and behavior – A
414 systematic review of influenza vaccine hesitancy. World Health Organization, 2016.
- 415 23. Goel A, Gupta L. Social Media in the Times of COVID-19. *Journal of clinical*
416 *rheumatology : practical reports on rheumatic & musculoskeletal diseases* 2020; 26:220-3.
- 417 24. Volkmer I. Social Media and COVID-19: A global study of digital crisis interaction
418 among Gen Z and millennials Melbourne, Australia: Wunderman Thompson, University of
419 Melbourne, Pollfish and the World Health Organization, 2021.
- 420 25. Barnhart B. Social Media Demographics to Inform Your Brand's Strategy in 2022.
421 Sprout Social: Sprout Social, 2022.
- 422 26. Edwards KM, Hackell JM. Countering Vaccine Hesitancy. *Pediatrics* 2016; 138.

- 423 27. Health IMo. Coronavirus in Israel - general situation
424 2022;<https://datadashboard.health.gov.il/COVID-19/general>.
- 425 28. Horiuchi S, Sakamoto H, Abe SK, Shinohara R, Kushima M, Otawa S, et al. Factors of
426 parental COVID-19 vaccine hesitancy: A cross sectional study in Japan. *PloS one* 2021;
427 16:e0261121.
- 428 29. Heller O. What we can learn from Israel's COVID-19 vaccination of children. The
429 Brookings Institution, 2021.
- 430 30. Heller O, Chun Y, Shlomo Y, Gewirtz-Meydan A, Acri M, Kulkarni S, et al. Public
431 perceptions and the willingness to get vaccinated against COVID-19: Lessons from Israel.
432 *Vaccine* 2022; 40:2053-61.
- 433 31. Alfieri NL, Kusma JD, Heard-Garris N, Davis MM, Golbeck E, Barrera L, et al. Parental
434 COVID-19 vaccine hesitancy for children: vulnerability in an urban hotspot. *BMC Public Health*
435 2021; 21:1662.
- 436 32. Robinson E, Jones A, Lesser I, Daly M. International estimates of intended uptake and
437 refusal of COVID-19 vaccines: A rapid systematic review and meta-analysis of large nationally
438 representative samples. *Vaccine* 2021; 39:2024-34.
- 439 33. Gorelik Y, Anis E, Edelstein M. Inequalities in initiation of COVID19 vaccination by age
440 and population group in Israel- December 2020-July 2021. *The Lancet regional health Europe*
441 2022; 12:100234.
- 442 34. Singh P, Dhalaria P, Kashyap S, Soni GK, Nandi P, Ghosh S, et al. Strategies to
443 overcome vaccine hesitancy: a systematic review. *Systematic reviews* 2022; 11:78.
- 444 35. Silver L, Taylor, K. . Smartphone Ownership Is Growing Rapidly Around the World, but
445 Not Always Equally. Pew Research Center, 2019.
- 446 36. Rhodes ME, Sundstrom B, Ritter E, McKeever BW, McKeever R. Preparing for A
447 COVID-19 Vaccine: A Mixed Methods Study of Vaccine Hesitant Parents. *Journal of Health*
448 *Communication* 2020; 25:831-7.
- 449

450




Figure 1: Social media outreach results, including initial post impressions, attrition, exclusion criteria, and conversion rate to clicks.