

Pattern and risk factors of menstrual regulation service use among ever-married women in Bangladesh: evidence from a nationally representative cross-sectional survey

Md. Rashed Alam¹, Md. Nuruzzaman Khan² and Yothin Sawangdee³

¹Department of Population Science and Human Resource Development, University of Rajshahi, Rajshahi Bangladesh. Email: mrasps29@gmail.com

²Department of Population Science, Jatiya Kabi Kazi Nazrul Islam University, Trisal, Mymensingh, Bangladesh. Email: sumonrupop@gmail.com

³Institute for Population and Social Research, Mahidol University, Thailand.
Email: yothin.saw@mahidol.ac.th

Short running title: Menstrual regulation service use among ever-married women in Bangladesh

Corresponding author

Yothin Sawangdee

³Institute for Population and Social Research, Mahidol University, Thailand.

Email: yothin.saw@mahidol.ac.th

Pattern and risk factors of menstrual regulation service use among ever-married women in Bangladesh: evidence from a nationally representative cross-sectional survey

Abstract

Background: Around 47% of the total conceptions in Bangladesh are unintended which leads to several adverse consequences, including maternal and child mortality. Availability of menstrual regulation (MR) service and its use can help women to end conception at an earlier stage, as such, reducing adverse consequences related to the unintended pregnancy. We explored the prevalence and determinants of MR service knowledge and its use among ever-married women in Bangladesh.

Methods: A total of 20 127 ever-married women data from the 2017 Bangladesh Demographic and Health Survey were analyzed. Knowledge about menstrual regulation (MR) and its use were our outcomes of interest. Several individual, household and community-level factors were considered as explanatory variables. The multilevel mixed-effects Poisson regression model was used to determine the factors associated with MR service knowledge and its use in Bangladesh.

Results: Around 71% of the total analyzed women reported they know about MR service while only 7% of them reported they used this service within three years of the survey date. MR service knowledge was found to be higher among women with increased age and education and engaged in income-generating employment. Knowledge about MR service was also found to be higher among women whose husbands were higher educated and engaged in physical work or business. Rural women and women who resided in the community with lower poverty and higher illiteracy were reported lower knowledge of MR service. MR service use was found higher among higher-aged women, women whose husbands were either physical workers or businessmen, women who have an increased number of children and inherent in the community with lower poverty. Lower use of MR service was found

among women who resided in the Chattogram, Khulna, and Mymensingh divisions and women who resided in the community with increased illiteracy.

Conclusion: Use of MR service is very low in Bangladesh. This could be responsible for higher adverse consequences related to unintended pregnancy including higher maternal and child mortality.

Policies and programs are important to aware women of MR.

Keywords: Menstrual regulation, Pattern, Risk, Maternal health, Bangladesh

Introduction

Bangladesh, the 8th largest populated country in the world, has around 20 million unintended conceptions every year that equates to around 47% of the country's total conceptions. Around half of them are ended before live births, mostly through unsafe abortion outside the formal healthcare facilities, as abortion is legally prohibited in Bangladesh unless it is required to save a woman's life. The remaining unintended conception are ended through live births, which contribute to around 25% of the total live births in the year in Bangladesh. Both unsafe abortion and live birth as a result of unintended conception are found to be associated with adverse birth and maternal health outcomes in low- and middle-income countries (LMICs), including Bangladesh (UNICEF 2019 & UN 2021). They include but not limited to pregnancy complications, stillbirths, preterm birth, low birth weight, neonatal, under-five, and maternal mortality. These are challenges to achieving the Sustainable Development Goals (SDGs) 3; health and wellbeing for all; particularly, to achieve the SDGs' targets to reduce neonatal (12 per 1,000), under-five (25 per 1,000) and maternal (73 per 10,000 live births) mortality by 2030 (UN-SDGs Goals 3).

Menstrual regulation (MR), a type of abortion that is performed in a very early stage of pregnancy, has been used in Bangladesh as an alternative of abortion to end pregnancy due to unintended conceptions. The government of Bangladesh in 1979 was approved MR as part of the family planning services and now available in governmental healthcare facilities. It combines manual vacuum aspiration and mifepristone/ misoprostol to regulate the menstrual cycle within 10/12 weeks of conception (Benson et al., 2011; Hossain et al., 2017 & Guttmacher Institute 2017).

Though MR service is available in the governmental healthcare facilities in free of cost or lower cost to access, however, such service use is still very low. The use of MR service is even now declining, alternative to abortion, which use is now increasing (Huda et al. 2010; S. Singh et al., 2017; Guttmarch

Institute, 2012). Consequently, abortion related adverse consequences, including maternal mortality, are very higher, which create a substantial burden to the healthcare system. For instance, abortion related maternal mortality rate was increased 35.5 per 1,000 live births in 2016 (7% of the total maternal mortality) 18 per 1,000 live births in 2010 (1% of the total maternal mortality) (Hossain A et al., 2014; M. Singh et al., 2019; Rana et al., 2019). Further policies and programs are therefore important to increase access of MR services. For this, proper understanding of the current level of knowledge of MR and the factors associated with MR service use are important. However, this understanding is still lacking. So far, only one research is published by using the nationally representative 2014 BDHS data which explored the basic socio-demographic factors associated with the MR service use. This is line of the other studies conducted in LMICs (Holmes K et al, 2021; Chae et al., 2017). However, as far know no research has been conducted yet in Bangladesh to current level of knowledge on MR service use. In addition, age-standardized prevalence knowledge of MR service and MR service use have not yet been investigated.

Therefore, the aims of this study were three-fold: (i) to determine the age-adjusted prevalence of MR service use and knowledge about MR, (ii) to explore the factors associated with MR service knowledge and (iii) to explore the factors associated with MR service use.

Methods

Study design and sample

The data analysed were extracted from the 2017 Bangladesh Demographic Health Survey (BDHS 2017). It is a nationally representative survey conducted in every three years by following a two-stage stratified random sampling method. The National Institute of Population Research and Training (NIPORT) conducted this survey. Developments partners, including the Macro International, ICF international, Mitra and Associates, US Agency for International Development (USAID), and Ministry of Health and Family Welfare, Bangladesh provided technical and financial supervision. In the first

stage of sampling, 675 PSUs were randomly selected from 293 579 PSUs which were generated by the Bangladesh Bureau of Statistics as part of conducting the 2011 National Population Census (BBS, 2014; BDHS, 2017). Data were collected from the 672 PSUs of them, whereas 20 160 households were selected for data collection with 30 households from each selected PSU in the second stage of sampling. Of them data collection was undertaken in 19 457 households with a 96.5% inclusion rate. There were 20 127 ever married women interviewed in from these selected households. Of them, we analysed 14 346 women data knowledge about MR and 1056 women use MR service in the 3 years preceding the survey (Figure. 1).

Outcome variable

Respondents' knowledge on MR and MR service use status within three years of the survey date were our outcomes of interest. Knowledge related data were collected by asking women "*have you ever heard of menstrual regulation (MR)?*". The woman who answered positively to this item was then asked "*have you ever used MR?*". Responses were recorded dichotomously for both questions (yes, no).

Predictor variables

The variables that were found significant for MR knowledge and MR service use were considered as predictor variables. For this, we first conducted an extensive literature search in five databases (PubMed, Embase, CINHALL, Web of Science and Goggle Scholar) and significant variables were listed. Their availability in the dataset we analysed were then checked and significance were tested. Finally, significant variables were kept in our analysis classified under three broad heading following the socio-ecological model of health: individual level factors, household level factors, and community level factors (Bronfenbrenner, U. 1994; Dahlberg & Krug, 2006). Individual level factors were women's age (15-19 years, 20-34 years and ≥ 35 years), education (no education, primary education, secondary education and higher education), age at marriage (below 18 years and 18 years and above) and working status (yes, no). Household level factors were partners' education (no education, primary

education, secondary education and higher education), occupation (agricultural, physical workers, services, business and others), number of children ever born (no child, 1-2 children and >2 children) and wealth quintile (poorest, poorer, middle, richer and richest). Place of residence (urban and rural), region of residence (Barisal, Chattogram, Dhaka, Khulna, Mymensingh, Rajshahi, Rangpur and Sylhet), community level poverty (high poverty, moderate poverty, low poverty and middle to richest community) and community level illiteracy (low, moderate and high) were considered as community level factors.

Statistical Analysis

Descriptive statistics were used to describe the characteristics of the respondents. Age-standardized prevalence of MR service knowledge and its use with their 95% confidence interval (95% CI) were also calculated. We calculated age-standardised estimates to the 2011 Census population of Bangladesh using the direct method $[\sum r_i X P_i / \sum P_i]$, where r_i is the prevalence of MR service knowledge or use in age group i and P_i is the population size in the i th age group. Predictors of MR service knowledge and its use status were determined using the multilevel mixed effect Poisson regression model. The reason of using this model were nested structure of the data we analysed and higher prevalence of MR use where conventional logistics regression model produces incorrect results ((Al-Balushi et al., 2020; Nelder, J. A, 1989). Five models were run separately for each outcome by following the progressive model building technique. Model 1 was null model determined the overall variability of MR service knowledge and its use across the PSUs. Individual level factors were adjusted with model 2. Household level factors were adjusted in model 3. Community level factors were considered in model 4. Model 5 was the final model where all individual, household and community level factors were adjusted. Sampling weight was considered in all analyses. Results were reported as Prevalence Ratio (PR) with

its 95% CI. All analyses were performed using statistical software packages Stata (version 13.0; Stata Corp LP, College Station, Texas).

Ethics statement

The study analyzed secondary data of BDHS. The BDHS is a part of the worldwide Demographic and Health Surveys (DHS) programme. The Bangladesh Ministry of Health and Family Welfare provided ethical approval for the surveys. The National Institute of Population Research and Training conducted the surveys with the support of the United States Agency for International Development (USAID). The survey protocol was approved by the National Research Ethics Committee in Bangladesh and ORC Macro (Macro International Inc.) Institutional Review Board. Informed consent was obtained from all participants.

Results

Of the 20 127 women data that were analysed in this study, 14 346 women reported they know about MR service which equates to 71% of the total respondents (Table 1). Of the women who reported positively, only 7% use MR service within three years of the survey date. The mean age of the respondents was 31.59 years at the time of the survey and mean years of education was 5.77 years.

Table 1: Background characteristics of the respondents

Variables	Mean (SE) or Frequency (%)
Women's age at the time of survey (mean±SD)	31.59 (±9.18)
Women's age at marriage	16.15 (±2.97)
Woman's education	5.77±4.14
Heard about menstrual regulation	
Yes	14346 (71.28)
No	5781 (28.72)
Use menstrual regulation within three years of the survey	
Yes	1056 (7.36)
No	13290 (92.64)

Of the women who used MR services, 62% resided in rural area (Figure 2). Moreover, of the urban women who know about MR service, only 9% used and about 93% of the rural women did not use of MR service within three years of the survey date. In contrary, nearly 7% of the total rural women who know about MR services used, MR services within three years of the survey date.

Prevalence of knowledge about Menstrual Regulation and MR service use

We calculated age-adjusted prevalence of MR service knowledge and use of MR service across individual-, household, and community level factors (Table 2). MR service knowledge and MR service use were found to be higher among respondents with increased age, education and who got marriage after their ages of 18 years or later. Women with formal engagement of employment were reported more use and knowledge of MR service. We found increase knowledge of MR and MR service use among women whose partners were educated and engaged in either services or business. Urban women reported more knowledge and use of MR. Both knowledge and use of MR were found higher among women who resided in the community with higher literacy and lower poverty.

Table 2: Age-adjusted prevalence of knowledge and use of menstrual regulation service across individual-, household-, and community level characteristics, Bangladesh.

Characteristics	Knowledge about menstrual regulation service % (95% CI)	Menstrual regulation service use % (95% CI)
Age of respondents		
15-19	55.34 (52.24 – 58.40)	1.56 (0.95 – 2.55)
20-34	74.10 (72.30 – 75.82)	6.53 (5.89 – 7.22)
≥35	71.67 (69.53 – 73.61)	9.82 (8.90 – 10.83)
Respondent educational level		
No education	61.46 (58.61 – 64.23)	6.90 (5.68 – 8.37)
Primary education	68.66 (66.46 – 70.78)	8.13 (7.21 – 9.14)
Secondary education	73.16 (71.25 – 74.98)	6.90 (6.24 – 7.63)
Higher education	84.80 (82.90 – 86.52)	7.49 (6.25 – 8.95)
Age at first marriage		
Early marriage (below 18 years)	69.75 (67.87 – 71.57)	7.51 (6.92 – 8.15)
Marriage 18 years and above	75.93 (73.88 – 77.87)	6.93 (5.99 – 7.99)

Respondent currently working status		
No	68.66 (66.47 – 70.81)	7.15 (6.42 – 7.95)
Yes	74.13 (72.31 – 75.86)	7.57 (6.86 – 8.35)
Husband educational status		
No education	64.96 (62.16 – 67.66)	6.50 (5.50 – 7.67)
Primary education	68.69 (66.48 – 70.82)	7.13 (6.25 – 8.12)
Secondary education	73.77 (71.78 – 75.66)	8.11 (7.17 – 9.15)
Higher education	82.96 (81.12 – 84.66)	7.95 (6.71 – 9.39)
Husband occupational status		
Agriculture	67.63 (65.08 – 70.07)	6.26 (5.41 – 7.23)
Physical workers	70.84 (68.82 – 72.77)	7.14 (6.37 – 7.99)
Service	84.84 (81.92 – 87.36)	8.69 (6.75 – 11.13)
Business	75.74 (73.34 – 77.99)	9.22 (8.10 – 10.46)
Others	63.74 (45.66 – 78.62)	6.32 (1.35 – 24.87)
Wealth Index		
Poorest	64.09 (61.04 – 67.03)	4.88 (3.97 – 5.99)
Poorer	68.41 (65.78 – 70.92)	6.71 (5.61 – 8.00)
Middle	70.74 (68.06 – 74.19)	7.57 (6.54 – 8.75)
Richer	71.62 (68.92 – 74.19)	7.17 (6.10 – 8.41)
Richest	80.59 (78.39 – 82.62)	9.62 (8.39 – 11.01)
Children ever born		
No Child	67.92 (65.75 – 70.00)	4.05 (3.41 – 4.79)
1-2 child	74.40 (72.57 – 76.15)	9.02 (8.24 – 9.87)
>2 Child	69.17 (66.71 – 71.53)	8.16 (7.00 – 9.49)
Place of Residence		
Urban	77.82 (75.23 – 80.21)	8.99 (7.91 – 10.20)
Rural	68.67 (66.46 – 70.81)	6.62 (6.01 – 7.29)
Place of regions		
Barisal	69.43 (64.16 – 74.24)	10.43 (8.45 – 12.82)
Chattogram	67.42 (63.43 – 71.17)	5.05 (4.11 – 6.19)
Dhaka	76.80 (72.29 – 80.77)	8.19 (6.80 – 9.84)
Khulna	63.87 (58.70 – 68.73)	7.01 (5.80 – 8.46)
Mymensingh	77.86 (72.51 – 82.42)	3.59 (2.74 – 4.69)
Rajshahi	69.36 (64.63 – 73.71)	7.59 (6.51 – 8.83)
Rangpur	74.42 (70.44 – 78.03)	10.64 (8.95 – 12.60)
Sylhet	65.23 (58.89 – 71.07)	5.76 (4.47 – 7.40)
Community level illiteracy		
Low	79.10 (74.73 – 82.88)	10.03 (8.49 – 11.80)
Moderate	70.72 (68.52 – 72.83)	7.24 (6.57 – 7.97)
High	67.94 (63.65 – 71.95)	5.72 (4.66 – 6.99)
Community level Poverty		
High poverty	70.87 (68.28 – 73.33)	6.33 (5.62 – 7.12)
Moderate poverty	68.48 (64.46 – 72.24)	6.47 (5.57 – 7.51)
Low poverty	69.16 (64.83 – 73.18)	9.61 (8.09 – 11.39)
Richest Community	78.69 (74.78 – 82.14)	9.54 (7.91 – 11.47)

The factors associated with the MR service knowledge and use of MR were determined using the multilevel Poisson regression model. Five different models were run separately. The summary of each model is presented in Table 3. The model with the lowest intra-class correlation and AIC and BIC are the best model. Full model, for both the MR service knowledge and use of MR service, met this condition. We found around 16% difference of MR service knowledge across clusters which reduced to 9% once individual, household and community level factors were adjusted. Similarly, there were 22.10% difference of MR service use was found across cluster which declined to 13.10% following the adjustment of individual, household, and community level factors.

Table 3: Results from random intercept model (measure of variation) for menstrual regulation knowledge and menstrual regulation use at cluster level using multilevel Poisson regression analysis.

	Null model	Individual level model	Household level model	Community level model	Full model
Menstrual regulation knowledge					
Intra-class correlation	16.10%	14.10%	13.28%	12.10%	9.40%
Variance of the random intercept	0.70	0.64	0.57	0.48	0.40
AIC	38186.7	37984.4	35100.4	38128.7	34973.7
BIC	38202.5	38056.01	35317.7	38246.7	35247.4
Menstrual regulation service use					
Intra-class correlation	22.10%	18.10%	16.12%	15.42%	13.20%
Variance of the random intercept	0.54	0.46	0.32	0.28	0.16
AIC	7551.04	7430.83	6964.96	7459.08	6854.04
BIC	7536.21	7499.09	7177.53	7572.86	7116.72

The results of the full models, for both the MR service knowledge and use of MR service, are presented in Table 4 while results of all models are presented in supplementary tables 1 and 2. The likelihoods of MR service knowledge were found to be higher among women aged 20-34 years (PR, 1.26, 95% CI, 1.20-1.33) and ≥ 35 years (PR, 1.30, 95% CI, 1.23-1.38) as compared to the women aged 15-19 years. Knowledge about MR service was found to be increased to around 6-16% among women with increased education from primary education. Alternatively, around 10% lower knowledge of MR service (PR, 0.90, 95% CI, 0.86-0.93) was found

among illiterate women. Similar associations were reported for women's partner education. We found around 4-9% lower knowledge of MR service among women with poorest and poorer socio-economic status as compared to the women with middle wealth quintile. Around 5-6% higher likelihoods of MR knowledge were found among women with increased number of children than among women who have no children. Rural women were 12% (PR, 0.88, 95% CI, 0.84-0.94) less likely to know about MR service as compared to the urban women. Women in the Mymensingh and Rangpur divisions were 10-17% more likely to know about MR service as compared to the women in the Barishal division. Knowledge about MR service was found to be increased with the increased level of illiteracy and decreased of community level poverty.

The MR service use was found to be increased to 3.13 times (95% CI, 1.85-5.28) and 4.23 times (95% CI, 2.44-7.32) among women aged 20-34 years and ≥ 35 years, respectively, as compared to the women aged 15-19 years. Likelihoods of MR service use was also found higher among women whose husbands were occupations were either physical worker (PR, 1.24, 95% CI, 1.03-1.50) or business (PR, 1.39, 95% CI, 1.13-1.70) as compared to the women whose husbands were engaged with agricultural work. As compared to the women with middle wealth quintile, women with poorest wealth quintile were 29% (PR, 0.71, 95% CI, 0.54-0.93) less likely to use MR service. Around 64-67% higher likelihoods of MR service use were found among women with increased number of children than women with no children. Women resided in the Khulna (PR, 0.68, 95% CI, 0.51-0.90) and Mymensingh (PR, 0.42, 95% CI, 0.30-0.58) divisions were less likely to use MR as compared to the women resided in the Barishal division. Use of MR service was also found to be increased with the decreased of community level poverty and increase of community level illiteracy.

Table 4: Factors associated with knowledge about menstrual regulation service and menstrual regulation service use using the multilevel Poisson regression model

Characteristics	Menstrual regulation knowledge, PR, (95% CI)	Menstrual regulation use, PR, (95% CI)
Age of respondents		
15-19 years	1.00	1.00
20-34 years	1.26 (1.20 – 1.33)**	3.13 (1.85 – 5.28)**
≥35 years	1.30 (1.23 – 1.38)**	4.23 (2.44 – 7.32)**
Respondent educational level		
No education	0.90 (0.86 – 0.93)**	0.85 (0.68 – 1.08)
Primary education	1.00	1.00
Secondary education	1.06 (1.03 – 1.09)**	0.85 (0.68 – 1.07)
Higher education	1.16 (1.11 – 1.20)**	0.86 (0.65 – 1.14)
Age at first marriage		
Marriage (10-17 years)	1.00	1.00
18 years and above	0.98 (0.97 – 1.01)	0.85 (0.73 – 1.01)
Respondent currently working status		
No	1.00	1.00
Yes	1.08 (1.05 – 1.11)**	1.07 (0.73 – 1.00)
Husband educational level		
No education	0.96 (0.94 – 0.99)**	0.90 (0.73 – 1.11)
Primary education	1.00	1.00
Secondary education	1.04 (1.01 – 1.06)**	1.18 (0.98 – 1.41)
Higher education	1.06 (1.02 – 1.09)**	1.05 (0.81 – 1.35)
Husband occupational status		
Agriculture	1.00	1.00
Physical workers	1.03 (1.00 – 1.05)*	1.24 (1.03 – 1.50)*
Service	1.04 (0.99 – 1.08)	1.33 (0.98 – 1.82)
Business	1.05 (1.00 – 1.08)**	1.39 (1.13 – 1.70)**
Others	1.00 (0.79 – 1.27)	0.97 (0.23 – 4.01)
Wealth Index		
Poorest	0.91 (0.88 – 0.95)**	0.71 (0.54 – 0.93)**
Poorer	0.96 (0.93 – 0.99)*	0.95 (0.75 – 1.21)
Middle	1.00	1.00
Richer	0.99 (0.96 – 1.02)	0.88 (0.70 – 1.29)
Richest	1.04 (0.99 – 1.08)	1.00 (0.78 – 1.29)
Children ever born		
No Child	1.00	1.00
1-2 child	1.06 (1.03 – 1.09)**	1.67 (1.37 – 2.04)**
>2 Child	1.05 (1.00 – 1.08)**	1.64 (1.27 – 2.14)**
Place of residence		
Urban	1.00	1.00
Rural	0.88 (0.84 – 0.94)**	0.96 (0.79 – 1.16)
Place of Regions		
Barisal	1.00	1.00
Chattogram	0.98 (0.91 – 1.04)	0.47 (0.35 – 0.62)**
Dhaka	1.12 (1.02 – 1.23)*	0.76 (0.57 – 1.01)

Khulna	0.93 (0.84 – 1.02)	0.68 (0.51 – 0.90)**
Mymensingh	1.17 (1.07 – 1.27)**	0.42 (0.30 – 0.58)**
Rajshahi	1.04 (0.94 – 1.13)	0.84 (0.65 – 1.07)
Rangpur	1.10 (1.01 – 1.19)**	1.25 (0.98 – 1.60)
Sylhet	1.01 (0.91 – 1.13)	0.63 (0.45 – 0.88)**
Community level Poverty		
High poverty	1.00	1.00
Moderate poverty	0.91 (0.86 – 0.97)**	0.99 (0.81 – 1.22)**
Low poverty	0.87 (0.81 – 0.94)**	1.41 (1.07 – 1.85)**
Middle and richest Community	0.86 (0.78 – 0.96)*	1.28 (0.92 – 1.79)
Community level illiteracy		
Low	1.00	1.00
Moderate	0.92 (0.87 – 0.98)**	0.82 (0.68 – 0.99)*
High	0.89 (0.82 – 0.97)**	0.67 (0.51 – 0.88)**

Discussion

In this study, we explored the knowledge and prevalence of MR service use in Bangladesh using most recent 2017 BDHS data. We found around 71.28% of the total women analysed know about this service while 7.36% of the women used this service within three years prior to the survey. Knowledge about MR service was found higher among higher aged women, higher educated women, women who engaged in income generating employment, women whose husbands were either work as physical worker or do business, women with lower socio-economic quintile and women with increased number of children. Rural women and women who resided in the community with low poverty and higher illiteracy were less likely to know about MR service. MR service use was found higher among higher aged women, women whose husbands were either physical workers or businessmen, women who having increased number of children and resided in the community with lower poverty. Lower use of MR service was found among women resided in the Chattogram, Khulna and Mymensingh divisions and women resided in the community with increased illiteracy. Findings will help the policy makers to make evidence-based policy and program to increase knowledge of MR service and its use whenever necessary.

We found higher knowledge and use of MR among increased aged women which are consistent with the findings of previous studies conducted in other LMICs (Alam & Sultan, 2019; Park et al., 2020; Reiss et al., 2019; Singh et al., 2017b). Lower aged women are more likely to be influenced by social stigma which might be linked with such lower use of MR. Moreover, increased aged women were less likely to use contraception or depend on less effective or traditional contraception's (Blake et al., 2007; Zahan & Feng, 2020). Also, they have already had desired number of children, so that any pregnancy is treated as unintended. Therefore, they consider MR service use (Bose S & Trent K 2006; Mote et al., 2010). Alternatively, if any unintended pregnancy occurred among lower aged women, who do not have required number of children, they may compromise with the time and continue the pregnancy to live births.

As like other studies in LMICs, in this study we found higher knowledge and use of MR among women with increased education (Alam & Sultan, 2019; Ankara, 2017; DaVanzo et al., 2013; Mote et al., 2010; Rana et al., 2019). Educated women were more likely to know about the reproductive and family planning services options either as part of their academic education, higher community level engagement, and higher exposure of mass media. Also, they have better decision-making autonomy over their reproductive life and know more about the importance of lower number of children (DaVanzo et al., 2013). This association is also justified by the higher use of MR service among with increased number of children as found in this study which is also similar to the previous studies in LMICs (Alam & Sultan, 2019; Aleni et al., 2020; Allotey et al., 2021; Dickson et al., 2018). Together these lead to higher use of MR service among women with increased education.

We also reported knowledge and use of MR services were different across divisions of Bangladesh and urban vs rural which are consistent with the available studies of LMICs (Chae et al., 2017; Kant

et al., 2015; Kapil Ahmed et al., 2005; Rana et al., 2019; Yassin K M 2000). These variations could be explained by community level social stigma, and community level religious and cultural norms. Beside these, proximity of healthcare facility is also important as found in a recent study of Bangladesh (Bearak et al., 2020; Rana et al., 2019).

Traditionally in Bangladesh, a community where people resided in is format with a combination of group of people with similar socio-economic status. Therefore, a majority or all of the people in lower poverty or illiterate community are either rich or educated. This direct higher knowledge and use of MR service among lower poverty and illiterate community as found in this study. In addition to these community level norms and social stigma, which are found more influential in the community with higher poverty and illiteracy, are also important for such associations (Ahmed & Ray, 2014; Ankara, 2017; Arambepola et al., 2016; Chae et al., 2017; Fusco et al., 2012; Johnston et al., 2010; SM Zahedul Islam, 2020; Vlassoff et al., 2012).

This study has several strengths and few limitations. In this study, we used most recent nationally representative data with a very higher sample size representative to all administrative divisions and urban and rural areas. We considered a range of individual-, household-, and community level factors and determined their linkages by using advanced statistical methods. Clusters level variations and sample weight were also considered in all analyses. Moreover, for the first time we calculated age-standardized prevalence of MR service knowledge and use. Therefore, the findings are mostly applicable for evidence-based decision and program making. However, the data analysed in this study were cross-sectional, therefore, the findings reported in this study were correlational only rather than casual. Moreover, since MR service use data were collected by asking directly to respondents without any validation, therefore, recall bias could be a major issue. Due to social and cultural norms, women may not give the accurate response on MR use status. However, any of such

bias is likely to be random. In addition, health facility level factors, including MR service availability are important predictors of MR service knowledge and use of MR. However, we could not adjust this variable in our analysis due to lack of data. Another important limitation of this study the BDHS authority did not mention in which places MR services use taken of the women.

Conclusion

Around 72% of the analysed women knew about MR while only 7% of them used MR within three years of the survey date. Several individual, household, and community level factors, including women's age, education, number of children, women's husbands' age, occupation, place of residence, division and community status in response to poverty and illiteracy were found to be associated with MR service knowledge and use of MR in Bangladesh. Findings suggest requirement of policy and program in the community level to increase knowledge of MR and its use as an effective family planning services to reduce burden of unintended conception in Bangladesh. This should include awareness building program at the community level to encourage people to access MR service when the pregnancy was undesired.

Acknowledgments

The authors thank MEASURE DHS for granting access to the BDHS 2017 data. The data is available on the website of the global DHS program, which could be accessed after getting registered with the MEASURES DHS. We wish to thank the director Aree Jampaklay , IPSR, MU, Thailand. The authors also thank to TWAS-IsDB and the editor and reviewers for their critical comments for the improvement of the manuscript.

Authors' contributions

MRA developed the study concepts. MRA and MNK analyzed the data. MRA drafted the manuscript. MRA, MNK and YS critically reviewed the manuscript. All authors have read and approved the final version of the paper.

Funding

The author(s) received no specific funding for this work.

Competing interests

The authors have declared that no competing interests exist.

Declaration of interests: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability: The datasets used and analyzed in this study are available from the Measure DHS website: <https://dhsprogram.com/data/available-datasets.cfm>

References

- Ahmed, S., & Ray, R. (2014). Determinants of pregnancy and induced and spontaneous abortion in a jointly determined framework: Evidence from a country-wide, district-level household survey in India. *Journal of Biosocial Science*, 46(4), 480–517. <https://doi.org/10.1017/S0021932013000369>
- Alam, M. Z., & Sultan, S. (2019). Knowledge and Practice of Menstrual Regulation (MR) in Bangladesh: Patterns and Determinants. *Journal of Population and Social Studies [JPSS]*, 27(3), 220–231.
- Al-Balushi, M. S., Ahmed, M. S., Islam, M. M., & Khan, M. H. R. (2020). Multilevel poisson regression modeling to identify factors influencing the number of children ever born to married women in Oman. *Journal of Statistics and Management Systems*, 23(8), 1357–1373. <https://doi.org/10.1080/09720510.2019.1709328>
- Aleni, M., Mbalinda, S., & Muhindo, R. (2020). Birth Intervals and Associated Factors among Women Attending Young Child Clinic in Yumbe Hospital, Uganda. *International Journal of Reproductive Medicine*, 2020, 1–11. <https://doi.org/10.1155/2020/1326596>
- Allotey, P., Ravindran, T. K. S., & Sathivelu, V. (2021). Trends in Abortion Policies in Low- and Middle-Income Countries. *Annual Review of Public Health*, 42(1), 505–518. <https://doi.org/10.1146/annurev-publhealth-082619-102442>
- Ankara, H. G. (2017). SOCIOECONOMIC VARIATIONS IN INDUCED ABORTION IN TURKEY. *Journal of Biosocial Science*, 49(1), 99–122. <https://doi.org/10.1017/S0021932016000158>
- Arambepola, C., Rajapaksa, L. C., Attygalle, D., & Moonasinghe, L. (2016). Relationship of family formation characteristics with unsafe abortion: Is it confounded by women’s socio-economic status? - A case-control study from Sri Lanka. *Reproductive Health*, 13(1), 75. <https://doi.org/10.1186/s12978-016-0173-5>
- BBS, B. (Ed.). (2014). *Bangladesh population and housing census, 2011. National report*. Bangladesh Bureau of Statistics, Statistics and Informatics Division, Ministry of Planning, Bangladesh, Government of the People’s Republic of Bangladesh.

BDHS. (2017). *Bangladesh Demographic and Health Survey 2017-18*.

<https://dhsprogram.com/publications/publication-FR344-DHS-Final-Reports.cfm>

Bearak, J., Popinchalk, A., Ganatra, B., Moller, A.-B., Tunçalp, Ö., Beavin, C., Kwok, L., & Alkema, L. (2020a). Unintended pregnancy and abortion by income, region, and the legal status of abortion: Estimates from a comprehensive model for 1990–2019. *The Lancet Global Health*, 8(9), e1152–e1161. [https://doi.org/10.1016/S2214-109X\(20\)30315-6](https://doi.org/10.1016/S2214-109X(20)30315-6)

Benson, J., Andersen, K., & Samandari, G. (2011). Reductions in abortion-related mortality following policy reform: Evidence from Romania, South Africa and Bangladesh. *Reproductive Health*, 8(1), 39.

<https://doi.org/10.1186/1742-4755-8-39>

Blake, S. M., Kiely, M., Gard, C. C., El-Mohandes, A. A. E., El-Khorazaty, M. N., & the NIH-DC Initiative.

(2007). Pregnancy Intentions and Happiness Among Pregnant Black Women at High Risk for Adverse Infant Health Outcomes. *Perspectives on Sexual and Reproductive Health*, 39(4), 194–205.

<https://doi.org/10.1363/3919407>

Bose, S., & Trent, K. (2006). Socio-demographic determinants of abortion in India: A north-South

comparison. *Journal of Biosocial Science*, 38(2), 261–282. <https://doi.org/10.1017/S0021932005026271>

Bronfenbrenner, U. (1994), B., U. (1994). Ecological models of human development. In T. Husen & T. N.

Postlethwaite (Eds.), *International Encyclopedia of Education* (2nd Ed., Vol. 3, pp. 1643– 1647).

Oxford, England: Pergamon Press. *Oxford: Elsevier*, 3(2), 37-.

Chae, S., Desai, S., Crowell, M., Sedgh, G., & Singh, S. (2017). Characteristics of women obtaining induced abortions in selected low- and middle-income countries. *PLOS ONE*, 12(3), e0172976.

<https://doi.org/10.1371/journal.pone.0172976>

Dahlberg, L. L., & Krug, E. G. (2006). Violence a global public health problem. *Ciência & Saúde Coletiva*,

11(2), 277–292. <https://doi.org/10.1590/S1413-81232006000200007>

DaVanzo, J., Rahman, M., Ahmed, S., & Razzaque, A. (2013). Influences on pregnancy-termination decisions in Matlab, Bangladesh. *Demography*, 50(5), 1739–1764. <https://doi.org/10.1007/s13524-013-0202-8>

- Dickson, K. S., Adde, K. S., & Ahinkorah, B. O. (2018). Socio – economic determinants of abortion among women in Mozambique and Ghana: Evidence from demographic and health survey. *Archives of Public Health*, 76(1), 37. <https://doi.org/10.1186/s13690-018-0286-0>
- Fusco, C. L. B., Silva, R. de S. e, & Andreoni, S. (2012). Unsafe abortion: Social determinants and health inequities in a vulnerable population in São Paulo, Brazil. *Cadernos De Saude Publica*, 28(4), 709–719. <https://doi.org/10.1590/s0102-311x2012000400010>
- Holmes, K., Curry, C., Sherry, Ferfolja, T., Parry, K., Smith, C., Hyman, M., & Armour, M. (2021). Adolescent Menstrual Health Literacy in Low, Middle and High-Income Countries: A Narrative Review. *International Journal of Environmental Research and Public Health*, 18(5), 2260. <https://doi.org/10.3390/ijerph18052260>
- Hossain, A., Maddow-Zimet, I., Ingerick, M., Bhuiyan, H. U., Vlassoff, M., & Singh, S. (n.d.). *Access to and Quality of Menstrual Regulation and Postabortion Care in Bangladesh: Evidence from a Survey Of Health Facilities, 2014*. 28.
- Hossain, A., Moseson, H., Raifman, S., Gerdt, C., Biswas, K. K., & Foster, D. G. (2016). ‘How shall we survive?’: A qualitative study of women’s experiences following denial of menstrual regulation (MR) services in Bangladesh. *Reproductive Health*, 13(1), 86. <https://doi.org/10.1186/s12978-016-0199-8>
- Huda, F. A., Ford, E., Deshpande, A., Haque, N. A., Quaiyum, M. A., Healy, J., Otsea, K., Johnston, H. B., & Ford, E. (2010). *Photographs*.
- Institute, G. (2017). *Menstrual Regulation and Unsafe Abortion in Bangladesh*. 2.
- Johnston, H. B., Oliveras, E., Akhter, S., & Walker, D. G. (2010). Health system costs of menstrual regulation and care for abortion complications in Bangladesh. *International Perspectives on Sexual and Reproductive Health*, 36(4), 197–204. <https://doi.org/10.1363/3619710>
- Kant, S., Srivastava, R., Rai, S. K., Misra, P., Charlette, L., & Pandav, C. S. (2015). Induced abortion in villages of Ballabgarh HDSS: Rates, trends, causes and determinants. *Reproductive Health*, 12(1), 51. <https://doi.org/10.1186/s12978-015-0040-9>

- Kapil Ahmed, M., van Ginneken, J., & Razzaque, A. (2005). Factors associated with adolescent abortion in a rural area of Bangladesh. *Tropical Medicine & International Health: TM & IH*, 10(2), 198–205.
<https://doi.org/10.1111/j.1365-3156.2004.01362.x>
- Menstrual Regulation and Unsafe Abortion in Bangladesh*. (2017, March 3). Guttmacher Institute.
<https://www.guttmacher.org/fact-sheet/menstrual-regulation-unsafe-abortion-bangladesh>
- Mote, CV., Larsen-Reindorf, R., Otupiri, E., & Hindin, M. (2010). FACTORS ASSOCIATED WITH INDUCED ABORTION AMONG WOMEN IN HOHOE, GHANA. *African Journal of Reproductive Health*, 14(4 0 0), 110–116.
- Nelder, J. A, McCullagh. P. (1989). *Generalized Linear Models 2nd Edition*. Chapman and Hall.
- Park, S., Yoon, T. W., Kang, D. R., & Chung, C. (2020). [Prevalence of Menstrual Disorders according to Body Mass Index and Lifestyle Factors: The National Health Insurance Service-National Health Screening Cohort in Korea, 2009~2016]. *Journal of Korean Academy of Nursing*, 50(3), 401–410.
<https://doi.org/10.4040/jkan.20054>
- Rana, J., Sen, K. K., Sultana, T., Hossain, M. B., & Islam, R. M. (2019). Prevalence and determinants of menstrual regulation among ever-married women in Bangladesh: Evidence from a national survey. *Reproductive Health*, 16(1), 123. <https://doi.org/10.1186/s12978-019-0785-7>
- Reiss, K., Andersen, K., Pearson, E., Biswas, K., Taleb, F., Ngo, T. D., Hossain, A., Barnard, S., Smith, C., Carpenter, J., Menzel, J., Footman, K., Keenan, K., Douthwaite, M., Reena, Y., Mahmood, H. R., Tabbassum, T., Colombini, M., Bacchus, L., & Church, K. (2019). Unintended Consequences of mHealth Interactive Voice Messages Promoting Contraceptive Use After Menstrual Regulation in Bangladesh: Intimate Partner Violence Results From a Randomized Controlled Trial. *Global Health, Science and Practice*, 7(3), 386–403. <https://doi.org/10.9745/GHSP-D-19-00015>
- Singh, M., Rajoura, O. P., & Honnakamble, R. A. (2019). Menstrual patterns and problems in association with body mass index among adolescent school girls. *Journal of Family Medicine and Primary Care*, 8(9), 2855–2858. https://doi.org/10.4103/jfmprc.jfmprc_474_19

- Singh, S., Hossain, A., Maddow-Zimet, I., Vlassoff, M., Bhuiyan, H. U., & Ingerick, M. (2017a). The Incidence of Menstrual Regulation Procedures and Abortion in Bangladesh, 2014. *International Perspectives on Sexual and Reproductive Health*, 43(1), 1–11. <https://doi.org/10.1363/43e2417>
- Singh, S., Hossain, A., Maddow-Zimet, I., Vlassoff, M., Bhuiyan, H. U., & Ingerick, M. (2017b). The Incidence of Menstrual Regulation Procedures and Abortion in Bangladesh, 2014. *International Perspectives on Sexual and Reproductive Health*, 43(1), 1–11. <https://doi.org/10.1363/43e2417>
- SM Zahedul Islam, C. (2020). Practices and Attitudes of Women Regarding Family Planning And Menstrual Regulation in The Sylhet Division of Bangladesh. *Global Journal of Infectious Diseases and Clinical Research*, 6(1), 014–028. <https://doi.org/10.17352/2455-5363.000030>
- Sustainable Development Goals | United Nations Development Programme*. (n.d.). UNDP. Retrieved December 8, 2021, from <https://www.undp.org/sustainable-development-goals>
- UNICEF, U. (2019). Child marriage-Bangladesh-2019. <https://www.ilo.org/surveyLib/index.php/catalog/3036/download/21390>
- United Nations. (2021). *United Nations 2021*. <https://worldpopulationreview.com/country-rankings/abortion-rates-by-country>
- Vlassoff, M., Hossain, A., Maddow-Zimet, I., Singh, S., & Bhuiyan, H. U. (2012). *Menstrual Regulation and Postabortion Care in Bangladesh: Factors Associated with Access to and Quality of Services*. <https://www.guttmacher.org/report/menstrual-regulation-and-postabortion-care-bangladesh-factors-associated-access-and-quality>
- Yassin, K. M. (2000). Incidence and socioeconomic determinants of abortion in rural Upper Egypt. *Public Health*, 114(4), 269–272.
- Zahan, R., & Feng, C. X. (2020). Bayesian spatial analysis of socio-demographic factors influencing pregnancy termination and its residual geographic variation among ever-married women of reproductive age in Bangladesh. *BMC Public Health*, 20(1), 1348. <https://doi.org/10.1186/s12889-020-09401-1>





Figure 1: Study sample selection process: Bangladesh Demographic and Health Survey 2017-18



medRxiv preprint doi: <https://doi.org/10.1101/2022.03.09.22272531>; this version posted March 18, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Figure 2: Menstrual regulation service use and knowledge by place of residence of BDHS 2017.