1 2 2	Estimating the burden of malaria and soil-transmitted helminth co- infection in sub-Saharan Africa: a geospatial study					
3 4 5	Short title: Malaria and helminth co-infections					
6	Muhammed O. Afolabi <sup>1</sup> , Adekola Adebiyi <sup>2</sup> , Jorge Cano <sup>3</sup> , Benn Sartorius <sup>1,4,5</sup> , Brian Greenwood <sup>1</sup> ,					
/ Q	<sup>1</sup> Department of Disease Control London School of Hygiene & Tropical Medicine, London, United					
9	Kingdom					
10 11	<sup>2</sup> College of Agriculture, Engineering and Environmental Design, Legacy University, Banjul, The Gambia					
12	<sup>3</sup> Expanded Special Project for Elimination of NTDs, World Health Organization Regional Office for					
13	Africa, Brazzaville, Republic of the Congo					
14	<sup>4</sup> Centre for Tropical Medicine and Global Health, University of Oxford, Oxford, United Kingdom					
15	<sup>5</sup> Department of Health Metric Sciences, University of Washington, Seattle, USA					
16	<sup>6</sup> Department of Mathematics, University of Manchester, Manchester, United Kingdom					
17	<sup>7</sup> Department of Infectious Diseases Epidemiology, London School of Hygiene & Tropical Medicine,					
18	London, United Kingdom					
19	<sup>8</sup> Vaccines and Immunity Theme, Medical Research Council Unit The Gambia at London School of					
20	Hygiene & Tropical Medicine, Fajara, The Gambia					
21						
22	Corresponding author: Muhammed.Afolabi@lshtm.ac.uk					
23						
24						

- 25 Abstract
- 26

27 Background: Limited understanding exists about the interactions between malaria and soil-

transmitted helminths (STH), their potential geographical overlap and the factors driving it. This study

29 characterised the geographical and co-clustered distribution patterns of malaria and STH infections

30 among vulnerable populations in sub-Saharan Africa (SSA).

31 Methodology/Principal findings: We obtained continuous estimates of malaria prevalence from the Malaria Atlas Project and STH prevalence surveys from the WHO-driven Expanded Special Project 32 33 for the Elimination of NTDs (ESPEN) covering 2000-2018 and used spatial autocorrelation methods to identify statistically significant clusters for both diseases across SSA. We used the inverse distance 34 weighted kriging (interpolation) methods to estimate STH prevalence. We calculated the population-35 weighted prevalence of malaria and STH co-infection, and used the bivariate local indicator of spatial 36 37 association (LISA analysis) to explore potential co-clustering of both diseases at the implementation 38 unit levels.

39 Our analysis shows spatial variations in the estimates of the prevalence of *Plasmodium falciparum*-

40 STH co-infections and identified hotspots across many countries in SSA with inter-and intra-country

41 variations. High *P. falciparum* and high hookworm co-infections were more prevalent in West and

42 Central Africa, whereas high P. falciparum, high Ascaris lumbricoides, high P. falciparum, and high

43 Trichuris trichiura co-infections were more predominant in Central Africa, compared to other sub-

44 regions in SSA.

45 Conclusions/Significance: Wide spatial heterogeneity exists in the prevalence of malaria and STH 46 co-infection within the regions and within countries in SSA. The geographical overlap and spatial co-47 existence of malaria and STH could be exploited to achieve effective control and elimination agendas 48 through the integration of the vertical programmes designed for malaria and STH into a more 49 comprehensive and sustainable community-based paradigm.

50 Keyword: malaria, geohelminths, spatial analysis, sub-Saharan Africa

## 52 Author Summary

53

Malaria and worms frequently co-exist together among children living in the poorest
countries of the world, but little is known about the specific locations of the combined
infections involving the two major parasitic diseases and how they interact and change over
the years.

We used open access data collected by two public registries, that is, the Malaria Atlas Project
and Expanded Special Project for the Elimination of NTDs, to understand the overlap of the
two diseases in different parts of Africa, where their burden are more predominant.

61 We found significant differences in the distributions of the combined diseases across different

62 parts of Africa, with large concentrations identified in Central and West Africa. For example,

63 double infections with malaria and hookworm were more common in West and Central

64 Africa, whereas malaria and roundworm, and malaria and whipworm were predominantly

65 found in Central Africa. A large collection of the dual infections was also found in some

66 localities within the countries which appeared to have low burden of the two diseases.

These findings provide a useful insight into the areas which could be serving as a reservoir to
propagating the transmission of the two diseases. The results of this study could also be used
to develop and implement integrated control programmes for malaria and worms, and this
could help to achieve the WHO NTD roadmap to ending the neglect to attain Sustainable
Development Goals by 2030.

72

73

74

## 76 Introduction

77

78 Due to environmental and host factors which favour transmission of multiple parasitic infections, malaria and soil-transmitted helminths (STH), including Ascaris lumbricoides, Trichuris trichiura, 79 and hookworms, co-exist in many parts of the world, predominantly in sub-Saharan Africa (SSA) [1], 80 81 where they are most prevalent. An estimated 241 million cases of malaria were reported in 2021 in 85 82 malaria endemic countries, 29 of which accounted for 96% of malaria cases globally, and six SSA countries (Nigeria, Democratic Republic of the Congo, Uganda, Mozambique, Angola and Burkina 83 84 Faso) carried more than half of this burden. [2] Similarly, a disproportionately high burden of STH has been reported in many countries in SSA with 85

large concentrations of moderate-to-heavy intensity infections in Nigeria, Democratic Republic of the 86 87 Congo, Ethiopia, Cameroon, Angola, Mozambique, Madagascar, Equatorial Guinea, and Gabon. [3] 88 In these malaria-STH co-endemic countries, estimating the national burden of the co-infection poses a 89 public health challenge because national surveillance systems are often sub-optimal.[4] For example, 90 deworming through mass drug administration (MDA) of anthelminthics is widely regarded as a cost-91 effective strategy. However, the impact of these MDA programmes on the changing prevalence and 92 intensity of STH in children, is often hindered by a lack of comprehensive baseline data before MDA 93 initiation, because of financial challenges or as a result of the administration of the drugs via undocumented channels.[3] Nevertheless, recent WHO reports [1, 4] indicate that considerable 94 95 progress has been made in reducing the overall burden of malaria and STH in many countries 96 including some in SSA. To sustain the gains of past decades and move the control efforts towards 97 elimination, WHO has recommended a paradigm shift from disease-specific to an integrated 98 approach.[4]

Empirical studies conducted across co-endemic countries have shown that variations in the prevalence
of malaria and STH co-infections differed significantly across geographic locations.[5-9] Limited
understanding exists about the spatial distribution of these co-infections in vulnerable populations.
Majority of the published studies targeting malaria-STH co-infection have focused on describing the

103 transmission and burden among affected populations, [5, 6, 9, 10] and have paid little attention to their concurrent spatial distribution and causes. Previous systematic reviews have also reported an 104 over-estimation of the relationship between malaria and helminths making it difficult to establish 105 conclusively the burden of malaria-STH dual infections. [11, 12] Consistent with findings of these 106 107 reviews, a recently conducted systemic review and meta-analysis of 55 studies which enrolled 37,559 children across low and middle- income countries (LMIC) found a wide variation in the prevalence of 108 109 malaria-helminth co-infections, ranging from 7-76% across LMICs.[13] These findings may be due to 110 the low sensitivity of the diagnostic methods employed for the detection of malaria-helminth co-111 infections in the primary studies included in the systematic review.

The WHO road map for the control and elimination of NTDs for the 2021-2030 period defines STH elimination as a public health problem (EPHP) when there is a prevalence of less than 2% of moderate-to-heavy intensity infections.[14] However, as STH control accelerates and new detection tools are developed, a significant number of people with clinical features outside high intensity infection hotspots are now being seen. [15]

Reliable estimates of malaria-STH morbidity hotspots are needed to comply with the NTD roadmap goal of intensifying cross-cutting approaches which integrate STH control with common delivery platforms for diseases that share similar epidemiology such as malaria. Also, empirical evidence shows that obtaining reliable estimates of the burden of STH-malaria co-infection would help address the critical gaps in planning and implementation of integration of STH control strategies with other interventions. [14]

We undertook a geospatial analysis on the datasets describing the prevalence of malaria and STH infections among adults and children aged 5-14 years, obtained from the geospatial open-access data generated from the Malaria Atlas Project (MAP)[16] and from the WHO-driven Expanded Special Project to Eliminate NTDs (ESPEN) [17], to characterise the geographical and co-clustered distribution patterns for malaria and STH co-infections among vulnerable populations in SSA.

128

# 129 Methods

130

#### 131 Study design

We analysed the spatial relationships between STH and malaria using spatial autocorrelation methods to identify statistically significant clusters over different regions in endemic countries in SSA. Based on the classifications of the United Nations Geoscheme for Africa[18], we categorised SSA into west, east, central and southern African sub-regions. These include 46 of Africa's 54 countries and territories that are fully or partially south of the Sahara, excluding Algeria, Djibouti, Egypt, Libya, Morocco, Somalia, Sudan and Tunisia.

138 We obtained datasets on the prevalence of *Plasmodium falciparum* from the MAP database (last 139 accessed September 2, 2021). MAP is a WHO collaborating platform for geospatial disease modelling which was created for the purpose of determining spatial limits, prevalence and endemicity of P. 140 falciparum and P. vivax. MAP provides a database to obtain data on the estimated prevalence of P. 141 falciparum and P. vivax at 5km resolution for each year from 2000 onwards. We focused on P. 142 falciparum in this study because it is the predominant causative agent for more than 95% of malaria 143 cases in SSA[19]. We also obtained datasets on the prevalence of STH (A. lumbricoides, T. trichiura 144 145 and hookworms) from the WHO-driven ESPEN portal. ESPEN portal contains survey datasets on NTDs in Africa. Survey datasets from both MAP and ESPEN were confined to surveys carried out from 2000 146 to 2018. 147

The *P. falciparum* prevalence data obtained from MAP were provided in a raster format for each year covering the entire African continent. To make them usable, we combined the data into one raster file using a mosaic tool. The description of the methodology used for generating the predicted prevalence is available at https://malariaatlas.org/api-docs/

The prevalence datasets on STH (*A. lumbricoides, T.trichiura*, and hookworms) extracted from the ESPEN portal were geolocated, that is, the data contains the geographic coordinates of the survey locations which can be either a school or village location. The data consist of 25,274 data points from 2000 to 2018, of which 817 (3.2%) were without spatial reference. Of the remaining 24,457 (96.8%) data points with spatial reference, 6,229 data points had prevalence rates above zero for *T. Trichiura*,
12,945 data points had prevalence rates above zero for hookworm and 9,728 for *A.lumbricoides*.
Incomplete data (those without spatial reference and explicit prevalence rates) were excluded from the
analyses.

160 Statistical analysis

161

### 162 Spatial auto-correlation

163 We used Moran's I statistics as the indicator of spatial auto-correlation. We plotted the prevalence of P. falciparum on the x-axis against the spatially lagged variables of hookworm, T. trichiura and A. 164 165 *lumbricoides* on the y-axis. All points with values above the mean (i.e. above zero) were sites with 166 high prevalence while all points with values below the mean (i.e. below zero) were sites with low 167 prevalence. This holds true for the prevalence of *P. falciparum* plotted on the x-axis. The scatter plot was divided into four quadrants. The values of P. falciparum and hookworm, T. trichiura and 168 169 A.lumbricoides that fell under the upper-right quadrant and the lower left quadrant were regarded as 170 high-high (hotspots) and low-low (coldspots) values, respectively. We used the bivariate Moran's I 171 algorithm in Geoda software (https://geodacenter.github.io/) to implement this analysis. **Spatial interpolation** 172 To produce the prevalence surface of STH from the point referenced data, we used the inverse distance 173 weighted kriging (interpolation) methods.[20] Prediction by kriging is based on the assumption that 174 175 prevalence estimates at locations close together are correlated. Therefore, it allowed us to estimate the value of STH prevalence at any point from the observed data. We carried out the Kriging using ArcGIS 176 version 10.3 software (https://arcview-gis.software.informer.com/10.3/). Our findings are virtualised 177 at implementation unit level (administrative level 2 for most of the SSA countries). By implementation 178 unit, WHO is considering the administrative division at which the preventive chemotherapy related 179

interventions are to be implemented. For most of countries in the African region, these areas correspondwith the second administrative level, although many countries have opted for using health district

182 division when this exists.

#### 184 Population-weighted prevalence by implementation unit

185 We calculated the population-weighted prevalence of malaria and STH for each implementation unit

- using the population density data obtained from the WorldPop database
- 187 (<u>https://www.worldpop.org/project/categories?id=14</u>) and the prevalence surfaces. Using 2020
- 188 Population raster of individual countries in SSA from the WorldPop database, estimated 165 million
- 189 people were in high-risk clusters of *P.falciparum*, hookworm and *A.lumbricoides*. For malaria-STH
- 190 co-infection, about 73 million, 21 million and 118 million people were in high risk clusters of
- 191 *P.falciparum* and hookworm; *P.falciparum* and *T. trichura; P.falciparum* and A. lumbricoides;
- 192 respectively. In most countries, administrative level 2 is the geographical area over which a particular
- treatment strategy or intervention is applied; this could be a district, local government area, county, or
- 194 province. There were a total of 5,935 administrative level 2 in the study areas reported in this paper.

#### 195 Bivariate spatial clustering

196 To identify the spatial clusters of an implementation unit, we used local Moran's I algorithm in Geoda

- 197 software (<u>https://geodacenter.github.io/</u>) to implement the bivariate local indicator of spatial
- association (LISA).[21] LISA is a spatial autocorrelation used to assess the influence of individual
- 199 locations on the magnitude of other locations. It gives an indication of the extent of significant spatial

200 clustering of similar values around that observation. We used LISA to identify the hotspots and

- 201 coldspots, and areas endemic to malaria-STH co-infection across the study areas. We considered areas
- with high malaria-STH co-infection and alpha level  $\leq 0.001$  as statistically significant hotspots.

## 203 Results

- 205 The spatial auto-correlation plot was a linear fit whose slope corresponds to Moran's I and its values
- for *P. falciparum* against hookworm, *T. trichiura* and *A. lumbricoides* were 0.302, 0.054 and 0.118,
- 207 p-value <0.05, respectively (Fig 1A, B, C). High P. falciparum and high hookworm (HW) co-
- 208 infection were more prevalent in the West and Central African sub-regions compared to other sub-
- regions of SSA (Figure 2A). The countries with high prevalence of *P. falciparum* and high HW co-
- 210 infections were Angola, Benin, Burundi, Cameroon, Central Africa Republic, Congo Brazzaville,

211 Cote d'Ivoire, Democratic Republic of the Congo (DRC), Ghana, Guinea, Kenya, Liberia, Mali,

212 Nigeria, Sierra Leone, South Sudan, Togo, and Uganda (Figure 2A). Within these countries with high

prevalence of *P. falciparum* and HW co-infection, there were specific areas (second administrative

214 unit) with statistically significant co-infection (p < 0.001) which can be considered as hotspots for the

co-infection.

216 The second administrative units with statistically significant P.falciparum and HW co-infection

217 (p<0.001) included north-west Angola; south-west Cameroon; west-coastal communities in Congo

218 Brazzaville; south-central and east-central Cote d'Ivoire; and Kongo central province of DRC (Figure

219 2B). Other hotspots for co-infection with *P.falciparum* and HW were communities in western Kenya;

south-western Liberia; eastern, southern and northern Sierra Leone; central and north-central Togo;

221 districts in the Lake Victoria region and eastern Uganda (Figure 2B). In Nigeria, hotspots for co-

222 infection with *P. falciparum* and HW co-infection included coastal communities in the southern and

south-eastern parts of the country (Figure 2B, Table S1).

224

225





Fig 1A: Moran's I statistic plot showing spatial auto-correlation for *P.falciparum* vs hookworm co-infections in sub-Saharan Africa, 2000-2018





Fig 1B: Moran's I statistic plot showing spatial auto-correlation for *P.falciparum* vs T. *trichiura* co-infections in sub-Saharan Africa, 2000-2018



- 252 Fig 2A: Local autocorrelation statistic interpretation of *P. falciparum* and hookworm co-
- 253 infection, SSA, 2000-2018



Fig 2B: Probability that a given implementation unit is a hotspot for *P. falciparum* and hookworm co-infection in SSA, 2000 – 2018



256	High <i>P.falciparum</i> and high <i>T. trichiura</i> co-infection were more prevalent in the Central African sub-
257	region compared to other sub-regions in SSA (Figure 3A). Angola, Cameroon, Congo Brazzaville,
258	DRC, Equatorial Guinea, Gabon, and coastal communities of southern Nigeria had a high prevalence
259	of <i>P. falciparum</i> and high <i>T. trichiura</i> co-infection (Figure 3A). Hotspots for <i>P. falciparum</i> and <i>T.</i>
260	trichiura co-infections included communities around the coast in Angola and several communities in
261	the central and southern Cameroon (Figure 3B). Other hotspots for co-infection with <i>P.falciparum</i>
262	and T. trichiura were communities spread across all districts of Congo Brazzaville; all districts in the
263	westernmost province of DRC; all regions in Equatorial Guinea and all districts except south-
264	eastern region of Gabon (Figure 3B, Table S1).
265	
266	
267	
268	
269	
270	
271	
272	
273	
274	
275	
276	

Fig 3A: Local autocorrelation statistic interpretation of P. falciparum and T. trichiura co-infection, SSA, 2000-2018



#### 

Fig 3B: Probability that a given implementation unit is a hotspot for *P.falciparum* and *T.trichiura* co-infection, SSA, 2000 – 2018



High *P.falciparum* and high *A. lumbricoides* (AS) co-infection was more prevalent in Central Africa
and parts of West Africa (Figure 4A). The countries with high prevalence of *P.falciparum* and high *A. lumbricoides* co-infection were Angola, Burundi , Cameroon, Central African Republic, Congo
Brazzaville, Cote d'Ivoire , DRC, Equatorial Guinea, Gabon, Kenya , Liberia, and Nigeria (Figure
3A). However, hotspots for *P.falciparum* and *A. lumbricoides* co-infection were identified in only
seven countries: Angola, Congo Brazzaville, DRC, Equatorial Guinea, Gabon, Liberia, and Nigeria
(Figure 4B).

Hotspots for *P.falciparum* and *A. lumbricoides* in Angola included communities in the northeast and
northern exclave of Angola, on the west (Atlantic) coast and north of the Congo River (Figure 4B).
Other hotspots for *P.falciparum* and *A. lumbricoides* co-infections in SSA included north-western
and south-western regions in DRC; all regions in Equatorial Guinea, the Littoral region and
south-eastern counties of Liberia; and almost all regions in Gabon. In Nigeria, hotspots for *P.falciparum* and *A. lumbricoides* co-infections were in 13 states located along the coastal lines of the
southern parts of the country (Figure 4B, Table S1).

Figure 4A: Local autocorrelation statistic interpretation of *P. falciparum* and *A. lumbricoides* co-infection, SSA, 2000-2018



### 303

<sup>304</sup> Figure 4B: Probability that a given implementation unit is a hotspot for *P. falciparum* and *A. lumbricoides* coinfection, SSA, 2000 – 2018



### 306 Discussion

307

308 Our analysis shows spatial variations in the estimates of prevalence of *P.falciparum*-STH co-309 infections and identified hotspots across many countries in SSA where the co-infections were high. 310 Consistent with previous studies [3, 12, 22-24], our findings also highlight inter- and intra-country variations in the prevalence of malaria-helminth co-infections. Across Central and West Africa sub-311 regions, a high prevalence of *P.falciparum* and high hookworm co-infection was predominant, while 312 the burden of P. falciparum-A. lumbricoides and P.falciparum-T-trichiura co-infections was higher in 313 Central Africa than in neighbouring sub-regions. Although, recent reports have shown that a 314 considerable gain has been made in the control of malaria and STH in the last two decades [1, 14], our 315 316 findings suggest that a high burden of falciparum malaria and moderate-to-heavy infections for at least one STH species are still occurring in the southern part of West Africa and extending to Central 317 Africa including Cameroon, Burundi and DRC. We observed similar patterns and trends in the 318 319 hotspot distributions for P. falciparum and any of the STH. While programmatic implementation of 320 preventive chemotherapy and seasonal malaria chemoprevention (SMC) have contributed to a significant reduction in the burden of STH and malaria in many countries, inherent challenges such as 321 322 logistical burden, medication adherence, and incomplete coverage have been reported to be 323 responsible for the persistently high burden of malaria and STH in the spatial areas identified in our 324 study.[14, 15]

325 Taken together the geographical overlap and spatial co-existence of malaria and STH, favoured 326 largely by environmental factors[1, 3], could be exploited to achieve effective control and elimination agendas through integration of the vertical programmes designed for malaria and STH into a more 327 comprehensive and sustainable community-based paradigm, that includes other preventive care such 328 329 as hand hygiene and water sanitation. The WHO 2030 NTD road map recommends concrete actions within integrated platforms for delivery of the interventions needed to improve the cost-effectiveness, 330 331 coverage and geographical reach of these integrated programmes.[14] A successful integration of the 332 vertical control programmes aligns well with the global trend towards integrated, non-disease specific 333 approaches which have become an increasingly recognised strategy recommended by WHO.[25]

334 However, limited evidence is currently available on the feasibility and effectiveness of integrating malaria with STH control at community level and evidence that this can be done effectively is needed. 335 Our study has some limitations. First, our study used secondary data obtained from two different 336 337 sources covering a period from 2000–2018, which may not fully capture the epidemiological status of the burden of the co-infection during the period under review. Our analysis also showed some 338 inconsistencies with findings of empirical studies [23, 26], especially for STH prevalence. For 339 example, we observed high uncertainty around estimates in Central Africa and some parts of West 340 Africa, which corresponds to areas with a high burden of infection. Lack of survey data is a plausible 341 342 reason for this observation.[3] Additionally, a substantial proportion of the data we used for our analysis was collected prior to the commencement of mass preventive treatments with albendazole 343 and SMC. Many countries have tailored the frequency of mass drug administration with preventive 344 345 chemotherapy to their STH prevalence, but annual SMC implementation during the period of high 346 malaria transmission is still being given as a single intervention.

347 The absence of survey data or inadequacy in definition of space-time contributes to the uncertainty observed in our finding and these may have also introduced bias leading to over-estimation or under-348 estimation of the prevalence of STH and malaria co-infection. Contributing further to the likely over 349 or under-estimation of the burden of STH and malaria could be empirical treatments of malaria and 350 351 STH outside the control programmes. Our findings could also be affected by the diagnostic approaches involving malaria rapid testing [27] and Kato-Katz method [28] used to generate the data, 352 as these have been widely reported to be less sensitive than PCR, especially in low-transmission 353 354 settings.

In conclusion, we have demonstrated wide spatial heterogeneity within the SSA and within countries, on the prevalence of malaria and STH confections. To consolidate on the encouraging progress made in the reduction of the prevalence of malaria and STH in some SSA countries, it is crucial that integrated control programmes target the regions with high-high burden of malaria and STH. Our study may also have critical implications for policy-making and resource allocations based on the needs of each region, as well as for the future of the implementation of integrated malaria-helminth

361	control programmes. Nevertheless, it is important that our findings are confirmed by further empirical
362	studies as this will provide the platform for research agenda that will lead to the establishment of the
363	much-needed platform for the implementation of the integrated programmes that are cost-effective,
364	and make optimum use of limited human resources frequently found in SSA.

Acknowledgement: We thank Mike Thorn of the Malaria Atlas Project and Ismaela Abubakar for
 supporting the data curation for this study

367 Author Contributions: MOA conceptualised the study. BS and JCO provided support in data

368 curation and analysis. AA and OJ performed the statistical analysis of the datasets. MOA, OW, AA

and OJ drafted different sections of the manuscript. BG revised the manuscript critically for important

intellectual content. All authors reviewed and approved the final draft of the manuscript.

### 371 Competing interests: None declared

372 Funding: This work was implemented as part of a career development fellowship awarded to MOA,

which is funded under the UK Research and Innovation Future Leaders Fellowship scheme

374 (MR/S03286X/1). The funders had no role in the study design, data collection and analysis, decision

375	to publish,	or preparation	of the manu	iscript
-----	-------------	----------------	-------------	---------

- 376
- 377

- 379
- 380
- 381
- 382
- 383

## 384 References

Brooker S, Akhwale W, Pullan R, Estambale B, Clarke SE, Snow RW, et al. Epidemiology of
 Plasmodium-Helminth Co-Infection in Africa: Populations at Risk, Potential Impact on Anemia, and
 Prospects for Combining Control. The American journal of tropical medicine and hygiene.
 2007;77(6 Suppl):88-98. doi: https://doi.org/10.4269/ajtmh.2007.77.88.

World malaria report 2021. Geneva: World Health Organization; 2021. Licence: CC BY-NC SA 3.0 IGO. Retrieved from <a href="https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2021">https://www.who.int/teams/global-malaria-programme/reports/world-</a>
 malaria-report-2021. Accessed 7 Dec 2021.

- 392 3. Sartorius B, Cano J, Simpson H, Tusting LS, Marczak LB, Miller-Petrie MK, et al.
- Prevalence and intensity of soil-transmitted helminth infections of children in sub-Saharan Africa,
  2000-18: a geospatial analysis. The Lancet Global health. 2021;9(1):e52-e60. Epub 2020/12/19. doi:
- 395 10.1016/s2214-109x(20)30398-3. PubMed PMID: 33338459; PubMed Central PMCID:
   396 PMCPMC7786448.
- 4. Ending the neglect to attain the Sustainable Development Goals: a road map for neglected
  tropical diseases 2021–2030. Geneva: World Health Organization; 2020. Licence: CC BY-NC-SA 3.0
  IGO.
- Mulu A, Legesse M, Erko B, Belyhun Y, Nugussie D, Shimelis T, et al. Epidemiological and
  clinical correlates of malaria-helminth co-infections in southern Ethiopia. Malaria Journal.
  2013;12(1):227. doi: 10.1186/1475-2875-12-227.
- Adedoja A, Tijani BD, Akanbi AA, 2nd, Ojurongbe TA, Adeyeba OA, Ojurongbe O. Coendemicity of Plasmodium falciparum and Intestinal Helminths Infection in School Age Children in
  Rural Communities of Kwara State Nigeria. PLoS neglected tropical diseases. 2015;9(7):e0003940-e.
  doi: 10.1371/journal.pntd.0003940. PubMed PMID: 26222743.
- Kepha S, Nuwaha F, Nikolay B, Gichuki P, Edwards T, Allen E, et al. Epidemiology of
  coinfection with soil transmitted helminths and Plasmodium falciparum among school children in
  Bumula District in western Kenya. Parasites & vectors. 2015;8:314-. doi: 10.1186/s13071-015-08915. PubMed PMID: 26063324.
- 8. Zeukeng F, Tchinda VHM, Bigoga JD, Seumen CHT, Ndzi ES, Abonweh G, et al. Coinfections of malaria and geohelminthiasis in two rural communities of Nkassomo and Vian in the
  Mfou health district, Cameroon. PLoS neglected tropical diseases. 2014;8(10):e3236-e. doi:
  10.1371/journal.pntd.0003236. PubMed PMID: 25329479.
- 415 9. Kinung'hi SM, Magnussen P, Kaatano GM, Kishamawe C, Vennervald BJ. Malaria and
  416 helminth co-infections in school and preschool children: a cross-sectional study in Magu district,
  417 north-western Tanzania. PloS one. 2014;9(1):e86510-e. doi: 10.1371/journal.pone.0086510. PubMed
  418 PMID: 24489732.
- Achidi EA, Apinjoh TO, Mbunwe E, Besingi R, Yafi C, Wenjighe Awah N, et al. Febrile
  status, malarial parasitaemia and gastro-intestinal helminthiases in schoolchildren resident at different
  altitudes, in south-western Cameroon. Annals of Tropical Medicine & Parasitology. 2008;102(2):10318. doi: 10.1179/136485908X252287.
- 11. Naing C, Whittaker MA, Nyunt-Wai V, Reid SA, Wong SF, Mak JW, et al. Malaria and soiltransmitted intestinal helminth co-infection and its effect on anemia: a meta-analysis. Transactions of
  The Royal Society of Tropical Medicine and Hygiene. 2013;107(11):672-83. doi:
- 426 10.1093/trstmh/trt086.
- 427 12. Degarege A, Erko B. Epidemiology of Plasmodium and Helminth Coinfection and Possible
  428 Reasons for Heterogeneity. Biomed Res Int. 2016;2016:3083568. Epub 2016/04/20. doi:
- 429 10.1155/2016/3083568. PubMed PMID: 27092310; PubMed Central PMCID: PMCPMC4820611.
- 430 13. Afolabi MO, Ale BM, Dabira ED, Agbla SC, Bustinduy AL, Ndiaye JLA, et al. Malaria and
  431 helminth co-infections in children living in endemic countries: A systematic review with meta-
- 432 analysis. PLOS Neglected Tropical Diseases. 2021;15(2):e0009138. doi:
- 433 10.1371/journal.pntd.0009138.
- 434 14. Degarege A, Veledar E, Degarege D, Erko B, Nacher M, Madhivanan P. Plasmodium
- falciparum and soil-transmitted helminth co-infections among children in sub-Saharan Africa: a
- 436 systematic review and meta-analysis. Parasites & vectors. 2016;9(1):344. Epub 2016/06/17. doi:

437 10.1186/s13071-016-1594-2. PubMed PMID: 27306987; PubMed Central PMCID:

**438** PMCPMC4908807.

- Hotez PJ, Molyneux DH. Tropical anemia: one of Africa's great killers and a rationale for
  linking malaria and neglected tropical disease control to achieve a common goal. PLoS Negl Trop
- 441 Dis. 2008;2(7):e270. Epub 2008/07/31. doi: 10.1371/journal.pntd.0000270. PubMed PMID:

442 18665256; PubMed Central PMCID: PMCPMC2474697.

- 16. Lyke KE, Dicko A, Dabo A, Sangare L, Kone A, Coulibaly D, et al. Association of
- Schistosoma haematobium infection with protection against acute Plasmodium falciparum malaria in
   Malian children. The American journal of tropical medicine and hygiene. 2005;73(6):1124-30. Epub
- 446 2005/12/16. PubMed PMID: 16354824; PubMed Central PMCID: PMCPMC2738948.
- 447 17. Expanded Special Project for Elimination of Neglected Tropical Diseases (ESPEN).
  448 Available online at https://espen.afro.who.int/. Accessed 12 June 2020.
- Available online at <u>https://cspell.ano.who.mit</u>. Accessed 12 June 2020.
  18. Adegnika AA, Kremsner PG. Epidemiology of malaria and helminth interaction: a review
- 450 from 2001 to 2011. Curr Opin HIV AIDS. 2012;7(3):221-4. Epub 2012/03/16. doi:
- 451 10.1097/COH.0b013e3283524d90. PubMed PMID: 22418449.
- Hartgers FC, Yazdanbakhsh M. Co-infection of helminths and malaria: modulation of the
  immune responses to malaria. Parasite Immunol. 2006;28(10):497-506. Epub 2006/09/13. doi:
- 454 10.1111/j.1365-3024.2006.00901.x. PubMed PMID: 16965285.
- 455 20. Nacher M. Interactions between worms and malaria: good worms or bad worms? Malar J.
- 456 2011;10:259. Epub 2011/09/14. doi: 10.1186/1475-2875-10-259. PubMed PMID: 21910854; PubMed
  457 Central PMCID: PMCPMC3192711.
- 458 21. Tao R, Thill J-C. BiFlowLISA: Measuring spatial association for bivariate flow data.
- 459 Computers, Environment and Urban Systems. 2020;83:101519. doi:
- 460 <u>https://doi.org/10.1016/j.compenvurbsys.2020.101519</u>.
- 461 22. Owada K, Lau CL, Leonardo L, Clements ACA, Yakob L, Nielsen M, et al. Spatial
- distribution and populations at risk of A. lumbricoides and T. trichiura co-infections and infection
- intensity classes: an ecological study. Parasites & vectors. 2018;11(1):535. Epub 2018/10/05. doi:
- 464 10.1186/s13071-018-3107-y. PubMed PMID: 30285906; PubMed Central PMCID:
- 465 PMCPMC6171148.
- Camara Y, Sanneh B, Joof E, Sanyang AM, Sambou SM, Sey AP, et al. Mapping survey of
  schistosomiasis and soil-transmitted helminthiases towards mass drug administration in The Gambia.
  PLoS neglected tropical diseases [Internet]. 2021 2021/07//; 15(7):[e0009462 p.].
- 469 24. Kabatereine NB, Standley CJ, Sousa-Figueiredo JC, Fleming FM, Stothard JR, Talisuna A, et
  470 al. Integrated prevalence mapping of schistosomiasis, soil-transmitted helminthiasis and malaria in
  471 lakeside and island communities in Lake Victoria, Uganda. Parasites & vectors. 2011;4(1):232. doi:
  472 10.1186/1756-3305-4-232.
- 473 25. Hotez PJ, Molyneux DH, Fenwick A, Ottesen E, Ehrlich Sachs S, Sachs JD. Incorporating a
  474 rapid-impact package for neglected tropical diseases with programs for HIV/AIDS, tuberculosis, and
  475 malaria. PLoS Med. 2006;3(5):e102. Epub 2006/01/27. doi: 10.1371/journal.pmed.0030102. PubMed
  476 PMID: 16435908; PubMed Central PMCID: PMCPMC1351920.
- 470 Finib. 10455508, Fublicate Central Finicity. Finicity (1551520).
  477 26. Pullan RL, Kabatereine NB, Bukirwa H, Staedke SG, Brooker S. Heterogeneities and
  478 consequences of Plasmodium analysis and healwarm existentiation is a nonvelation head study in
- 478 consequences of Plasmodium species and hookworm coinfection: a population based study in
  479 Uganda. The Journal of infectious diseases. 2011;203(3):406-17. Epub 2010/12/27. doi:
- Uganda. The Journal of Infectious diseases. 2011;203(3):406
   10 1093/infdis/iia063. PubMed PMID: 21187338
- 480 10.1093/infdis/jiq063. PubMed PMID: 21187338.
- 481 27. Mahende C, Ngasala B, Lusingu J, Yong TS, Lushino P, Lemnge M, et al. Performance of
  482 rapid diagnostic test, blood-film microscopy and PCR for the diagnosis of malaria infection among
  483 febrile children from Korogwe District, Tanzania. Malar J. 2016;15(1):391. Epub 2016/07/28. doi:
  484 10.1186/s12936-016-1450-z. PubMed PMID: 27459856; PubMed Central PMCID:
- 485 PMCPMC4962396.
- 486 28. Cools P, Vlaminck J, Albonico M, Ame S, Ayana M, José Antonio BP, et al. Diagnostic
- 487 performance of a single and duplicate Kato-Katz, Mini-FLOTAC, FECPAKG2 and qPCR for the
- 488 detection and quantification of soil-transmitted helminths in three endemic countries. PLOS
- 489 Neglected Tropical Diseases. 2019;13(8):e0007446. doi: 10.1371/journal.pntd.0007446.
- 490