Conditions of Confinement in U.S. Carceral Facilities during COVID-19: Individuals Speak: Incarcerated during the COVID-19 Epidemic (INSIDE)

# Short Running Title: Conditions of Confinement in U.S. Carceral Facilities during COVID-19

# Alternative: Confinement in Carceral Facilities during COVID-19 (Characters w/ spaces: 50/50)

Nicole Cassarino, BA<sup>\*1</sup>, Harika Dabbara, BS<sup>\*1,2</sup>; Carla B. Monteiro, MSW<sup>3,4</sup>; Arthur Bembury<sup>5</sup>; Leslie Credle, BA<sup>6,7,8</sup>; Uma Grandhi<sup>9</sup>, Samantha White, AB<sup>10</sup>, Monik C. Jiménez, ScD<sup>1</sup>

\*Both authors contributed equally to the preparation of the manuscript and are listed in alphabetical order

<sup>1</sup>Division of Women's Health, Brigham and Women's Hospital, Boston, MA, USA
<sup>2</sup>Boston University School of Medicine, Boston, MA, USA
<sup>3</sup>Brigham Health Bridge Clinic, Boston, MA, USA
<sup>4</sup>Cape Verdean Social Workers Association, Boston, MA, USA
<sup>5</sup>Partakers Organization - College Behind Bars, Auburndale, MA, USA
<sup>6</sup>National Council for Incarcerated and Formerly Incarcerated Women and Girls
<sup>7</sup>Justice 4 Housing, Boston, MA
<sup>8</sup>Families for Justice as Healing, Boston, MA
<sup>9</sup>University of California Santa Cruz, Santa Cruz, CA, USA
<sup>10</sup>Harvard University, Cambridge, MA, USA

#### **Corresponding Author:**

Monik C. Jiménez, ScD, SM, FAHA Division of Women's Health Brigham and Women's Hospital, Harvard Medical School 1620 Tremont St., 3-034, Boston, MA 02120 Telephone: 617-525-7516; Fax: 617-525-7746 mjimenez11@partners.org

Keywords (4-6): Incarceration, COVID-19, Word count: 2995/3000 References: 30/100 Tables: 4/4 Figures: 0/4

Acknowledgements: Dr. Jiménez is supported by the Brigham and Women's Hospital Richard Nesson Fellowship. The authors would like to thank Massachusetts State Senator Rebecca Rausch, communications director Evan Berry, and intern Sandy Mabee for their contributions to this study.

**Author contributions:** MCJ initiated the study and led framing of the study, contributed to the analysis and drafted the manuscript; AB, CM, HD, LC and NC conceptualized the study design and contributed to drafting the manuscript; NC led and conducted the data analysis; HD led manuscript preparation; HD and NC jointly led data interpretation. AB, CM, HD, LC, MJ, NC, SW and UG contributed to interpreting the results and all approved the final version of the submitted manuscript.

#### Abstract (word count: 230/250, incl headers)

**Objectives**: We aimed to describe conditions of confinement among people incarcerated in the United States during the coronavirus disease 2019 (COVID-19) pandemic and assess the feasibility of a community-science data collection approach.

**Methods**: We developed a web-based survey with community partners to collect information on confinement conditions (COVID-19 safety, basic needs, support). Formerly incarcerated adults released after March 1, 2020, or non-incarcerated adults in communication with an incarcerated person (proxy) were recruited through social media from July 25, 2020, through March 27, 2021. Descriptive statistics were estimated in aggregate and separately by proxy or formerly incarcerated status. Additionally, we compared responses between proxy and formerly incarcerated respondents using chi-square or Fisher's exact tests as appropriate based on alpha=0.05.

**Results**: Of 378 responses, 94% were by proxy, and 76% reflected state prison conditions. Participants reported inability to physically distance ( $\geq$ 6ft at all times) (92%), inadequate access to soap (89%), water (46%), toilet paper (49%) and showers (68%). Among people who received mental healthcare before the pandemic, 75% reported reduced care. We found that responses were consistent between formerly incarcerated people and proxy-respondents.

**Conclusions**: Our findings suggest that a community-science approach to data collection is feasible. Based on these findings, COVID-19 safety and basic needs were not sufficiently addressed within some carceral settings. Thus, we recommend the lived experiences of incarcerated individuals should be included to make informed and equitable policy decisions.

# INTRODUCTION

Over a year after the initial global response to coronavirus disease 2019 (COVID-19), the United States (U.S.) sits at the top of two lists: the highest rate of incarceration, and the highest rate of COVID-19 deaths.<sup>1,2</sup> In 2020, COVID-19 case rates among individuals in U.S. federal and state departments of corrections (DOC) were 4.8-5.5 times higher than the general population.<sup>3-5</sup> Previous outbreaks of influenza,<sup>6-8</sup> Legionnaires' disease,<sup>9</sup> H1N1,<sup>10,11</sup> and tuberculosis<sup>12</sup> were harbingers of carceral facilities' vulnerability to SARS-CoV-2. Conditions such as overcrowding, inadequate ventilation, poor sanitation, and difficulty accessing personal protective equipment (PPE) have created environments primed for outbreaks. Additionally, highly prevalent conditions like diabetes, hypertension, and asthma place incarcerated populations at greater risk of COVID-19-related morbidity and mortality.<sup>13</sup>

Physical distancing, mask wearing, handwashing, and testing have been key public health COVID-19 mitigation strategies.<sup>14</sup> However, anecdotal reports indicate that mitigation strategies reported by state-level DOCs are irregularly implemented and highly punitive. Incarcerated individuals report inadequate access to soap, water, and disinfectant supplies,<sup>15</sup> and extensive periods of lockdowns (suspension of activities and confinement restricted to housing areas) or solitary confinement (housing with minimal to rare contact with others) to achieve physical distancing.<sup>15-17</sup> The punitive nature of these measures may have unknown consequences for physical and mental health.<sup>18</sup>

While anecdotal evidence is compelling, robust data on the lived experience of incarcerated people during the pandemic is limited with only one quantitative peer reviewed study to date.<sup>19</sup> Moreover, participants were only included from prisons in three Midwestern or Southeastern states. Due to the decentralized response to COVID-19 across state carceral

3

systems, data from more states and facility types are needed to understand the lived experience of incarceration during the COVID-19 pandemic.

Therefore, we partnered with community advocates to design and disseminate a webbased survey to collect detailed data on the conditions of confinement and determine the feasibility of collecting data through outreach to community members such as loved ones, formerly incarcerated individuals (FII), and professional partners.

#### METHODS

#### **Instrument Development**

In May 2020, a Community Advisory Board (CAB) was convened to design a survey to examine condition experienced by individuals incarcerated during the pandemic. The CAB included FII advocates from local and national organizations, Families for Justice as Healing, Justice 4 Housing, the National Council of Incarcerated and Formerly Incarcerated Women and Girls, and the Partakers Organization. Each member described their commitment to their communities below.

"My work and experience with the criminal justice system has been the impetus that motivates me to seek solutions that can help reduce recidivism by providing essential resources to formerly incarcerated individuals ... I believe in working with the gatekeepers to ensure that funding is directed to critical areas that are needed most, and will have a significant impact on the population and the communities we work with." —Arthur Bembury "Because of the things I saw during my 19 years of incarceration I have become an advocate for the women I had to leave behind and the women that will sleep in a prison bed tomorrow." —CAB Member

"Surviving the Federal carceral punishment system is the motivation that drives the work I do today. At Justice 4 Housing we advocate for the abolishment of discriminatory housing policies and ending incarceration of women." —Leslie Credle

At study onset, there were no validated instruments to examine confinement conditions among incarcerated individuals during the pandemic; therefore, we developed a novel survey across four domains: COVID-19 safety, basic needs, support, and demographics factors. Each domain was developed based on carceral system mitigation strategies, Center for Disease Control and Prevention (CDC) guidelines,<sup>14</sup> community-informed concerns of CAB members,<sup>20</sup> publicly available anecdotal evidence, and surveys targeting the general population.

The instrument was piloted among twelve individuals who had been incarcerated and released after March 1, 2020, or were in communication with an incarcerated individual. The pilot survey included questions regarding estimated time of completion, accessibility, necessity of questions asked, and inclusion of additional questions to better capture lived experiences. In partnership we reviewed the pilot findings, implemented changes, and planned survey dissemination.

The final instrument consisted of 42 questions, (37 multiple choice and 5 open response) (Online Supplement), of which three were administered only to proxies and seven only to FII. Data were collected anonymously using REDCap and the survey was distributed using a

5

snowball sampling method through Twitter and Facebook and by email to community-partner organizations, medical and legal professionals, and local politicians. After deployment, proxy respondents were allowed to indicate their medical or legal professional status.

#### **Study Sample**

Participants were required to be aged  $\geq 18$  years and either (1) a formerly incarcerated individual (FII), released after March 1, 2020 (beginning of the U.S. response to the pandemic), or (2) in contact with someone currently incarcerated (proxy). Currently incarcerated individuals were not recruited. To maximize reported data questions could be skipped. Proxies in contact with multiple incarcerated individuals were asked to submit a separate survey for each incarcerated person, unless proxies were legal or medical professionals, who were prompted to report the number of people for whom the information was applicable.

#### STATISTICAL ANALYSIS

All primary analyses were descriptive and thus did not include statistical hypothesis testing. Frequencies and relative frequencies were calculated for each question based on available data for all respondents in aggregate, with a missing indicator for skipped questions. Geographic regions were defined according to the U.S. Census Bureau classifications.<sup>21</sup>

We assessed internal validity by comparing the distribution of responses (excluding missing values) by proxies to FII respondents for questions regarding COVID-19 safety, basic needs, and support. Comparisons were conducted using two-sided Fisher's Exact Tests based on an  $\alpha$ =0.05. In addition, we conducted sensitivity analyses to examine differences over time and

state of incarceration for any variables with inconsistencies between proxy- and FII respondents. All analyses were conducted using Stata/IC (version 16.1, College Station, TX).

### **STATEMENT OF ETHICS**

This study was approved by the institutional review board of Mass General Brigham and deemed exempt because no personal identifiers were collected, and the research posed minimal risk. All procedures were in accordance with institutional guidelines. However, we acknowledge that these data represent the lived experiences of individuals who have been incarcerated and those who know them personally.

## RESULTS

Of 378 responses collected between July 28, 2020 and March 27, 2021, 94.4% (n=357) were proxies who completed the survey based on experiences shared by someone currently incarcerated (Table 1). The remaining 5.6% (n=21) were FII who were released after March 1, 2020 (Table 1). Most respondents provided data on facilities in the Northeast (37.0%), followed by the West (28.7%), South (26.1%), and Midwest (8.2%). In addition, 75.7% of respondents reflected conditions in state prisons, followed by jails (13.5%), federal prisons (5.8%), and other facilities including Immigration and Customs Enforcement (ICE) and juvenile detention centers (5.0%) (Table 1). Among respondents who reported gender of the impacted person, 39.6% identified as female, 60.1% were White, followed by Black (21.5%), and Hispanic/Latinx (12.9%).

#### **COVID-19 Safety**

Responses about COVID-19 safety measures and access to basic needs are reported for proxies and FII in aggregate (Table 2). Most participants (93.6%) reported experiencing lockdown procedures, of which 86.6% reported lockdowns >20 hours/day, and 48.8% reported a duration of >3 months. Over 92% of respondents indicated inability to always keep the CDC-recommended six-foot distance from others. Sixteen percent of respondents reported individual housing, while 53.7% reported housing with one other person, and 30% with two or more people. While 81.6% reported being provided PPE, (primarily face masks), 85.3% reported inconsistent use of PPE by staff. Over one-third of respondents (35.5%) reported that individuals who experienced COVID-19 symptoms were returned to their cells regardless of whether their cellmate also experienced symptoms. In addition, 53.5% reported that individuals were unable to take their belongings when being moved due to COVID-19. Lastly, nearly one-third (32.0%) of respondents reported that no information about COVID-19 was shared with individuals during incarceration. Among those who received information, the number of facility-wide COVID-19 cases was the most reported (32.8%).

#### **Basic Needs, Medical Care, and Support**

Overall, respondents indicated that individuals had inadequate access to basic necessities including personal hygiene items (Table 2). Although 67.9% received free soap, only 11.5% reported it was enough for their needs. Forty-six percent indicated insufficient access to water, toilet paper (48.5%) or daily showers (67.8%). In addition, 86.0% of respondents indicated insufficient food supply.

Medical care and support resources were also affected by the pandemic response (Table 2). A delay beyond normal delivery of medical care for flu-like symptoms was reported by

81.3% of respondents. Similarly, of those receiving mental healthcare before the pandemic,74.7% reported a decrease access. Access to legal services were also reduced; 79.1% reportedreductions in access to legal aid, and 36.0% indicated a delay of parole hearings.

#### **Questions only answered by Formerly Incarcerated Respondents**

Questions about PPE replacement, worry and added stress, and receipt of financial support were asked only to formerly incarcerated respondents (n=21; Table 3). In total, 81.8% indicated new PPE was provided less than once per week. In addition, 87.5% reported that they worried about contracting COVID-19 "most of the time" and experienced "a lot of added stress" (93.8%). Most FII respondents (62.5%) received less financial support during the pandemic and only 16.7% reported that they received facility-level COVID-19 updates. Nearly two-thirds (64.3%) of FII expressed interest in getting a COVID-19 vaccine if available.

#### **Questions only answered by Proxy Respondents**

Questions about recency of communication with incarcerated individuals, concerns about contracting COVID-19 from FII when released or through visitation, and increased expenses to support incarcerated individuals were posed specifically to proxies. Most had communicated with the individual within the past week (85%). A high proportion (70.1%) of proxies reported concerns about contracting the virus when facilities reopened for visitation or from the incarcerated individual if released (49.8%). Nearly a quarter of respondents (24.1%) reported needing a rapid COVID-19 test within a week to feel safe if the incarcerated individual were released tomorrow. Additionally, 81.9% reported increased expenses to support the individual.

Since respondents were allowed to skip questions, missing data ranged from 4.7% for facility lockdown to 59% for access to mental healthcare.

#### **Internal Validity**

The distribution of responses between proxies and FII were compared for key metrics and were found to be consistent (p>0.05) for all but two questions. First, proxies were significantly more likely to report a facility being on lockdown (94.4%) than FII (79.0%, p=0.03). Additionally, fewer proxies (30.0%) reported that incarcerated individuals had access to daily showers compared to 63.2% of FII (p=0.01).

### DISCUSSION

To our knowledge, this is the first study designed in partnership with FII to collect data on confinement conditions among individuals incarcerated across all four regions of the U.S. Our findings suggest that at least in some facilities, individuals were unable to always physically distance from others, despite extensive periods of lockdown, and lacked access to basic needs (i.e., soap, water, toilet paper, and showers). Respondents also indicated limited access to healthcare and legal aid.

One other quantitative study has reported confinement conditions based on 327 individuals incarcerated in three U.S. states.<sup>19</sup> The COVID-19 Questionnaire for Correctional Populations (CQCP) survey was used to collect data on mitigation strategies observed by staff and incarcerated individuals. Our findings were similar to those reported by the CQCP study for some conditions, such as provision of soap (INSIDE: 67.9%; CQCP: 70.6%). However, we observed discrepancies for questions regarding physical distancing (INSIDE: 7.9%; CQCP: 66.4%), which may be due to wording differences. While the CQCP asked if participants "Stayed 6 feet away from others if possible," we explicitly asked, "Were you allowed to keep your distance from others by staying 6 feet away at all times?". Our study complements and expands this previous work by collecting nationwide data, reflecting a longer time-period (July 2020-March 2021 versus April 2020-November 2020) and includes questions on lockdown, basic needs, and mental healthcare access.

Our findings also paralleled several studies from the gray literature. The Essie Justice Group surveyed 729 people who were in contact with an incarcerated individual.<sup>22</sup> Overall, 62% reported that their incarcerated loved ones feared losing their lives, similar to 87.5% of FII from INSIDE who worried about getting COVID-19 "most of the time". Similarly, a Physicians for Human Rights survey among 50 individuals formerly detained by ICE, reported that 80% were not able to maintain a 6-foot distance from others while eating, consistent with the 92% of our respondents who reported an inability to distance  $\geq$ 6-feet at all times.<sup>15</sup>

Our data suggest that implementation of COVID-19 mitigation strategies may not meet CDC-outlined practices and may lead to physical and psychological distress. For example, numerous studies have shown solitary confinement's association with psychological harm<sup>23-25</sup> including anxiety, depression, aggression, self-harm, and increased mortality post-release.<sup>23</sup> Yet, lockdowns which result in extended confinement and isolation have been prioritized as a strategy to reduce transmission. One INSIDE FII reported,

"Being locked in a cell for 23 hours and 40 minutes a day for weeks at a time (whenever there was a report of a positive case among staff/incarcerated citizens was extremely

stressful. I often considered hurting other prisoners or the guards simply because I was angry. I am not a violent person. I was incarcerated for a drug offense."

In addition, in the general population, post-pandemic delayed or restricted care have likely contributed to the 18% increase in mortality (2020 versus 2019).<sup>26</sup> Specifically, mortality increases in diabetes (15%), unintentional injury (11%), Alzheimer's (10%), and heart disease (5%) are suggestive of collateral pandemic consequences.<sup>26</sup> In our study, prolonged time to receive medical care (81.3%) and reduced access to mental health care (74.7%) may be associated with increased risk of death or morbidity either from COVID-19 directly or chronic conditions. One proxy respondent reported,

"My loved one recently got infected with COVID-19 and is also suffering for underlying medical conditions (chronic Hepatitis B and asthma). He didn't receive any medical treatments other than having his inhaler with him... The only "treatment" that was given to some individuals were over-the-counter medications such as Tylenol to reduce some flu-like symptoms. My loved one is still suffering from some lingering symptoms such as wheezing and coughing."

Our findings support recommendations from public health professionals and advocacy groups to reform crisis responses strategies by centering (1) population decompression, (2) improvement of living conditions through external oversight, and (3) incorporation of input from incarcerated people and their families.<sup>27,28</sup> Importantly, the American Public Health Association has called for wide-spread population decompression due to the well-documented harms associated with incarceration which have only been exacerbated by the pandemic.<sup>27</sup> Their

comprehensive recommendations include use of diversion programs, sentencing, misdemeanor and bail reform. Additional tools for population decompression include expansion of compassionate release for those with chronic medical conditions and the elderly, home confinement programs, and expedited release of those nearing their release date.<sup>29</sup>

Our data also demonstrated some facilities' failure to provide basic needs, medical and mental healthcare, and implement safety measures to mitigate infectious disease transmission. However, there is no central external oversight body for US prisons and jails, resulting in limited accountability, inconsistent crisis response, and a lack of standardized protections.<sup>30</sup> In order to minimize harm to incarcerated people, transparent external oversight is necessary to ensure humane adherence to CDC guidelines. Lastly, our study provides a model for how directly impacted people and their loved ones can provide critical evidence regarding implementation of crisis response strategies, while highlighting the need for anonymous collection of firsthand experiences to foster transparency and ensure compliance.

To our knowledge, this study is the only study to quantitatively document confinement conditions during the pandemic with data from all four U.S. regions and across a variety of facility types. Importantly, this study was conducted in partnership with a CAB and driven by community-identified needs. Lastly, we demonstrate the feasibility and validity of gathering data on the lived experiences of incarcerated individuals via proxies.

We acknowledge that generalizability of our findings may be limited using a convenience sample. However, we aimed to increase geographic and carceral system diversity and recognized that obtaining required approvals from each independent facility to directly survey incarcerated individuals would hinder the collection of time-sensitive data. Despite this limitations, general comparisons with other studies and anecdotal reports support the patterns we observed. In

13

addition, generalizability of our sample by race and ethnicity is uncertain due to missing data for these variables. Although our sample is likely to overrepresent conditions in female facilities given our CAB member's networks; there is no evidence to suggest that conditions would be improved for incarcerated males or individuals of color. Furthermore, our novel data collection methods were developed early in the pandemic based on public health priorities at that time which have changed with respect to sanitation procedures over the course of the pandemic. Furthermore, data from proxies may not be optimally reported for all variables due to the proportion of "I don't know" or "Skip" responses to some questions, such as mental healthcare access. Lastly, two of our variables (facility on lockdown and ability to shower when wanted) demonstrated low internal validity; significantly more FII reported ability to shower daily if wanted, compared to proxies, while more proxies reported lockdown conditions compared to FII. However, 15.8% of FII who answered the questions regarding showering and lockdown were released before May 2020, prior to widespread response in all states. Additionally, all were incarcerated in Florida, which may reflect variability in different states' responses. However, we observed high internal validity for other variables, suggesting that proxy reports largely matched those of incarcerated individuals.

# **HEALTH EQUITY IMPLICATIONS**

Our data demonstrate that at least some incarcerated people experienced inadequate implementation of safety protocols, difficulty accessing basic needs, and were subjected to long periods of restricted confinement. Importantly, our data highlight the need to prioritize the lived experiences of incarcerated individuals in policy design and oversight, to ensure appropriate and humane policies and implementation. Our data were collected through robust community

partnerships, and we urge practitioners to engage with community organizations to develop

meaningful research questions, leverage community strengths, and shape policy.

# REFERENCES

States of Incarceration: The Global Context 2021. Prison Policy Initiative, 2021.
 (Accessed November 21, 2021, at <a href="https://www.prisonpolicy.org/global/2021.html">https://www.prisonpolicy.org/global/2021.html</a>.)

2. WHO Coronavirus (COVID-19) Dashboard. 2021. (Accessed February 10, 2022, at https://covid19.who.int/.)

3. Saloner B, Parish K, Ward JA, DiLaura G, Dolovich S. COVID-19 Cases and Deaths in Federal and State Prisons. JAMA 2020.

 Jiménez MC, Cowger TL, Simon LE, Behn M, Cassarino N, Bassett MT. Epidemiology of COVID-19 among incarcerated individuals and staff in Massachusetts jails and prisons.
 JAMA Netw Open 2020;3:e2018851.

Hagan LM, Williams SP, Spaulding AC, et al. Mass Testing for SARS-CoV-2 in 16
 Prisons and Jails - Six Jurisdictions, United States, April-May 2020. MMWR Morb Mortal Wkly
 Rep 2020;69:1139-43.

 Robinson S, Smith P, Stephen D, Shubert J, Reed C, Manning S. Influenza Outbreaks at Two Correctional Facilities - Maine, March 2011. Morbidity and Mortality Weekly Report 2012;61:229-32.

 Besney J, Moreau D, Jacobs A, et al. Influenza outbreak in a Canadian correctional facility. J Infect Prev 2017;18:193-8.

8. Young LC, Dwyer DE, Harris M, et al. Summer outbreak of respiratory disease in an Australian prison due to an influenza A/Fujian/411/2002(H3N2)-like virus. Epidemiol Infect 2005;133:107-12.

9. Lucas KD, Wheeler C, McLendon P, Leistikow BN, Mohle-Boetani JC. Outbreak of Legionnaires' disease associated with cooling towers at a California state prison, 2015. Epidemiol Infect 2018;146:297-302.

10. Chao WC, Liu PY, Wu CL. Control of an H1N1 outbreak in a correctional facility in central Taiwan. J Microbiol Immunol Infect 2017;50:175-82.

Turner KB, Levy MH. Prison outbreak: pandemic (H1N1) 2009 in an Australian prison.
 Public Health 2010;124:119-21.

12. Lambert LA, Armstrong LR, Lobato MN, Ho C, France AM, Haddad MB. Tuberculosis in Jails and Prisons: United States, 2002-2013. Am J Public Health 2016;106:2231-7.

13. Nowotny KM, Rogers RG, Boardman JD. Racial disparities in health conditions among prisoners compared with the general population. SSM Popul Health 2017;3:487-96.

14. COVID-19 and Your Health. U.S. Department of Health & Human Services 2022.

(Accessed February 20, 2022, at <u>https://www.cdc.gov/coronavirus/2019-ncov/your-health/need-</u>to-know.html.)

15. Peeler K, Erfani P, Lee CH, et al. Praying for Hand Soap and Masks: Health and Human Rights Violations in U.S. immigration Detention during the COVID-19 Pandemic. New York, NY2021.

16. What Happens When More Than 300,000 Prisoners Are Locked Down? The MarshallProject, 2020. (Accessed February 10, 2022, at

https://www.themarshallproject.org/2020/04/15/what-happens-when-more-than-300-000prisoners-are-locked-down.)

Association AB. ABA Standards for Criminal Justice: Treatment of Prisoners. Standard
 23-10. Washington, D.C.: American Bar Association; 2011.

 Cloud DH, Ahalt C, Augustine D, Sears D, Williams B. Medical Isolation and Solitary Confinement: Balancing Health and Humanity in US Jails and Prisons During COVID-19. J Gen Intern Med 2020;35:2738-42.

19. Pettus-Davis C, Kennedy SC, Veeh CA. Incarcerated individuals' experiences of COVID-19 in the United States. Int J Prison Health 2021;ahead-of-print.

20. Berk J, Rich JD, Brinkley-Rubinstein L. What are the greatest health challenges facing people who are incarcerated? We need to ask them. Lancet Public Health 2021;6:e703-e4.

21. 2010 Census Regions and Divisions of the United States. U.S. Census Bureau, 2010.

(Accessed February 10, 2022, at https://www2.census.gov/geo/pdfs/maps-

data/maps/reference/us\_regdiv.pdf.)

22. Clayton-Johnson G, Karefa-Johnson R, Rasaki T. Lives on the Line: Women with incarcerated loved ones and the impact of COVID-19 behind bars. Oakland, CA: Essie Justice Group, ; 2020.

23. Brinkley-Rubinstein L, Sivaraman J, Rosen DL, et al. Association of Restrictive Housing During Incarceration With Mortality After Release. JAMA Netw Open 2019;2:e1912516.

24. Kaba F, Lewis A, Glowa-Kollisch S, et al. Solitary confinement and risk of self-harm among jail inmates. Am J Public Health 2014;104:442-7.

 Haney C. Restricting the use of solitary confinement. Annual Review of Criminology 2018;1:285-310.

26. Ahmad FB, Cisewski JA, Minino A, Anderson RN. Provisional Mortality Data - United States, 2020. MMWR Morb Mortal Wkly Rep 2021;70:519-22.

27. Advancing Public Health Interventions to Address the Harms of the Carceral System.
2021. (Accessed November 22, 2021, 2021, at <a href="https://www.endingpoliceviolence.com">https://www.endingpoliceviolence.com</a>.)

28. Natoli LJ, Vu KL, Sukhija-Cohen AC, et al. Incarceration and COVID-19:

Recommendations to Curb COVID-19 Disease Transmission in Prison Facilities and

Surrounding Communities. Int J Environ Res Public Health 2021;18.

29. ACLU. COVID-19 Model Finds Nearly 100,000 More Deaths Than Current Estimates,

Due to Failures to Reduce Jails2020.

30. Deitch M. But who oversees the overseers?: The status of prison and jail oversight in the United States. American Journal of Criminal Law 2020;47:207-73.

N = 378	n	%
Role		
Proxy Respondent	357	94.4%
Formerly Incarcerated	21	5.6%
Age		
18-30 years	27	14.8%
30-44	90	49.5%
45-64 years	58	31.9%
> 65 years	7	3.9%
<b>Region of Carceral Facility</b> <sup>a</sup>		
Southern U.S.	98	26.1%
Northeastern U.S.	139	37.0%
Western U.S.	108	28.7%
Midwestern U.S.	31	8.2%
Type of Carceral Facility		
Jail	51	13.5%
State Prison	286	75.7%
Federal Prison	22	5.8%
ICE Detention Center	11	2.9%
Other	8	2.1%

# Table 1. Facility and Demographic Information of Incarcerated People Represented

## **Racial/ethnic Background**

American Indian/Alaska Native	0	0.0%
Asian	4	2.5%
Black	35	21.5%
Hispanic/Latinx	21	12.9%
Native Hawaiian/Pacific Islander	1	0.6%
White	98	60.1%
Multiracial	4	2.5%
Sex and Gender		
Male	106	58.2%
Female	72	39.6%
Trans Man	1	0.6%
Trans Woman	1	0.6%
Non-binary	2	1.1%

ICE: Immigration and Customs Enforcement

All responses exclude "Don't know" and skipped questions.

<sup>a</sup>Southern: Alabama, Arkansas, Delaware, DC, Florida, Georgia, Kentucky, Louisiana,

Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, West Virginia

Northeastern: Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York,

Pennsylvania, Rhode Island, Vermont

Western: Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, Wyoming

Midwestern: Indiana, Illinois, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, Wisconsin

N = 378	n	0⁄0ª		
Lockdown				
Any lock-down due to COVID-19 <sup>b</sup>	335	93.6%		
Duration of lock-down				
<2 weeks	17	5.8%		
>2 weeks - <1 month	51	17.5%		
1-2 months	47	16.2%		
2-3 months	34	11.7%		
> 3 months	142	48.8%		
Hours per day on lock-down				
$\leq$ 20hrs/day	35	13.4%		
>20hrs/day	227	86.6%		
COVID-19 Safety				
Physical distancing at all times	26	7.9%		
Number of people in cell				
Single	55	16.1%		
Double	183	53.7%		
>2/barracks/dormitory	103	30.2%		
Disinfection of common items				
No	147	58.6%		

# Table 2. Distribution of Reported Conditions of Confinement

More than daily	30	12.0%
At least daily	74	29.5%
Staff wearing PPE		
None	19	6.5%
Some staff	230	78.8%
All staff	43	14.7%
Type of PPE worn by staff <sup>b</sup>		
Face Mask	259	94.9%
Gloves	79	28.9%
Face Shield	19	7.0%
Incarcerated Person Given any PPE	244	81.6%
Type of PPE received <sup>b</sup>		
Face Mask	240	98.4%
Gloves	1	0.4%
Homemade	6	2.5%
Action taken if someone reported symptoms		
Isolation/Quarantine	147	58.6%
Returned to cell, IF cellmate ALSO has symptoms	15	6.0%
Returned to cell, EVEN IF cellmate has <i>NO</i> symptoms	89	35.5%

Able to reject COVID test without being punished	32	20.8%		
Type of information about COVID-19 received <sup>b</sup>				
# of people with COVID in facility	124	32.8%		
# of people tested for COVID in facility	87	23.0%		
# of deceased from COVID in facility	77	20.4%		
Plan for dealing with COVID	51	13.5%		
How to protect self	45	11.9%		
None	121	32.0%		
Basic Needs				
Received free soap from the facility				
No	78	32.1%		
Yes and enough for needs	28	11.5%		
Yes but NOT enough for needs	137	56.4%		
Unable to access water when wanted	107	46.1%		
Unable to access to enough toilet paper	96	48.5%		
Unable to shower every day if wanted	187	67.8%		
Delayed medical care for people with flu-like symptoms	191	81.3%		
Allowed to take possessions if moved within facility	87	46.5%		
Type of meals received				
1 hot meal/day and 2 bag lunches	120	62.5%		

Only bag lunches	72	37.5%		
Quantity of food received				
Enough	33	14.0%		
Not enough	203	86.0%		
Support		·		
Mental healthcare changes <sup>c</sup>				
More care received	7	4.6%		
Less care received	115	74.7%		
No care received	32	20.8%		
Given more stamps, telephone or video calls <sup>b</sup> Able to use increased telephone or video calls	136	36.0%		
Able to use for full allotted time	47	37.3%		
Able to use but not for full allotted time	52	41.3%		
Not able to use	27	21.4%		
Changes in receipt of legal aid				
More aid	5	2.6%		
Less aid	155	79.1%		
No change	36	18.4%		
Changes in parole hearings				
Paused/delayed	50	36.0%		
Limited access to parole hearings	38	27.3%		

Remote/full	12	8.6%
No change	39	28.1%

<sup>a</sup>Excluding "don't know" and "other" answers

<sup>b</sup>Binary Y/N question. Yes was used as reference and reported. Responses are not mutually exclusive.

<sup>c</sup>*Changes in mental healthcare were collected among those who were currently receiving mental healthcare.* 

N = 378	n	<b>%</b> <sup>a</sup>		
Questions for Formerly Incarcerated Respondents Only				
Month Released				
March 2020 - May 2020	10	47.6%		
June 2020 - August 2020	8	38.1%		
September 2020 - November 2020	3	14.3%		
Received COVID-19 facility updates	3	16.7%		
Frequency of PPE replacement				
Not daily but more than once per week	2	18.2%		
Less than once per week	9	81.8%		
Worried about getting COVID-19				
Most of the time	14	87.5%		
Sometimes/Not much/none	2	12.5%		
Added stress because of COVID-19				
A lot of added stress	15	93.8%		
Some stress/A little/no added stress	1	6.3%		
Decreased financial support <sup>a</sup>	10	62.5%		
Interested in getting a vaccine <sup>a</sup>	9	64.3%		
Questions for Proxy Respondents Only				

# Table 3. Questions Specifically for Formerly Incarcerated or Proxy Respondents

Last communication with incarcerated person

Less than 1 week ago	303	84.9%
Between 1-2 weeks ago	33	9.2%
Between 2 weeks to one month ago	13	3.6%
More than one month ago	8	2.2%
Concerned about contracting COVID-19 from incarcerated individual if released	126	49.8%
Concerned about getting COVID-19 if facility opened visitation	169	70.1%
Which of the following would proxy need to feel safe from contracting COVID-19 from incarcerated individual? <sup>b</sup>		
A rapid COVID test within a week	86	24.1%
Knowing that they had been tested in past month	27	7.6%
Place in your home where they could quarantine	52	14.6%
Increased expenses in order to support incarcerated	185	81.9%
person		

<sup>a</sup>Yes was used as reference and reported.

<sup>b</sup>Binary Y/N question. Yes was used as reference and reported. Responses are not mutually exclusive.

Question	Proxy Respondents	Formerly Incarcerated Respondents	Two-sided Fisher's exact test p-value
Facility on any type of lockdown	94.4% (320)	79.0% (15)	0.03
Hours spent in lockdown (>20 hours/day)	86.2% (213)	93.3% (14)	0.70
Lockdown time period exceeding 3 months	49.6% (137)	33.3% (5)	0.29
Ability to maintain 6-foot distance at all times	7.8% (24)	10.0% (2)	0.67
Routine disinfection of commonly used surfaces	40.6% (95)	52.9% (9)	0.32
Adequate access to toilet paper	50.3% (90)	63.2% (12)	0.34
Ability to shower every day, if wanted	30.0% (77)	63.2% (12)	0.01
Adequate access to water	53.5% (114)	57.9% (11)	0.81
Enough free soap for needs	10.3% (23)	26.3% (5)	0.05
Isolated/Quarantined from cellmates if experiencing COVID-19 symptoms	59.2% (142)	45.5% (5)	0.37
Longer waits than normal to access medical care	81.2% (177)	82.4% (14)	1.00
Increased access to mental healthcare	4.1% (6)	11.1% (1)	0.35

# **Table 4. Internal Validity Results**

All responses are dichotomized; values represent responses to "yes."