

Acceptance of COVID-19 vaccine among healthcare workers in Katsina state, Northwest Nigeria

Ahmed Tijani Abubakar^{1,2} Kabir Suleiman³ Ahmad Suleiman Idris^{*4,5} Shamsuddeen Yahaya Suleiman³ Umar Bello Ibrahim⁶ Suleiman Bello Abdullahi^{4,7} Ahmed Suleiman Haladu⁵ Ahmad Ibrahim Al-Mustapha^{2,8} Musa Imam Abubakar⁹

¹African Center for Disease Prevention and Control, Abuja, Nigeria

²Department of Veterinary Services, Kwara State Ministry of Agriculture and Rural Development, Ilorin, Kwara State, Nigeria

³Katsina State Primary Healthcare Agency

⁴Nigeria Field Epidemiology and Laboratory Training Program

⁵Africa Field Epidemiology Network

⁶Katsina State Ministry of Health

⁷Federal Medical Centre Katsina

⁸Faculty of Pharmaceutical Sciences, Universite de Tours, Tours, France

⁹REDISSE Project, Federal Ministry of Agriculture and Rural Development, Abuja, Nigeria

*Corresponding Author: Dr Ahmad Suleiman Idris

Abstract

High acceptance of COVID-19 vaccines is crucial to ending the COVID-19 pandemic. Healthcare workers (HCWs) are frontline responders in the fight against COVID-19, they were prioritized to receive the COVID-19 vaccine in Nigeria. This study assessed the acceptance of the COVID-19 vaccine among HCWs in Katsina State using an online structured questionnaire and predicted variables that could increase the acceptance of the vaccine among HCWs using logistic regression analysis.

A total of 793 HCWs were included in this study. Of these, 65.4% (n=519) of them were male and 36.2% (n=287) were aged between 30-39 years. Eighty percent (80%, n=638) of the HCWs have been tested for the SARS-CoV-2 out of which 10.8% (n=65) of them tested positive. The majority of the HCWs (97.3%, n=765) believed that the COVID-19 vaccine was safe and 90% (n=714) of the HCWs have received the first dose of the COVID-19 vaccine. Our findings showed that the age of the HCW, their COVID-19 testing status, and the type of health facility they work (either public or private) were the main predictors for the acceptance of the COVID-19 vaccine among HCWs in Katsina State. HCWs between the age of 30-39 years were more likely (OR:7.06; 95% CI: 2.36, 21.07; $p < 0.001$) to accept the vaccine than others. In the same vein, HCWs that have been tested for COVID-19 were more likely (OR:7.64; 95% CI: 3.62, 16.16; $p < 0.001$) to accept the vaccine than those that have not been tested. In addition, HCWs in public health facilities were more likely (OR: 2.91; 95% CI: 1.17, 6.11; $p = 0.094$) to accept the COVID-19 vaccine than their counterparts in private HFs.

There was a high acceptance of the COVID-19 vaccines among HCWs in Katsina State. More emphasis should be paid on adherence to non-pharmaceutical interventions and availability of vaccines for HCWs in private hospitals.

Keywords: COVID-19, HCWs, Vaccine Acceptance, Nigeria.

Introduction

The coronavirus disease 2019 (COVID-19) is a major public health threat that has affected over 220 countries worldwide. As of June 15th 2021, Nigeria has recorded 167, 078 COVID-19 cases and 2,117 COVID-19 related deaths out of the 176 million cases and 3.8 million deaths recorded globally [1,2].

To curb the spread of the virus, save lives, and flatten the epidemic curve, several public health measures such as social distancing, wearing of face mask, frequent hand washing, avoidance of crowded places, self-isolation, and quarantine were instituted globally. However, these measures were not sufficient [3]. Thus, humanity had the urgent need to establish herd immunity through alternative interventions such as vaccinations [4,5]. The collaborative effort of the international scientific community led to the rapid development of effective and safe vaccines against COVID-19.

Nigeria (with a population of 214 million inhabitants) prioritized the frontline healthcare workers (HCWs), elder statesmen, and the elderly persons (>60 years) as the first recipients of the 4 million doses of the AstraZeneca COVID-19 vaccine delivered to the country [6]. The uptake of the COVID-19 vaccine has however faced challenges that ranged from vaccine hesitancy and lack of trust in vaccine safety and effectiveness even among HCWs [3,7]. HCWs are crucial to Nigeria's primary healthcare system as they provide reliable information about vaccines and vaccine-preventable diseases especially at the grassroots [8,9]. Hence, their attitude towards the COVID-19 preventive practices and their acceptance of the COVID-19 vaccine could influence the acceptance of the vaccine by the general population [10,11].

Katsina state has recorded 2,011 COVID-19 positive cases and 35 COVID-19 associated deaths since its index case on the 7th of April 2020. The highest daily COVID-19 incidence in Katsina state was 70 (17th December 2020) [1]. The state received 107,504 doses of the COVID-19 vaccine and has since started vaccinating its priority individuals [12]. To the best of our knowledge, this is the first study evaluating the acceptance of the COVID-19 vaccine among HCWs in Northern Nigeria. Hence, this study assessed the acceptance of the COVID - 19 vaccine among HCWs in Katsina State, Nigeria.

Materials and Methods

Study Area

The study was conducted in Katsina State, Northwestern Nigeria. The state has 33 local government areas and a population of 5,801,584 inhabitants [13]. The state is mostly dominated by Hausa and Fulani ethnic tribes.

Survey methodology

The survey instrument was designed to assess the acceptance of the COVID-19 vaccine by HCWs in Katsina State. The questionnaire had two sections: Section A focused on the demographics of the HCWs whereas Section B assessed the acceptance of the vaccine by HCWs. The survey instrument was validated by two independent academic reviewers to examine its content and face validity as well as its ease of administration. The questionnaire was administered as an electronic survey (via open data kit) as part of the COVID-19 operational research in Katsina State between 1st and 22nd May 2021. Participants were recruited by the non-probability sampling (snowballing) technique. We shared the questionnaire via social media platforms such as WhatsApp, Twitter, etc. to capture all the health facilities registered on the DHIS2 (except a few that could not be reached).

The inclusion criteria were that the participant had to be a healthcare worker. Since there were no published studies that evaluated COVID-19 vaccine acceptance in Katsina state, we assumed that 50% of the HCWs would accept the vaccine. Hence, a sample size of at least 384 respondents was needed. Study participants were selected by simple random sampling based on their willingness to participate in the study.

Statistical analysis

The data generated from this study were analyzed using Statistical Package for Social Sciences v. 26 (Armonk, New York, USA). Descriptive statistics of the data were presented as frequencies and proportions for qualitative data as well as the mean and standard deviation for quantitative data. We conducted a univariable logistic regression analysis to evaluate the effect of the independent variables (socio-demographic variables) on the binary outcome variable (acceptance of the COVID-19 vaccine). All the significant variables at $p \leq 0.05$ were

retained and included in the final model of the multivariable logistic regression analysis using the logit function and α at 0.25.

Ethical approval

The ethical approval of this study was obtained from the Katsina State Ministry of Health (Reference number: MOH/KS/EHC/77/98). The objective of the study was briefly explained to the participants before a written consent form was administered to each participant. Only participants that consented were included in this survey. Participants were free to withdraw from the study at any point and no personal identifying information was obtained from any of the study participants.

Results

A total of 793 HCWs were included in this study. Of these, 519 of them (65.4%) were male and 36.2% (n=287) were aged between 30-39 years. Most of the respondents (87.5%, n=694) were from public health facilities (Table 1). The majority of HCWs complied and practiced the laid down COVID-19 preventive measures such as regular handwashing with soap and water or using hand sanitizers. However, only around $\frac{3}{4}$ of the HCWs frequently used nose masks, maintained social distancing, and avoided crowded places (Table 2).

Table 1. Structure of health care workers who participated

Eighty percent (80%, n=638) of the HCWs in both the public and private health facilities (HFs) have been tested for the COVID-19 disease either via polymerase chain reaction (PCR) or via rapid diagnostic test kits (RDT kits). Of the HCWs that have been screened for the COVID-19, 10.8% (n=65) tested positive and were managed appropriately. The majority of the HCWs (97.3%, n=765) believed that the AstraZeneca COVID-19 vaccine received by Nigeria (from the global COVAX system) was safe and some of them (74.7%, n=573) opined that the vaccine should be mandated on all Nigerians. Ninety percent (n=714) of all HCWs who participated in this survey have received the first dose of the AstraZeneca COVID-19 vaccine. Only 4% (n=32) of all respondents were not satisfied with the government's handling of the pandemic.

Table 2. Vaccine acceptance among HCWs in Katsina State (n=793)

Our findings showed that the age of the HCW, their COVID-19 testing status, and the type of HF were the main predictors for the acceptance of the Covid-19 vaccine among HCWs in Katsina state. Older HCWs between the age of 30-39 were more likely (OR:7.06; 95% CI: 2.36, 21.07; $p < 0.001$) to accept the vaccine than others. A similar higher likelihood ratio was predicted for other groups as against young HCWs between 20-29 years. In the same vein, HCWs that have been tested for COVID-19 were more likely (OR:7.64; 95% CI: 3.62, 16.16; $p < 0.001$) to accept the vaccine than those that have not been tested. In addition, HCWs in public HFs were more likely (OR: 2.91; 95% CI: 1.17, 6.11; $p = 0.094$) to accept the COVID-19 vaccine than their counterparts in private HFs.

Table 3. Logistic regression analysis of the factors that could predict the acceptance of the AstraZeneca COVID-19 vaccine among HCWs in Katsina State, Nigeria.

Discussion

For centuries, public health interventions such as vaccination have been crucial to the prevention and control of infectious diseases [14-16]. The arrival of the COVID-19 vaccine into Nigeria marked another phase in the control of the pandemic. Our findings showed that the majority of HCWs complied and practiced the laid down COVID-19 preventive measures. HCWs complied more with regular handwashing with soap and water or using hand sanitizers as against other COVID-19 preventive practices (Table 2).

As the pandemic fatigue worsens, only around 75% of the HCWs frequently used nose masks, maintained social distancing, and avoided crowded places. This fatigue could have been further complicated by the availability of vaccines among other factors [17]. However, HCWs should strictly comply and adhere to the COVID-19 preventive practices despite the availability of COVID-19 vaccines. Moore et al., [18] and Yu et al., [19] reported that non-pharmaceutical interventions are still needed to control the pandemic despite the global COVID-19 vaccine rollout.

Our findings showed that the prevalence of COVID-19 among HCWs in Katsina State was 10.8%. However, it remains to be ascertained whether HCWs contracted the SARS-CoV-2 via community transmission or through contact with infected patients in their respective HFs. Reports by Alasia and Maduka, [20] as well as Ogboghodo *et al.*, [21] had reported a higher (15.3%) and lower (5.3%) COVID-19 prevalence rate among HCWs in Rivers and Edo states of Nigeria respectively.

Our study showed that 90% of HCWs in Katsina state have received the first dose of the vaccine. This is higher than 77% and 51% vaccine uptake reported by Maltezou *et al.*, [11] among Physicians and Nursing staff respectively in Greece. The high vaccine uptake may not be unconnected to the perception of the vaccine safety by HCWs in Katsina state as 93.7% of the HCWs believed the vaccine is safe and effective. The high uptake of the vaccine by the HCW will play a great role in increasing acceptance and uptake of the vaccine by the general population since HCWs serve as trusted source of information to the general public and their advice can be a determinant of the patient's decision to take the vaccine [3]. Similarly, HCWs highly rated the government's handling of the pandemic as only 4% were not satisfied with the government's handling of the pandemic. These high ratings coupled with the good perception of the COVID-19 vaccine could be responsible for the high acceptance rate [22].

Our findings showed that the main predictors of increased acceptance of the COVID-19 vaccine among HCWs in Katsina were age, COVID-19 testing status, and the type of HF. HCWs aged between 30-39 years were likely to accept the vaccine than others. This may be due to the higher percentage of this age bracket among the health sector workforce in Nigeria. A similar trend was observed by Shekhar *et al.*, [23] who reported that COVID-19 vaccine acceptance increased with increasing age.

Similarly, an increased acceptance of the COVID-19 vaccine could be predicted by the testing status of HCW. Eighty percent (80%) of the study participants have been tested by either PCR or RDT kits. The willingness to get tested reflects the level of education and knowledge of the HCWs, which are additional factors that could influence vaccine acceptance by HCW. This agrees with the findings of Maltezou *et al.*, [11] and Shekhar *et al.*, [23]. Vaccine acceptance was lower in private HFs compared to public HF. This may be due to the increased risk perceptions of COVID-19 among HCW in public HF compared to those in private HF. Perception of risk is significant for vaccine decision-making [3]. Another factor that may explain the disparity in vaccine uptake among HCW in public and private HF is the prioritization of vaccines to the frontline HCW which are majorly found in the public HF where COVID-19 cases are mostly managed, and vaccines are administered. It is however imperative to increase vaccine uptake among HCW in private HF by including them among those prioritized for vaccination and making the private HF a site of vaccination for the public. This will generally increase the number of vaccine uptake among HCW, prevent infection and reduce loss of essential workforce to infection and death [23]. By extension, it will have a positive impact on vaccine uptake among the public because of the strong association identified by Shekhar *et al.*, [23] among HCW who may intend to get vaccinated and recommend the vaccine to others.

The main strength of the study was the sample size and the timeliness of this survey. The main limitations were the sampling bias because some HFs could not access the online survey.

Conclusion

This study reports the high acceptance of the COVID-19 vaccine among HCWs in Katsina State, Northwest Nigeria. The main predictors of an increased vaccine acceptance among HCWs were the age and the testing status of the HCW. HCWs must adhere more to COVID-19 preventive practices.

Declarations

Acknowledgment: We acknowledge acknowledge Dr Muftau Oyewo for reviewing the manuscript.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflicts of Interest: The authors declare that they have no competing interests.

Author Contributions: Authors contributed equally to the conception, analysis, and writing of the manuscript.

Data Availability: The dataset generated during the survey is available on reasonable request.

REFERENCES

1. Nigeria Centre for Disease Control (2021). NCDC Coronavirus Disease (COVID -19) Dashboard. Available at: <https://covid19.ncdc.gov.ng/>. Accessed June 16, 2021
2. World Health Organization (2021). WHO Coronavirus Disease (COVID-19) Dashboard. Geneva: World Health Organization. Available at: <https://covid19.who.int/>. Accessed June 16, 2021.
3. Nzaji, M. K., Ngombe, L. K., Mwamba, G. N., Ndala, D. B. B., Miema, J. M., Lungoyo, C. L., Mwimba, B. L., Bene, A. C. M. and Musenga, E. M. (2020). Acceptability of vaccination against COVID -19 among health care workers in the Democratic Republic of the Congo. *Pragmatic and Observational Research*, 11: 103 – 109.
4. Fu, C., Wei, Z., Pei, S., Li, S., Sun, X. and Liu, P. (2020). Acceptance and preference for COVID-19 vaccination in health-care workers (HCWs). *medRxiv*, 2962(548).
5. Paterson, P., Paterson, P., Meurice, F., Stanberry, L.R., Glismann, S. and Rosenthal, S.L. (2016). Vaccine hesitancy and healthcare providers vaccine hesitancy and healthcare providers. *Vaccine*, 34:6700–6706.
6. World Health Organization, Africa (2021). COVID – 19 vaccines shipped by COVAX arrive Nigeria. Available at: <https://www.afro.who.int/news/covid-19-vaccines-shipped-covax-arrive-nigeria>. Accessed June 29, 2021
7. Macdonald, N.E. (2015) Vaccine hesitancy: definition, scope and determinants. *Vaccine*, 33: 4161–4164.
8. Danchin, M., Biezen, R., Manski-Nankervis, J.A., Kaufman, J. and Leask, J. (2020). Preparing the public for COVID-19 vaccines: how can general practitioners build vaccine confidence and optimise uptake for themselves and their patients? *Australian Journal of General Practice*, 49:625-9
9. Maltezou, H.C., Theodora, M., Lytras, T., Fotiou, A., Nino, E. and Rodolakis, A. (2020). Knowledge, attitudes and practices about vaccine-preventable diseases and vaccinations of children among pregnant women in Greece. *Vaccine*, 38:7654-8.
10. Agyekum, M. W., Afrifa-Anane, G. F., Kyei-Arthur, F. and Addo, B. (2021). Acceptability of COVID – 19 vaccination among health care workers in Ghana. *Advances in Public Health*, 2021.
11. Maltezou, H. C., Pavli, A., Dedoukou, X., Georgakopoulou, T., Raftopoulos, V., Drositis, I., Bolikas, E., Ledda, C., Adamis, G., Spyrou, A., Karantoni, E.,

- Gamaletsou, M. N., Koukou, D. M., Lourida, A., Moussas, N., Petrakis, V., Panagopoulos, P., Hatzigeorgiou, D., Theodoridou, M., Lazanas, M., Gargalianos, P. and Sipsas, N. V. (2021). Determinants of intention to get vaccinated against COVID – 19 among healthcare personnel in hospitals in Greece. Research paper available at <http://www.journals.elsevier.com/infectiondisease-and-health/>. Accessed June 16, 2021.
12. The Guardian. Katsina, three others get 276,564 doses of COVID-19 vaccine. Guardian.ng. 2021 [cited 5 August 2021]. Available from: <https://guardian.ng/news/katsina-three-others-get-276564-doses-of-covid-19-vaccine/>.
 13. Katsina State Government. Katsina State [cited 5 July 2021]. Available from: <https://www.katsinastate.gov.ng/about-katsina/>.
 14. Wilder-Smith A, Longini I, Zuber PL, Bärnighausen T, Edmunds WJ, Dean N, Spicher VM, Benissa MR, Gessner BD. The public health value of vaccines beyond efficacy: methods, measures and outcomes. *BMC medicine*. 2017 Dec;15(1):1-9.
 15. Paul E, Steptoe A, Fancourt D. Attitudes towards vaccines and intention to vaccinate against COVID-19: Implications for public health communications. *The Lancet Regional Health-Europe*. 2021 Feb 1;1:100012.
 16. Folb PI, Bernatowska E, Chen R, Clemens J, Dodoo AN, Ellenberg SS, Farrington CP, John TJ, Lambert PH, MacDonald NE, Miller E. A global perspective on vaccine safety and public health: the Global Advisory Committee on Vaccine Safety. *American journal of public health*. 2004 Nov;94(11):1926-31.
 17. GAVI. 10 reasons why pandemic fatigue could threaten global health in 2021 [Internet]. Gavi.org. 2021 [cited 5 August 2021]. Available from: <https://www.gavi.org/vaccineswork/10-reasons-why-pandemic-fatigue-could-threaten-global-health-2021>.
 18. Moore S, Hill EM, Tildesley MJ, Dyson L, Keeling MJ. Vaccination and non-pharmaceutical interventions for COVID-19: a mathematical modelling study. *The Lancet Infectious Diseases*. 2021 Jun 1;21(6):793-802.
 19. Liu Y, Morgenstern C, Kelly J, Lowe R, Jit M. The impact of non-pharmaceutical interventions on SARS-CoV-2 transmission across 130 countries and territories. *BMC medicine*. 2021 Dec;19(1):1-2.
 20. Alasia DD, Maduka O. Prevalence and Pattern of COVID-19 among Healthcare Workers in Rivers State Nigeria. *Occupational Diseases and Environmental Medicine*. 2021 Jan 7;9(1):20-32.
 21. Ogboghodo EO, Osaigbovo II, Obarisiagbon OO, Okwara BU, Obaseki DE, Omo-Ikrodah OT, Ehinze ES, Adio F, Nwaogwugwu JC, Esegibe EF. Facility-Based Surveillance Activities

- for COVID-19 Infection and Outcomes among Healthcare Workers in a Nigerian Tertiary Hospital. *The American Journal of Tropical Medicine and Hygiene*. 2021 Mar;104(3):1034.
22. Lazarus JV, Ratzan SC, Palayew A, Gostin LO, Larson HJ, Rabin K, Kimball S, El-Mohandes A. A global survey of potential acceptance of a COVID-19 vaccine. *Nature medicine*. 2021 Feb;27(2):225-8.
23. Shekhar R, Sheikh AB, Upadhyay S, Singh M, Kottewar S, Mir H, Barrett E, Pal S. COVID-19 vaccine acceptance among health care workers in the United States. *Vaccines*. 2021 Feb;9(2):119.

Appendices
 Appendix 1
 Tables

Table 1. Structure of health care workers who participated

Variable	Frequency (%)
Age (years)	
<20	22 (2.8)
20-29	213 (26.8)
30-39	287 (36.2)
40-49	190 (24)
50-59	72 (9.1)
>60	9 (1.1)
Gender	
Female	274 (34.6)
Male	519 (65.4)
Designation	
Health Information technology	27 (3.4)
Medical laboratory scientist/technicians	89 (11.2)
Medical student	117 (14.8)
Midwives	73 (9.2)
Nurse	160 (20.1)
Community health workers	215 (27.1)
Physician	92 (11.6)
Health facility	
Private health facility	99 (12.5)
Public health facility	694 (87.5)

Table 2. Vaccine acceptance among HCWs in Katsina State (n=793).

Variable	Frequency (%)
1. Do you practice any of the following?	
a) Regular handwashing	764 (96.3)
b) Frequent use of nose masks	601 (75.8)
c) Social distancing	623 (78.6)
d) Avoid crowded places	641 (80.8)
2a. Have you been tested for the COVID-19?	
No	155 (19.6)
Yes	638 (80.4)
2b. If yes in 2 above, was it positive?	
No	573 (89.2)
Yes	65 (10.8)
3. Do you think we need a COVID-19 vaccine to curb the pandemic in Nigeria?	
No	26 (3.3)
Yes	767 (96.7)
4. Have you taken the COVID-19 vaccine?	
No	79 (10)
Yes	714 (90)
5. Do you think that the COVID-19 vaccine brought to Nigeria is safe?	
No	21 (2.7)
Yes	765 (97.3)
6. Do you think that the COVID-19 vaccine should be mandated on all Nigerians?	
Maybe	171 (21.6)
No	29 (3.7)
Yes	593 (74.7)
7. What is your rating of the government's handling?	
Poor	8 (1.0)
Fair	24 (3.0)
Average	91 (11.5)
Good	211 (26.6)
Excellent	459 (57.9)

Table 3. Logistic regression analysis of the factors that could predict the acceptance of the AstraZeneca COVID-19 vaccine among HCWs in Katsina State, Nigeria.

Variable	Referent		Univariable analysis				Multivariable analysis			
			Odds Ratio (95% CI)	X^2	DF	<i>P-value</i>	Adjusted Odds Ratio (95% CI)	X^2	DF	<i>P-value</i>
Age	20-29	30-39	7.68 (2.61, 22.52)	25.91	4	<0.001	7.06 (2.36, 21.07)	20.57	4	<0.001
		40-49	6.72 (1.99, 22.88)				5.05 (1.46, 17.49)			
		50-59	2.50 (0.73, 8.56)				1.75 (0.49, 6.30)			
		>60	0.87 (0.10, 7.25)				1.01 (0.11, 9.49)			
Have you been tested for COVID-19?	No	Yes	8.63 (4.17, 17.87)	34.93	1	<0.001	7.64 (3.62, 16.16)	29.59	1	<0.001
		Tested positive to COVID-19	Yes				2.15 (0.28, 16.26)			
Gender	Male	Female	1.52 (0.76, 3.05)	1.39	1	0.239	-	-	-	-
Type of health facility	Private HFs	Public HFs	2.67 (1.21, 5.91)	5.10	1	0.024	2.91 (1.17, 6.11)	5.27	1	0.097

Table 4 Logistic regression analysis of the factors that could predict the completion of the *Kanamycin (KANTH-04)* course among HIV-1 in Kaduna State, Nigeria

Variable	Reference		Univariable analysis				Multivariable analysis			
			Odds Ratio (95% CI)	OR	95%	P-value	Adjusted Odds Ratio (95% CI)	OR	95%	P-value
Age	20-29	30-39	1.00(1.04, 21.02)	21.04	1	<0.001	0.99(1.24, 11.81)	20.17	1	<0.001
		40-49	0.65(1.05, 21.98)				2.05(1.45, 17.93)			
		50-59	2.50(0.21, 3.20)				1.15(0.75, 6.88)			
		60-69	0.17(0.28, 1.22)				1.33(0.21, 20.85)			
Have you been treated for TB (KANTH-04)	No	Yes	5.71(4.17, 37.82)	14.04	1	<0.001	7.08(3.47, 14.15)	20.19	1	<0.001
Treated prior to KANTH-04	No	Yes	0.14(0.08, 0.24)	0.49	1	<0.001	-	-	-	-
Gender	Male	Female	1.81(0.78, 2.49)	1.25	1	<0.205	-	-	-	-
Type of health facility	Private Lab	Public Lab	2.87(1.21, 3.92)	2.30	1	<0.028	1.91(1.10, 4.18)	2.27	1	<0.001

Table 7. Mean responses given using WOTWOT (Online Survey) (n=200)

Variable	Frequency (%)
1. Do you practice any of the following?	
a) Regular handwashing	774 (78.7%)
b) Frequent use of face masks	603 (61.3%)
c) Social distancing	625 (63.5%)
d) Avoid crowded places	688 (70.4%)
2a. Have you been tested for the COVID-19?	
No	117 (11.8%)
Yes	888 (90.2%)
2b. If yes, in 2 weeks, was it positive?	
No	101 (10.2%)
Yes	66 (6.7%)
3. Do you think we need a COVID-19 vaccine to curb the pandemic in Nigeria?	
No	28 (2.8%)
Yes	767 (78.7%)
4. How you value the COVID-19 vaccine?	
No	78 (7.9%)
Yes	714 (73%)
5. Do you think that the COVID-19 vaccine brought in Nigeria is safe?	
No	28 (2.8%)
Yes	767 (78.7%)
6. Do you think that the COVID-19 vaccine should be mandated on all Nigerians?	
Maybe	118 (11.9%)
No	28 (2.8%)
Yes	649 (66.3%)
7. What is your rating of the government's handling?	
None	26 (2.6%)
Very	28 (2.8%)
Average	86 (8.7%)
Good	218 (22.2%)
Excellent	428 (43.7%)

Table 1. Structure of health care workers who participated

Variable	Frequency (%)
Age (years)	
<20	22 (2.8)
20-29	213 (26.8)
30-39	287 (36.2)
40-49	190 (24)
50-59	72 (9.1)
>60	9 (1.1)
Gender	
Female	274 (34.6)
Male	519 (65.4)
Designation	
Health Information technology	27 (3.4)
Medical laboratory scientist/technicians	89 (11.2)
Medical student	117 (14.8)
Midwives	73 (9.2)
Nurse	160 (20.1)
Community health workers	215 (27.1)
Physician	92 (11.6)
Health facility	
Private health facility	69 (8.7)
Public health facility	694 (87.5)