

Socioeconomic status and immune aging in older US Adults in the Health and Retirement Study

Eric T. Klopack, Ph.D.*
Leonard Davis School of Gerontology
University of Southern California
klopack@usc.edu
<https://orcid.org/0000-0002-0019-1332>

Bharat Thyagarajan, Ph.D.
Department of Laboratory Medicine and Pathology
University of Minnesota
thya0003@umn.edu

Jessica D. Faul, Ph.D.
Survey Research Center
University of Michigan
jfaul@umich.edu

Helen C. S. Meier, Ph.D.
Survey Research Center
University of Michigan
hspink@umich.edu

Ramya Ramasubramanian, M.P.H.
Division of Epidemiology and Community Health
University of Minnesota School of Public Health
ramas063@umn.edu

Jung Ki Kim, Ph.D.
Leonard Davis School of Gerontology
University of Southern California
jungk@usc.edu

Eileen M. Crimmins, Ph.D.
Leonard Davis School of Gerontology
University of Southern California
crimmin@usc.edu

*Corresponding author

Funding Statement

Support was provided by grants from the National Institute on Aging (P30AG017265 and R01AG060110).

The Health and Retirement Study is supported by the National Institute on Aging (NIA U01AG009740) and the Social Security Administration with grants to the University of Michigan.

Author Contributions

EK conceived the study, completed the analyses, and wrote the initial draft. BT helped in interpretation of findings and editing the manuscript. JF helped in interpretation of findings and editing the manuscript. HM helped in interpretation of findings and editing the manuscript. RR helped in development of measures. JK helped with coding and data visualization. EC conceived the study, developed the data, secured the funding, supervised the study, helped in interpretation of the findings, and helped in writing.

Conflict of Interest Disclosure

The authors report no conflict of interest.

Ethics Approval Statement

This research was approved by the University of Southern California IRB.

Data Availability Statement

All data are publicly available: <https://hrs.isr.umich.edu/>

Abstract

Background: Life course socioeconomic and demographic factors including educational attainment, race and ethnicity, and childhood (SES) are very powerful predictors of large inequalities in aging, morbidity, and mortality. Immune aging, including accumulation of late-differentiated, senescent-like lymphocytes and lower level of naïve lymphocytes, may play a role in the development of the age-related health inequalities. However, there has been little research investigating association between socioeconomic status and immune aging, particularly age-related changes in lymphocyte percentages.

Methods: This study used nationally representative data from more than 9000 US adults from the Health and Retirement Study to investigate associations between educational attainment, race and ethnicity, and childhood SES and lymphocyte percentages.

Results: Respondents with lower educational attainment, Hispanic adults, and those who had a parent with less than high school education had lymphocyte percentages consistent with more highly accelerated immune aging compared to those with greater educational attainment, non-Hispanic White adults, and respondents who had parents with a high school education, respectively. Associations between education, Hispanic ethnicity, and parents' education and late differentiated senescent-like T lymphocytes (TemRA) and B cells were largely driven by cytomegalovirus (CMV) seropositivity.

Conclusions: Results suggest that CMV is a major driver of observed SES inequalities in immunosenescence and may therefore be an important target for interventions. Naïve T lymphocytes may be particularly affected by socioeconomic position and may therefore be of particular interest to research interested in inequalities in health and aging.

Keywords: race and ethnicity; lymphocytes; cytomegalovirus; immunosenescence

Introduction

Age-related changes in immune function has been associated with increased risk cardiovascular disease, cancer, and other chronic conditions as well as reduced efficacy in fighting acute infections and reduced response to vaccines¹⁻⁵. The aging immune system is characterized by lower percentages of naïve T and B lymphocytes, higher percentages of late differentiated lymphocytes, increases in cytotoxic (CD8⁺) relative to helper (CD4⁺) T cells, and decreases in naïve helper T cells relative to memory helper T cells. As the thymus involutes with age, fewer naïve cells are available to respond to novel viral and vaccine exposures⁶. Late differentiated cells accumulate over time and are less effective in viral response and other senescent cell and cancerous cell clearance and produce inflammatory cytokines and display a senescence-associated secretory phenotype⁷⁻¹⁰. Though other immune cells are involved in the immune aging, past research suggests that lymphocytes—and T lymphocytes in particular—are most affected by age-related immune changes^{6,11}. Though age is a very powerful predictor of these changes⁴, there is substantial individual variation within age groups. Uncovering the sources of this variance is important for understanding and combating age-related health problems and functional decline.

A robust body of research links socioeconomic status (SES; e.g., wealth, education, income, occupational prestige) with various age-related health outcomes generally¹² and with immune dysfunction particularly¹³. Indeed, past research suggests that lower income is associated with age-related cell type percentages¹⁴. Education may affect immune cell percentages by increasing exposure to health risks like social stressors that can accelerate immunosenescence^{2,15}.

Lower educational attainment is also associated with increased risk of human cytomegalovirus infection (CMV) infection^{13,16}. CMV is a common herpesvirus that is typically asymptomatic. CMV remains latent (i.e., once it is acquired, it is never cleared) and can be reactivated throughout life. This ongoing exposure to low levels of CMV infection results in an increasing proportion of the immune cells being dedicated to the management of CMV, leading to the exhaustion of naïve lymphocytes and the accumulation of memory T cells devoted to the virus (so called memory inflation)^{8,17,18}. Thus, CMV infection is a strong determinant of cell percentages that characterize an aging immune system.

Race and ethnicity are highly related to SES in the US and are also robustly associated with age-related health problems¹⁹. Most past research on immune aging has utilized small or all White samples, making comparisons across race and ethnicity difficult. There has been mixed evidence from research comparing European and African populations²⁰ and comparing White, Black, and Hispanic adults in the US²¹ regarding racial and ethnic differences in CD4⁺ and CD8⁺ T cell percentages, and to our knowledge, no research focusing on racial and ethnic differences in cell types specifically relevant to the immune aging (e.g., naïve and late differentiated T and B cells). However, there is good reason to expect that non-White and Hispanic US adults would have percentages of these cells indicative of accelerated immune aging. Black and Hispanic US adults have higher CMV seroprevalence and higher CMV antibody levels among CMV seropositive individuals^{13,22}. Racial and ethnic minority groups in the US are structurally disadvantaged and therefore are exposed to more environmental toxins and social stressors and have limited access to socioeconomic opportunity¹⁹, all of which are associated with the immune age phenotype^{23,24}. Racial and ethnic minorities are also exposed to particular stressors like racial discrimination that over time weather individuals²⁵, leading to immune dysfunction and

accelerated aging^{26,27}. We therefore expect that non-White US adults will have lymphocyte percentages more typical of accelerated immune aging.

SES in childhood is also a major determinant of adult health, independent of adult SES²⁸⁻³⁰. Low childhood SES is associated with adverse health behaviors^{31,32}, greater adversity and toxic stress³³, lower adult SES attainment^{34,35}, and may directly alter developmental trajectories toward a more accelerated aging phenotype^{36,37}. Past research in small samples has found that early life adversity increases risk of CMV infection and T cell senescence¹⁷.

Though understanding the drivers of the immune age phenotype may be important for understanding health changes with age, relatively little research has investigated the roles of socioeconomic factors and race and ethnicity in immune aging. What research has been conducted on SES and immune age phenotype generally has utilized limited community samples. This study builds on this past research by utilizing nationally representative data on over 9000 US adults from the Health and Retirement Study. This study represents the first time that detailed immunophenotyping has been conducted in a large US population representative sample of older adults.

We investigated first whether educational attainment, race/ethnicity, and childhood SES (assessed by parental education) were associated with percentages of age-related lymphocytes: percentages of naïve CD4⁺ and CD8⁺ T cells, late differentiated (so called TemRA) CD4⁺ and CD8⁺ T cells, the CD4:CD8 cell ratio, the CD4 naïve:CD4 total memory cell ratio, naïve B cells, and late memory B cells. We expect higher educational attainment, being non-Hispanic White, and higher parental education to be associated with higher percentages of naïve lymphocytes, lower percentages of late differentiated (e.g., TemRA and late memory) lymphocytes, and higher

CD4:CD8 and CD4 naïve:CD4 total memory cell ratios. We next examined whether these effects differed by CMV seropositivity. Finally, we examined potential interactions among these factors.

Methods

Sample

This study utilizes the Health and Retirement Study (HRS) 2016 Venous Blood Study (N = 9,934). HRS is a national panel survey of older Americans, which when weighted is designed to be representative of community dwelling older Americans older than 50. The VBS consisted of panel members interviewed before 2016 at least once who consented to a venous blood draw. 78.5% of eligible respondents consented and 82.9% of those respondents complete a blood draw. Sample characteristics are described in the results section. More information about this study is available elsewhere³⁸. Absolute counts and percentages of 24 different types of immune cells were assessed using Multiparameter flow cytometry using the standardized protocol by the Human Immunology Project³⁹ with minor modifications performed on an LSRII or a Fortessa X20 flow cytometer (BD Biosciences, San Diego, CA) More detailed methods for this sample are published elsewhere^{38,40,41}.

Measures

Lymphocyte percentages were assessed following the protocol described above. In the current study, we focus on the percentage of naïve CD4⁺ (CD4⁺/CD3⁺/CD19⁻/CD45RA⁺/CCR7⁺/CD28⁺) and CD8⁺ cells (CD8⁺/CD3⁺/CD19⁻/CD45RA⁺/CCR7⁺/CD28⁺), TemRA CD4⁺ (CD4⁺/CD3⁺/CD19⁻/CD45RA⁺/CCR7⁻/CD28⁻) and CD8⁺ cells (CD8⁺/CD3⁺/CD19⁻/CD45RA⁺/CCR7⁻/CD28⁻), the ratio of CD4⁺ to CD8⁺ T cells using the quotient of counts of these compartments, the ratio of naïve to total memory CD4⁺ T cells (sum of central memory (CD4⁺/CD3⁺/CD19⁻/CD8⁻/CD45RA⁻/CCR7⁺/CD28⁺), effector memory

(CD4⁺/CD3⁺/CD19⁻/CD8⁻/CD45RA⁻/CCR7⁻/CD28⁻) and TemRA cells, naïve B cells (CD3⁻/CD19⁺/IgD⁺/CD27⁻), and late memory B cells⁴² (CD3⁻/CD19⁺/IgD⁻/CD27⁺). Variables that showed substantial skewness (skewness > 2; viz., CD4⁺ TemRA, the CD4:CD8 ratio, CD4 naïve:CD4 total memory cell ratio, and late memory B cells) were natural log transformed to approximate a normal distribution. These variables were standardized to have a mean of 0 and variance of 1 to ease comparison.

CMV seropositivity was assessed using IgG antibodies to CMV using the Roche e411 immunoassay analyzer (Roche Diagnostics Corporation, Indianapolis, IN). Cutoffs were used to identify non-reactive (<0.5 COI; coded 0), borderline (0.5 to <1.0 COI; coded 0), and reactive (>1.0 COI; coded 1) individuals. Non-reactive and borderline groups were combined and were used as the reference group. More information is available elsewhere³⁸.

Educational attainment was assessed by self-reported years of education. Respondents were categorized as 0-11 years (reference group), 12 years, 13-15 years, and 16 or more years.

Race/ethnicity was assessed by self-report. Respondents were categorized as non-Hispanic White (reference group), Hispanic, non-Hispanic Black, and non-Hispanic other race.

Parental low education was assessed using respondents' reports of their parents' educational attainment from the RAND HRS longitudinal file. If respondents reported on both parents, they were coded as 1 if either parent had less than high school education. If respondents only reported on one parent, they were coded as 1 if that parent had less than high school education. Respondents who only reported on their parents' education at Wave 2H reported only whether their parent had attended school for 8 or more years. These respondents were assigned 7.5 years if less than 8 years and 8.5 is assigned if 8 or more. We also ran these models

excluding participants who only reported on their parents' education at Wave 2H. Results for these models were highly similar and had an identical pattern of significant results.

We control for age in years and sex (male as reference) in all models. The weighted sample had a mean age of 67.66 years, ranging from 50 to 107. The sample was 55.02% female.

Plan of Analysis

Of the initial 9,934 respondents, 201 were excluded because they were cohort ineligible (e.g., were not a member of one of the birth cohorts represented in HRS) or were living in a nursing home. An additional 85 respondents were excluded because they were missing data on CMV seropositivity. 548 were excluded because they were missing on all flow cytometry measures in the current study. Sample weights, strata, and PSUs from the HRS tracker file were used to weight the sample. Respondents missing Venous Blood Study specific weights were assigned their core 2016 weight. The resulting sample size for this study is 9100. However, some individuals were missing data on specific cell types or predictor variables. To retain information, we used all respondents with complete data on each cell type for analyses using that cell type as the dependent variable. Additionally, there were too few participants in the non-Hispanic other race category to make meaningful comparisons, so those 277 participants were excluded in analyses with race and ethnicity as the focal predictors. E-table 1 shows final sample sizes for analyses for each model.

A series of generalized linear models were estimated regressing each cell type percentage/ratio on educational attainment or race/ethnicity, age and sex and then on educational attainment or race/ethnicity, CMV seropositivity, age, and sex. We then estimated a final set of models regressing each cell type percentage/ratio on race/ethnicity, education, CMV seropositivity, age, and sex. Because educational attainment and parental education highly

covaried, we also estimated a model in which participants were categorized as high or low in each (not shown). This model produced a similar pattern of results. All analyses were conducted in R 4.1.1⁴³ using the survey package⁴⁴.

Results

The weighted sample had a mean age of 68 years, ranging from 50 to 107. The sample was 55% female, 78.12% non-Hispanic White, 9.74% non-Hispanic Black, 9.01% Hispanic, and 3.13% non-Hispanic other race. 14.23% had 0-11 years of education, 29.15% had 12 years of education, 25.32% had 13-15 years of education, and 31.30% had 16 or more years of education. 62.97% of participants were CMV seropositive.

Figure 1 shows the weighted percent of participants who were CMV positive by age and sociodemographic factors (also see e-table 2). Higher education was associated with lower CMV seropositivity in our sample. Participants with fewer than 12 years of education had the highest percentage of seropositivity before age 80 and the percentage of CMV seropositive respondents remained stable across age (Figure 1, Panel A). Higher educated participants tended to have lower seropositivity at all ages, and seropositivity increased with age. Non-Hispanic White respondents had lower CMV seropositivity at all ages compared to both non-Hispanic Black and Hispanic respondents (Figure 1, Panel B). CMV seropositivity was also associated with parents' education such that participants whose parent(s) had less than a high school education had higher percentages of CMV seropositivity (Figure 1, Panel C). Finally, when considering education and race/ethnicity, CMV seropositivity tended to be lower for people with higher educational attainment across racial/ethnic groups (Figure 1, Panel D). To assess whether the association between educational attainment and CMV serostatus differed by race/ethnicity we estimated a

model predicting CMV seropositivity with interactions between educational attainment and race/ethnicity (not shown), and no interaction term was significant.

Results indicating educational differences from models regressing cell type percentages and ratios on education and controls are shown in the left panel of Figure 2. Compared to participants with the lowest level of education, i.e. fewer than 12 years, having 12 years of education, 13-15 years of education, or 16 or more years of education was associated with a greater percentage of naïve CD4⁺ T cells, CD8⁺ T cells, higher CD4:CD8 T cell ratio and CD4 naïve:CD4 total memory cell ratios, and lower percentages of CD4⁺ and CD8⁺ TemRA and late memory B cells. Participants with 12 and 16 or more years of education had a greater percentage of naïve B cells compared to those with less than 12 years of education.

The right panel of Figure 2 shows the same models additionally controlling for CMV seropositivity. Compared to having fewer than 12 years of education, having 12 years of education was still associated with a greater percentage of naïve CD4⁺, a higher CD4 naïve:CD4 total memory cell ratio, and fewer late memory B cells, but all other associations were attenuated and no longer statistically significant. Having 13-15 years of education was also associated with fewer CD8⁺ TemRA cells, a greater percentage of naïve CD4⁺ T and CD8⁺ T cells, and a higher CD4 naïve:CD4 total memory cell ratio but was no longer associated with CD4:CD8 ratio or with CD8⁺ TemRA and late memory B cells. Finally, having 16 or more years of education was associated with a greater percentage of naïve CD4⁺ T and CD8⁺ T cells, and a higher CD4 naïve:CD4 total memory cell ratio, and fewer late memory B cells.

Results from models regressing cell type percentages and ratios on race and ethnicity and controls are shown in the left panel of Figure 3. Compared to non-Hispanic White participants, non-Hispanic Black participants had fewer naïve CD4⁺ T cells, fewer naïve B cells, and lower

CD4:CD8 and CD4 naïve:CD4 total memory cell ratios. Black participants had a higher percentage of CD4⁺ TemRA cells, CD8⁺ naïve cells, CD8⁺ TemRA cells, and late memory B cells. Compared to non-Hispanic White participants, Hispanic participants had fewer naïve CD4⁺ T and CD8⁺ T cells, lower CD4:CD8 and CD4 naïve:CD4 total memory cell ratios, and a higher percentage of CD4⁺ and CD8⁺ TemRA cells and late memory B cells.

The right panel of Figure 3 shows the same models additionally controlling for CMV seropositivity. After controlling for CMV seropositivity, non-Hispanic Black participants no longer differed from non-Hispanic White participants on CD4:CD8 ratio or late memory B cell percentage. The non-Hispanic Black-White CD8⁺ TemRA cell percentage difference changed direction after controlling for CMV seropositivity. All differences between Hispanic participants and non-Hispanic White participants except on naïve T cell percentages and the CD4 naïve:CD4 total memory cell ratio were no longer statistically significant after controlling for CMV.

We further investigated the sign change in CD8⁺ TemRA cells for Black participants after controlling for CMV seropositivity noted in Figure 3 by looking at cell type percentages by race/ethnicity and CMV status (see e-table 3). Among CMV non-reactive and among CMV reactive individuals, non-Hispanic Black participants had a lower percentage of CD8⁺ TemRA cells compared to non-Hispanic White participants. These results are further discussed below.

Results of models regressing cell type percentages and ratios on parental education and controls are shown in the left panel of Figure 4. Participants who had at least one parent with less than high school education had a lower percentage of naïve CD4⁺ T, CD8⁺ T, lower CD4:CD8 T and CD4 naïve:CD4 total memory cell ratios, and higher percentages of CD4⁺ and CD8⁺ TemRA and late memory B cells. After controlling for CMV seropositivity, participants who had at least one parent with less than high school education had lower percentages of CD4⁺ and CD8⁺ naïve

cells and lower CD4:CD8 T and CD4 naïve:CD4 total memory cell ratios compared to other participants.

Next, all sociodemographic factors were included in one model, adjusting for covariates, to determine whether the estimates of associations were altered with mutual adjustment (shown in e-table 4). Controlling for childhood SES, race/ethnicity, and CMV serostatus, having 12 of years of education was associated with more CD4⁺ and CD8⁺ naïve cells compared to having less than 12 years of education. Having 12 years of education is no longer associated with the CD4 naïve:CD4 total memory cell ratio after including all covariates. Having 12 years of education is associated with a greater percentage of naïve B cells (a non-significant association in model shown in Figure 2) after including all covariates. Having 13-15 years of education was associated with Fewer CD4⁺ TemRA cells, more CD4⁺ and CD8⁺ naïve cells, and a higher CD4 naïve:CD4 total memory cell ratio. These associations were all also significant in results shown in Figure 2. Similar to results shown in Figure 2, having 16 or more years of education was associated with lower percentages of CD4⁺ and CD8⁺ naïve cells, and a higher CD4 naïve:CD4 total memory cell ratio, but was no longer with a lower percentage of late memory B cells. After controlling for all covariates, having 16 or more years of education was also associated with a higher percentage of naïve B cells, a non-significant association in model shown in Figure 2.

Similar to results shown in Figure 3, compared to non-Hispanic White participants, non-Hispanic Black participants had more CD4⁺ TemRA cells, fewer CD4⁺ naïve cells, fewer CD8⁺ TemRA cells, more CD8⁺ naïve cells, a lower CD4 naïve:CD4 total memory cell ratio, and fewer naïve B cells. Similar to results shown in Figure 3, adjusting for childhood SES and educational attainment, compared to non-Hispanic White participants, Hispanic participants had fewer naïve CD4⁺ cells and a lower CD4 naïve:CD4 total memory cell ratio. Unlike results shown in Figure 3,

Hispanic respondents no longer differed from White respondents in percentage of naïve CD8⁺ T cells. Also different from results shown in Figure 3, after controlling for all covariates, being Hispanic was associated with a higher CD4:CD8 ratio.

Similar to results shown in Figure 4, parental low education was associated with fewer CD4⁺ and CD8⁺ naïve cells, and a lower CD4:CD8 ratio adjusting for educational attainment and race/ethnicity. Unlike results shown in Figure 4, after controlling for all covariates, low parental education was not associated with the CD4 naïve:CD4 total memory cell ratio. Because educational attainment and parental education highly covaried, we also estimated a model in which participants were categorized as high or low in each (results not shown). Results from this model were similar to those in the model presented above, leading to the same conclusions.

Finally, we assessed interactions among these factors by regressing each cell type on all predictors and interactions among focal predictors (not shown). Non-Hispanic Black participants appear to benefit more from 13-15 and 16+ years of education in terms of naïve B cell percentage compared to non-Hispanic White participants. Non-Hispanic Black participants also appear to be less negatively affected by low parental education in terms of CD4⁺ naïve and B naïve cells compared to non-Hispanic White participants.

Discussion

In this nationally representative sample of US adults over the age of 50, we found that cell type percentages and ratios indicative of immune aging were associated with SES and demographic factors. Higher educational attainment, non-minority status, and higher levels of parental education were associated less “aged” lymphocyte percentages. Though there has been little research on the association between immune aging and SES, these findings are consistent with evidence that income is associated with age-related immune change¹⁴ and that SES is

associated with immune functioning broadly¹³. Our findings are also consistent with past research showing that early life SES is associated with CMV exposure and altered immune functioning¹⁷.

Controlling for CMV attenuated many of the associations with sociodemographic factors observed in this study, which underscores the importance of CMV in later life immune function. CMV infection is patterned by SES and race/ethnicity in the US²². Of note, individuals of low SES and minorities are more likely to acquire CMV earlier in life than high SES individuals and Non-Hispanic Whites⁴⁵. Since the suppression of CMV requires substantial immunological resources, leading to memory inflation and the expansion of late-differentiated T cells^{8,18}, having to control the virus for additional years may be relevant for the sociodemographic patterning of decline in immune function observed with aging⁴⁶.

In this study, higher percentages of naïve T cells were consistently indicative of higher SES and non-minority status independent of CMV serostatus. It is possible that lifestyle factors associated with SES (e.g., diet and exercise) may slow thymic involution, leading to a relatively larger percentage of new naïve T cells being produced⁷. Thus, naïve T cells may be an immunological bellwether for socioeconomic and structural forces of interest to researchers interested in health inequalities, however further research is needed to refine this hypothesis.

Ratios among lymphocytes may be particularly useful for measuring age-related changes in immune cell percentages. Higher educational attainment was positively associated with CD4/CD8 ratio, indicating stronger immune function, however, all observed associations were attenuated and no longer statistically significant after adjusting for CMV. Thus, the differential prevalence of CD4⁺ and CD8⁺ T lymphocytes associated with socioeconomic position may be driven by differential exposure to and reactivation of CMV^{22,47,48}. Alternatively, the ratio of

CD4⁺ naïve to CD4⁺ memory cells was statistically significantly associated with educational attainment, race and ethnicity, and parental education after controlling for CMV seropositivity. Therefore, CD4⁺ naïve/CD4⁺ memory cell ratio may be particularly indicative of age-related immune phenotype that is affected by socioeconomic position.

Non-Hispanic Black-White differences in CD8⁺ naïve T cell percentages became more extreme after controlling for CMV and CD8⁺ TemRA cell differences changed direction. This was because non-Hispanic Black participants had a lower percentage of CD8⁺ TemRA cells and higher percentage of CD8⁺ naïve T cells compared to non-Hispanic White participants within CMV seropositivity category. These differences were masked or reversed at the population level, suggesting this may be an example of Simpson's paradox. That is, were it not for differential CMV exposure, Black US adults might have less accelerated immunosenescence compared to White US adults. Differential exposure to CMV may be a major driver of racial differences in immunosenescence and potentially in racial health disparities, though further research is needed to confirm this. Interventions targeting exposure risk factors (e.g., number of individuals in the household, nativity outside of the US, region of residence, day care attendance, and sexual behaviors²²) or potentially CMV vaccination may help in addressing racial health disparities. Future work on nutritional, psychological stress, inflammation, and morbidity and mortality prediction of immune aging is warranted.

This study has limitations. First, CMV seroprevalence and flow cytometry were only assessed at one time. Longitudinal assessments are needed to establish temporal ordering of CMV infection and age-related cell type percentages and ratios. Second, this study is limited to US adults 50 years and older. Further research in international samples with different educational systems and CMV seroprevalence profiles are needed to understand place-based SES and

infection influences on later life immune function. Despite these limitations, this study has many strengths. The age-related cell type percentages and ratios examined expands on previous research into the immune risk profile^{49,50}, immune parameters associated with mortality in very old individuals, to include age-related changes to immune function in a population representative study of older US adults. This study provides evidence that higher socioeconomic status may be protective against immunosenescence. Future research on nutritional, psychosocial stress, and inflammatory drivers of cell type percentages and ratios, as well as the relationship between cell type percentages and ratios and morbidity and mortality are warranted. Understanding the aging immune system may illuminate the biological processes underlying inequalities in age-related morbidity and mortality.

References

1. Huff WX, Kwon JH, Henriquez M, Fetcko K, Dey M. The Evolving Role of CD8+CD28– Immunosenescent T Cells in Cancer Immunology. *International Journal of Molecular Sciences*. 2019;20(11):2810. doi:10.3390/ijms20112810
2. Aiello A, Farzaneh F, Candore G, et al. Immunosenescence and Its Hallmarks: How to Oppose Aging Strategically? A Review of Potential Options for Therapeutic Intervention. *Front Immunol*. 2019;10. doi:10.3389/fimmu.2019.02247
3. Pawelec G. Age and immunity: What is “immunosenescence”? *Exp Gerontol*. 2018;105:4-9. doi:10.1016/j.exger.2017.10.024
4. Thyagarajan B, Faul J, Vivek S, et al. Age-related differences in T cell subsets in a nationally representative sample of people over age 55: Findings from the Health and Retirement Study. *The Journals of Gerontology: Series A*. 2021;(glab300). doi:10.1093/gerona/300
5. Adriaensen W, Pawelec G, Vaes B, et al. CD4:8 Ratio Above 5 Is Associated With All-Cause Mortality in CMV-Seronegative Very Old Women: Results From the BELFRAIL Study. *GERONA*. Published online December 7, 2016:glw215. doi:10.1093/gerona/300
6. Pangrazzi L, Weinberger B. T cells, aging and senescence. *Experimental Gerontology*. 2020;134:110887. doi:10.1016/j.exger.2020.110887
7. Palmer S, Albergante L, Blackburn CC, Newman TJ. Thymic involution and rising disease incidence with age. *PNAS*. 2018;115(8):1883-1888. doi:10.1073/pnas.1714478115
8. Bektas A, Schurman SH, Sen R, Ferrucci L. Human T cell immunosenescence and inflammation in aging. *Journal of Leukocyte Biology*. 2017;102(4):977-988. doi:10.1189/jlb.3RI0716-335R

9. Callender LA, Carroll EC, Beal RWJ, et al. Human CD8+ EMRA T cells display a senescence-associated secretory phenotype regulated by p38 MAPK. *Aging Cell*. 2018;17(1):e12675. doi:10.1111/accel.12675
10. Covre LP, De Maeyer RPH, Gomes DCO, Akbar AN. The role of senescent T cells in immunopathology. *Aging Cell*. 2020;19(12):e13272. doi:10.1111/accel.13272
11. Adriaensen W, Derhovanessian E, Vaes B, et al. CD4:8 Ratio >5 Is Associated With a Dominant Naive T-Cell Phenotype and Impaired Physical Functioning in CMV-Seropositive Very Elderly People: Results From the BELFRAIL Study. *The Journals of Gerontology: Series A*. 2015;70(2):143-154. doi:10.1093/gerona/glu018
12. Kristenson M, Eriksen HR, Sluiter JK, Starke D, Ursin H. Psychobiological mechanisms of socioeconomic differences in health. *Social Science & Medicine*. 2004;58(8):1511-1522. doi:10.1016/S0277-9536(03)00353-8
13. Dowd JB, Aiello AE. Socioeconomic Differentials in Immune Response. *Epidemiology*. 2009;20(6):902-908. doi:10.1097/EDE.0b013e3181bb5302
14. Aiello AE, Feinstein L, Dowd JB, et al. Income and Markers of Immunological Cellular Aging. *Psychosomatic Medicine*. 2016;78(6):657-666. doi:10.1097/PSY.0000000000000320
15. Braveman PA, Cubbin C, Egerter S, Williams DR, Pamuk E. Socioeconomic Disparities in Health in the United States: What the Patterns Tell Us. *Am J Public Health*. 2010;100(S1):S186-S196. doi:10.2105/AJPH.2009.166082
16. Hoes J, Boef AGC, Knol MJ, et al. Socioeconomic Status Is Associated With Antibody Levels Against Vaccine Preventable Diseases in the Netherlands. *Frontiers in Public Health*. 2018;6:209. doi:10.3389/fpubh.2018.00209

17. Elwenspoek MMC, Sias K, Hengesch X, et al. T Cell Immunosenescence after Early Life Adversity: Association with Cytomegalovirus Infection. *Front Immunol.* 2017;8.
doi:10.3389/fimmu.2017.01263
18. Derhovanessian E, Maier AB, Hähnel K, et al. Infection with cytomegalovirus but not herpes simplex virus induces the accumulation of late-differentiated CD4+ and CD8+ T-cells in humans. *Journal of General Virology.* 2011;92(12):2746-2756.
doi:10.1099/vir.0.036004-0
19. Williams DR. Miles to go before we Sleep: Racial inequities in health. *Journal of Health and Social Behavior.* 2012;53(3):279-295. doi:10.1177/0022146512455804
20. Achhra AC, Amin J. Race and CD4+ T-cell count in HIV prognosis and treatment. *Future Virology.* 2012;7(2):193-203. doi:http://dx.doi.org/10.2217/fvl.11.143
21. Nowicki MJ, Karim R, Mack WJ, et al. Correlates of CD4+ and CD8+ Lymphocyte Counts in High-Risk Immunodeficiency Virus (HIV)-Seronegative Women enrolled in the Women's Interagency HIV Study (WIHS). *Human Immunology.* 2007;68(5):342-349.
doi:10.1016/j.humimm.2007.01.007
22. Dowd JB, Aiello AE, Alley DE. Socioeconomic disparities in the seroprevalence of cytomegalovirus infection in the US population: NHANES III. *Epidemiol Infect.* 2009;137(1):58-65. doi:10.1017/S0950268808000551
23. Reed RG. Stress and immunological aging. *Current Opinion in Behavioral Sciences.* 2019;28:38-43. doi:10.1016/j.cobeha.2019.01.012
24. Prather AA, Epel ES, Portela Parra E, et al. Associations between chronic caregiving stress and T cell markers implicated in immunosenescence. *Brain, Behavior, and Immunity.* 2018;73:546-549. doi:10.1016/j.bbi.2018.06.019

25. Geronimus AT. The weathering hypothesis and the health of African-American women and infants: Evidence and speculations. *Ethnicity & Disease*. 1992;2(3):207-221.
26. Simons RL, Lei MK, Klopack ET, Beach SRH, Gibbons FX, Philibert RA. The effects of social adversity, discrimination, and health risk behaviors on the accelerated aging of African Americans: Further support for the weathering hypothesis. *Social Science & Medicine*. 2021; Advance online publication. doi:10.1016/j.socscimed.2020.113169
27. Schmeer KK, Tarrence J. Racial-ethnic disparities in inflammation: Evidence of weathering in childhood? *Journal of Health and Social Behavior*. 2018;59(3):411-428.
doi:10.1177/0022146518784592
28. Yang Y, Gerken K, Schorpp K, Boen C, Harris KM. Early-life socioeconomic status and adult physiological functioning: A life course examination of biosocial mechanisms. *Biodemography & Social Biology*. 2017;63(2):87-103.
doi:10.1080/19485565.2017.1279536
29. Liu Y, Croft JB, Chapman DP, et al. Relationship between adverse childhood experiences and unemployment among adults from five US states. *Soc Psychiatry Psychiatr Epidemiol*. 2013;48(3):357-369. doi:10.1007/s00127-012-0554-1
30. Font SA, Maguire-Jack K. Pathways from childhood abuse and other adversities to adult health risks: The role of adult socioeconomic conditions. *Child Abuse & Neglect*. 2016;51:390-399. doi:10.1016/j.chiabu.2015.05.013
31. Wickrama KAS, Kwon JA, Oshri A, Lee TK. Early socioeconomic adversity and young adult physical illness: The role of body mass index and depressive symptoms. *J Adolesc Health*. 2014;55(4):556-563. doi:10.1016/j.jadohealth.2014.04.006

32. Pudrovska T, Logan ES, Richman A. Early-life social origins of later-life body weight: The role of socioeconomic status and health behaviors over the life course. *Social Science Research*. 2014;46:59-71. doi:10.1016/j.ssresearch.2014.02.007
33. McEwen CA, McEwen BS. Social structure, adversity, toxic stress, and intergenerational poverty: An early childhood model. *Annual Review of Sociology*. 2017;43(1):445-472. doi:10.1146/annurev-soc-060116-053252
34. Pudrovska T, Anikputa B. Early-Life Socioeconomic Status and Mortality in Later Life: An Integration of Four Life-Course Mechanisms. *The Journals of Gerontology: Series B*. 2014;69(3):451-460. doi:10.1093/geronb/gbt122
35. Willson AE, Shuey KM. Life course pathways of economic hardship and mobility and midlife trajectories of health. *Journal of Health and Social Behavior*. 2016;57(3):407-422. doi:10.1177/0022146516660345
36. Nettle D, Frankenhuys WE, Rickard IJ. The evolution of predictive adaptive responses in human life history. *Proceedings of the Royal Society B-Biological Sciences*. 2013;280(1766):9. doi:10.1098/rspb.2013.1343
37. Duggal NA, Pollock RD, Lazarus NR, Harridge S, Lord JM. Major features of immunesenescence, including reduced thymic output, are ameliorated by high levels of physical activity in adulthood. *Aging Cell*. 2018;17(2):e12750. doi:10.1111/accel.12750
38. Crimmins EM, Faul JD, Thyagarajan B, Weir DR. *Venous Blood Collection and Assay Protocol in the 2016 Health and Retirement Study 2016 Venous Blood Study (VBS).*; 2017. <https://hrsdata.isr.umich.edu/sites/default/files/documentation/data-descriptions/HRS2016VBSDD.pdf>

39. Maecker HT, McCoy JP, Nussenblatt R. Standardizing immunophenotyping for the Human Immunology Project. *Nat Rev Immunol*. 2012;12(3):191-200. doi:10.1038/nri3158
40. Thyagarajan B, Barcelo H, Crimmins E, et al. Effect of delayed cell processing and cryopreservation on immunophenotyping in multicenter population studies. *Journal of Immunological Methods*. 2018;463:61-70. doi:10.1016/j.jim.2018.09.007
41. Barcelo H, Faul J, Crimmins E, Thyagarajan B. A Practical Cryopreservation and Staining Protocol for Immunophenotyping in Population Studies. *Current Protocols in Cytometry*. 2018;84(1):e35. doi:10.1002/cpcy.35
42. Frasca D, Diaz A, Romero M, Blomberg BB. Human peripheral late/exhausted memory B cells express a senescent-associated secretory phenotype and preferentially utilize metabolic signaling pathways. *Experimental Gerontology*. 2017;87:113-120. doi:10.1016/j.exger.2016.12.001
43. R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing; 2021. <https://www.R-project.org/>
44. Lumley T. Analysis of Complex Survey Samples. *Journal of Statistical Software*. 2004;9(1):1-19. doi:10.18637/jss.v009.i08
45. Colugnati FA, Staras SA, Dollard SC, Cannon MJ. Incidence of cytomegalovirus infection among the general population and pregnant women in the United States. *BMC Infectious Diseases*. 2007;7(1):71. doi:10.1186/1471-2334-7-71
46. Solana R, Tarazona R, Aiello AE, et al. CMV and Immunosenescence: from basics to clinics. *Immun Ageing*. 2012;9(1):1-9. doi:10.1186/1742-4933-9-23
47. Reed RG, Presnell SR, Al-Attar A, Lutz CT, Segerstrom SC. Perceived stress, cytomegalovirus titers, and late-differentiated T and NK cells: Between-, within-person

associations in a longitudinal study of older adults. *Brain, Behavior, and Immunity*.


2019;80:266-274. doi:10.1016/j.bbi.2019.03.018

48. Bauer ME, Fuente MD la. The role of oxidative and inflammatory stress and persistent viral infections in immunosenescence. *Mechanisms of Ageing and Development*. 2016;158:27-37. doi:10.1016/j.mad.2016.01.001
49. Wikby A, Johansson B, Ferguson F, Olsson J. Age-related changes in immune parameters in a very old population of Swedish people: A longitudinal study. *Experimental Gerontology*. 1994;29(5):531-541. doi:10.1016/0531-5565(94)90036-1
50. Pawelec G. The human immunosenescence phenotype: does it exist? *Semin Immunopathol*. 2020;42(5):537-544. doi:10.1007/s00281-020-00810-3


Figures

Figure 1. Percent CMV seropositive by SES variables and age


A




B



C




D




Note: HS = high school; 95% confidence intervals shown.

Figure 2. Educational attainment associated with cell type percentages and ratios




Note: cell type percentages and ratios have been standardized to have a mean of 0 and standard deviation of 1; regression coefficients and 95% confidence intervals from GLMs regressing cell type percentages/ratios on educational attainment; reference group is less than 12 years of education.

Figure 3. Race/ethnicity associated with cell type percentages and ratios



Note: NH = non-Hispanic; cell type percentages and ratios have been standardized to have a mean of 0 and standard deviation of 1; regression coefficients and 95% confidence intervals from GLMs regressing cell type percentages/ratios on race/ethnicity; reference group is non-Hispanic White.

Figure 4. Parents' Education associated with cell type percentages and ratios



Note: cell type percentages and ratios have been standardized to have a mean of 0 and standard deviation of 1; regression coefficients and 95% confidence intervals from GLMs regressing cell type percentages/ratios on parents' education; reference group is low parental education