

Association of COVID-19 with risks of hospitalization and mortality from other disorders post-infection: A study of the UK Biobank

Yong XIANG¹, Ruoyu Zhang¹, Jinghong QIU¹, Hon-Cheong SO^{1-7*}

¹School of Biomedical Sciences, The Chinese University of Hong Kong, Shatin, Hong Kong

²KIZ-CUHK Joint Laboratory of Bioresources and Molecular Research of Common Diseases, Kunming Institute of Zoology and The Chinese University of Hong Kong, China

³CUHK Shenzhen Research Institute, Shenzhen, China

⁴Department of Psychiatry, The Chinese University of Hong Kong, Shatin, Hong Kong

⁵Margaret K.L. Cheung Research Centre for Management of Parkinsonism, The Chinese University of Hong Kong, Shatin, Hong Kong

⁶Brain and Mind Institute, The Chinese University of Hong Kong, Shatin, Hong Kong

⁷Hong Kong Branch of the Chinese Academy of Sciences Center for Excellence in Animal Evolution and Genetics, The Chinese University of Hong Kong, Hong Kong SAR, China

**Corresponding author*

Correspondence to: Hon-Cheong So, MBBS, PhD. Lo Kwee-Seong Integrated Biomedical Sciences Building, The Chinese University of Hong Kong, Shatin, Hong Kong. Tel: +852 3943 9255; E-mail: hcsso@cuhk.edu.hk

Submitted on 23 Mar 2022

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Abstract

Background More than 0.4 billion cases of COVID-19 have been reported worldwide. The sequelae of COVID-19 remains unclear, especially whether it may be associated with increased hospitalization and mortality from other diseases.

Methods We leveraged a large prospective cohort of the UK biobank (UKBB) ($N=412,096$; current age 50-87) to identify associations of COVID-19 with hospitalization and mortality due to different diseases post-infection. We conducted a comprehensive survey on disorders from all systems (up to 135 disease categories). Multivariable Cox and Poisson regression was conducted controlling for main confounders. For sensitivity analysis, we also conducted separate analysis for new-onset and recurrent cases, and other sensitivity analysis such as the prior event rate adjustment (PERR) approach to minimize effects of unmeasured confounders.

Results Compared to individuals with no known history of COVID-19, those with severe COVID-19 (requiring hospitalization) exhibited higher hazards of hospitalization and/or mortality due to multiple disorders (median follow-up=608 days), including disorders of respiratory, cardiovascular, neurological, gastrointestinal, genitourinary and musculoskeletal systems. Increased hazards of hospitalizations and/or mortality were also observed for injuries due to fractures, various infections and other non-specific symptoms. These results remained largely consistent after sensitivity analyses. Severe COVID-19 was also associated with increased all-cause mortality (HR=14.700, 95% CI: 13.835-15.619).

Mild (non-hospitalized) COVID-19 was associated with modestly increased risk of all-cause mortality (HR=1.237, 95% CI 1.037-1.476) and mortality from neurocognitive disorders (HR=9.100, 95% CI: 5.590-14.816), as well as hospital admission from a few disorders such as aspiration pneumonitis, musculoskeletal pain and other general signs/symptoms.

Conclusions In conclusion, this study revealed increased risk of hospitalization and mortality from a wide variety of pulmonary and extra-pulmonary diseases after COVID-19, especially for severe infections. Mild disease was also associated with increased all-cause mortality. Further studies are required to replicate our findings.

Introduction

There are more than 400 million¹ documented cases of COVID-19 worldwide, and accumulating evidence have shown that the disease may be associated with various sequelae. For example, respiratory symptoms and disorders were reported frequently in the post-infection period, such as dyspnoea and cough², lung function impairment³, and bronchiectasis⁴. Relatively less attention has been paid on other organ systems. For example, studies have reported that neurological manifestations could persist in patient with COVID-19⁵⁻⁷. Increased risks of cardiovascular diseases have also been observed⁸. Several other studies have also investigated the health consequences of COVID-19 across different systems⁹⁻¹². Nevertheless, the full spectrum of sequelae from COVID-19 remains unclear, and there are still limited studies which have conducted a comprehensive analysis on potential sequelae from *all* body systems.

Given the very large number of people affected by COVID-19 worldwide, additional studies are still very much needed to delineate clearly the sequelae of COVID-19. There are also some limitations of existing studies. For example, some studies^{11,12} are of relatively small sample sizes, which limits the power to detect moderate associations. Another study⁹ examined the risks of a wide range of sequelae; however the study was restricted to veterans, who are predominantly males and the findings may not be readily generalizable to the general population¹³. In addition, very few works (except e.g. ref¹⁴) have studied how vaccination may change the risk of sequelae from COVID-19.

Here we leveraged a large prospective cohort of the UK biobank (UKBB) to identify associations of COVID-19 with hospitalization and mortality due to different diseases post-infection. We conducted a comprehensive survey on disorders from all systems, and a series of additional analysis, for example on the risk of hospitalization from new-onset and recurrent diseases, risk of sequelae across different follow-up periods, and how vaccination (before or after the infection) may alter the risk of sequelae,

Methods

UK Biobank sample

The UK Biobank (UKBB) is a large prospective cohort comprising ~500,000 UK participants with continuous follow-up. The current age of subjects in this study ranged from 50 to 87 years. The details of the UKBB sample are described in ref¹⁵. As not every participant is linked to hospital inpatient records, our analysis was restricted to those subjects (sample size, $N= 412,096$) with at least one hospitalization record and alive as at 31-Jan-2020, the date for the first confirmed case of COVID-19 in UK. The current study was conducted under the project number 28732.

Study Outcomes

Our main study outcomes were hospitalization and mortality due to various disorders. Hospitalization records were extracted from the Hospital Episode Statistics (HES) data of UKBB. Detailed descriptions of HES can be found in <https://biobank.ndph.ox.ac.uk/ukb/ukb/docs/HospitalEpisodeStatistics.pdf>. The hospital inpatient ‘core’ and diagnoses datasets were updated to 30 Sep 2021 (accessed 21 Dec 2021). The diagnosis codes and corresponding dates were summarized based on each participant’s inpatient record. All the diagnosis codes were converted to 3-character ICD-10 codes. We followed the mapping strategy as described in https://biobank.ndph.ox.ac.uk/showcase/showcase/docs/first_occurrences_outcomes.pdf. The mortality records of UKBB contain the cause of death (ICD10-coded) and were linked to national death registries. Details can be found in <https://biobank.ctsu.ox.ac.uk/crystal/crystal/docs/DeathLinkage.pdf>. Mortality data were updated to 18 Oct 2021 (accessed 21 Dec 2021).

All ICD-10 codes were covered in analysing the sequelae of COVID-19 infection. We employed the latest version (2022.1) of Clinical Classifications Software Refined (CCSR)¹⁶ to classify ICD-10 codes to a smaller number of clinically meaningful categories of diagnoses. Totally 320 clinical categories covering 21 body systems were included as outcomes.

Only primary causes of admission/mortality were considered. All diagnosis given before the study start-date (i.e. time0; the date a subject entered the cohort study) were regarded as medical history of comorbidities; diagnoses given after time0 were treated as hospitalization/mortality due to ‘new-onset’ diseases. For better statistical power, we primarily present the results from *any* hospitalization or mortality from diseases, without consideration of past medical history. Nevertheless, we also performed additional stratified analysis on new-onset (subjects with no known prior history) and recurrent/relapsed diseases (subjects with known history of the disease).

Covariates

We performed multivariable regression analysis with adjustment for covariates including basic demographics (age, sex, ethnicity), major risk factors of cardiovascular disorders (lipid and glucose levels, HbA1c), immunological disorders (autoimmune diseases, immunodeficiency, history of receiving immunosuppressants), major comorbidities (coronary artery disease[CAD], stroke, Type 2 diabetes mellitus[T2DM], hypertension[HTN], atrial fibrillation[AF], chronic obstructive pulmonary disease[COPD], dementia, history of cancer, chronic kidney disease[CKD] and pneumonia), overall health status and indicators of multi-morbidities (number of non-cancer illnesses, number of medications prescribed by GP and number of hospitalizations in the past year), indicators of obesity (body mass index [BMI], waist circumference), socioeconomic status (Townsend Deprivation index) and smoking status. Covariates were picked based on their potential relevance to COVID-19 infection and/or its complications. Please refer to Table S1 for the full list and distribution of covariates.

To define history of relevant diseases, we included information from primary care and hospital inpatient data, as well as ICD-10 diagnoses (code 41270) and self-reported illnesses (code 20002). The strategy of integrating all diagnoses were based on instructions from UKBB (https://biobank.ndph.ox.ac.uk/showcase/showcase/docs/first_occurrences_outcomes.pdf).

Missing covariate data

Missing values of features were imputed with “missRanger”. The program is based on missForest, an iterative imputation approach using random forest. The program has been widely used and has been shown to produce low imputation errors and good performance in predictive models¹⁷. The missing rate of each covariate and the OOB (out-of-bag error) of imputation are listed in Table S2.

COVID-19 infection status

COVID-19 testing data were downloaded from the UKBB data portal (please refer to <https://biobank.ndph.ox.ac.uk/showcase/exinfo.cgi?src=COVID19>). Briefly, the latest COVID test results were downloaded on 21 Dec 2021 (last update on 30 Oct 2021). COVID-19 test results after the study end-date (30 Sep 2021 for hospital admission and 18 Oct 2021 for mortality) were ignored. COVID-19 infection status was defined by any positive test results, or a diagnosis of COVID-19 (ICD-10 code: U071) from inpatient/mortality records or the code “Y2a3b” within the TPP GP records (<https://biobank.ndph.ox.ac.uk/showcase/coding.cgi?id=8708&nl=1>) in the follow-up period.

“Severe” COVID-19 was defined by any case who is hospitalized or whose mortality is due to U07.1. Other non-hospitalized and non-fatal cases are considered as ‘mild’. For hospitalized cases, we required both test result and test origin to be 1 (positive test and inpatient origin). For a small number of cases with initial outpatient origin and positive test result, but changed to inpatient origin within 2 weeks, we still considered these subjects as inpatient cases (i.e., we shall assume the hospitalization was related to COVID-19).

Four models of analysis

Four sets of analysis were performed for hospitalization and mortality as outcome separately (Table 1). For model A, it was restricted to test-positive subjects, while for other models, they were based on the whole population. For Model A, we compared severe vs mild COVID-19 cases for post-infection hospital admission and mortalities. For model B, we compared severe COVID-19 cases against those with no known history of infection. For model C, all infected cases were compared against those with no known infection. Finally, ‘mild COVID-19’ were compared against subjects with no known infection in model D.

Statistical analysis

Cox and Poisson regression were employed for analysis. Cox regression models time to development of event, while Poisson regression models the incidence rate (with consideration of follow-up time). Briefly, the time to hospitalization or mortality from the studied diseases was treated as outcome in Cox regression, controlling for other covariates. For Poisson regression, presence/absence of the event of interest was considered as outcome, with ‘offset’ specified (=number of days of follow-up) to account for the differences in duration of follow-up for each subject.

For subjects with COVID-19 infection, the first positive date was treated as the start-date (time0) of this cohort study. For subjects free of COVID-19 infection, 31-Jan-2020 were assigned as time0. The end of follow-up was set as the time a participant developed an event of interest or 30 Sep 2021 (for analysis of hospitalization) or 18 Oct 2021 (for analysis of mortality). All statistical analysis were performed by R (version number 3.6.1). The false discovery rate (FDR) approach by Benjamini and Hochberg¹⁸ was performed to control for multiple testing, which controls the expected proportion of false-positives in the rejected hypotheses.

To avoid convergence issues when large number of predictors are included, we followed the methodology described by Zhao et al.¹⁹ to further select covariates for inclusion. We first included several ‘default’ variables, including age, sex and general health status (number of hospitalization and medications received in the past year) into the regression model. For the rest of the covariates, univariate testing was performed with each predictor; those with nominally significant associations ($p < 0.05$) with the outcome were selected into the final model. This approach¹⁹ was shown by theory and simulations to control the proportion of false positive predictors.

Prior event rate ratio (PERR) adjustment

Although many possible confounders have been adjusted for, the chance of residual confounding remains. The PERR approach exploits a before-and-after design to minimize the effects of unmeasured confounders²⁰⁻²². The basic idea is to compare the differences in hazard ratios between the exposed and unexposed groups before and after the COVID-19 pandemic, as differences in hospitalization/mortality risks prior to the pandemic can be due to baseline differences (unmeasured confounders) between the two groups.

We may assume that *before* the onset of the pandemic in UK (31 Jan 2020), the hazard ratio of event (e.g. hospitalization due to a specific disease) between the exposed and unexposed groups is HR_{before} . Since neither group would have COVID-19 before our study baseline, HR_{before} is assumed to reflect baseline differences owing to unmeasured confounders, independent of the effect of COVID-19. Similarly, we define the differences between groups *after* the pandemic onset to be HR_{after} .

The PERR-adjusted HR, calculated as $HR_{\text{after}}/HR_{\text{before}}$, provides an estimate of the effect of COVID-19 infection accounting for both unmeasured and measured confounders. The evaluation of $HR_{\text{after}}/HR_{\text{before}}$ is equivalent to evaluating the difference in regression coefficients before and after treatment, i.e. $\beta_{\text{diff}} = \beta_{\text{after}} - \beta_{\text{before}}$. A non-parametric bootstrap procedure (re-sampling observations with replacement) was used to estimate the standard error (SE) of β_{diff} ²². The z-statistic can then be computed by $\beta_{\text{diff}}/SE(\beta_{\text{diff}})$. We performed further PERR adjustment for all significant associations (unadjusted p-value < 0.05) in the original analysis. Two hundred bootstraps were performed.

The PERR approach has several assumptions. It is assumed that the confounders (measured and unmeasured) have the same effect in both pre- and post-COVID-19 periods. Also, the probability of (severe) COVID-19 infection is assumed to be unaffected by events (hospitalization due to diseases) in the prior period.

Prescription-time distribution matching (PTDM)

In this cohort study, the start-date (time0) for the exposed group is the first date of being tested positive, while an artificial time0 was assigned to the unexposed group. Imbalance of ‘prescription time’ (here ‘prescription’ refers to being infected by COVID-19) distribution may lead to a typical bias known as ‘survival bias’. Survival bias may be generated when individuals who die shortly after the start of follow-up might not have a probability of being infected and would only be classified into the ‘unexposed group’ by definition. As a result, events in the unexposed group may be over-estimated, and this bias is likely more prominent for the study of mortalities (or hospitalizations due to a lethal disease, e.g. stroke).

To reduce such bias, several methods have been proposed (see²³). Here we employed the PTDM approach as it is computationally fast and has been shown to effectively reduce survival bias²³. In brief, for each person without COVID-19 infection, instead of using the same start-date (31 Jan 2020), an alternative time0 was generated by randomly choosing from the sets of time0 of the COVID-19-exposed group. Subjects who are unaffected by COVID-19 but had an event before the assigned time0 are excluded from the analysis. The objective is to ensure a similar distribution of the time0 across both groups.

Note that in this study, we may expect more associations of COVID-19 with sequelae after PTDM adjustment, as under PTDM time0 is pushed to a later date for the unexposed group and fewer events (or lower incidence rate) would be observed in that group.

Other analysis to check for robustness of findings

We performed a variety of different analyses to verify the robustness of our findings under different modeling strategies and assumptions:

- (1) While we primarily focused on all hospitalizations, we also performed stratified analysis for new-onset and recurrent diseases, as mentioned earlier;
- (2) for counting the follow-up end date, we primarily consider the last date of available hospitalization/mortality records as the end of follow-up. However, since vaccination may change the risk of sequelae, we also performed our analysis limited to the period when no vaccines were yet available, i.e. setting 8 Dec 2020 (the date of first vaccination) as our study end-date;

Association with hospital admission/mortality stratified by vaccination status and follow-up period

We also conducted further (exploratory) analysis stratified by vaccination status. Briefly, we compared hospitalization rates for subjects who have received at least one dose of vaccine *before* infection, those who have received vaccine *after* the infection, those who are vaccinated both before and after the infection, those who are infected but not vaccinated and those with no known COVID-19. One of our main questions is whether prior vaccination (and to a lesser extent, vaccination after infection) may ameliorate the risk of sequelae, when compared to infected subjects who are not vaccinated.

In another additional analysis, we examined the effect of COVID-19 on hospitalization risks as stratified by the follow-up period. Three periods were specified, namely <3 months, 3-6 months and >6 months. We hypothesized that the risk would be most prominent in the initial months and might wane afterwards.

Results

Overview

In this study, we found that severe COVID-19 was associated with elevated hazards ratios (HR) of hospital admission and mortality due to various diseases involving many body systems. These patterns were consistent in a series of sensitivity analyses, and after PTDM and PERR adjustment.

The four sets of analysis are summarized in Table 1 and the number of subjects for each model is presented in Table 2.

Compared to subjects with no known history of COVID-19, association of hospitalization with severe COVID-19 (severe disease vs uninfected)

Severe infection was associated with higher hazards of hospitalization and mortality due to a broad array of diseases (median follow-up = 608 days). The main results are listed in Table 3 and full results in Table S3. Here we primarily present the results based on Cox regression model, without further adjustments by PTDM or PERR.

Considering all hospitalizations (regardless of history), severe COVID-19 was significantly associated with elevated hazards of hospitalization involving multiple body systems, as described below.

Respiratory diseases Elevated hazards of hospitalization due to respiratory conditions after severe COVID-19 infection were observed. Increased risk of hospital admission was observed for pneumonia (except that caused by tuberculosis) (Hazard ratio [HR]=2.886 (95% Confidence Interval: 2.381-3.498); FDR-adjusted p-value (p.adj)=2.70e-25) (figures in the parentheses organized in the same way below), chronic obstructive pulmonary disease and bronchiectasis (3.013 (2.206-4.114); p.adj=1.06e-10), aspiration pneumonitis (7.332 (5.142-10.453); p.adj=3.06e-26), respiratory failure/insufficiency/arrest (7.442 (4.109-13.480); p.adj=8.58e-10), pneumothorax (12.799 (5.485-29.867); p.adj=7.07e-08), and other specified and unspecified lower respiratory disease (2.694 (1.941-3.739); p.adj=6.07e-08), which includes interstitial pulmonary disorder, hypostatic pneumonia without specified organism, rheumatoid lung disease, etc.

Cardiovascular diseases We observed increased hazards of hospital admission from several cardiovascular/cerebrovascular conditions, including acute myocardial infarction (AMI) (2.369 (1.606-3.494); p.adj=1.34e-04), nonspecific chest pain (1.843 (1.133-2.998); p.adj=4.68e-02), cardiac dysrhythmias (2.162 (1.524-3.066); p.adj=1.45e-04), heart failure (2.797 (1.831-4.273); p.adj=2.30e-05), cerebral infarction (3.028 (2.034-4.508); p.adj=7.99e-07), acute hemorrhagic cerebrovascular disease (4.244 (2.524-7.138); p.adj=8.17e-07), peripheral and visceral vascular disease (3.166 (1.913-5.240); p.adj=7.52e-05) and hypotension (3.512 (2.415-5.109); p.adj=1.16e-09).

Genitourinary diseases There was evidence for increased hazards of hospitalization from genitourinary disorders, including the following conditions: nephritis/nephrosis/renal sclerosis (6.508 (2.731-15.507); p.adj=2.14e-04), acute and unspecified renal failure (5.285 (4.089-6.832); p.adj=5.07e-35), chronic kidney disease (CKD) (2.711 (1.727-4.256); p.adj=1.40e-04), urinary tract infection (UTI) (3.262 (2.635-4.037); p.adj=1.44e-25), calculus of urinary tract (2.284 (1.403-3.716); p.adj=4.63e-03), other specified and unspecified diseases of kidney and ureters (e.g. hydronephrosis, obstructive and reflux uropathy, cyst) (2.693 (1.597-4.540); p.adj=1.34e-03).

Nervous system disorders Increased hazards of hospitalization from neurological disorders were evident and included Parkinson's disease (3.877 (2.056-7.312); p.adj=2.49e-04), headache (including migraine) (1.968 (1.150-3.366); p.adj=4.60e-02), neurocognitive disorders (4.364 (3.098-6.146); p.adj=1.39e-15), transient cerebral ischemia (2.649 (1.565-4.484); p.adj=1.79e-03), coma, stupor and brain damage (9.848 (4.045-23.972); p.adj=6.03e-06) and nerve and nerve root disorders (2.804 (1.301-6.046); p.adj=3.14e-02).

Digestive conditions Our results showed increased hazards of hospitalization due to digestive disorders after severe COVID-19, including oesophageal disorders (1.626 (1.100-2.405); p.adj=4.98e-02), pancreatic disorders (excluding diabetes) (2.980 (1.731-5.132); p.adj=6.20e-04), hemorrhage of digestive tract (1.979 (1.414-2.770); p.adj=5.40e-04), infectious and non-infectious diseases of digestive tract (2.052 (1.176-3.581); p.adj=3.99e-02). Higher hazards of hospitalization due to other specified and unspecified gastrointestinal disorders (2.895 (1.612-5.199); p.adj=2.24e-03) was also observed.

Musculoskeletal disorders Severe COVID-19 were associated with higher hazards of hospitalization from osteomyelitis (5.095 (2.524-10.283); p.adj=5.79e-05), musculoskeletal pain (2.593 (1.955-3.439); p.adj=8.77e-10), low back pain (2.444 (1.615-3.701); p.adj=2.15e-04). Higher hospitalization risks from initial encounters of fracture were observed, including fracture of the spine and back (3.643 (1.893-7.010); p.adj=7.80e-04), the upper limb (2.526 (1.631-3.912); p.adj=2.85e-04), the lower limb (except hip (3.368 (1.769-6.413); p.adj=1.44e-03) and the neck of the femur (hip), (3.708 (2.604-5.282); p.adj=1.12e-11). High hazards of hospital admission were also seen in subsequent encounter of above-mentioned fracture and other injury sequelae, like dislocation, sprain, etc.

Neoplasms History of severe COVID-19 was associated with increased hazards of hospitalization from several neoplasms, including stomach cancers (3.787 (1.820-7.879); p.adj=2.22e-03), colorectal cancers (2.664 (1.842-3.853); p.adj=2.78e-06), brain cancers (5.443 (2.345-12.636); p.adj=6.11e-04), secondary malignancies (2.370 (1.540-3.646); p.adj=6.42e-04) and neoplasms of unspecified nature or uncertain behaviour (2.912 (1.727-4.908); p.adj=4.83e-04).

Other infections History of severe COVID-19 was associated with increased hazards of hospitalization from septicemia (4.089 (3.238-5.164); p.adj=2.71e-30), bacterial infections (4.040 (3.275-4.983); p.adj=7.66e-37), as well as parasitic and other unspecified infections (2.559 (1.793-3.653); p.adj=3.20e-06).

Other sequelae We also observed increased hazards of hospitalization from nutritional anemia (1.568 (1.151-2.137); p.adj=1.81e-02), aplastic anemia (1.887 (1.364-2.611); p.adj=8.82e-04), metabolic disorders, like fluid and electrolyte disorders (2.746 (1.817-4.150); p.adj=1.96e-05), depressive disorders (5.329 (2.380-11.929); p.adj=3.93e-04), and general poor wellbeing, such as syncope (2.455 (1.722-3.500); p.adj=8.58e-06), fever (3.343 (1.840-6.073); p.adj=5.70e-04), and other general signs and symptoms (7.383 (6.797-8.019); p.adj<9.60e-231), abnormal findings without diagnosis (3.605(2.352-5.526); p.adj=7.56e-08).

When we restricted the outcome to hospitalization from *new-onset or recurrent* diseases, most of our results were similar. Higher risks of hospitalization from a few additional disorders were found when we performed analysis on hospitalizations from new-onset diseases, including coronary atherosclerosis and other heart diseases

(2.248 (1.195-4.229); p.adj=4.19e-02), benign neoplasms (1.576 (1.121-2.215); p.adj=3.24e-02), and other specified hereditary and degenerative nervous system conditions (4.763 (1.851-12.257); p.adj=6.10e-03).

We also repeated the analysis using Poisson regression which models the incidence rate of hospitalization. The results were similar to those from Cox regression, with similar significant associations observed (Table S3).

Compared to subjects with mild COVID-19, association of hospitalization with severe COVID-19 (severe vs mild infection)

Here we compared the risk of hospitalization of severe vs mild cases within the infected individuals (Table 4). Compared with mild COVID-19, higher hazards of hospitalization from multiple diseases were observed after severe infection (median follow-up: 248 days), including for example, hospitalization from aplastic anaemia (hazard ratio [HR]= 3.606 (95% CI: 1.854-7.014); FDR-adjusted p-value (p.adj)= 1.08e-03), acute myocardial infarction [AMI] (HR=3.674 (1.788-7.549); p.adj=2.36e-03), cardiac dysrhythmias (HR=4.364 (2.232-8.531); p.adj=1.54e-04), cerebral infarction (HR=6.307 (2.331-17.061); p.adj= 1.79e-03), hypotension (HR=3.965 (1.867-8.423); p.adj= 2.07e-03), acute and unspecified renal failure (HR=7.009 (3.744-13.119); p.adj= 2.38e-08), chronic kidney disease [CKD] (HR=4.769 (1.970-11.542); p.adj= 3.01e-03), urinary tract infections [UTI] (HR=4.533 (2.915-7.048); p.adj= 4.85e-10), calculus of urinary tract (HR=2.992 (1.321-6.777); p.adj= 3.16e-02), neurocognitive disorders (HR=10.081 (4.201-24.193); p.adj= 3.21e-06), pneumonia (except that caused by tuberculosis) (HR=5.226 (3.446-7.924); p.adj= 2.46e-13), chronic obstructive pulmonary disease and bronchiectasis [COPD] (HR=3.633 (1.904-6.935); p.adj= 6.71e-04), and aspiration pneumonitis (HR=3.573 (1.921-6.644); p.adj= 4.66e-04). We also observed increased risk of hospitalization from several digestive disorders, infectious conditions, injury and general signs/symptoms, which are listed in Table 4.

Compared to subjects with no known history of COVID-19, association of hospitalization with mild COVID-19 (mild infection vs uninfected)

We found from this analysis (model D) that history of ‘mild’ (i.e. non-hospitalized) COVID-19 were associated increased hazards of hospital admission due to musculoskeletal pain (excluding low back pain) (HR=1.937 (1.463-2.566); p.adj=4.29e-05), aspiration pneumonitis (HR=2.933 (1.716-5.012); p.adj=6.25e-04), and other general signs and symptoms (like edema, abnormal weight loss, anorexia, and etc.) (HR=1.353 (1.163-1.574); p.adj= 6.59e-04), as shown in Table S3.

Higher hazards of all-cause mortality (1.237 (1.037-1.476); p.adj=2.27e-02) and mortality due to neurocognitive disorders (9.100 (5.590-14.816); p.adj=7.33e-18) were also observed.

Compared with subjects with no known history of COVID-19, association of hospitalization with any COVID-19 infection (any infection vs uninfected)

The results from this analysis (model C) were largely similar to those from model B. Details can be found in Table S3. In addition to findings from model B, we also found increased hazards of hospitalization from essential hypertension (3.061 (1.397-6.705); p.adj=2.09e-02), obesity (6.609 (2.263-19.298); p.adj=3.10e-03), hematuria (9.131 (3.184-26.186); p.adj=3.27e-04), osteoarthritis (1.515 (1.267-1.813); p.adj=5.75e-05), muscle disorders (6.564 (2.390-18.022); p.adj=1.66e-03), lung disease due to external agents (4.211 (1.558-11.384); p.adj=1.88e-02), abdominal pain and other digestive/abdomen signs and symptoms (1.591 (1.098-2.306); p.adj=4.79e-02) and circulatory signs and symptoms (1.645 (1.104-2.451); p.adj=4.88e-02).

Association of hospitalization with severe COVID-19, with PTDM adjustment

Results with PTDM adjustment are listed in Table S3. Since PTDM pushed the start-date of the unexposed to a later date, the number of events was smaller for the unexposed. We relaxed our results presenting criteria (cell count \geq 3 for at least one cell). Under the PTDM adjustment, increased risks of hospitalization were still observed for multiple pulmonary disorders, such as pneumonia (except that caused by tuberculosis) (3.612 (2.944-4.432); p.adj=1.75e-32), COPD (4.325 (3.102-6.029); p.adj=4.13e-16), aspiration pneumonitis (7.796 (5.303-11.460); p.adj=2.00e-23), respiratory failure; insufficiency; arrest (8.559 (4.559-16.070); p.adj=8.64e-10), pneumothorax (19.199 (7.665-48.088); p.adj=9.23e-09), lung disease due to external agents (8.660 (2.210-33.939); p.adj=1.55e-02).

In addition, increased hazards of hospital admission from cardiovascular and several other diseases were observed, such as AMI (2.167 (1.459-3.217); p.adj=1.56e-03), other and ill-defined heart disease (10.610 (2.696-41.752); p.adj=7.15e-03), cardiac dysrhythmias (2.195 (1.539-3.130); p.adj=2.27e-04), heart failure (2.688 (1.731-4.173); p.adj=1.74e-04), cerebral infarction (3.274 (2.167-4.945); p.adj=4.94e-07), acute hemorrhagic cerebrovascular disease (4.251 (2.480-7.288); p.adj=3.36e-06), peripheral and visceral vascular disease (3.986 (2.354-6.747); p.adj=5.93e-06), hypotension (3.002 (2.034-4.431); p.adj=8.06e-07), acute phlebitis; thrombophlebitis and thromboembolism (8.753 (3.037-25.224); p.adj=7.91e-04), nephritis; nephrosis; renal sclerosis (7.471 (2.967-18.814); p.adj=3.07e-04), acute and unspecified renal failure (5.321 (4.051-6.989); p.adj=5.60e-31), chronic kidney disease (5.915 (3.766-9.291); p.adj=6.69e-13); mental disorders (schizophrenia spectrum and other psychotic disorders (9.199 (3.108-27.224); p.adj=8.14e-04)), and rheumatoid arthritis/related disease (2.992 (1.512-5.923); p.adj=1.34e-02). Elevated risks of hospitalization due to some other infectious diseases, injury, neoplasms, and general poor well-being were also observed.

Association of hospitalization with severe COVID-19 with PERR adjustment

The primary results with PERR adjustment (with prior exposure period restricted to 30-Jan-2018 to 30-Jan-2020) are listed in Table 5.

With the PERR adjustment, compared to those with no known history of COVID-19, severe COVID-19 was significantly associated with increased hazards of hospital admission due to pulmonary disorders, like pneumonia (except that caused by tuberculosis) (2.528 (1.725-3.705); $p= 1.98e-06$), aspiration pneumonitis (2.521 (1.099-5.786); $p= 2.91e-02$), respiratory failure; insufficiency; arrest (5.496 (1.593-18.966); $p= 7.01e-03$); and extra-pulmonary system diseases, like AMI (2.432 (1.045-5.658); $p= 3.91e-02$), cardiac dysrhythmias (2.219 (1.249-3.941); $p= 6.57e-03$), cerebral infarction (4.120 (1.989-8.535); $p= 1.39e-04$), peripheral and visceral vascular disease (3.331 (1.129-9.825); $p= 2.93e-02$), acute and unspecified renal failure (4.941 (2.918-8.364); $p= 2.73e-09$), UTI (3.569 (2.228-5.716); $p= 1.20e-07$), neurocognitive disorders (2.786 (1.160-6.689); $p= 2.19e-02$), transient cerebral ischemia (2.829 (1.080-7.409); $p= 3.42e-02$), osteomyelitis (8.640 (1.909-39.096); $p= 5.12e-03$), and low back pain (2.023 (1.097-3.728); $p= 2.40e-02$). Under PERR adjustment, we also found that higher hospitalization risks due to various infections, endocrine disorders, fever, and other general signs.

When compared with mild (non-hospitalized) COVID-19, severe COVID-19 illness (i.e. model A) were associated with higher hazards of hospitalization due to pneumonia (except that caused by tuberculosis) (2.170 (1.140-4.129); $p= 1.83e-02$), cardiac dysrhythmias (3.524 (1.288-9.646); $p= 1.42e-02$), acute and unspecified renal failure (4.025 (1.260-12.854); $p= 1.87e-02$), UTI (2.296 (1.142-4.614); $p= 1.96e-02$), injury and intestinal infection.

Considering any (severe/mild) infection (model C), it was also observed that COVID-19 increased the hazards of hospitalization of COPD (1.635 (1.010-2.648); $p= 4.55e-02$), hypotension (2.069 (1.142-3.751); $p= 1.66e-02$), osteoarthritis (1.388 (1.101-1.750); $p= 5.50e-03$), and other general respiratory symptoms, in addition to most associations found under model B.

We note that the PERR adjustment tends to be conservative in general in our case. By the assumptions of PERR, the prior events should have no effect on the risk of exposure²¹; however here the history of many diseases could increase the risk of (severe) COVID-19 infection. As a result, the hazard ratio of hospitalization (for infected vs non-infected groups) will be higher than expected even in the pre-COVID-19 period, since the (severely) infected group are more likely to be hospitalized for diseases even in the pre-COVID period. This will lead to a conservative bias, as also shown in previous simulation studies²¹.

Association of mortality after severe COVID-19

The primary results of associations with mortality after COVID-19 (without PTDM adjustment) are listed in Table 6 and full results in Table S4. We found that severe COVID-19 was associated with increased all-cause mortality (HR=14.700 (13.835-15.619); p.adj<9.6e-231); significantly increased risk of mortality was also observed for mild disease, albeit with an attenuated effect size (1.237 (1.037-1.476); p.adj= 2.27e-02).

Considering cause-specific mortality (without considering the past history of disease), severe infection was associated with mortality due to pneumonia (except that caused by tuberculosis) (3.404 (1.434-8.083); p.adj= 7.55e-03), COPD (8.448 (5.434-13.134); p.adj= 3.23e-20), AMI (2.658 (1.432-4.931); p.adj= 2.89e-03), coronary atherosclerosis and other heart disease (2.700 (1.549-4.707); p.adj= 8.25e-04), Parkinson's disease (3.540 (1.947-6.439); p.adj= 7.37e-05), multiple sclerosis (19.784 (7.588-51.582); p.adj= 4.76e-09), neurocognitive disorders (13.406 (8.729-20.588); p.adj= 4.28e-31), and death due to several cancers (see Table 6).

Associations with mortality after any infection (model C) are largely similar. When we only considered fatalities from those with a history of the disease, most associations were non-significant, likely due to the small number of events and lower statistical power. Associations with mortality with PTDM adjustment is similar to those without PTDM adjustment (Table S4).

Analysis restricted to the pre-vaccination period

Association of all hospitalization/mortality after COVID-19 before the launch of vaccination (i.e. prior to 8 Dec 2020) are presented in Tables S5 and S6. The spectrum of sequelae remained similar. As for mortality, increased hazards of all-cause mortality and mortality due to specific common causes such as coronary atherosclerosis and other heart disease, acute hemorrhagic cerebrovascular disease, COPD, and other cancers were also observed when the analysis was limited to the 'pre-vaccination' period.

Associations results stratified by vaccination status and follow-up period

The associations with sequelae stratified by vaccination status are presented in Table S7. We will focus on the comparison between subjects with vaccination before (and to a lesser extent, after) the infection against those infected but have not received any vaccination. A few results were nominally significant. For the analysis on breakthrough infections vs infection without vaccination, we observed lower risks of hospitalization from pneumonia, unspecified lower respiratory diseases and other general signs/symptoms from the former group. We observed increased rates of hospitalization from a few non-specific causes, including low back pain, non-specific chest pain and cataracts. However, such results may not be straightforward to interpret as the exact reason for admission is known (e.g. admission could be for an elective operation). The findings should be considered

tentative as most of the above findings are marginally significant except for general signs and symptoms ($p=1.83e-15$).

We then compared those vaccinated after infection against those infected and unvaccinated. The former group had lower hazards of hospitalization from injuries, pneumonia, bacterial infections, septicaemia and other general signs and symptoms, under the original model without PTDM adjustment.

Nevertheless, for subjects who received vaccinations after being infected, we only consider events occurring after vaccination, when comparing with infected but unvaccinated subjects (for whom we will consider all events after infection). As such, the PTDM adjustment may produce more reasonable results. No results were nominally significant under PTDM except for urinary tract infections which was marginally significant.

As for the analysis stratified by follow-up period, we observed larger effects from COVID-19 infection and more statistically significant associations when restricting analysis of outcome to the first 3 months, and in general the effects of infection attenuated over time (Table S8). Nevertheless, it is noteworthy that even after 6 months, infections or severe infections in particular, were still associated with elevated hospitalization risks from numerous conditions. These conditions included, for example, aspiration pneumonitis, acute renal failure, injuries and fractures, and various infections such as urinary tract infections, septicaemia, pneumonia and other bacterial infections (Table S8).

Discussions

Overview

Overall, we found that COVID-19, especially severe disease, were associated with increased risks of hospitalization and/or mortality due to pulmonary, cardiovascular, digestive, neurological, genitourinary and musculoskeletal disorders in the post-infection period. These results were largely consistent and robust to multiple sensitivity analysis, and after PERR and PTDM adjustment.

Interpretation of findings in the context of previous studies

We noted that a recent study by Al-Aly et al. has characterized the post-acute sequelae of COVID-19 in US veterans⁹ for a range of outcomes, however the study was on veterans only which may affect its generalizability to the general population and females. The focus of Al-Aly et al. was also somewhat different from ours, as they focused on incident (new) diagnoses post-infection, while we studied *hospitalization* due to new or recurrent diseases. As such, a person with history of a disorder (e.g. stroke) who was admitted for the same disease post-

infection will be considered by our study but not in Al-Aly et al. We also studied mortality due to all disorders covering all systems.

In this study we utilized a large prospective cohort of UKBB to delineate the septum of hospital admission and mortality cause up to 20 months after COVID-19. We have conducted a very comprehensive set of analyses, which includes all hospitalization and mortality from diseases covering all systems, as well as stratified analysis on new-onset and recurrent diseases. We also performed other sensitivity analyses such as PTDM and PEER adjustments. The PEER adjustment for example can help to reduce residual confounding and help evaluate causal relationships between COVID-19 and possible sequelae, which has not been employed in the above study⁹.

A number of studies have been performed on the longer-term sequelae of COVID-19 (mostly focused on specific organ systems), and have revealed elevated risks for a variety of disorders, which corroborate with our findings in general. For example, impaired lung function and respiratory conditions are among the most commonly reported sequelae of COVID-19^{3,4}. Two previous studies on US veterans^{8,9} have revealed that COVID-19 was associated with a wide range of cardiovascular disorders including dysrhythmias, inflammatory and ischemic heart diseases, thrombotic disorders, heart failure, cerebrovascular conditions and others. The current study also revealed a similar spectrum of cardiovascular in a general population cohort based on an independent sample. Cerebrovascular disorders were also commonly reported as complications of COVID-19, as reviewed elsewhere^{31,32}. Also, renal involvement is not uncommon during the acute phase of infection, and subclinical inflammation may persist for many months^{33,34}. Deterioration of renal function following acute infection has been reported¹¹. Although the mechanisms remain poorly understood, neurocognitive impairment was observed for patients after recovery from COVID-19³⁵⁻³⁷. In a large-scale retrospective cohort study³⁸, it was reported that the incidence of most neurological and psychiatric disorders were raised after COVID-19, compared to those with influenza or other respiratory tract infections, especially for the most severe COVID-19 cases requiring intensive care support.

We also observed that COVID-19 patients were more likely to be admitted for other bacterial or viral infections post-infection. Elevated risk of hospitalization was not only observed for pneumonia, but also other infective conditions such as septicemia, peritonitis and intra-abdominal abscess, intestinal infection, urinary tract infections (UTI), osteomyelitis, and other bacterial infections. The exact mechanisms remain unclear, however for post-influenza infections, it has been suggested that secondary infections could be due to disruption of airway epithelium owing to immune-mediated damage and dysfunctional innate immune responses^{39,40}. Since corticosteroids are often prescribed for severe COVID-19 infections, it may be worthwhile to investigate whether steroid use is associated with increased risks of other infections⁴¹.

Another interesting finding is that we observed increased risk of hospitalization from several musculoskeletal problems, especially fractures. Increased risks of fall may be one important contributing factor, which in turn could be due to myriad reasons such as malaise and fatigue, musculoskeletal pain, muscle weakness, neurocognitive impairment, poor general well-being and the increased incidence of other disorders that may increase fall risk post-infection^{9,42}. Another contributing factor may be increased risks of osteoporosis due to COVID-19 itself⁴³ and/or its treatment (e.g. steroids), but further clinical studies are required. We also noted increased risks of hospitalization from several neoplasms, which may be due to, for example, deconditioning and increased risk of secondary infections after the acute COVID-19 infection, given that it is unlikely that COVID-19 would affect the incidence of neoplasms.

While this is an observational study, it is interesting to note that Mendelian randomization studies, which are less prone to confounding, have also shown that COVID-19 may be a causal risk factor for various cardiometabolic and neuropsychiatric disorders^{44,45} .

The mechanisms underlying COVID-19 sequelae are still largely unknown, and different symptoms/disorders may also involve different pathophysiologies. It has been postulated that, for example, persistent reservoirs of the virus, on-going inflammation, dysregulated immune response and re-activation of pathogens may all contribute to the long-term sequelae^{46,47} .

Clinical implications

Our findings may have important clinical implications. We observed that COVID-19, especially severe infections requiring hospitalization, is associated with a wide variety of pulmonary and extra-pulmonary sequelae. Alarming, elevated risk of admission and mortality was observed for multiple disorders spanning many organ systems. Patients who have recovered from especially severe COVID-19 may therefore require appropriate follow-up, monitoring and risk assessment for different complications. It remains to be investigated what types of interventions may help to reduce the risk of sequelae.

From a public health perspective, when determining policies related to the pandemic (such as lifting restrictions), one should consider the total burden caused by COVID-19 that includes its possible complications, and the disabilities and mortalities due to these complications. Our findings also call for allocation of resources to prevent and treat the relevant complications.

Another alarming finding is that even mild (non-hospitalized) COVID-19 is associated with modestly increased risk of all-cause mortality (HR=1.237, 95% CI 1.037-1.476) and mortality from neurocognitive disorders (HR=9.100, 95% CI: 5.590-14.816), as well as hospital admission from a few disorders such as aspiration pneumonitis,

musculoskeletal pain and other general signs/symptoms. Such findings are in line with previous studies^{31,32} which also showed elevated risks of cardiovascular and other sequelae even for mild COVID-19 cases. Given the huge number of people infected by COVID-19, even a modest rise in the hazard for mortality would imply a large (absolute) number of fatalities. This suggests containment of infection is still important together with efforts to reduce the number of severe cases.

Strength and limitation

The major strength of this study included large sample size of the UK Biobank with detailed health records and sufficient period of follow-up. This study is very comprehensive and covers most clinically relevant categories of diagnosis. In addition to any hospitalization/mortality, we also performed additional analysis on hospitalization with/without relevant history of disease (i.e. new-onset and recurrent diseases). Apart from these, we also made use of advanced statistical methods to evaluate whether our findings are consistent to different modeling strategies. For example, the PERR adjustment was employed to minimize unmeasured confounding.

There are several limitations of this study. Firstly, this is an observational study. Although we have included many known confounders in the model and used other methods (e.g. PERR) to minimize residual confounding, confounding effects cannot be excluded completely. However, many results remained significant after PERR adjustment which provides further support to our findings. Secondly, subclinical disorders may be underestimated and some disorders may be too rare to be included for further analysis. In addition, due to the relatively small number of events for some disorders, absence of significant associations can also be due to inadequate statistical power. We also note that mild or asymptomatic infections may not be detected and considered as having no history of COVID-19; however this is likely to result in a conservative bias. Variant information is not known and thus cannot be analyzed, although we expect the alpha and delta variants are likely the predominant variants. Further studies are required to investigate the sequelae from other variants, such as Omicron. Finally, UKBB participants are likely healthier and of higher socioeconomic status⁴⁸, and generalizability of current findings to other populations (e.g. other ethnicities and those aged <50) may require further studies.

Conclusions

In conclusion, this study revealed increased risk of hospitalization and mortality from a wide variety of pulmonary and extra-pulmonary diseases after COVID-19, especially for those with severe disease. On the other hand, we also observed increased all-cause mortality and mortality due to neurocognitive disorders after mild infections. The findings may have important clinical and public health implications given the huge number of people infected by COVID-19. Proper monitoring and assessments for the risk of sequelae may be warranted for patients recovered from severe COVID-19 in particular. Further studies are required to replicate the current findings and to investigate the mechanisms underlying the sequelae, and interventions for prevention and treatment.

Author Contributions

Conception and design: HCS (lead), with input from YX. Study supervision: HCS. Funding acquisition: HCS. Methodology: HCS (lead), YX. Data curation: YX (lead), RZ, JQ. Data analysis: YX (lead), RZ, JQ. Data interpretation: HCS, YX, RZ, JQ. Preparation of first draft of manuscript: YX, HCS, with input from RZ and JQ.

Acknowledgements

This work was supported partially a National Natural Science Foundation China (NSFC) grant (81971706) and the Lo Kwee Seong Biomedical Research Fund from The Chinese University of Hong Kong and the KIZ-CUHK Joint Laboratory of Bioresources and Molecular Research of Common Diseases, Kunming Institute of Zoology and The Chinese University of Hong Kong, China. We thank Prof. Pak Sham for support on data access.

Supplementary Information

All supplementary Tables and notes are available at the journal's website and at <https://drive.google.com/drive/folders/1Mqy5Qfu8QSe1RAYgy11fcqkY0M-XzZtJ?usp=sharing>

Conflicts of interest

The authors declare no conflict of interest.

References

1. Dong, E., Du, H. & Gardner, L. An interactive web-based dashboard to track COVID-19 in real time. *The Lancet infectious diseases* **20**, 533-534 (2020).
2. Daines, L., Zheng, B., Pfeffer, P., Hurst, J.R. & Sheikh, A. A clinical review of long-COVID with a focus on the respiratory system. *Curr Opin Pulm Med* (2022).
3. Torres-Castro, R. *et al.* Respiratory function in patients post-infection by COVID-19: a systematic review and meta-analysis. *Pulmonology* **27**, 328-337 (2021).
4. José, R.J., Manuel, A., Gibson-Bailey, K. & Lee, L. Post COVID-19 bronchiectasis: a potential epidemic within a pandemic. Vol. 14 1183-1184 (Taylor & Francis, 2020).
5. Wang, F., Kream, R.M. & Stefano, G.B. Long-Term Respiratory and Neurological Sequelae of COVID-19. *Medical science monitor : international medical journal of experimental and clinical research* **26**, e928996-e928996 (2020).
6. Fiani, B., Covarrubias, C., Desai, A., Sekhon, M. & Jarrah, R. A Contemporary Review of Neurological Sequelae of COVID-19. *Frontiers in Neurology* **11**(2020).
7. Taquet, M., Geddes, J.R., Husain, M., Luciano, S. & Harrison, P.J. 6-month neurological and psychiatric outcomes in 236 379 survivors of COVID-19: a retrospective cohort study using electronic health records. *The Lancet Psychiatry* **8**, 416-427 (2021).
8. Xie, Y., Xu, E., Bowe, B. & Al-Aly, Z. Long-term cardiovascular outcomes of COVID-19. *Nature Medicine* **28**, 583-590 (2022).
9. Al-Aly, Z., Xie, Y. & Bowe, B. High-dimensional characterization of post-acute sequelae of COVID-19. *Nature* **594**, 259-264 (2021).
10. Nalbandian, A. *et al.* Post-acute COVID-19 syndrome. *Nature Medicine* **27**, 601-615 (2021).

11. Huang, C. *et al.* 6-month consequences of COVID-19 in patients discharged from hospital: a cohort study. *The Lancet* **397**, 220-232 (2021).
12. Carfi, A., Bernabei, R. & Landi, F. Persistent symptoms in patients after acute COVID-19. *Jama* **324**, 603-605 (2020).
13. Agha, Z., Lofgren, R.P., VanRuiswyk, J.V. & Layde, P.M. Are Patients at Veterans Affairs Medical Centers Sicker?: A Comparative Analysis of Health Status and Medical Resource Use. *Archives of Internal Medicine* **160**, 3252-3257 (2000).
14. Kuodi, P. *et al.* Association between vaccination status and reported incidence of post-acute COVID-19 symptoms in Israel: a cross-sectional study of patients tested between March 2020 and November 2021. *medRxiv*, 2022.01.05.22268800 (2022).
15. Sudlow, C. *et al.* UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS medicine* **12**, e1001779-e1001779 (2015).
16. Cost, H. Utilization Project (HCUP) Agency for healthcare research and quality, Rockville, MD. *Clin Classifications Software Refined (CCSR)* (2020).
17. Waljee, A.K. *et al.* Comparison of imputation methods for missing laboratory data in medicine. *BMJ Open* **3**, e002847 (2013).
18. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)* **57**, 289-300 (1995).
19. Zhao, S.D. & Li, Y. Principled sure independence screening for Cox models with ultra-high-dimensional covariates. *Journal of Multivariate Analysis* **105**, 397-411 (2012).
20. Rodgers, L.R. *et al.* Prior event rate ratio adjustment produced estimates consistent with randomized trial: a diabetes case study. *Journal of Clinical Epidemiology* **122**, 78-86 (2020).
21. Uddin, M.J. *et al.* Performance of prior event rate ratio adjustment method in pharmacoepidemiology: a simulation study. *Pharmacoepidemiol Drug Saf* **24**, 468-77 (2015).
22. Weiner, M.G., Xie, D. & Tannen, R.L. Replication of the Scandinavian Simvastatin Survival Study using a primary care medical record database prompted exploration of a new method to address unmeasured confounding. *Pharmacoepidemiol Drug Saf* **17**, 661-70 (2008).
23. Zhou, Z., Rahme, E., Abrahamowicz, M. & Pilote, L. Survival bias associated with time-to-treatment initiation in drug effectiveness evaluation: a comparison of methods. *Am J Epidemiol* **162**, 1016-23 (2005).
24. Beumont, P., Al-Alami, M. & Touyz, S. Relevance of a standard measurement of undernutrition to the diagnosis of anorexia nervosa: Use of Quetelet's Body Mass Index (BMI). *International Journal of Eating Disorders* **7**, 399-405 (1988).
25. Han, T.S. & Lean, M.E. A clinical perspective of obesity, metabolic syndrome and cardiovascular disease. *JRSM cardiovascular disease* **5**, 2048004016633371 (2016).
26. Ascaso, J.F. *et al.* Diagnosing insulin resistance by simple quantitative methods in subjects with normal glucose metabolism. *Diabetes care* **26**, 3320-3325 (2003).
27. Grundy, S.M. *et al.* 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA Guideline on the Management of Blood Cholesterol: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation* **139**, e1082-e1143 (2019).
28. NICE, N. National Institute for health and care excellence. *Rivaroxaban for the prevention of venous thromboembolism after total hip or total knee replacement in adults [Internet]* (2009).
29. Hawkins, W., Speck, E. & Leonard, V.G. Variation of the hemoglobin level with age and sex. *Blood* **9**, 999-1007 (1954).
30. Askari, M. & Namayandeh, S.M. The Difference between the Population Attributable Risk (PAR) and the Potential Impact Fraction (PIF). *Iranian journal of public health* **49**, 2018-2019 (2020).
31. Jarrahi, A. *et al.* Neurological consequences of COVID-19: what have we learned and where do we go from here? *Journal of neuroinflammation* **17**, 286-286 (2020).
32. Choi, Y. & Lee, M.K. Neuroimaging findings of brain MRI and CT in patients with COVID-19: A systematic review and meta-analysis. *European journal of radiology* **133**, 109393-109393 (2020).
33. Yende, S. & Parikh, C.R. Long COVID and kidney disease. *Nature Reviews Nephrology* **17**, 792-793 (2021).
34. Long, J.D., Strohbehn, I., Sawtell, R., Bhattacharyya, R. & Sise, M.E. COVID-19 Survival and its impact on chronic kidney disease. *Transl Res* **241**, 70-82 (2022).
35. Mukaetova-Ladinska, E.B., Kronenberg, G. & Raha-Chowdhury, R. COVID-19 and neurocognitive disorders. *Current opinion in psychiatry* **34**, 149-156 (2021).

36. Dąbrowska, E., Galińska-Skok, B. & Waszkiewicz, N. Depressive and Neurocognitive Disorders in the Context of the Inflammatory Background of COVID-19. *Life (Basel, Switzerland)* **11**, 1056 (2021).
37. Woo, M.S. *et al.* Frequent neurocognitive deficits after recovery from mild COVID-19. *Brain communications* **2**, fcaa205-fcaa205 (2020).
38. Asriati, S.Q. 6-month neurological and psychiatric outcomes in 236 379 survivors of COVID-19: a retrospective cohort study using electronic health records. *Jurnal Latihan* **1**, 1-12 (2021).
39. Metzger, D.W. & Sun, K. Immune dysfunction and bacterial coinfections following influenza. *The Journal of Immunology* **191**, 2047-2052 (2013).
40. McCullers, J.A. The co-pathogenesis of influenza viruses with bacteria in the lung. *Nature Reviews Microbiology* **12**, 252-262 (2014).
41. Garcia-Vidal, C. *et al.* Incidence of co-infections and superinfections in hospitalized patients with COVID-19: a retrospective cohort study. *Clin Microbiol Infect* **27**, 83-88 (2021).
42. Wilkinson, E. UK Government urged to recognise post-COVID-19 cancer backlog. *The Lancet Oncology* **22**, 910 (2021).
43. Awosanya, O.D. *et al.* Osteoclast-mediated bone loss observed in a COVID-19 mouse model. *Bone* **154**, 116227 (2022).
44. Xiang, Y., Chau, C.K.-L., Qiu, J., Rao, S. & So, H.-C. Exploring causal relationships between COVID-19 and cardiometabolic disorders: A bi-directional Mendelian randomization study. *medRxiv*, 2021.03.20.21254008 (2021).
45. Xiang, Y. *et al.* Neuropsychiatric disorders as risk factors and consequences of COVID-19: A Mendelian randomization study. *medRxiv*, 2021.06.29.21259609 (2021).
46. Mehandru, S. & Merad, M. Pathological sequelae of long-haul COVID. *Nature Immunology* **23**, 194-202 (2022).
47. Ortona, E. & Malorni, W. Long COVID: to investigate immunological mechanisms and sex/gender related aspects as fundamental steps for tailored therapy. *The European respiratory journal* **59**, 2102245 (2022).
48. Fry, A. *et al.* Comparison of Sociodemographic and Health-Related Characteristics of UK Biobank Participants With Those of the General Population. *Am J Epidemiol* **186**, 1026-1034 (2017).

Table 1 The four sets of analyses performed in this study

Model	Cohort 1	Cohort 2
A	Hospitalized or fatal (U07.1)	Non-hospitalized infection
B	Hospitalized or fatal (U07.1)	UKBB subjects without COVID-19 Dx or tested negative
C	All subjects with infection	UKBB subjects without COVID-19 Dx or tested negative
D	Non-hospitalized infection	UKBB subjects without COVID-19 Dx or tested negative

Model A was restricted to infected patients. Other models also involves UKBB subjects with no known history of infection. U07.1 is the code for fatal (laboratory-confirmed) COVID-19 infection based on the latest ICD coding. Dx, diagnosis.

Table 2 Number of available subjects in models A to D

Model	Cohort 1	Cohort 2	Total
A	6484	18976	25460
B	6484	386636	393120
C	25460	386636	412096
D	18976	386636	405612

Table 3. Compared with subjects with no known history of COVID, the associations of severe COVID-19 with hospitalization due to various disorders, without PTDM adjustment (i.e. model B)

Outcome	coef	exp(coef)	se(coef)	z	Pr(> z)	p.adj	2.5.%	97.5.%	CCSR Category Description
BLD001	0.450	1.568	0.158	2.848	4.40E-03	1.81E-02	1.151	2.137	Nutritional anemia
BLD003	0.635	1.887	0.166	3.836	1.25E-04	8.82E-04	1.364	2.611	Aplastic anemia
BLD010	1.343	3.832	0.472	2.848	4.39E-03	1.81E-02	1.520	9.659	Other specified and unspecified hematologic conditions
CIR009	0.862	2.369	0.198	4.347	1.38E-05	1.34E-04	1.606	3.494	Acute myocardial infarction
CIR012	0.612	1.843	0.248	2.464	1.37E-02	4.68E-02	1.133	2.998	Nonspecific chest pain
CIR017	0.771	2.162	0.178	4.324	1.53E-05	1.45E-04	1.524	3.066	Cardiac dysrhythmias
CIR019	1.029	2.797	0.216	4.755	1.98E-06	2.30E-05	1.831	4.273	Heart failure
CIR020	1.108	3.028	0.203	5.456	4.87E-08	7.99E-07	2.034	4.508	Cerebral infarction
CIR021	1.446	4.244	0.265	5.450	5.04E-08	8.17E-07	2.524	7.138	Acute hemorrhagic cerebrovascular disease
CIR026	1.152	3.166	0.257	4.482	7.39E-06	7.52E-05	1.913	5.240	Peripheral and visceral vascular disease
CIR031	1.256	3.512	0.191	6.570	5.02E-11	1.16E-09	2.415	5.109	Hypotension
DIG001	1.001	2.721	0.160	6.272	3.57E-10	7.95E-09	1.990	3.720	Intestinal infection
DIG004	0.486	1.626	0.200	2.436	1.49E-02	4.98E-02	1.100	2.405	Esophageal disorders
DIG012	0.890	2.435	0.253	3.522	4.28E-04	2.52E-03	1.484	3.996	Intestinal obstruction and ileus
DIG014	0.641	1.899	0.222	2.889	3.86E-03	1.63E-02	1.229	2.935	Hemorrhoids
DIG015	0.761	2.140	0.228	3.329	8.70E-04	4.55E-03	1.367	3.349	Anal and rectal conditions
DIG016	1.450	4.264	0.271	5.354	8.60E-08	1.30E-06	2.508	7.251	Peritonitis and intra-abdominal abscess
DIG017	1.091	2.977	0.238	4.581	4.63E-06	5.02E-05	1.867	4.749	Biliary tract disease
DIG020	1.092	2.980	0.277	3.938	8.22E-05	6.20E-04	1.731	5.132	Pancreatic disorders (excluding diabetes)
DIG021	0.683	1.979	0.172	3.981	6.86E-05	5.40E-04	1.414	2.770	Gastrointestinal hemorrhage
DIG022	0.719	2.052	0.284	2.530	1.14E-02	3.99E-02	1.176	3.581	Noninfectious gastroenteritis
DIG025	1.063	2.895	0.299	3.558	3.74E-04	2.24E-03	1.612	5.199	Other specified and unspecified gastrointestinal disorders
END011	1.010	2.746	0.211	4.793	1.65E-06	1.96E-05	1.817	4.150	Fluid and electrolyte disorders
END015	1.096	2.991	0.261	4.193	2.75E-05	2.44E-04	1.792	4.991	Other specified and unspecified endocrine disorders

EYE002	0.402	1.495	0.155	2.602	9.26E-03	3.37E-02	1.104	2.024	Cataract and other lens disorders
FAC010	0.661	1.937	0.223	2.965	3.03E-03	1.32E-02	1.251	2.997	Other aftercare encounter
GEN001	1.873	6.508	0.443	4.228	2.36E-05	2.14E-04	2.731	15.507	Nephritis; nephrosis; renal sclerosis
GEN002	1.665	5.285	0.131	12.713	5.02E-37	5.07E-35	4.089	6.832	Acute and unspecified renal failure
GEN003	0.997	2.711	0.230	4.336	1.45E-05	1.40E-04	1.727	4.256	Chronic kidney disease
GEN004	1.182	3.262	0.109	10.862	1.76E-27	1.44E-25	2.635	4.037	Urinary tract infections
GEN005	0.826	2.284	0.248	3.324	8.88E-04	4.63E-03	1.403	3.716	Calculus of urinary tract
GEN006	0.991	2.693	0.266	3.718	2.01E-04	1.34E-03	1.597	4.540	Other specified and unspecified diseases of kidney and ureters
INF002	1.408	4.089	0.119	11.825	2.89E-32	2.71E-30	3.238	5.164	Septicemia
INF003	1.396	4.040	0.107	13.043	7.00E-39	7.66E-37	3.275	4.983	Bacterial infections
INF009	0.940	2.559	0.182	5.176	2.27E-07	3.20E-06	1.793	3.653	Parasitic, other specified and unspecified infections
INJ002	1.293	3.643	0.334	3.871	1.08E-04	7.80E-04	1.893	7.010	Fracture of the spine and back, initial encounter
INJ004	0.927	2.526	0.223	4.152	3.30E-05	2.85E-04	1.631	3.912	Fracture of the upper limb, initial encounter
INJ005	1.214	3.368	0.329	3.695	2.20E-04	1.44E-03	1.769	6.413	Fracture of the lower limb (except hip), initial encounter
INJ006	1.311	3.708	0.180	7.264	3.75E-13	1.12E-11	2.604	5.282	Fracture of the neck of the femur (hip), initial encounter
INJ039	1.293	3.643	0.334	3.871	1.08E-04	7.80E-04	1.893	7.010	Fracture of the spine and back, subsequent encounter
INJ041	0.927	2.526	0.223	4.152	3.30E-05	2.85E-04	1.631	3.912	Fracture of the upper limb, subsequent encounter
INJ042	1.214	3.368	0.329	3.695	2.20E-04	1.44E-03	1.769	6.413	Fracture of lower limb (except hip), subsequent encounter
INJ043	1.311	3.708	0.180	7.264	3.75E-13	1.12E-11	2.604	5.282	Fracture of the neck of the femur (hip), subsequent encounter
INJ073	1.094	2.985	0.076	14.342	1.19E-46	1.57E-44	2.571	3.466	Injury, sequela
MBD002	1.673	5.329	0.411	4.069	4.73E-05	3.93E-04	2.380	11.929	Depressive disorders
MUS002	1.628	5.095	0.358	4.544	5.51E-06	5.79E-05	2.524	10.283	Osteomyelitis
MUS010	0.953	2.593	0.144	6.617	3.68E-11	8.77E-10	1.955	3.439	Musculoskeletal pain, not low back pain
MUS038	0.894	2.444	0.212	4.223	2.41E-05	2.15E-04	1.615	3.701	Low back pain
NEO013	1.331	3.787	0.374	3.562	3.69E-04	2.22E-03	1.820	7.879	Gastrointestinal cancers - stomach
NEO015	0.980	2.664	0.188	5.204	1.95E-07	2.78E-06	1.842	3.853	Gastrointestinal cancers - colorectal
NEO048	1.694	5.443	0.430	3.943	8.05E-05	6.11E-04	2.345	12.636	Nervous system cancers - brain

NEO070	0.863	2.370	0.220	3.925	8.66E-05	6.42E-04	1.540	3.646	Secondary malignancies
NEO072	1.069	2.912	0.266	4.012	6.03E-05	4.83E-04	1.727	4.908	Neoplasms of unspecified nature or uncertain behavior
NVS004	1.355	3.877	0.324	4.187	2.83E-05	2.49E-04	2.056	7.312	Parkinson`s disease
NVS010	0.677	1.968	0.274	2.472	1.34E-02	4.60E-02	1.150	3.366	Headache; including migraine
NVS011	1.473	4.364	0.175	8.429	3.50E-17	1.39E-15	3.098	6.146	Neurocognitive disorders
NVS012	0.974	2.649	0.269	3.629	2.85E-04	1.79E-03	1.565	4.484	Transient cerebral ischemia
NVS013	2.287	9.848	0.454	5.039	4.68E-07	6.03E-06	4.045	23.972	Coma; stupor; and brain damage
NVS017	1.031	2.804	0.392	2.630	8.53E-03	3.14E-02	1.301	6.046	Nerve and nerve root disorders
RSP002	1.060	2.886	0.098	10.799	3.49E-27	2.70E-25	2.381	3.498	Pneumonia (except that caused by tuberculosis)
RSP008	1.103	3.013	0.159	6.939	3.96E-12	1.06E-10	2.206	4.114	Chronic obstructive pulmonary disease and bronchiectasis
RSP010	1.992	7.332	0.181	11.008	3.50E-28	3.06E-26	5.142	10.453	Aspiration pneumonitis
RSP012	2.007	7.442	0.303	6.623	3.53E-11	8.58E-10	4.109	13.480	Respiratory failure; insufficiency; arrest
RSP014	2.549	12.799	0.432	5.896	3.71E-09	7.07E-08	5.485	29.867	Pneumothorax
RSP016	0.991	2.694	0.167	5.926	3.11E-09	6.07E-08	1.941	3.739	Other specified and unspecified lower respiratory disease
SYM001	0.898	2.455	0.181	4.965	6.86E-07	8.58E-06	1.722	3.500	Syncope
SYM002	1.207	3.343	0.305	3.963	7.40E-05	5.70E-04	1.840	6.073	Fever
SYM010	0.919	2.508	0.181	5.075	3.87E-07	5.09E-06	1.758	3.577	Nervous system signs and symptoms
SYM013	0.745	2.107	0.244	3.051	2.28E-03	1.05E-02	1.305	3.401	Respiratory signs and symptoms
SYM014	0.978	2.660	0.393	2.492	1.27E-02	4.39E-02	1.232	5.741	Skin/Subcutaneous signs and symptoms
SYM016	1.999	7.383	0.042	47.386	0.00E+00	0.00E+00	6.797	8.019	Other general signs and symptoms
SYM017	1.282	3.605	0.218	5.883	4.03E-09	7.56E-08	2.352	5.526	Abnormal findings without diagnosis

Table 4. Compared to subjects with mild (non-hospitalized) COVID, the associations of severe COVID-19 with hospitalization due to various disorders, without PTDM adjustment (i.e. model A)

Outcome	coef	exp(coef)	se(coef)	z	Pr(> z)	p.adj	2.5.%	97.5.%	CCSR Category Description
BLD003	1.283	3.606	0.339	3.779	1.57E-04	1.08E-03	1.854	7.014	Aplastic anemia
CIR009	1.301	3.674	0.367	3.542	3.98E-04	2.36E-03	1.788	7.549	Acute myocardial infarction
CIR017	1.473	4.364	0.342	4.307	1.65E-05	1.54E-04	2.232	8.531	Cardiac dysrhythmias
CIR020	1.842	6.307	0.508	3.627	2.87E-04	1.79E-03	2.331	17.061	Cerebral infarction
CIR031	1.378	3.965	0.384	3.584	3.39E-04	2.07E-03	1.867	8.423	Hypotension
DIG001	1.656	5.237	0.330	5.011	5.42E-07	6.91E-06	2.740	10.008	Intestinal infection
DIG016	1.489	4.432	0.486	3.065	2.17E-03	1.01E-02	1.711	11.483	Peritonitis and intra-abdominal abscess
DIG017	1.361	3.899	0.449	3.031	2.44E-03	1.10E-02	1.617	9.400	Biliary tract disease
DIG020	1.428	4.172	0.471	3.030	2.44E-03	1.10E-02	1.656	10.508	Pancreatic disorders (excluding diabetes)
DIG021	0.768	2.156	0.271	2.838	4.54E-03	1.86E-02	1.268	3.665	Gastrointestinal hemorrhage
DIG022	1.504	4.500	0.527	2.854	4.32E-03	1.81E-02	1.602	12.643	Noninfectious gastroenteritis
END015	1.375	3.955	0.556	2.474	1.33E-02	4.58E-02	1.331	11.756	Other specified and unspecified endocrine disorders
FAC009	1.719	5.582	0.548	3.137	1.71E-03	8.30E-03	1.906	16.342	Implant, device or graft related encounter
GEN002	1.947	7.009	0.320	6.088	1.14E-09	2.38E-08	3.744	13.119	Acute and unspecified renal failure
GEN003	1.562	4.769	0.451	3.464	5.32E-04	3.01E-03	1.970	11.542	Chronic kidney disease
GEN004	1.511	4.533	0.225	6.711	1.94E-11	4.85E-10	2.915	7.048	Urinary tract infections
GEN005	1.096	2.992	0.417	2.627	8.62E-03	3.16E-02	1.321	6.777	Calculus of urinary tract
GEN006	1.678	5.353	0.481	3.486	4.91E-04	2.81E-03	2.084	13.748	Other specified and unspecified diseases of kidney and ureters
INF002	1.798	6.039	0.268	6.709	1.96E-11	4.85E-10	3.571	10.212	Septicemia
INF003	1.758	5.801	0.237	7.434	1.06E-13	3.46E-12	3.649	9.222	Bacterial infections
INF009	1.693	5.438	0.374	4.533	5.81E-06	6.05E-05	2.615	11.310	Parasitic, other specified and unspecified infections
INJ006	1.703	5.489	0.392	4.349	1.37E-05	1.34E-04	2.548	11.824	Fracture of the neck of the femur (hip), initial encounter
INJ043	1.703	5.489	0.392	4.349	1.37E-05	1.34E-04	2.548	11.824	Fracture of the neck of the femur (hip), subsequent encounter
INJ073	1.184	3.269	0.131	9.020	1.87E-19	8.49E-18	2.527	4.228	Injury, sequela

MUS010	0.593	1.810	0.215	2.757	5.83E-03	2.34E-02	1.187	2.760	Musculoskeletal pain, not low back pain
NEO015	1.279	3.593	0.318	4.023	5.74E-05	4.66E-04	1.927	6.701	Gastrointestinal cancers - colorectal
NEO070	1.628	5.095	0.439	3.706	2.11E-04	1.40E-03	2.153	12.056	Secondary malignancies
NVS011	2.311	10.081	0.447	5.173	2.30E-07	3.21E-06	4.201	24.193	Neurocognitive disorders
RSP002	1.654	5.226	0.212	7.786	6.93E-15	2.46E-13	3.446	7.924	Pneumonia (except that caused by tuberculosis)
RSP008	1.290	3.633	0.330	3.912	9.15E-05	6.71E-04	1.904	6.935	Chronic obstructive pulmonary disease and bronchiectasis
RSP010	1.273	3.573	0.317	4.023	5.74E-05	4.66E-04	1.921	6.644	Aspiration pneumonitis
RSP011	1.516	4.552	0.573	2.645	8.17E-03	3.07E-02	1.481	13.996	Pleurisy, pleural effusion and pulmonary collapse
RSP016	1.270	3.559	0.307	4.137	3.52E-05	3.00E-04	1.950	6.495	Other specified and unspecified lower respiratory disease
SYM001	0.933	2.542	0.297	3.140	1.69E-03	8.24E-03	1.420	4.550	Syncope
SYM010	0.729	2.073	0.279	2.614	8.95E-03	3.27E-02	1.200	3.582	Nervous system signs and symptoms
SYM016	2.108	8.231	0.090	23.440	1.65E-121	3.62E-119	6.901	9.817	Other general signs and symptoms
SYM017	1.192	3.294	0.379	3.146	1.65E-03	8.13E-03	1.567	6.921	Abnormal findings without diagnosis

Table 5. The associations of any hospitalization with severe COVID-19 due to various disorders, with *PERR* adjustment

Outcome	Model	beta diff	boot sd	zval final	pval final	HR	CI lower	CI upper	no.boot	CCSR.Category.Description
CIR009	B	0.889	0.431	2.063	3.91E-02	2.432	1.045	5.658	200	Acute myocardial infarction
CIR017	A	1.260	0.514	2.452	1.42E-02	3.524	1.288	9.646	200	Cardiac dysrhythmias
CIR017	B	0.797	0.293	2.718	6.57E-03	2.219	1.249	3.941	200	Cardiac dysrhythmias
CIR020	B	1.416	0.372	3.810	1.39E-04	4.120	1.989	8.535	200	Cerebral infarction
CIR026	B	1.203	0.552	2.180	2.93E-02	3.331	1.129	9.825	200	Peripheral and visceral vascular disease
CIR031	C	0.727	0.303	2.396	1.66E-02	2.069	1.142	3.751	200	Hypotension
DIG001	A	1.157	0.469	2.466	1.37E-02	3.180	1.268	7.974	200	Intestinal infection
DIG001	B	0.996	0.302	3.301	9.63E-04	2.707	1.499	4.889	200	Intestinal infection
DIG001	C	0.460	0.197	2.339	1.94E-02	1.584	1.077	2.329	200	Intestinal infection
DIG016	B	1.505	0.511	2.944	3.24E-03	4.503	1.653	12.262	200	Peritonitis and intra-abdominal abscess
DIG016	C	0.793	0.330	2.401	1.64E-02	2.211	1.157	4.225	200	Peritonitis and intra-abdominal abscess
DIG017	B	0.903	0.421	2.148	3.17E-02	2.468	1.082	5.627	200	Biliary tract disease
DIG021	B	0.742	0.316	2.345	1.90E-02	2.100	1.130	3.904	200	Gastrointestinal hemorrhage
DIG021	C	0.417	0.180	2.313	2.07E-02	1.517	1.066	2.160	200	Gastrointestinal hemorrhage
DIG025	B	1.448	0.500	2.896	3.78E-03	4.255	1.597	11.336	200	Other specified and unspecified gastrointestinal disorders
END015	B	1.672	0.584	2.864	4.19E-03	5.323	1.695	16.716	200	Other specified and unspecified endocrine disorders
EYE002	C	0.380	0.133	2.866	4.16E-03	1.463	1.128	1.898	200	Cataract and other lens disorders
GEN002	A	1.393	0.592	2.351	1.87E-02	4.025	1.260	12.854	200	Acute and unspecified renal failure
GEN002	B	1.597	0.269	5.947	2.73E-09	4.941	2.918	8.364	200	Acute and unspecified renal failure
GEN002	C	1.172	0.218	5.367	8.00E-08	3.229	2.105	4.955	200	Acute and unspecified renal failure
GEN004	A	0.831	0.356	2.334	1.96E-02	2.296	1.142	4.614	200	Urinary tract infections
GEN004	B	1.272	0.240	5.294	1.20E-07	3.569	2.228	5.716	200	Urinary tract infections
GEN004	C	0.759	0.162	4.683	2.83E-06	2.135	1.554	2.933	200	Urinary tract infections
INF002	B	1.004	0.224	4.491	7.10E-06	2.730	1.761	4.231	200	Septicemia
INF002	C	0.712	0.185	3.852	1.17E-04	2.037	1.418	2.926	200	Septicemia
INF003	B	1.284	0.199	6.439	1.20E-10	3.610	2.442	5.335	200	Bacterial infections

INF003	C	0.887	0.164	5.396	6.80E-08	2.427	1.759	3.350	200	Bacterial infections
INF009	B	0.719	0.359	2.001	4.54E-02	2.052	1.015	4.148	200	Parasitic, other specified and unspecified infections
INJ004	C	0.486	0.229	2.121	3.39E-02	1.626	1.038	2.548	200	Fracture of the upper limb, initial encounter
INJ041	C	0.486	0.239	2.031	4.22E-02	1.626	1.017	2.599	200	Fracture of the upper limb, subsequent encounter
INJ073	A	0.836	0.205	4.069	4.72E-05	2.306	1.542	3.449	200	Injury, sequela
INJ073	B	0.850	0.130	6.516	7.21E-11	2.339	1.812	3.021	200	Injury, sequela
INJ073	C	0.445	0.097	4.580	4.66E-06	1.561	1.290	1.888	200	Injury, sequela
MUS002	B	2.156	0.770	2.800	5.12E-03	8.640	1.909	39.096	200	Osteomyelitis
MUS002	C	1.616	0.626	2.581	9.86E-03	5.033	1.475	17.173	200	Osteomyelitis
MUS006	C	0.328	0.118	2.776	5.50E-03	1.388	1.101	1.750	200	Osteoarthritis
MUS010	B	0.929	0.225	4.128	3.65E-05	2.531	1.629	3.934	200	Musculoskeletal pain, not low back pain
MUS010	C	0.674	0.130	5.185	2.16E-07	1.962	1.521	2.531	200	Musculoskeletal pain, not low back pain
MUS038	B	0.704	0.312	2.258	2.40E-02	2.023	1.097	3.728	200	Low back pain
NEO015	B	0.859	0.351	2.450	1.43E-02	2.360	1.187	4.692	200	Gastrointestinal cancers - colorectal
NVS010	C	0.598	0.272	2.195	2.82E-02	1.818	1.066	3.102	200	Headache; including migraine
NVS011	B	1.025	0.447	2.293	2.19E-02	2.786	1.160	6.689	200	Neurocognitive disorders
NVS012	B	1.040	0.491	2.117	3.42E-02	2.829	1.080	7.409	200	Transient cerebral ischemia
RSP002	A	0.775	0.328	2.359	1.83E-02	2.170	1.140	4.129	200	Pneumonia (except that caused by tuberculosis)
RSP002	B	0.927	0.195	4.755	1.98E-06	2.528	1.725	3.705	200	Pneumonia (except that caused by tuberculosis)
RSP002	C	0.579	0.146	3.981	6.86E-05	1.785	1.342	2.374	200	Pneumonia (except that caused by tuberculosis)
RSP008	C	0.492	0.246	2.000	4.55E-02	1.635	1.010	2.648	200	Chronic obstructive pulmonary disease and bronchiectasis
RSP010	B	0.925	0.424	2.182	2.91E-02	2.521	1.099	5.786	200	Aspiration pneumonitis
RSP010	C	1.045	0.360	2.905	3.68E-03	2.842	1.405	5.752	200	Aspiration pneumonitis
RSP012	B	1.704	0.632	2.696	7.01E-03	5.496	1.593	18.966	200	Respiratory failure; insufficiency; arrest
RSP012	C	1.431	0.666	2.151	3.15E-02	4.184	1.135	15.421	200	Respiratory failure; insufficiency; arrest
RSP016	B	0.646	0.265	2.440	1.47E-02	1.908	1.136	3.207	200	Other specified and unspecified lower respiratory disease
RSP016	C	0.535	0.213	2.519	1.18E-02	1.708	1.126	2.590	200	Other specified and unspecified lower respiratory disease
SYM002	B	1.305	0.624	2.092	3.64E-02	3.688	1.086	12.524	200	Fever
SYM010	B	0.890	0.337	2.644	8.20E-03	2.436	1.259	4.714	200	Nervous system signs and symptoms
SYM010	C	0.515	0.220	2.335	1.95E-02	1.673	1.086	2.577	200	Nervous system signs and symptoms

SYM013	C	0.767	0.304	2.522	1.17E-02	2.153	1.186	3.906	200	Respiratory signs and symptoms
SYM016	A	1.763	0.181	9.739	2.05E-22	5.828	4.088	8.310	200	Other general signs and symptoms
SYM016	B	2.009	0.090	22.267	7.69E-110	7.456	6.248	8.899	200	Other general signs and symptoms
SYM016	C	1.268	0.067	19.062	5.22E-81	3.553	3.119	4.048	200	Other general signs and symptoms
SYM017	B	2.059	0.414	4.971	6.65E-07	7.835	3.480	17.640	200	Abnormal findings without diagnosis
SYM017	C	1.315	0.289	4.552	5.32E-06	3.726	2.115	6.566	200	Abnormal findings without diagnosis

Table 6. Association of mortality with severe COVID-19, without PTDM adjustment

Outcome	coef	exp(coef)	se(coef)	z	Pr(> z)	p.adj	2.5.%	97.5.%	Model	Analysis	CCSR Category Description
All-cause Mortality	3.055	21.229	0.094	32.536	3.27E-232	9.60E-231	17.661	25.520	A	All-cause mortality	
All-cause Mortality	2.688	14.700	0.031	86.872	0.00E+00	0.00E+00	13.835	15.619	B	All-cause mortality	
All-cause Mortality	2.035	7.655	0.029	69.261	0.00E+00	0.00E+00	7.226	8.109	C	All-cause mortality	
All-cause Mortality	0.213	1.237	0.090	2.360	1.83E-02	2.27E-02	1.037	1.476	D	All-cause mortality	
NEO015	1.871	6.493	0.537	3.487	4.89E-04	8.30E-04	2.269	18.583	A	Any death	Gastrointestinal cancers - colorectal
NVS011	0.771	2.162	0.311	2.475	1.33E-02	1.70E-02	1.174	3.980	A	Any death	Neurocognitive disorders
CIR009	0.978	2.658	0.315	3.100	1.94E-03	2.89E-03	1.432	4.931	B	Any death	Acute myocardial infarction
CIR011	0.993	2.700	0.284	3.504	4.59E-04	8.25E-04	1.549	4.707	B	Any death	Coronary atherosclerosis and other heart disease
GEN004	2.627	13.832	0.407	6.461	1.04E-10	5.70E-10	6.235	30.687	B	Any death	Urinary tract infections
NEO013	1.619	5.050	0.484	3.344	8.25E-04	1.34E-03	1.955	13.048	B	Any death	Gastrointestinal cancers - stomach
NEO015	1.396	4.040	0.256	5.448	5.08E-08	1.94E-07	2.445	6.676	B	Any death	Gastrointestinal cancers - colorectal
NEO018	2.026	7.587	0.484	4.189	2.80E-05	6.16E-05	2.940	19.582	B	Any death	Gastrointestinal cancers - bile duct
NEO025	2.100	8.168	0.493	4.256	2.08E-05	4.82E-05	3.105	21.487	B	Any death	Skin cancers - melanoma
NEO039	1.826	6.208	0.252	7.233	4.72E-13	3.19E-12	3.785	10.181	B	Any death	Male reproductive system cancers - prostate
NEO043	1.940	6.962	0.396	4.898	9.68E-07	2.94E-06	3.203	15.133	B	Any death	Urinary system cancers - bladder
NEO048	2.003	7.411	0.401	5.001	5.71E-07	1.93E-06	3.380	16.249	B	Any death	Nervous system cancers - brain
NEO051	1.317	3.732	0.331	3.984	6.79E-05	1.39E-04	1.952	7.134	B	Any death	Endocrine system cancers - pancreas
NEO071	2.118	8.314	0.439	4.828	1.38E-06	3.92E-06	3.519	19.646	B	Any death	Malignant neoplasm, unspecified
NVS004	1.264	3.540	0.305	4.143	3.43E-05	7.37E-05	1.947	6.439	B	Any death	Parkinson's disease
NVS005	2.985	19.784	0.489	6.105	1.03E-09	4.76E-09	7.588	51.582	B	Any death	Multiple sclerosis
NVS011	2.596	13.406	0.219	11.858	1.95E-32	4.28E-31	8.729	20.588	B	Any death	Neurocognitive disorders
RSP002	1.225	3.404	0.441	2.777	5.49E-03	7.55E-03	1.434	8.083	B	Any death	Pneumonia (except that caused by tuberculosis)
RSP008	2.134	8.448	0.225	9.479	2.57E-21	3.23E-20	5.434	13.134	B	Any death	Chronic obstructive pulmonary disease and bronchiectasis

CIR011	0.718	2.050	0.251	2.855	4.31E-03	6.02E-03	1.252	3.355	C	Any death	Coronary atherosclerosis and other heart disease
EXT020	1.495	4.459	0.498	2.999	2.71E-03	3.97E-03	1.679	11.844	C	Any death	External cause codes: intent of injury, accidental/unintentional
EXT029	1.412	4.105	0.493	2.866	4.15E-03	5.90E-03	1.563	10.781	C	Any death	External cause codes: subsequent encounter
EXT030	1.412	4.105	0.493	2.866	4.15E-03	5.90E-03	1.563	10.781	C	Any death	External cause codes: sequela
GEN004	2.221	9.217	0.389	5.706	1.15E-08	4.83E-08	4.298	19.766	C	Any death	Urinary tract infections
NEO013	1.095	2.990	0.442	2.477	1.33E-02	1.70E-02	1.257	7.113	C	Any death	Gastrointestinal cancers - stomach
NEO015	0.867	2.380	0.226	3.840	1.23E-04	2.35E-04	1.529	3.704	C	Any death	Gastrointestinal cancers - colorectal
NEO018	1.377	3.963	0.442	3.117	1.83E-03	2.77E-03	1.667	9.421	C	Any death	Gastrointestinal cancers - bile duct
NEO025	1.254	3.504	0.486	2.582	9.82E-03	1.31E-02	1.353	9.078	C	Any death	Skin cancers - melanoma
NEO039	1.345	3.837	0.242	5.560	2.70E-08	1.08E-07	2.389	6.165	C	Any death	Male reproductive system cancers - prostate
NEO043	1.268	3.552	0.387	3.277	1.05E-03	1.68E-03	1.664	7.582	C	Any death	Urinary system cancers - bladder
NEO048	1.039	2.827	0.376	2.762	5.75E-03	7.79E-03	1.352	5.912	C	Any death	Nervous system cancers - brain
NEO071	1.529	4.614	0.389	3.935	8.32E-05	1.66E-04	2.154	9.881	C	Any death	Malignant neoplasm, unspecified
NVS004	1.008	2.741	0.253	3.991	6.57E-05	1.38E-04	1.671	4.497	C	Any death	Parkinson's disease
NVS005	2.157	8.645	0.479	4.506	6.61E-06	1.62E-05	3.383	22.092	C	Any death	Multiple sclerosis
NVS011	1.404	4.072	0.172	8.159	3.37E-16	2.96E-15	2.906	5.706	C	Any death	Neurocognitive disorders
RSP002	0.957	2.603	0.385	2.485	1.29E-02	1.70E-02	1.224	5.535	C	Any death	Pneumonia (except that caused by tuberculosis)
RSP008	1.647	5.190	0.214	7.706	1.30E-14	1.04E-13	3.414	7.890	C	Any death	Chronic obstructive pulmonary disease and bronchiectasis
NVS011	2.208	9.100	0.249	8.880	6.66E-19	7.33E-18	5.590	14.816	D	Any death	Neurocognitive disorders