

1           **The scope and needs related to training, research, and mentorship among**  
2           **early-career addiction medicine professionals: an online global survey**  
3           **followed by expert group discussions**  
4

5 Roshan Bhad <sup>1,2</sup>, Sophia Achab <sup>2,3</sup>, Parnian Rafei <sup>2,4</sup>, Preethy Kathiresan <sup>5</sup>, Hossein  
6 Mohaddes Ardabili <sup>2,6</sup>, Jenna L. Butner <sup>2,7</sup>, Laura Orsolini <sup>2,8</sup>, Katrine Melby <sup>2,9</sup>, Mehdi  
7 Farokhnia <sup>2,10</sup>, Venkata Lakshmi Narasimha <sup>2,11</sup>, Kelly Ridley <sup>2,12</sup>, Serenella Tolomeo <sup>2,13</sup>,  
8 Mitika Kanabar <sup>2,14</sup>, Beatrice Matanje <sup>2,15</sup>, Paolo Grandinetti <sup>2,16,17</sup>, ISAM NEXt Consortium  
9 <sup>18\*</sup>, Marc N. Potenza <sup>7</sup>, Hamed Ekhtiari <sup>19,20</sup>, Alexander Baldacchino <sup>21</sup>

- 10  
11 1. Department of Psychiatry & National Drug Dependence Treatment Centre (NDDTC), All India  
12 Institute of Medical Sciences (AIIMS), New Delhi, India  
13 2. ISAM NEXt (New Professionals Exploration, Training & Education Committee, International  
14 Society of Addiction Medicine (ISAM)  
15 3. Psychological and Sociological Research Unit, Faculty of Medicine, University of Geneva, Geneva,  
16 Switzerland  
17 4. Department of Psychology, Faculty of Psychology and Education, University of Tehran, Tehran,  
18 Iran  
19 5. Department of Psychiatry, All India Institute of Medical Sciences (AIIMS), Jodhpur, Rajasthan, India  
20 6. Psychiatry and Behavioural Sciences Research Center, Mashhad University of Medical Sciences,  
21 Mashhad, Iran  
22 7. Yale School of Medicine, New Haven, CT, USA  
23 8. Unit of Clinical Psychiatry, Department of Neurosciences/DIMSC, School of Medicine and Surgery,  
24 Polytechnic University of Marche, Ancona, Italy  
25 9. Department of Clinical Pharmacology, St. Olav University Hospital, Trondheim, Norway  
26 10. National Institute on Drug Abuse and National Institute on Alcohol Abuse and Alcoholism,  
27 National Institutes of Health (NIH), Baltimore, Maryland, USA  
28 11. Department of Psychiatry, All India Institute of Medical Sciences (AIIMS), Deoghar, Jharkhand,  
29 India  
30 12. The Rural Clinical School of Western Australia, University of Western Australia, Albany, Australia  
31 13. Singapore Agency for Science, Technology and Research & National University of Singapore,  
32 Singapore  
33 14. Southern California Permanente Medical Group, California, USA  
34 15. Partners In Health/Abwenzi Pa Za Umoyo, Neno, Malawi  
35 16. SerD Teramo, Department of Territorial Assistance, ASL Teramo, Teramo, Italy  
36 17. Network of Early Career Professionals working in the area of Addiction Medicine (NECPAM)  
37 18. Consortia included 26 members across 22 countries \*  
38 19. Laureate Institute for Brain Research, Tulsa, OK, USA  
39 20. Department of Psychiatry, University of Minnesota, MN, USA  
40 21. Division of Population and Behavioural Science, Medical School, University of St Andrews, St  
41 Andrews, UK

42  
43 *(Members includes 1.Ankita Chattopadhyay; 2.Bezzina Gianluca; 3.Chia-Chun Hung; 4.Danielle Jackson; 5.Edem*  
44 *Sallah; 6.Enjeline Hanafi; 7.Erna Gunnthorsdottir; 8.Francina Fonseca; 9.Gayatri Bhatia; 10.Georgios Tzeferakos;*  
45 *11.Hillary Selassi Nutakor; 12.Hussein Elkholy; 13.Irfan Ullah; 14.Jiang Long; 15.Jibril Handuleh; 16.Leonardo*  
46 *E.Allagoa; 17.Mandana Sadeghi; 18.Mehrnoush Vahidi; 19.Mohammed Aljenibi; 20.Mohammadreza Shalbafan;*  
47 *21.Paxton Bach; 22.Sung Young Huh; 23.Surajuddin Abdulkadir; 24.Takeo Toyoshima; 25.Vicky Phan;*  
48 *26.Wafaa Elsayy)*

49  
50           **NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.**

51 **Abstract:**

52 Addiction medicine is a developing field, with many young professionals opting for a career in  
53 this area. However, globally, early-career professionals often face challenges in this field,  
54 such as lack of competency-based training due to a shortage of trainers, low availability of  
55 institutions with appropriate infrastructure, and limited resources for adequate training,  
56 particularly in developing countries. On the other hand, in developed countries, early career  
57 professionals may struggle with mentorship, limited job opportunities, and challenges with  
58 establishing a suitable research area.

59 The International Society of Addiction Medicine (ISAM) New Professionals Exploration,  
60 Training & Education (NExT) committee, a global platform for early-career addiction medicine  
61 professionals (ECAMPs), conducted an online survey using a modified Delphi-based  
62 approach among ECAMPs across 56 countries to assess and understand the need and  
63 scope for standardized training, research opportunities, and mentorship. The survey was  
64 conducted in 2 phases. A total of 110 respondents participated in Phase I (online key  
65 informant survey), and 28 respondents participated in Phase II (online expert group  
66 discussions on the three themes identified in Phase I). Most participants agreed with the  
67 lack of standardized training, structured mentorship programmes, research funding, and  
68 research opportunities in addiction medicine for ECAMPs. There is a need for standardized  
69 training programmes, improving research opportunities, and effective mentorship  
70 programmes to promote the next generation of addiction medicine professionals and further  
71 development to the entire field. The efforts of ISAM-NExT are well-received and give a  
72 template of how this gap can be addressed.

73

74 **Keywords:** Addiction medicine; Addiction psychiatry; Early career professionals;  
75 Mentorship; Research; Substance-Related Disorders; Training

76

77

78

79 **1. Introduction**

80

81 Addiction medicine is a relatively new field of medicine, with a growing number of early  
82 career professionals (ECPs) opting for a career in this area. However, there are several  
83 challenges in terms of lack of well-structured training shortage of institutes with infrastructure  
84 for adequate training and trainers, i.e., formally trained mental health and medical  
85 professionals with experience in addiction medicine (1, 2). These limitations prevent ECPs in  
86 several countries from pursuing a career in addiction medicine. There are limited resources  
87 and training opportunities for ECPs in upper-middle (UMICs) and lower-middle-income  
88 countries (LMICs). In high-income countries (HICs), where there is no or less dearth of  
89 experts and infrastructure, the challenges include receiving appropriate mentorship and  
90 choosing a suitable research area (3–5). Therefore, there is a need for a global platform  
91 helping early-career addiction medicine professionals (ECAMPs), including trainees, and  
92 connecting them with each other and with trainers and mentors worldwide. There is also a  
93 need to facilitate the launch and implementation of standardized training programmes,  
94 creating research and education opportunities, as well as fellowships and mentorship  
95 programmes in each subspecialty of the addiction medicine field (6). These needs become  
96 extremely necessary mainly due to the significant variability in the standards and quality of  
97 training programmes in the field of addiction and/or psychiatry globally, which is a major  
98 challenge for many ECAMPs in many countries (7).

99 Similarly, the assessment of training in addiction medicine and/or psychiatry is limited. For  
100 example, the International Certification in Addiction Medicine by the International Society of  
101 Addiction Medicine (ISAM) is one of the few examples of a well-established association able  
102 to provide global standards in validating and certifying knowledge in addiction medicine for  
103 professionals. However, currently, the examination by ISAM includes assessment for theory  
104 (based on multiple-choice questions (MCQs), without any practical exam or real case  
105 vignettes (8, 9). Hence, it is essential first to identify and clearly understand the needs and  
106 the demand for a standardized assessment of training in addiction medicine and/or

107 psychiatry to develop and implement a universal curriculum in addiction training  
108 programmes.

109 Building research collaboration across the globe and developing a practice-based research  
110 network is of high importance, given the eclectic nature of the field of addiction medicine and  
111 its significance is emphasized by policymakers and various global organizations, as well.  
112 Unfortunately, there are hardly any international organizations and networks in addiction  
113 medicine that address the need for researchers on a global platform for research  
114 collaboration. The National Institute on Drug Abuse (NIDA) is an organization that provides  
115 research support for early and mid-career addiction professionals; however, support is often  
116 limited to United States-based researchers and institutes (10). Assessing the need and  
117 scope for research opportunities exclusively for ECAMPs will inform policymakers regarding  
118 various issues and challenges. Quality mentoring and strategic planning, along with a  
119 favorable environment, are some of the elements that should be combined to create a  
120 successful career in research (11). Moreover, there is a need to assess deficiencies in  
121 training, research interest, and need for mentoring among early-career addiction  
122 professionals and address important issues that may help them in career development to  
123 mid-career. This may motivate and encourage ECAMPs to take up addiction medicine as an  
124 informed career choice since they can see the career trajectory and growth prospects ahead.  
125 The ISAM NExT (New Professionals Exploration, Training & Education) committee was  
126 established in 2020 with the primary objective of increasing and improving the capacity of  
127 addiction medicine training and other educational activities among ECAMPs. The committee  
128 constitutes 30 early career addiction professional members, including a chair and two co-  
129 chairs from 22 countries. We conducted a two-phase global cross-sectional online survey  
130 among ECAMPs to understand the need and scope for standardized training, research  
131 opportunities, and mentorship in the field of addiction medicine.

132  
133  
134  
135

136 **2. Methods**

137

138 **2.1. Study design**

139

140 A two-phase global online survey was conducted using a mixed-method, modified Delphi-  
141 based approach (12, 13). The first phase of the survey was carried out using an online  
142 survey in which 270 ECAMPs were approached. The first round of the survey took place  
143 from October 2020 to March 2021, and the results were finalized in April 2021. The second  
144 phase of the study comprised three focus group discussions to obtain consensus on key  
145 themes elicited in the first phase.

146 An online Google survey tool was prepared by the research team for phase I (available as  
147 supplementary material). Eligible participants (ECAMPs) were identified (sample of  
148 convenience) across different regions of the world using membership directory of  
149 professional societies in the field and social media/research networks, i.e., ResearchGate  
150 and LinkedIn enrolled. ECAMPs (n=270) were then invited to participate in the study via  
151 email. Total of 125 respondents provided written informed consent for participation in study  
152 on email in the first phase. Written informed consent were also obtained from participants  
153 prior to enrollment in the second phase of the study. to Subsequently, the data were  
154 analyzed, and the recommendations were compiled based on feedback from a core group of  
155 13 collaborators of the ISAM NExT expert committee for the research project.

156 **2.2. Inclusion and exclusion criteria**

157 For the purpose of this study, the operational definition of “Early Career Addiction Medicine  
158 Professionals (ECAMPs)” has been used (Table 1) (14) who were clinicians, scholars,  
159 resident doctors, and professionals working in or with an interest in the field of addiction  
160 medicine within 10 years of obtaining MD/MSc/Equivalent degree OR within 5 years of  
161 obtaining PhD degree depending on national context) and were aged between 25-45 years.  
162 All participants who gave informed consent to participate in online surveys and expert group  
163 discussions were included in the study. A sample size of a minimum of 100 respondents for  
164 phase I (online Google Form<sup>LM</sup> survey) across at least ten countries in the world and a

165 minimum of twenty respondents from phase I for phase II (online expert group discussion)  
166 were decided based on feasibility, time constraints, and their COVID-19 pandemic  
167 circumstances.

### 168 **2.3. Recruitment strategy**

169 For the first phase of the study, potential participants were identified using the membership  
170 directory of organizations working in the field of addiction medicine {e.g., International  
171 Society of Addiction Medicine (ISAM), International Society of Substance Use Prevention  
172 and Treatment Professionals (ISSUP), World Psychiatric Association (WPA), Indian  
173 Psychiatry Society (IPS), and social media (LinkedIn) and research network (ResearchGate).  
174 We ensured that our sample included at least ten participants from each World Health  
175 Organization (WHO) region (African Region, Region of the Americas, South-East Asia  
176 Region, European Region, Eastern Mediterranean Region, and Western Pacific Region) in  
177 order to increase diversity and global representation. For the second phase of the study, all  
178 participants who took part in the first phase were randomized per WHO regions and were  
179 invited to participate in the in-depth interviews (within online expert group discussions) on the  
180 themes that emerged on training, research, and mentorship, using stratified random  
181 sampling strategy. A random sample of participants was engaged in 3 sessions of  
182 discussion, each comprising 8-11 respondents, for a duration of 2 hours, in April 2021. The  
183 participants were contacted in advance through email with enclosed information about the  
184 questionnaire, an expert group discussion guide, the rules of engagement in the discussion,  
185 the participant information sheet, and a consent form. Upon receiving written consent, a link  
186 for an online meeting was shared.

187 The online expert group discussions (Training, Research, and Mentorship) were facilitated by  
188 collaborator members from the ISAM NEXt. The moderator (an ISAM NEXt Member)  
189 guided the participants with questions and facilitated the discussion. The meetings were  
190 video recorded (with permission from the participants) and later transcribed for thematic  
191 analysis.

192

193

#### 194 **2.4. Ethics approval and consent:**

195 The online survey was conducted according to the principles of good scientific practice (15).  
196 Ethical approval for the study was sought and granted by the Institutional Ethics Committee  
197 (IEC) at the All India Institute of Medical Sciences (AIIMS), New Delhi, India (September 16,  
198 2020, reference number IEC-888/04.09.2020). Participants provided a written online  
199 informed consent to participate in this study before filling out the self-administered survey,  
200 voluntary and anonymously.

201

#### 202 **2.5. Analysis:**

203 Data of phase I was analyzed using the Software Package for Social Sciences for Windows  
204 v. 24.0 (SPSS 24) (IBM Corp, Armonk, NY). Categorical variables were summarized as n  
205 (%), and continuous variables as means [standard deviation (SD)]. Focus group discussions  
206 data were analyzed using thematic analysis of the video recorded expert group discussions  
207 transcribed to the word document by the independent researchers (PK, KR). A core group of  
208 13 collaborators from the ISAM NExT committee reviewed the data before publication.

209

### 210 **3. Results**

211

#### 212 **3.1. Phase I: Online Survey**

213 Out of the 270 potential respondents approached, a total of 125 responses were received  
214 from across 56 countries and 6 WHO regions during six months of the data collection period  
215 (response rate 46.3%). Figure 1 shows the geographical distribution of the survey  
216 respondents (Phase 1) and the WHO regions. Fifteen responses were excluded as 10  
217 respondents did not fulfill eligibility criteria and 5 responses were duplicates; therefore, data  
218 were analyzed for a total of 110 respondents for the phase I survey (Table 1). The mean age  
219 of respondents was 35.66 (SD=4.97) years. About half of the respondents were from the  
220 Southeast Asia and Africa region. Respondents had been working in the field of addiction  
221 medicine or psychiatry profession for an average of 5.78 years (ranging from 1-17 years).

222 Out of these respondents, 56% were psychiatrists, with some exclusively practicing addiction  
 223 psychiatry. Around 27% of respondents were addiction medicine physicians; psychologists  
 224 (4.5%), nursing professionals (3.6%), social workers (3.6%), and post-doctoral fellows (4.5%)  
 225 constituted the remaining professionals. Table 2 overviews the responses to the survey  
 226 questions concerning key themes related to addiction medicine training, mentorship, and  
 227 research shortages and needs.

228  
 229 **Table 1: Phase I Survey - Socio-demographic and addiction medicine training-related**  
 230 **information (n=110)**  
 231

<b>Variables</b>	<b>Mean (SD)/Frequency (%)</b>
<b>Age (in years)</b>	35.66 (4.97)
<b>Gender</b> <ul style="list-style-type: none"> <li>● Male</li> <li>● Female</li> </ul>	68 (61.8 %) 42 (38.2 %)
<b>World-Health Organization (WHO) Regions</b> <ul style="list-style-type: none"> <li>● Africa</li> <li>● Americas</li> <li>● Eastern Mediterranean</li> <li>● European</li> <li>● Southeast Asia</li> <li>● Western Pacific</li> </ul>	27 (24.5%) 14 (12.8%) 12 (10.9%) 16 (14.6%) 28 (25.4%) 13(11.8%)
<b>Type of professionals</b> <ul style="list-style-type: none"> <li>● General/Addiction Psychiatrist</li> <li>● General/Addiction Physician</li> <li>● Clinical Psychologist</li> <li>● General/Psychiatrist Nursing</li> <li>● Psychiatric Social Worker</li> <li>● Doctoral (PhD) student/Post-Doctoral Fellow</li> </ul>	62 (56.5%) 30 (27.3%) 5 (4.5%) 4 (3.6%) 4 (3.6%) 5 (4.5%)
Years of clinical/research experience in Addiction Medicine	5.78 (9.35)



Deficiency in existing addiction training programme/curriculum for ECAMPs	Yes – 95 (86.4%) No – 15 (13.6%)
---	-------------------------------------

232  
233

**Table 2: Phase I Survey – key themes for addiction medicine training needs**

Key problems in lack of standardized training programmes	<ol style="list-style-type: none"> <li>1. Exposure to specialty clinics – 23 (24.2%)</li> <li>2. Lack of trained faculty in subject – 18 (18.9%)</li> <li>3. Less rotation/duration of posting in addiction medicine during postgraduate training – 18 (18.9%)</li> <li>4. Lack of proper assessment of training – 14 (14.8%)</li> <li>5. Lack of standard textbook – 14 (14.8%)</li> <li>6. Less patient load – 8 (8.4%)</li> </ol>
Lack of research opportunities for ECAMPs	Yes - 98 (89.1%) No – 12 (10.9%)
Ways to improve research opportunities for ECAMPs	<ol style="list-style-type: none"> <li>1. Research project compulsory part of training – 28 (28.6%)</li> <li>2. Workshops on how to conduct research – 26 (26.5%)</li> <li>3. Workshops on how to get funding/grant for research – 20 (20.4%)</li> <li>4. Funding (Intramural/Extramural) – 16 (16.3%)</li> <li>5. Interdisciplinary team – 8 (8.2%)</li> </ol>
Key recommendations for a successful mentorship programme	<ol style="list-style-type: none"> <li>1. Trained addiction professionals as mentors – 35 (31.8%)</li> <li>2. Networking opportunities – 19 (17.3%)</li> <li>3. Mentorship programme as an integral part of training – 18 (16.4%)</li> <li>4. Institutional support – 17 (15.4%)</li> <li>5. No recommendation – 21 (19.1%)</li> </ol>

234

235 **Figure 1 — the geographical distribution of survey respondents (Phase I) and their affiliated WHO regions**  
236 **illustrated on a world map**

237 **3.2. Phase II: Online expert group discussions (n=28)**

238

239 All participants of the phase I study were invited to participate in group discussions on  
240 training, mentorship, or research needs. A total of 110 participants were randomized based  
241 on the WHO regions (stratified) into groups of 37, 37, and 36. Subsequently, these groups  
242 were assigned one of the three themes (Training, Mentorship, and Research), and an email  
243 invitation along with questions on that theme was sent 10-14 days before the online expert  
244 group discussions. A total of 28 respondents participated in phase II (overall response rate

245 25.4 %). The response rate was 8/36 (22.2%) for the training theme, 8/37 (21.6%) for  
 246 mentorship theme, and 12/37 (32.4%) for the research theme. The geographical distribution  
 247 and the number of representatives in expert group discussions are depicted in figure 2.

248 The thematic analysis results of the expert group discussions are as shown in Tables 3, 4,  
 249 and 5.

250 **Figure 2 — the number and geographical distribution of representatives who contributed to expert group**  
 251 **discussions illustrated on a world map**

252 **Table 3A: Expert group discussion on training (n=8)**  
 253

Part A:	Responses to the questions for advance distribution (frequency)
Countries (Participants)	Italy (2), India (2), Ghana (2), Indonesia (1), Iran (1)
Questions	Responses (n)
1. Current phase of training in the field of Addiction Medicine/Psychiatry?	<ul style="list-style-type: none"> <li>● Post-graduate/super-specialty Trainee Resident (3/8)</li> <li>● Junior faculty member/Consultant (3/8)</li> <li>● Nursing professional (2/8)</li> </ul>
2. Availability of standard training programme for Addiction Medicine/Psychiatry in medical school/institute/university?	<ul style="list-style-type: none"> <li>● Yes (6/8)</li> <li>● No (2/8)</li> </ul>
3. Duration of training in Addiction Medicine/Psychiatry in respondent's country	<ul style="list-style-type: none"> <li>● Two years or less (1/6)</li> <li>● Three years or less (4/6)</li> <li>● More than three years (1/6)</li> </ul>
4. How satisfied respondent is with the training programme in addiction medicine/psychiatry	<ul style="list-style-type: none"> <li>● Very unsatisfied (1/6)</li> <li>● Neutral (1/6)</li> <li>● Satisfied (3/6)</li> <li>● Very satisfied (1/6)</li> </ul>

5. Main strengths of the training programme	<ul style="list-style-type: none"> <li>● Exposure to the management of different substances/specialty clinics (4/6)</li> <li>● Availability of eminent addiction professionals as mentors (2/6)</li> </ul>
6. Main limitations of the training programme	<ul style="list-style-type: none"> <li>● Lack of addiction specialists as trainers (4/6)</li> <li>● Lack of specialty clinics and exposure to multiple substance use problems (1/6)</li> <li>● Difficulties in obtaining needed materials for training (1/6)</li> </ul>
7. Number of patients consulted under supervision/year during training?	<ul style="list-style-type: none"> <li>● Less than 10 patients/year (1/6)</li> <li>● 20-50 patients/year (1/6)</li> <li>● More than 50 patients/year (4/6)</li> </ul>
8. Exposure/posting/rotation to Addiction Medicine/Psychiatry during medicine or psychiatry training?	<ul style="list-style-type: none"> <li>● Yes (8/8)</li> </ul>
9. Duration of posting/rotation in Addiction Medicine/Psychiatry	<ul style="list-style-type: none"> <li>● One month or less (2/8)</li> <li>● 1-3 Months (3/8)</li> <li>● 3-6 months (3/8)</li> </ul>
10. Exposure/clinical experience in the following subspecialties/special treatment facilities of Addiction Medicine/Psychiatry	<ul style="list-style-type: none"> <li>● Dual diagnosis clinic, Opioid agonist treatment, Behavioural addiction clinic (7/8)</li> <li>● Consultation liaison (6/8)</li> <li>● Tobacco cessation clinic, Adolescent clinic (5/8)</li> <li>● Community addiction treatment clinic, Injecting drug users clinic (4/8)</li> <li>● Pain clinic, New psychoactive substances (NPS) clinic (1/8)</li> </ul>

254  
255  
256  
257  
258

**Table 3B: Thematic analysis of expert group discussion (Training)**

<b>Part B:</b>	<b>Expert group discussion</b>
----------------	--------------------------------

<b>Question</b>	1. What do you think about issues faced by early-career addiction medicine professionals in terms of standardized training, deficiencies, suggestions for improving the same?	
<b>Themes</b>	<b>Examples/Consensus</b>	<b>Remarks</b>
Entry/eligibility to join addiction medicine training/course	<p><i>“Both physicians and psychiatrists can practice addiction medicine in Italy.”</i> General psychiatrists manage addiction problems in Iran. The eligibility for training is defined and limited to only psychiatrists in India. Only a few institutes offer addiction medicine training in Indonesia and India. <i>“The training is heterogeneous and varies across institutes in India.”</i> There is an MCQ-based entrance examination for entry into addiction training in India. Training is available for nursing professionals in addiction treatment.</p> <p>Consensus: There is a variation in entry into the addiction training programme across countries.</p>	No national training programmes in addiction medicine in Italy and Ghana
Duration of posting/rotation in addiction medicine training	<p>No exposure for undergraduate students. Postgraduate students are posted for 3–6 months. <i>“The training duration varies from 1–3 years and super-specialty courses available in India where trainees are posted for 21-14 months in core specialty and remaining months in Neurology, Gastroenterology, Child Psychiatry, etc.”</i> There is no specialized training in Italy, and psychiatry residents are posted for 3 months–2 years. There is a 3-month duration of posting for a psychiatrist in addiction medicine in Iran, which is not enough. <i>“There is no rotation in addiction medicine in Ghana, and there are no trained addiction medicine doctors.”</i> In Indonesia, there is a 3–4 weeks posting for undergraduate medical students and 1ne month posting in addiction psychiatry for psychiatry residents. <i>“Two years of experience in the addiction treatment facility is needed to practice as a specialist.”</i> Recently a subspecialty programme in addiction medicine started in one university in Indonesia.</p> <p>Consensus: There is a variation in the duration of training/posting in addiction medicine across countries.</p>	Addiction medicine training was recently established in Indonesia, and the curriculum is yet to develop.

Nature of training	<p>An adequate number of patients with substance use disorders across countries for training needs. Both Buprenorphine and Methadone are available as OAT in Italy, India, and Indonesia. In Iran, tincture opium is also available for treatment. <i>“There is also a separate 3-month course in opioid agonist treatment including Methadone for physicians in Iran”</i>. Treatment options are limited to tramadol in Ghana. Medications are available for the treatment of alcohol use disorders in all countries. In-patient treatment facilities are available in all countries.</p> <p>Consensus: Outpatient treatment facilities are key places for clinical exposure.</p>	ECAMPs are exposed to clinical training in outpatient treatment services in all participating countries
Exposure to specialty clinics/sub-specialties	<p><i>“In Iran, exposure to various substance use problems is diverse, and supervision by expert faculty is available.”</i> Specialty clinics are still in the budding stage across countries, and some countries have behavioral addiction clinics. In Italy, trainees are able to see dual diagnosis patients as a part of a training programme. No NPS clinic in India; other specialty clinics are available. No specialty clinics in Ghana. In Indonesia, exposure to behavioral addiction and tobacco cessation clinics are available as a part of training in limited institutes.</p> <p>Consensus: There is limited exposure to specialty clinics across countries.</p>	Trained faculty in addiction medicine are available in Iran
Assessment at the end of the training	<p>Six monthly assessments of training in Italy. End semester comprehensive exams, including theory and clinical case presentation, is conducted in India for super specialty trainees. No separate assessment in Iran and Indonesia for MD (Psychiatry) residents. <i>“10 % marks of OSCE are dedicated to addiction psychiatry in the MD (Psychiatry) final exam in India.”</i> In India, the assessment is predominantly summative with some elements of formative assessment. Clinical assessments are conducted for nursing professionals specialized in addiction treatment in Ghana.</p> <p>Consensus: Training assessments are conducted in all the countries. However, the method of assessment varies.</p>	End of the training standard assessment is conducted in India

259  
260  
261  
262  
263

**Table 4A: Expert group discussion on mentorship (n=8)**

<b>Part A:</b>	<b>Responses to the questions sent for advance distribution</b>
----------------	---

Countries (Participants)/ Questions	Canada (1), Iran (1), India (1), Nigeria (1) Singapore (1), Somalia (1), USA (2)
<b>Questions</b>	<b>Responses (n)</b>
1. How would you describe your current phase of career in the field of addiction medicine?	<ul style="list-style-type: none"> <li>● Post-graduate/super-specialty Trainee Resident (3/8)</li> <li>● Junior faculty member (2/8)</li> <li>● Doctoral/Fellowship Student (1/8)</li> <li>● Post-doctoral Student/Fellow (1/8)</li> <li>● Associate Physician (1/8)</li> </ul>
2. Do you have a formal mentorship programme in your medical school/institute/university?	<ul style="list-style-type: none"> <li>● Yes (3/8)</li> <li>● No (5/8)</li> </ul>
3. Do you consider your current supervisor/thesis guide/principal investigator as your mentor?	<ul style="list-style-type: none"> <li>● Yes (5/8)</li> <li>● No (2/8)</li> <li>● No, but I have a mentor from outside my medical school/institute/university (long-distance mentor) (1/8)</li> </ul>
4. If a mentorship programme was available, then how satisfied were you with the mentorship programme?	<ul style="list-style-type: none"> <li>● Unsatisfied (1/3)</li> <li>● Neutral (1/3)</li> <li>● Satisfied (1/3)</li> </ul>
5. Can you describe two reasons for effectiveness/non-effectiveness (if any) for the available mentorship programme?	<p>Effectiveness</p> <ul style="list-style-type: none"> <li>● Similar experiences, life goals, and professional aims to mine, an ability to relate (2/3)</li> <li>● Genuine concern for my wellbeing and success (1/3)</li> </ul> <p>Non-effectiveness</p> <ul style="list-style-type: none"> <li>● Limited time and busy schedule, and lack of protected time for mentoring and research (2/3)</li> <li>● Mentors possessed biased views against non-psychiatrist's ability to practice in the field of addiction (1/3)</li> </ul>

6. Ideally, how many mentors should a mentee have?	<ul style="list-style-type: none"> <li>● Single mentor (5/8)</li> <li>● Multiple mentors (2/8)</li> <li>● Multiple mentors with a single senior mentor in charge (1/8)</li> </ul>
7. According to your experience, which phase of a mentorship programme* is the most important to you? * Zachary's The Mentor's Guide, 2000 (15)	<ul style="list-style-type: none"> <li>● Initiation or Preparation (1/8)</li> <li>● Cultivation or Negotiation &amp; Enabling or Protégé (5/8)</li> <li>● Separation or Closing or Break up (1/8)</li> <li>● Redefinition or Lasting friendship (1/8)</li> </ul>
8. Do you have an office for delivering needed guidance on career development/training/post-doctoral while training at your medical school/institute/university?	<ul style="list-style-type: none"> <li>● Yes (1/8)</li> <li>● No (7/8)</li> </ul>
9. If the ideal mentorship programme is available, then what sort of guidance and assistance would you need from a mentor as a mentee (in order of preference)?	<ul style="list-style-type: none"> <li>● Long-term career planning (3/8)</li> <li>● Research design (1/8)</li> <li>● Networking nationally and internationally (1/8)</li> <li>● Balancing personal &amp; Professional demands (1/8)</li> <li>● Developing a research portfolio (1/8)</li> <li>● Addressing burnout during training (1/8)</li> </ul>
10. By the end of an effective mentorship programme, what kind of abilities should a mentee obtain?	<ul style="list-style-type: none"> <li>● Ability to conduct research ethically independently and responsibly in the given area (4/8)</li> <li>● Ability to achieve career progression financial independence through satisfactory job opportunities in the given area (3/8)</li> <li>● Ability to become a mentor and run an effective mentorship programme in the given area (1/8)</li> </ul>

264  
265  
266  
267

**Table 4B: Thematic analysis of expert group discussion (Mentorship)**

<b>Part B:</b>	<b>Expert group discussion</b>	
<b>Questions</b>	1. In Your university, in your city, in your academic field, do you have a mentorship programme in the field of addiction medicine or general psychiatry? If so, how satisfying is the programme?	
<b>Themes</b>	<b>Examples/Consensus</b>	<b>Remarks</b>

<p>Availability of mentorship programme</p>	<p><i>“There is no formal mentorship programme in some institutes. But anyone can identify someone with similar research interests as a mentor. There is a mentorship programme for Buprenorphine training to get a special waiver to prescribe Buprenorphine in USA (Called as PCSS-O), but it is not associated with any university, and it is not that satisfying.”</i></p> <p>Consensus: The mentorship programme is not available widely in developed countries and non-existent in developing countries</p>	<p>Mentorship programmes for early-career addiction medicine professionals are not available in most countries</p>
<p>Nature of mentorship programme</p>	<p><i>“In Canada, there is mentorship programme only for current fellows in addiction medicine, for the duration they are in the programme to provide them primarily with career advice.”</i></p> <p>Consensus: There are no formal mentorship programmes for addiction medicine in most institutes, and among the available programmes, only a few are satisfactory.</p>	
	<p>2. Can you describe reasons for effectiveness and non-effectiveness for mentorship programmes?</p>	
<p>Effectiveness/non-effectiveness of the mentorship programme</p>	<p><i>“Mismatch about expectations is a very critical thing. There should be flexibility in the programme for people or prospective mentees to be specific in what they are looking for and choose mentors based on their expectations. The relationship between mentor and mentee is another important reason for failure or success of a programme.”</i></p> <p>Consensus: The programme should be interdisciplinary, and mentees should be able to choose their mentor based on their needs or interests; The mentor should have adequate emotional intelligence, should be available and accessible, and be able to relate to the mentees, and there should be SMART goals. The programme should not be short-term but should be continuous or long-term.</p>	
	<p>3. What are your suggestions for successful mentorship programmes, especially for early-career addiction medicine professionals?</p>	



Appropriate time and duration for the mentorship programme	<p><i>“Exposing medical students early in their career, preferably at the undergraduate level even before they decide to go into psychiatry or surgery specialties, etc. Identifying those who are passionate about addiction. For graduates or those in private practice, etc., ISAM or MSR can create networks. There is a need for continuity instead of being short-lived. Information about mentorship programmes should be given to trainees during the last year of training.”</i></p>	
Career-based mentorship programme	<p><i>“We need to have early career mentoring programmes. Successful mentoring programmes should have time-bound SMART goals. Also, some kind of career-based mentoring on how to maintain a balance between personal and professional aspects of life. We should have some wellness programmes on how to balance lifestyle and work in addiction.”</i></p> <p>Consensus: For a successful mentorship programme, it should be introduced at the medicine level (undergraduate level), residency as well as early-career level. There should be flexibility keeping mentees in the center of everything, and mentees should be able to choose their mentors. There should be SMART goals. The programme should not be short-lived but should be continuous and comprehensive. ISAM can help in international collaboration for mentorship. Also, there should be a career-based mentoring programme to discuss work-life balance issues.</p>	Continuous mentorship programme as per the stage of career is needed for its effectiveness in shaping the career of ECAMPs

268  
269  
270

**Table 5: Expert group discussion on research needs (n=12)**

<b>Thematic analysis of expert group discussion (Research Needs)</b>	
<b>Countries/participants</b>	Australia (2), China (1), Iceland (1), India (2)*, Iran (1), Nigeria (1), Norway (1), Pakistan (1)*, South Korea (1), Switzerland (1), USA (2) <i>*two participants left the group discussion halfway, eleven participants attended the full discussions</i>
<b>Questions</b>	<b>1. What is the scope of research in Addiction Medicine in your country?</b>

Themes	Examples/Consensus	Remarks
Research themes covered worldwide	<p><i>“In Korea, many are interested in research in gaming disorders and other behavioral addictions. The government encourages the researchers to do various pieces of research in the field of behavioral addictions and alcohol use disorder.”</i></p> <ul style="list-style-type: none"> <li>•China and Korea are more focused on behavioral addictions.</li> <li>•Australia: psychedelics, methamphetamine, amphetamine-type stimulants, newer psychoactive substances, as well as tobacco and alcohol.</li> <li>•USA: all types of illicit substances</li> <li>•Iceland, all types of research</li> <li>•In Nigeria, there is a need for research in addiction.</li> </ul>	<p>Heterogeneity in the scope of research in addiction medicine among countries:</p> <p>Research topics orientation based on a specific drug or behavior</p>
Determinants of research themes conducted	<ul style="list-style-type: none"> <li>· Cultural influences (politics, religion) on research areas investigated</li> </ul> <p>USA: “research on all types of drugs are being conducted and all varieties of work including controversial researches with political and religious influences.”</p> <ul style="list-style-type: none"> <li>· Economic influence (funding opportunities) on the scope of research themes investigated.</li> </ul> <p>Iceland: “ there is a small population, but accessible and adequate funding opportunities for research.”</p>	
2. Are there enough research opportunities, either funded or non-funded, for early-career addiction medicine professionals in your institution?		
Impact of funding on research opportunities for early careers in addiction medicine	<p>Inadequate funding for addiction research compared to other psychiatric disorders and non-communicable diseases. In Australia, there are enough research opportunities. In China, there is difficulty in getting funding for research, although the government is advocating for conducting research on behavioral addictions as it is a public health problem. In the USA, Iceland, and Norway, plenty of funding opportunities are available. “There is no sufficient funding for research in Nigeria, Pakistan.”</p>	<p>Some jurisdictions, totally (e.g., African countries) or partially (e.g., China) resulted in a lack of opportunities for research in addiction medicine</p>

Impact of mentorship on research opportunities for early careers in addiction medicine	Nigeria: <i>“There are no sufficient mentorship opportunities for research.”</i> Some mentorship opportunities are available in most other countries.	
Impact of clinical workload on research opportunities for early careers in addiction medicine	USA: <i>“It is difficult to manage time between research and clinical work.”</i>	
3. How independently the research (meaning not funded by industries with competing interests like alcohol, tobacco, gambling, and gaming industry) could be conducted in your country?		
Availability of infrastructure for independent research	<i>“I have never heard about competing for interest institution ever participating or having influence in addiction research in Iceland. We have access to independent research.”</i> Consensus: Conducting research projects independently is possible in most countries.	
4. How easy is it for you to consolidate research, clinical, and training activities?		
Multitasking	Consensus: In Australia, it depends on papers and position (academic/clinical/combined). It is easier to combine research activities with training activities. However, it is harder to make systematic reviews or other things like that during business hours. It is easier to do teamwork. It is difficult to combine clinical work with research work in the USA and China, given the time, energy burnout, etc.	
5. How do you manage time, in between clinical and training work for research work, in terms of applying for funding, writing protocol, research projects, etc.?		

<p>Dedicated time for research</p>	<p><i>“It is difficult to manage clinical and training work for research work. I do not have the expertise or skills for applying for a larger-scale grant - mentorship and training are needed for this purpose. Also, I have to spend more time outside my clinical and teaching hours. It is a struggle in terms of balance.” A respondent from the USA</i></p> <p>Consensus: In the USA, it is a struggle to manage timings for clinical and training work for research. In Australia and China, Grant writing is the biggest time suck – We put in a lot of effort outside usual working hours, but in the end, you may not get funding.</p> <p>It is easier to be part of a research group in the USA and Nigeria than doing independent research.</p>	
<p>6. Do you have a form of a research programme in your medical school/institute/university for a research programme in addiction medicine? And if there is, are you satisfied with the programme? (Very unsatisfied to very satisfied)</p>		
<p>Availability of research programme</p>	<p>Consensus: In Australia, there is a research programme as part of specialty training in addiction and PhD opportunities in addiction. Research activity satisfaction depends on your personal interest in research or the project.</p> <p>In the USA, the satisfaction of research programmes depends on the location of the practice. Some people say that it is pretty satisfying.</p> <p>In China and Korea, there are satisfying research programmes.</p> <p>In Nigeria, there is no research programme.</p> <p>In Iceland, there are no addiction psychiatry research programmes.</p>	
<p>7. Can you describe any two reasons for the effectiveness or non-effectiveness of the available research programmes you just described in your country?</p>		

*“The primary reason for effectiveness as well as non- effectiveness of Australian research programme is the fact that it is mandatory. It is effective because you have to do it. It is ineffective as it is forcing people into research, when they may be not interested at that time of training, producing opposite effects.”*

Consensus: In Australia, research is mandatory in research programmes. This is a reason for effectiveness as people do it, as well as non-effectiveness, forcing people who may not be interested in doing research at that point in life.

In the USA, the reason for the effectiveness of research usually involves specific people and when specific research is done. The reason for non-effectiveness is that it is difficult for people to enter into research from clinical work.

In China and Korea, the programmes are free-styled and depend on the style of mentors, which is the reason for non-effectiveness. There is a need for standardization or regularization for the programme and the mentors.

In Iceland and Nigeria, there is no programme, and there is a need for the same

8. Would you be interested in workshops on scientific work? And if any, what would you be interested in (workshop on finding donors, scientific writing, or any other topics)?

<p>Modes of improving research programme</p>	<ul style="list-style-type: none"> <li>• In Australia, there is a need for workshops on the structure of research papers, writing papers in an efficient way, and grant writing.</li> <li>• In the USA, workshops are not necessary, as most programmes have research-related guidance built-in. There is a need for pairing with a mentor and support groups for paper rejection.</li> <li>• Australia also reported that some didactic teaching could be repetitive as universities often have online guides or webinars on academic writing, literature review, etc.</li> <li>• In China, the participant wanted training on the whole process of publication – how to get a paper published how to cope with every step, and how to cope with rejection.</li> <li>• In Korea, there was a need for a workshop on the structure of research paper</li> <li>• In Nigeria, there was a need for finding suitable Journals.</li> </ul>	<p>There is demand for workshops on scientific writing, obtaining grants among ECAMPs</p>
<p>9. Is there an addiction research group community or network where you work which would be interested in participating in digital meetings with other PhD students, postgraduates, etc., in other countries?</p>		
<p>Professionals groups engaged in research</p>	<ul style="list-style-type: none"> <li>• In Australia, the USA, China, and Korea, addiction research groups or communities exist. All expressed their interest in participating in digital meetings.</li> <li>• In Iceland, there is an addiction research group and community. The representative stated that she was interested in participating only when she got into a residency programme or similar qualification, as she felt it would be pointless without active interaction.</li> </ul>	
<p>10. What are your suggestions for the improvement of research activities in the field of addiction medicine?</p>		

<p>Ways to improve research opportunities</p>	<p><i>“It would be wonderful if there was a mechanism that can provide some guidance or huge pass for research. One association of international organization can provide the junior researchers with a road map for research -what we can do, what we can research on, what we can start - More than mentorship - We can gather a lot of eminence or important researchers, cutting edge directions for research in the future for next 5 years or 10 years” (Respondent from China)</i></p> <p>Consensus: In Australia, all the options plus mentorship plus listing of research skills desired in the project, so that jobless or homeless people can choose the project based on their skills.</p> <p>In the USA, having a website or page or resource listing opportunities can help to collaborate and/or work regionally as well as internationally.</p> <p>In China, a mechanism to provide guidance or huge pass for research and organization of training provides more resources and more platforms for junior researchers who do not speak English.</p> <p>In Korea, all the suggestions are listed as fellowship opportunities.</p> <p>In Iceland, International collaboration.</p> <p>In Nigeria, there is a need for workshops for writing and applying for grants and international collaboration.</p>	<p>There is a need for more funding for ECAMPs and a roadmap for developing/improving their research skills.</p>
---	--	--

271

272

273 **4. Discussion**

274

275 This study was one of the most extensive surveys conducted among early-career addiction  
 276 medicine professionals assessing the need and scope for standardized training, mentorship  
 277 programmes, and research opportunities — the online survey methodology allowed for a  
 278 broad representation of participants from 56 countries.

279 There is wide variability in entry requirements for addiction medicine training globally. In  
 280 countries such as the USA and the UK, both family physicians and psychiatrists can practice  
 281 as addiction specialists. In others, such as in Italy, both physicians with a specialty in  
 282 internal medicine or other medical subspecialties together with psychiatrists and

283 pharmacologists can treat patients with addiction. In most other countries, only psychiatrists  
284 can train in addiction medicine as a specialty. Developing countries allow nursing  
285 practitioners and social workers to pursue addiction medicine training due to a shortage of  
286 specialty physicians.

287 Training and exposure to addiction medicine also differ considerably across different  
288 countries. With respect to undergraduate training, exposure to addiction medicine as a  
289 specialty is minimal (about one week in the USA, 7 hours in the UK) or absent in most  
290 countries (6, 16–20). The biopsychosocial model of addiction is taught as a part of the  
291 theory in undergraduate medical school (21). Addiction medicine is an integral part of  
292 psychiatry and family medicine residencies in some countries like the USA, India, and Iran  
293 with respect to postgraduate training. Indonesia and India offer additional certificate courses  
294 after postgraduate training. There is considerable variation in curriculum and duration of the  
295 training placement of psychiatry residents in addiction psychiatry across countries.

296 In the USA & Australia, addiction psychiatry is a separate specialty as a postgraduate  
297 programme. In India and Indonesia, addiction psychiatry specialty is in the early developing  
298 stage (18,20,22). Lack of a structured curriculum is an important issue highlighted by the  
299 present survey participants. Content of the addiction medicine curriculum varies due to the  
300 nature of local substance use, availability of specialty clinics, opioid agonist treatments and  
301 other pharmacotherapy options available in different countries, and availability of trained  
302 faculty members for teaching. A structured curriculum can improve the knowledge of  
303 addiction medicine among internal medicine residents and hence need to be developed  
304 updated in different countries in order to improve the delivery of quality addiction treatment  
305 services (23). The quality of training is an issue for both internal medicine residents in the  
306 USA and also among psychiatry trainees across European countries as per a recent survey,  
307 which makes the call for a structured curriculum ever more important and urgent (24,25).

308 A survey in China revealed that doctors involved in drug treatment are not well prepared or  
309 experienced and have negative attitudes toward substance use disorders and afflicted  
310 patients (26). The low number and level of professional addiction experts are the potential



311 outcomes of inadequate addiction medicine training for medical students and residents in the  
312 USA, which is highlighted previously (9). From a trainee point of view, there is a demand for  
313 standardized training as emphasized in past reviews and found in the present study (27,28).  
314 The evaluation of standardized, structured short term training is also found to be an effective  
315 tool for addiction medicine training (29, 30) . The next generation of addiction treatment  
316 providers needs to be trained adequately to deal with emerging substance use problems  
317 across the globe.

318 There is a standard exit exam after completion of addiction medicine training in some  
319 countries like the USA, Australia, and India. There are regular mid-term assessments (6  
320 monthly or yearly) that are also conducted in countries like Italy, India, Iran, and Ghana for  
321 the trainees. In this regard, efforts by the ISAM to successfully conduct International  
322 Certification in Addiction Medicine exams for global trainees for the past 10 years need to be  
323 acknowledged (8).

324 The availability of mentorship programmes and needs were assessed in the present study.  
325 We found there are limited mentorship programmes available for early-career addiction  
326 medicine professionals. Such programmes are limited to developed countries like the USA &  
327 Australia. The mentorship programme is non-existent in most African and Asian countries  
328 like Ghana, Nigeria, China, India, Indonesia etc. Most participants recognized their training  
329 programme supervisor, thesis guide as a mentor. A single mentor was desired by most,  
330 although some participants expressed the need for multiple mentors depending upon the  
331 need in particular areas of interest and the stage of their career. The barriers identified for  
332 quality mentorship programmes were lack of time, funding, and trained faculty members (31).  
333 Most participants favored a continuous mentorship programme in different stages of their  
334 careers and were not limited to only the training duration. Mentorship programmes are vital  
335 for the development of the career of ECAMPs, and there is a need to facilitate mentorship  
336 programmes across nations. There are limited research studies on understanding the  
337 challenges faced with mentorship problems for ECAMPs. Among the available programmes,  
338 The Learning for Early Careers in Addiction & Diversity (LEAD) Programme, funded by the

339 National Institute of Drug Abuse, uses a team mentoring approach. Each LEAD Programme  
340 scholar works with a Clinical Trial Network (CTN) primary mentor while also receiving  
341 guidance from a UCSF University of California San Francisco (UCCSF) mentor and a  
342 nationally regarded diversity advisor (32). Other similar programmes are run by addiction  
343 medicine societies like the American Academy of Addiction Psychiatry (AAAP) & ISAM  
344 (International Society of Addiction Medicine) (33). There is a tremendous need to develop  
345 the culture of mentorship for strengthening academic medical centers engaged in addiction  
346 medicine training. Innovative methods like co-training with general physicians can facilitate  
347 mentorship programmes in such centres. The mentoring need is now even greater with the  
348 expansion of addiction medicine as a specialty and many young professionals joining their  
349 respective training programmes (11, 34, 35).

350

351 Most of the study participants reported there are limited research opportunities for ECAMPs.  
352 There are many challenges like clinical workload, funding, few suitable research mentors,  
353 obtaining research grants, and publishing the research. The challenges are existent even in  
354 developed countries like the USA & Australia. The research capacity has to be more  
355 developed during the training programme and is effective when mandatory for the completion  
356 of training. There is an unmet demand for grant writing, workshops for conducting research,  
357 and writing papers among ECAMPs. The research grants available from NIDA are mostly  
358 limited to USA Residents/Citizens (10). There are limited opportunities from addiction  
359 medicine societies. The United Nations Office on Drugs & Crime (UNODC), with support  
360 from the Drug Abuse Prevention Center (DAPC), started offering grants for early career  
361 researchers for projects related to prevention and promotion activities recently (36).

362 Combining clinical training and research would be a step ahead in improving addiction  
363 medicine training programmes and creating research opportunities for ECAMPs (37).

364 Developing research capacity among ECAMPs from low and lower-middle-income countries  
365 by conducting workshops with the support of facilitators from high-income countries can be a  
366 solution for the problem. Other allied addiction medicine professionals can also be engaged

367 in such training programmes to develop workforce and build more capacity (35, 38, 39). The  
368 main challenges encountered for conducting research by ECAMPs in the European survey  
369 (n=258) were lack of time as a large proportion of participants (87.2%) reported conducting  
370 research after regular working hours or partly during and after working hours. Only one-tenth  
371 ever received a grant for their work. Lack of funding is an important hurdle in conducting  
372 research in spite of ECAMPs being motivated to conduct the research(40). Global societies  
373 and institutes working in the field of addiction medicine need to provide adequate research  
374 opportunities as there is a risk of early-career addiction medicine professionals falling prey to  
375 predatory publishing and industry-sponsored research in the early stage of their career,  
376 which may bias their subsequent research projects (41–43).

377

378 The results from the present study suggest that there is variation in eligibility, the content of  
379 the curriculum, and assessments for addiction training across the globe. It is essential to  
380 develop a standard curriculum and training content that is competency-based, culturally  
381 sensitive, and can include local jurisdictional norms with substance use disorders. Flexibility  
382 is needed in the curriculum to account for the possibility of various medical professionals  
383 starting addiction medicine as a career. The study findings emphasized the need for  
384 mentorship programmes and more research opportunities for ECAMPs as a vital component  
385 of addiction medicine training.

386

387 A major strength of the present study is the perspective from more than 50 countries and  
388 covering all 6 WHO regions. We used a robust methodology with an online two-phase  
389 survey with systematic randomization for the second phase. The second phase, i.e., the  
390 qualitative part of the study using expert group discussions, adds perspective on attitudes  
391 and opinions of survey participants and hence adds more meaning and depth to the data  
392 collected using the online survey. Limitations of our study include self-reported data and a  
393 relatively small sample size. The study was conducted during the COVID-19 pandemic,  
394 where there was disruption of training programmes and shift to online teaching, which may

395 have influenced some of the findings in the study. The generalizability of the data is another  
396 limitation, as only participants who were members of professional societies and available on  
397 social media were approached. Future studies should address these limitations using  
398 randomized control trials for studying models of training, innovative techniques of training,  
399 and longitudinal study design to study mentorship needs in long-term career growth. The  
400 study findings emphasize the need for standardized training programmes, improving  
401 research opportunities and collaboration, and effective mentorship programmes for the next  
402 generation of addiction medicine professionals.

403

404

405 **Authors' Contributions:** RB & PK drafted the protocol. RB, MK, PR, and VLN recruited the  
406 phase I survey participants. PR sent the invitations and arranged online expert group  
407 discussion meetings. SA, MF, KM, PR developed questions and animated the discussions  
408 for expert group discussion research theme. RB, SA, MF, KM, PR, HMA, JLB, ST, MK, LO,  
409 VLN were collaborators for expert group discussions and facilitated the sessions. PK was  
410 note taker for expert group discussion sessions, transcribed and analyzed research &  
411 mentorship expert group discussions. KR validated the expert group discussions data. PR  
412 illustrated the figures and maps. ISAM NExT Consortia members participated in both phases  
413 of the survey. MNP, HE & AB reviewed the manuscript and mentored the group. All authors  
414 contributed to the revision and final edits of the manuscript.

415

416 **Acknowledgments:** We would like to thank Dr Shalini Arunogiri, Training Officer, ISAM; and  
417 Dr Cor De Jong Co-chair, Training Committee, ISAM; Dr Susana Galea Singer, Education  
418 Officer, ISAM; Dr Nady el-Guebaly, Chief Examiner, ISAM for their guidance and Ms Marilyn  
419 Dorozio, ISAM NExT office for technical support in conducting online meetings.

420

421 **Funding Statement:** This research received no specific grant from any funding agency,  
422 commercial or not-for-profit sectors.

423

424 MF is supported by NIDA and NIAAA intramural research funding (ZIA-DA000635 and ZIA-  
425 AA000218). The content of this article is solely the responsibility of the authors and does not  
426 necessarily represent the official views of the National Institutes of Health.

427

428 **Declaration of interest statement:** The authors have no conflict of interest to declare.

429 **References**

430

431 1. Smith DE. The Evolution of Addiction Medicine as a Medical Specialty. *AMA J Ethics*. 2011  
432 December 1;13(12):900–5.

433 2. Soyka M, Gorelick DA. Why should addiction medicine be an attractive field for young  
434 physicians? *Addict Abingdon Engl*. 2009 Feb;104(2):169–72.

435 3. Ayu A, Schellekens A, Iskandar S, Pinxten L, Jong C. Effectiveness and Organization of  
436 Addiction Medicine Training Across the Globe. *Eur Addict Res*. 2015 May 5;21:223–39.

437 4. Klimas J, Small W, Ahamad K, Cullen W, Mead A, Rieb L, et al. Barriers and facilitators to  
438 implementing addiction medicine fellowships: a qualitative study with fellows, medical students,  
439 residents and preceptors. *Addict Sci Clin Pract*. 2017 Sep 20;12(1):21.

440 5. De Jong CAJ, Van De Wetering BJM. Addiction medicine is an attractive field for young  
441 physicians—a call for a European initiative for the training in addiction medicine. *Addict Abingdon Engl*.  
442 2009 Jul;104(7):1258–9.

443 6. Ayu AP, el-Guebaly N, Schellekens A, De Jong C, Welle-Strand G, Small W, et al. CORE  
444 ADDICTION MEDICINE COMPETENCIES FOR DOCTORS: AN INTERNATIONAL CONSULTATION  
445 ON TRAINING. *Subst Abuse*. 2017;38(4):483–7.

446 7. Haber PS. International Perspectives in Postgraduate Medical Training in Addiction Medicine.  
447 *Subst Abuse*. 2011 April 13;32(2):75–6.

448 8. el-Guebaly N, Violato C. The international certification of addiction medicine: validating clinical  
449 knowledge across borders. *Subst Abuse*. 2011 Apr;32(2):77–83.

450 9. Rasyidi E, Wilkins JN, Danovitch I. Training the next generation of providers in addiction  
451 medicine. *Psychiatr Clin North Am*. 2012 Jun;35(2):461–80.

452 10. Abuse NI on D. Early & Mid-Investigator Career Development (K) Awards [Internet]. National  
453 Institute on Drug Abuse. 2020 [cited 2020 Jul 16]. Available from: [https://www.drugabuse.gov/about-](https://www.drugabuse.gov/about-nida/organization/offices/office-nida-director-od/office-research-training-diversity-disparities-ortdd/office-diversity-health-disparities-odhd/early-mid-investigator-career-development-k-awards)  
454 [nida/organization/offices/office-nida-director-od/office-research-training-diversity-disparities-](https://www.drugabuse.gov/about-nida/organization/offices/office-nida-director-od/office-research-training-diversity-disparities-ortdd/office-diversity-health-disparities-odhd/early-mid-investigator-career-development-k-awards)  
455 [ortdd/office-diversity-health-disparities-odhd/early-mid-investigator-career-development-k-awards](https://www.drugabuse.gov/about-nida/organization/offices/office-nida-director-od/office-research-training-diversity-disparities-ortdd/office-diversity-health-disparities-odhd/early-mid-investigator-career-development-k-awards)

456 11. Alford DP, Carney BL, Jackson AH, Brett B, Bridden C, Winter M, et al. Promoting addiction  
457 medicine teaching through functional mentoring by co-training generalist chief residents with faculty  
458 mentors. *Subst Abuse*. 2018;39(3):377–83.

459 12. McMillan SS, King M, Tully MP. How to use the nominal group and Delphi techniques. *Int J Clin*  
460 *Pharm*. 2016 Jun;38(3):655–62.

461 13. Niederberger M, Spranger J. Delphi Technique in Health Sciences: A Map. *Front Public Health*.  
462 2020;8:457.

463 14. Rayhan RU, Zheng Y, Uddin E, Timbol C, Adewuyi O, Baraniuk JN. Administer and collect  
464 medical questionnaires with Google documents: a simple, safe, and free system. *Appl Med Inform*.  
465 2013;33(3):12–21.

466 15. The Mentor's Guide: Facilitating Effective Learning Relationships, 2nd Edition | Wiley [Internet].  
467 Wiley.com. [cited 2021 Dec 25]. Available from: <https://www.wiley.com/en->

- 468 [us/The+Mentor%27s+Guide%3A+Facilitating+Effective+Learning+Relationships%2C+2nd+Edition-p-](https://doi.org/10.2196/jmir.2042)  
469 [9780470907726](https://doi.org/10.2196/jmir.2042)
- 470 16. Eysenbach G. Improving the quality of Web surveys: the Checklist for Reporting Results  
471 of Internet E-Surveys (CHERRIES) [published correction appears in doi:10.2196/jmir.2042]. *J*  
472 *Med Internet Res*. 2004;6(3):e34. Published 2004 Sep 29. doi:10.2196/jmir.6.3.e34  
473
- 474 17. Iannucci R, Sanders K, Greenfield SF. A 4-year curriculum on substance use disorders for  
475 psychiatry residents. *Acad Psychiatry J Am Assoc Dir Psychiatr Resid Train Assoc Acad Psychiatry*.  
476 2009 Feb;33(1):60–6.
- 477 18. Tontchev GV, Housel TR, Callahan JF, Kunz KB, Miller MM, Blondell RD. Specialized training on  
478 addictions for physicians in the United States. *Subst Abuse*. 2011 Apr;32(2):84–92.
- 479 19. Pinxten WJL, De Jong C, Hidayat T, Istiqomah AN, Achmad YM, Raya RP, et al. Developing a  
480 competence-based addiction medicine curriculum in Indonesia: the training needs assessment. *Subst*  
481 *Abuse*. 2011 Apr;32(2):101–7.
- 482 20. O'Brien S, Cullen W. Undergraduate medical education in substance use in Ireland: a review of  
483 the literature and discussion paper. *Ir J Med Sci*. 2011 Dec;180(4):787–92.
- 484 21. Tripathi R, Singh S, Bhad R. The need, scope, challenges, and potential solutions for enhancing  
485 addiction psychiatry training in India. *Indian J Psychiatry*. 2020;62(6):728–31.
- 486 22. Carroll J, Goodair C, Chaytor A, Notley C, Ghodse H, Kopelman P. Substance misuse teaching  
487 in undergraduate medical education. *BMC Med Educ*. 2014 Feb 17;14:34.
- 488 23. Das S, Roberts LW. Addiction Training: Striving to Fill an Unmet Need. *Acad Psychiatry*. 2016  
489 June 1;40(3):451–3.
- 490 24. Brown AT, Kolade VO, Staton LJ, Patel NK. Knowledge of addiction medicine among internal  
491 medicine residents and medical students. *Tenn Med J Tenn Med Assoc*. 2013 Mar;106(3):31–3.
- 492 25. Wakeman SE, Baggett MV, Pham-Kanter G, Campbell EG. Internal medicine residents' training  
493 in substance use disorders: a survey of the quality of instruction and residents' self-perceived  
494 preparedness to diagnose and treat addiction. *Subst Abuse*. 2013;34(4):363–70.
- 495 26. Orsolini L, Rojnić Palavra I, Papanti GD, Potočan M, Quattrone D, Martens M, et al. Psychiatry  
496 Trainees' Attitudes, Knowledge, and Training in Addiction Psychiatry—A European Survey. *Front*  
497 *Psychiatry*. 2021 January 8;11:585607.
- 498 27. Tang Y, Wiste A, Mao P, Hou Y. Attitudes, knowledge, and perceptions of Chinese doctors  
499 toward drug abuse. *J Subst Abuse Treat*. 2005 Oct;29(3):215–20.
- 500 28. Klimas J, Kelly D, Adam A, Arya S, Ruiz BII, Krupchanka D, et al. What Are the Self-Assessed  
501 Training Needs of Early Career Professionals in Addiction Medicine? A BEME Focused Review.  
502 *MedEdPublish [Internet]*. 2020 Apr 3 [cited 2021 May 19];9. Available from:  
503 <https://www.mededpublish.org/manuscripts/2843#bibliographyreferences>
- 504 29. Kelly D, Adam A, Arya S, Indave I, Krupchanka D, Cullen W, et al. What are the training needs of  
505 early career professionals in addiction medicine? A BEME scoping review. :15.

- 506 30. Ayu AP, Schellekens AFA, Iskandar S, Pinxten L, De Jong CAJ. Effectiveness and Organization  
507 of Addiction Medicine Training Across the Globe. *Eur Addict Res.* 2015;21(5):223–39.
- 508 31. Barron R, Frank E, Gitlow S. Evaluation of an experiential curriculum for addiction education  
509 among medical students. *J Addict Med.* 2012 Jun;6(2):131–6.
- 510 32. Kahan M, Midmer D, Wilson L, Liu E. Faculty Rating of Learning Objectives for an Undergraduate  
511 Medical Curriculum in Substance Abuse. *Subst Abuse.* 2001 Dec;22(4):257–63.
- 512 33. Learning for Early Careers in Addiction & Diversity (LEAD) Programme. :20.
- 513 34. AAAP Mentoring Programme - AAAP [Internet]. American Academy of Addiction Psychiatry.  
514 [cited 2021 Dec 25]. Available from: <https://www.aaap.org/membership/aaap-mentoring/>
- 515 35. Choi AMK, Moon JE, Steinecke A, Prescott JE. Developing a Culture of Mentorship to  
516 Strengthen Academic Medical Centers. *Acad Med.* 2019 May;94(5):630–3.
- 517 36. Masson C, Sorensen J. Mentoring Emerging Scientists for Careers in Substance Use Research.  
518 [cited 2021 Dec 25]; Available from: <https://grantome.com/grant/NIH/R25-DA035163-06>
- 519 37. DAPC Grant [Internet]. United Nations : Office on Drugs and Crime. [cited 2021 Dec 25].  
520 Available from: [//www.unodc.org/unodc/en/prevention/youth-initiative/dapc-grant.html](http://www.unodc.org/unodc/en/prevention/youth-initiative/dapc-grant.html)
- 521 38. Klimas J, McNeil R, Ahamad K, Mead A, Rieb L, Cullen W, et al. Two birds with one stone:  
522 experiences of combining clinical and research training in addiction medicine. *BMC Med Educ.* 2017  
523 January 23;17(1):22.
- 524 39. Merritt C, Jack H, Mangezi W, Chibanda D, Abas M. Positioning for success: building capacity in  
525 academic competencies for early-career researchers in sub-Saharan Africa. *Glob Ment Health*  
526 [Internet]. 2019 ed [cited 2021 Dec 25];6. Available from:  
527 [https://www.cambridge.org/core/journals/global-mental-health/article/positioning-for-success-building-  
528 capacity-in-academic-competencies-for-earlycareer-researchers-in-subsaharan-  
529 africa/0C590397A6380629B5835434A5F2B39E](https://www.cambridge.org/core/journals/global-mental-health/article/positioning-for-success-building-capacity-in-academic-competencies-for-earlycareer-researchers-in-subsaharan-africa/0C590397A6380629B5835434A5F2B39E)
- 530 40. McCarty KN, McDowell YE, Sher KJ, McCarthy DM. Training Health Services Psychologists for  
531 Research Careers in Addiction Science. *Train Educ Prof Psychol.* 2020 Feb;14(1):70–7.
- 532 41. Koelkebeck K, Andlauer O, Asztalos M, Jovanovic N, Kazakova O, Naughton S, et al. Research  
533 by Psychiatric Trainees and Early Career Psychiatrists—Results of a Survey From 34 Countries in  
534 Europe. *Front Psychiatry.* 2021 Sep 10;12:718669.
- 535 42. Bhad R, Hazari N. Predatory journals in psychiatry: A note of caution. *Asian J Psychiatry.* 2015  
536 Aug;16:67–8.
- 537 43. Forero DA, Oermann MH, Manca A, Deriu F, Mendieta-Zerón H, Dadkhah M, et al. Negative  
538 Effects of “Predatory” Journals on Global Health Research. *Ann Glob Health.* 84(4):584–9.
- 539 44. Mitchell G, McCambridge J. The ‘snowball effect’: short and long-term consequences of early  
540 career alcohol industry research funding. *Addict Res Theory.* 2021 July 22;0(0):1–7.

541  
542  
543  
544




Figure 1






Figure 2