1 Key topics in pandemic health risk communication: A qualitative

2 study of expert opinions and knowledge

3 Siv Hilde Berg^{1*}, Marie Therese Shortt¹, Jo Røislien¹, Daniel Adrian Lungu¹, Henriette

4 Thune^{1,} Siri Wiig¹

- 5 1. Centre for Resilience in Healthcare, Department of Quality and Health Technology,
- 6 Faculty of Health Sciences, University of Stavanger, Stavanger, Norway

7 *Corresponding author

- 8 Email: <u>Siv.h.berg@uis.no</u>(SHB)
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19

20 Abstract

21	Background. Science communication can provide people with more accurate information on
22	pandemic health risks by translating complex scientific topics into language that helps people
23	make more informed choices on how to protect themselves and others. During pandemics,
24	experts in medicine, science, public health, and communication are important sources of
25	knowledge for science communication. This study uses the COVID-19 pandemic to explore
26	these experts' opinions and knowledge of what to communicate to the public during a
27	pandemic. The research question is: What are the key topics to communicate to the public
28	about health risks during a pandemic?
29	Method. We purposively sampled 13 experts in medicine, science, public health, and
30	communication for individual interviews, with a range of different types of knowledge of
31	COVID-19 risk and communication at the national, regional and hospital levels in Norway.
32	The interview transcripts were coded and analysed inductively in a qualitative thematic
33	analysis.
34	Results. The study's findings emphasise three central topics pertaining to communication
35	about pandemic health risk during the first year of the COVID-19 pandemic in Norway: 1)
36	how the virus enters the human body and generates disease; 2) how to protect oneself and
37	others from being infected; and 3) pandemic health risk for the individual and the society.
38	Conclusion. The key topics emerging from the expert interviews relate to concepts
39	originating from multiple disciplinary fields, and can inform frameworks for interprofessional
40	communication about health risks during a pandemic. The study highlights the complexity of
41	communicating pandemic messages, due to scientific uncertainty, fear of risk amplification,
42	and heterogeneity in public health and scientific literacy. The study contributes with insight
43	into the complex communication processes of pandemic health risk communication.
44	
45	Key words: Science communication, mental models' theory, expert study, risk
46	communication, health communication, pandemic response.

48 1 Introduction

In a pandemic, a virus capable of infecting humans spreads and crosses international
borders, usually affecting a large number of people (1). In the past century, there have been
four pandemics: influenza (1918–1919), AIDS (1966-), H1N1 "swine flu" (2009–2010), and
COVID-19 (2019-).

53 A pandemic associated with an unknown virus brings uncertainty that challenges 54 politicians' and policy makers' decision making. Scientific uncertainty gives rise to a difficult 55 balancing act in public communication: being transparent about the limits of human 56 knowledge, yet still communicating clear messages (2). The challenge is reinforced by poor 57 public health literacy, which impedes people's ability to understand the rationale for 58 governmental recommendations and to anticipate the consequences of their health choices (3). 59 To strengthen peoples' scientific-and health literacy, science communication should convey 60 accurate information on health risks in a language that enables people to make independent 61 and informed choices that will protect both themselves and others (4). 62 Scientific uncertainty, combined with the difficulty of delivering timely but complex 63 messages to the population in a competitive media market, highlights the importance of health 64 risk communication being not only correct and trustworthy, but also engaging. What to 65 communicate to the public about staying safe during a pandemic – and how to communicate it 66 - thus needs to be done with tremendous care (5). 67 Experts in medicine, risk, public health, and communication are important sources of 68 knowledge in science communication during public health crisis (4, 6). This study uses the 69 COVID-19 pandemic to explore experts' opinions and knowledge of what to communicate to 70 the public during a pandemic. The research question is: What are the key topics to 71 communicate to the public about health risks during a pandemic?

72 1.1 Health risk communication

73	Public communication during health emergencies aims to improve health outcomes by
74	influencing, engaging, and reaching out to different at-risk audiences (7, 8). Public
75	communication about a pandemic is, as with all communication, a multifaceted field in which
76	the contents of the message, the messenger and the qualities of the audience attributes affect
77	the reception of the message itself (8-11). In this section, we present four topics which are
78	relevant to this study's findings regarding the experts' communication of key topics: 1) trust
79	and communicating uncertainty; 2) tradeoffs in decision making; 3) risk amplification; and 4)
80	public engagement.

81 1.1.1 Trust and communicating uncertainty

82 Communicating honestly about uncertainty can build public trust and increase the 83 legitimacy and credibility of the decision making process (12). Public trust in the messengers 84 of pandemic health risk information is associated with their willingness to comply with 85 measures to prevent infection (13, 14). The World Health Organisation (WHO) advises health 86 authorities to be candid about risk, events and interventions and to be forthcoming about what 87 is known and what is not (7). However, the different types of uncertainty that have been 88 communicated have different effects on trust. Gustafson and Rice (15) reviewed the 89 experimental literature and found that communicating disagreements or conflict in science 90 (consensus uncertainty) were most often associated with reduced credibility. Communicating 91 technical uncertainty (e.g., probabilities) had positive or no effect on perceived credibility. 92 The effect was mixed in relation to communicating scientific uncertainty, which had both 93 positive and negative effects on perceived credibility.

94 1.1.2 Tradeoffs in decision making

95 Several safety science theorists have outlined *tradeoffs* as an inevitable balancing act 96 among all kinds of pressures and safety in hazardous systems (16-18), including healthcare 97 (19). Tradeoffs involve ambiguity over what constitutes the right answer. According to 98 Gregory et al. (20) tradeoffs are usually difficult to communicate. Changes in one endpoint 99 are often achieved at the expense of other goals, and the consequences of tradeoffs are often 100 unclear to the decision makers. According to Gregory et al. (20) tradeoffs reflect issues that 101 science cannot resolve; they depend on values in making decisions about risk. Little is known 102 about the public behavioral effect of communicating tradeoffs on decision making in 103 pandemics, yet Gregory et al. (20) and Norheim et al. (21) argued that tradeoffs should be 104 transparent to foster trust.

105 1.1.3 Risk amplification

106 The Social Amplification of Risk Framework (SARF) suggests that most health risk 107 messages are sent through transmitters (e.g. media, messengers) which alter the original 108 message by intensifying or attenuating some signals before passing them on (10). SARF 109 proposes that risk deemed less urgent by experts may be *amplified* by other transmitters; 110 however, risk that experts consider less urgent may be *attenuated*. According to SARF, 111 silence from experts and decision makers may breed fear and suspicion among those at risk 112 and lead to risk amplification. Nevertheless, a moderate level of fear can mitigate the risk of a 113 pandemic by encouraging behaviour that protects public health (22). To minimise risk 114 amplification, health risk communicators should avoid sensationalism, speculation and 115 disturbing images (23).

116 1.1.4 Public engagement

117	In the early 1980s, "lay" perspectives were perceived as fallible and subjective by
118	communication experts and policy makers, and communication interventions aimed to align
119	lay perspectives with those of the experts in a one-way approach. As the field of health and
120	risk communication evolves, multi-way approaches to communication emphasising public
121	engagement have largely replaced the one-way approach to communication (9). The WHO
122	advises health authorities to involve the public in decision making to ensure that interventions
123	are collaborative and contextually suitable, thereby constituting a two-way approach (7).
124	There is growing awareness of the benefits of including patients and the public in healthcare.
125	However, involvement needs to be conducted at an even wider system level, e.g., involvement
126	in the design of public health communication (24).
127	

128 2 Method

129 The present study applies a qualitative design using individual interviews with 130 Norwegian experts in medicine, science, public health, and communication guided by the 131 mental models' approach to risk communication (4, 25). The mental models' approach to risk 132 communication involves both experts' knowledge and opinions of what people should know 133 to make informed decisions about a topic (normative research) and the ability of the intended 134 audience (descriptive research) to create tailored risk communication interventions 135 (prescriptive research) (4). This study reports on the normative research phase (26). See Berg 136 et al. (27) for results from the audience analysis, and Røislien at al. (28) for the design of the 137 intervention.

Expert opinions are valuable components of communication designs when the scientific evidence is missing, is inconsistent or incomplete, as is often the case with new viruses such as the coronavirus (SARS-COV-2) (25). Prior research of expert mental models' have used

- 141 health scientists' and medical practitioners' knowledge of infectious diseases to design health
- 142 risk communications (29, 30). To our knowledge, no mental models' studies of infectious
- 143 diseases have so far included the knowledge of communication experts.

144 2.1 Participants and sampling strategy

145 Communicating key topics during a pandemic is a dynamic process in which a range of

146 interprofessional issues of pandemic health risk are attributed different temporal weights.

147 Infection control was the Norwegian government's and policy makers' main focus in the first

148 wave of the pandemic, while mental health and public health became increasingly important

as the pandemic evolved (31). Communication science is an important field of knowledge to

150 pandemic responses (7), however it is not necessarily emphasised as part of the public health

151 risk communication field. We therefore sought informants with expertise in medicine,

- science, public health, and communication to cover the variety of expertise relevant to
- 153 pandemic risk communication.

154 We purposively sampled 13 experts in medicine, medical/health research, public health,

and communication for individual interviews, all with extensive knowledge of COVID-19 and

156 risk communication at the hospital, regional and national levels in Norway. Participants who

157 met the inclusion criteria were identified and recruited by invitation from the research group.

158 One participant dropped out of the study due to time issues. Table 1 displays sample

159 characteristics and the participants' field of expertise, professional role during the COVID-19

160 pandemic and in public communication.

	Field of expertise and sample category	Professional role	Public communication role related to COVID-19	Gender
1	Public health management, nurse (Public health)	Head of department for community healthcare service. Infection tracing and training of healthcare professionals	Advisor of municipal communication to the public	F

Table 1 Sample Characteristics

2	Specialist in community medicine (medicine)	Municipal chief physician and head of COVID-19 emergency room. Infection prevention at individual and group level	Municipal communication to the public	М
3	Public emergency and risk management (public health)	Head of municipal emergency management and community risk communication	Municipal communication to the public	М
4	Marketing (communication)	Audience analysis, targeted communication to adolescents and young adults	Advisor of municipal communication to the public	F
5	Clinical psychologist, PhD in psychology, journalism (public health)	Psychologist in primary mental healthcare	Municipal communication to the public	F
6	Communication and journalism (communication)	Audience analysis. Targeted communication related to motivation and mental health	Advisor of municipal communication to the public	F
7	Professor and infection medicine specialist (medicine)	Head of infection medicine in a Regional Hospital. Infection prevention at individual and group level	Communication to the public at regional level	М
8	Infection medicine specialist (medicine)	Infection prevention at hospital level. Microbiology, virus transmission and treatment of COVID patients	Science dissemination in medical journals	F
9	Infection medicine specialist (medicine)	Head of infection medicine in a Regional Hospital. Infection prevention at individual and group level	Communication to the public at regional level	М
10	Risk communication and ethnography (communication)	Director of nationwide health risk communication	Strategy level, nationwide health risk communication	F
11	Professor of statistics (medical research)	Biostatistician. Mathematical modelling of the R-value and the spread of COVID-19.	Communication to the public at national level	М
12	Professor of health communication and cell-biologist (health research)	Health communication researcher. Health literacy in the general public	Science dissemination in national media	М
13	PhD and infection medicine specialist (medical research)	Infection medicine researcher. Virus transmission, immunity and antibodies related to SARS-COV-2	Communication in national media	F

161

162 2.2 Interviews

163 A semi-structured interview guide was developed to explore key topics related to

164 communication about COVID-19 virus transmission and exposure, mitigation management,

165 and potential effects of infection (S1 File). The interview guide also explored key topics

166	related to pandemic health risk generally and those in each expert's specialty. The interview
167	guide was pilot tested with participant number one. The interviewer had no prior established
168	relationships with the participants, and the participants were informed about the reasons for
169	this research (exploring key topics in pandemic communication and their experiences with
170	communication). Due to national COVID-19 measures to keep physical distance, the
171	participants were interviewed individually by SHB using a video conferencing software
172	program (Zoom). Mean duration of interviews was 59 (range 46-80 min) minutes. Interviews
173	were audio recorded and transcribed. Ethical approval was obtained from the Norwegian
174	Centre for Research Data (Ref nr. 583192). All participants gave written, voluntary and
175	informed consent. The interviews were conducted between February and April 2021.

176 2.3 Context

177 The data material in this study presents important topics for a first-year pandemic

178 response. At the time of data collection, individuals in Norway with the highest risk of severe

179 consequences of COVID-19 were being vaccinated. The UK variant of the coronavirus

180 (B.1.1.7) had started to spread in the Norwegian population and led to the third wave of

181 COVID-19.

182 In the first year of the pandemic, the number of positive cases and deaths per million due

to COVID-19 were much lower in Norway than in other European countries (32). Since the

184 first wave of COVID-19 struck Norway in March-April 2020, the government strategy has

185 been to limit the spread of infection by using testing, isolation, contact tracing, and

186 quarantine, and by occasionally closing schools and kindergartens, and cancelling cultural

187 events (33). The goal has been to control the spread of infection so that the infection is

188 manageable and does not exceed the capacity of the health and care service and the municipal

189 health service (34).

190	Norway has a strong public healthcare system; most hospitals are funded and owned by
191	the state. The welfare state is well-developed and founded on the principles of universal
192	access, decentralisation and free choice of provider (35). Compared to other OECD countries,
193	Norway is a high-trust society (36); in a recent study 96% of the participants reported that
194	they trusted the health authorities in general (37). Trust in the health authorities has remained
195	high and stable during the pandemic; between 80 and 90% of the population expressed a high
196	level of trust in the health authorities (31), and especially in the Norwegian Institute of Public
197	Health (NIPH) (38). Norway has a population of 5.4 million. Although the country has the
198	third lowest population density in Europe, at the beginning of the COVID-19 outbreak,
199	Norway was second only to Italy in the number of confirmed cases per capita. However, by
200	April 2020 Norway had been able to "flatten the curve" and ease the competition for hospital
201	beds (38). A strategy used by the Norwegian government during the first pandemic response
202	was to appeal to people's sense of solidarity by using the term <i>dugnad</i> , an old cultural
203	practice that describes volunteer activities that will benefit the community (39).

204 2.4 Analysis

205 The expert interview transcripts were coded and analysed inductively in a qualitative 206 thematic analysis (40). The unit of analysis was "what people should know", and the analysis 207 followed the five phases of systematic thematic analysis as outlined by Braun and Clark (40). 208 To become familiar with the data, all transcripts were read by SHB and MT (phase 1), who 209 discussed first impressions of the material with SW. SHB inductively coded data across the 210 entire data set related to the meaning units (phase 2). The initial codes were extracted to 211 tables, which were used to search for meaningful patterns, themes, and levels of themes. All 212 co-authors reviewed the initial organisation of codes. The codes were then compiled under 36 213 key messages, which were then abstracted into seven sub-themes and three themes (phase 3).

214	In phase 4, a table with thematically organised codes was made to validate the themes in
215	relation to the codes and the entire data set. The themes were refined and renamed to generate
216	clear terms for each theme (phase 5). The fifth and final phase consisted of writing out the
217	results with codes and extract examples, which all co-authors read and discussed.
218	Thematic mapping is used in thematic analysis to find and display relationships between
219	codes and themes (40) and influence diagrams are often used to visualise experts mental
220	models (25, 26). However, these visualisation techniques are one-dimensional and do not
221	allow for multi-layered information to be displayed. A bubble chart was thus created; a
222	visualisation method developed by Shortt et al. (41) for visualising qualitative interview data,
223	inspired by thematic mapping and influence diagrams (25, 40). The bubble chart displays
224	topics emphasised by experts in medicine, medical/health researchers (science), public health,
225	and communication, with colour coding that enables visual comparison of topics emphasised
226	by the different experts. Topics with disagreement whether they should be communicated to
227	the public or due to being perceived as not relevant, to complicated or too uncertain to
228	communicate to the public, are highlighted.

3

Results

229

230

231 The first impression of the data material was linked to how seriously the medical and 232 research experts took their responsibility for communicating pandemic risk to the public, yet 233 they emphasised not being trained in public communication. The medical and research 234 experts were reflexive in crafting each message to each audience, and in maintaining their 235 ethics in public communication. They expressed personal uncertainty in their role as 236 communicators to the public. Although the communication experts had been trained, none of 237 them had ever had to issue communications about such an ambiguous situation, especially one

238	of the scale of the	COVID-19 pande	mic. All of the expert	ts had to learn a	about this type of

- 239 communication during a pandemic by doing it, despite scientific uncertainty and constantly
- changing evidence.
- 241 The analysis of expert opinions and knowledge identified 36 key messages in
- 242 pandemic health risk communication to the public which were organised into seven sub-
- themes as part of three main themes: how the virus enters the human body and produces
- 244 disease; how to protect oneself and others from infection; and pandemic health risk for the
- 245 individual and the society (Table 2).

Themes	How the virus gets into the human body and generates disease	How to protect oneself and others from being infected	Pandemic health risk for the individual and the society
Sub- themes	Modes of virus transmission	Infection prevention at the individual level	Solidarity
	Virus and immunity	Infection prevention at group level	Control of the spread
			Risk tradeoffs


Table 2 Key Topics to Communicate to the Public Related to Pandemic Risk

246

247 How each of the expert groups emphasised each of the themes and sub-themes is visualised in

Figure 1. An animated version of the bubble chart reveals the multi-layer organisation of the

themes and sub-themes according to the various expert groups (S2 File).




251

252 **Figure 1** Bubble chart visualising concept mapping of expert opinions.

253 3.1 How the virus gets into the human body and generates disease

- 254 The theme "how the virus gets into the body and generates disease" has two sub-
- themes: modes of virus transmission, and virus and immunity (Figure 2).



256

257 *Figure 2* Bubble chart representation of information relating to the theme 'How the virus gets into the human body and generates disease'

259

- 260 3.1.1 Modes of virus transmission
- 261 The experts' emphasis on modes of virus transmission is related to a potential increase
- 262 in people's comprehension of why they should comply with recommendations to protect
- themselves from COVID-19. One necessary message to communicate was how droplets
- 264 containing the virus could reach people within a metre of the person emitting the droplets.
- 265 This message helps to reinforce the need to keep physical distance.

266 Experts from several fields insisted that the public needed to comprehend the concepts

267 of *incubation period* and *asymptomatic transmission*, which they expressed as: "You can be

contagious even if you don't have symptoms. Non-symptomatic people can carry the disease
without being aware of it. You cannot tell if a person is infected or not, so everyone needs to
keep distance".

271	At the time of data collection there was scientific uncertainty about the modes of
272	virus transmission of the corona virus. Specialists in infectious disease highlighted that it was
273	important to address this uncertainty by communicating the most dominant route of
274	transmission to help people distinguish possible from the most probable way of transmission.
275	Some of the experts used the terms contact/surface transmission, droplet transmission
276	and airborne transmission when communicating with the public. The experts agreed that
277	there was not a clear distinction but rather a continuum between droplet and airborne
278	transmission. Although one expert in infectious diseases believed it was important to illustrate
279	a continuum of droplets with different sizes and reach, another expert communicated how
280	temperature and humidity affected how long the droplets stayed in the air and thus increased
281	the probability of virus survival, and therefore the chance of being infected.
282	Contact transmission was found to be difficult to explain, as people who received the
282 283	Contact transmission was found to be difficult to explain, as people who received the message interpreted the same information in different ways:
283 284 285 286 287 288 289	
283 284 285 286 287 288	message interpreted the same information in different ways: We try to communicate that the main infection path is air and other people. If you practice good hand hygiene then you are reasonably safe, even if you touch a door handle covered with virus. But people perceive that information very differently. Some people perceive it as deadly and go to extremes, but others will perceive this as something they don't have to think
283 284 285 286 287 288 289 290	 message interpreted the same information in different ways: We try to communicate that the main infection path is air and other people. If you practice good hand hygiene then you are reasonably safe, even if you touch a door handle covered with virus. But people perceive that information very differently. Some people perceive it as deadly and go to extremes, but others will perceive this as something they don't have to think about, so it does not matter how you deliver the message (No. 12).
283 284 285 286 287 288 289 290 291	 message interpreted the same information in different ways: We try to communicate that the main infection path is air and other people. If you practice good hand hygiene then you are reasonably safe, even if you touch a door handle covered with virus. But people perceive that information very differently. Some people perceive it as deadly and go to extremes, but others will perceive this as something they don't have to think about, so it does not matter how you deliver the message (No. 12). Some of the infectious disease experts did not communicate airborne transmission to

air. The health communication expert described how creating mental images of the airspace

around people by using analogies to water have been used successfully as a pedagogic tool:

297 298 299 300 301 302	People have rather vague ideas about the medium of air, and what it means to breathe at a distance of one metre. It spreads in the same way, as if you dip a teabag in a cup of water. Because there is no hole here, it's compact matter These particles are carried through matter in the same way as if you had dropped a small wooden plank on the water surface (No. 12)
303	Experts who did not use the terms droplets, airborne and contact/surface
304	transmission, believed they were too technical and ambiguous. Instead, they prioritised the
305	conveying of clear messages; how people could protect themselves by washing their hands
306	and keeping their distance, as exemplified by a communication expert for nationwide health
307	risk communication:
308 309 310 311 312 313 314 315 316	We have not been completely oblique about how the virus infects, and new virus variants change how the virus behaves We have not used the terms airborne, droplet and contact transmission so often. We have been more concerned with explaining what the best precautions to take to avoid being infected by the virus areWe have noticed that there is a great demand among the general public for information about the type of infection, but people are just as concerned about how to handle the infection. They want to know what to do (No. 10).
317	The marketing expert pointed out that modes of virus transmission were not
318	mentioned in their target group analysis of young people's concerns and values, thus the topic
319	was not included in their targeted communication to adolescents.
320	3.1.2 Virus and immunity
321	The infectious diseases experts believed that it was important to explain how a virus
322	attaches itself to the cells and creates disease. Such mechanisms were important to increase
323	peoples understanding of variant virus' changes in transmission. The variants of the virus had

324 different causes for their increased contagiousness; for example, an infectious diseases expert

325	explained how a virus which can be in the upper respiratory tract one to two days before the
326	disease breaks out is extremely contagious. The health communication expert believed that
327	explanations of virus autonomy and multiplication were too complicated for most people to
328	understand. The communication expert for nationwide health risk communication noted that
329	they had included this information when communicating to the public, despite its complexity:
 330 331 332 333 334 335 336 337 	We have tried to explain that here is a part of this virus, this spike that attached better [SARS-COV-2 Spike Protein]. It is complex information, but we have really thought that it is important to try to give it anyway, so that people understand why we still ask them to keep distance, and maybe even keep more distance now, because the contagiousness in the virus variations is different to the first virus variant that we are familiar with. We have changed the information both about the virus itself and also about the advice we give (No. 10)
338	
339	Infectious diseases experts experienced that the public tended to perceive mutations of
340	viruses as dangerous, and considered it important to communicate the message that mutations
341	are a natural part of the virus's life cycle. The virus spreads by becoming more contagious
342	without killing its host. Thus, by mutating the virus becomes less lethal.
343	Experts from diverse fields considered it important to inform about the effects of
344	vaccination on immunity and how vaccination affected the individual and the society. At the
345	individual level, it was important to convey the message that "you can still be infected and
346	infect others even if you are vaccinated, however the vaccine may limit the lethality of
347	infection". At the society level, reducing the risk of infection through herd immunity through
348	mass vaccination was a relevant concept to communicate.
349	3.2 How to protect oneself and others from being infected
350	The theme "how to protect oneself and others from being infected" pertained to two
251	sub the mass infection provention of the individual level, and at the group level (Figure 2)

351 sub-themes: infection prevention at the individual level, and at the group level (Figure 3).





Figure 3 Bubble chart representation of information from interviews relating to the theme 'How to protect oneself and others from being infected'

355

356 3.2.1 Infection prevention at the individual level



-		
	^	7
٦	n	- 1
~	v	

307	
368	However, the rules were difficult to enforce if people did not understand the
369	underlying principles. Many of the experts insisted that the rationales for infection prevention
370	measures were important to explain to the public. Explaining why people had to follow the
371	rules, and being open about the evidence and the pandemic decision making, were considered
372	important in the creation of trust. The marketing expert emphasised that adolescents did not
373	understand the rationale for the rules, so they produced videos to describe not only how to
374	protect themselves, but also why.
375	The infectious diseases experts considered it important to teach the public about
376	probabilities, and to convey the message that "there are no clear boundaries between safe and
377	unsafe distance, the larger distance you keep, the less probability you have for being exposed
378	to infection". According to the research and infectious diseases expert:
379 380 381 382 383 384 385	In communication with the public, it is important to emphasise that when you set a two-metre limit, it is not because there is a black-and-white limit that if you are 2 metres and 10 cm from someone, you are safe, and if you are 1 metre and 90 cm, then you are not. That's silly. So, you have to trust people to use their common sense and understand that it is a question of probabilities in all of the rules and guidelines that exist (No.13).
386	3.2.2 Infection prevention at group level
387	While the scientific evidence for infection prevention was more definitive at the
388	individual level, higher scientific uncertainty was related to infection prevention at the group
389	level, which often led to public debates in the media over whether or not the evidence
390	supported lockdowns of specific activities. The infectious disease experts emphasised the

391 need to communicate the risk of being physically close to others and engaging in activities


that increased the expulsion of droplets containing the virus (e.g., singing, going to pubs and

393 gyms) to increase people's understanding of why those activities had been stopped. As the

394 nationwide health risk communication expert emphasised:

395 396 397 398	It is not about whether it is correct to close that particular school or shopping centre or not, it's about reducing the total contact frequency among individuals, and these choices will be implemented with a high level of uncertainty regarding their efficiency (No. 10).
399	
400	To reduce the stigma of contracting COVID-19, the communication director
401	communicated that "this virus does not differentiate between people". However, some of the
402	experts at community and regional hospitals reported feeling trapped between safeguarding
403	individuals' right to privacy and satisfying the media's need for information.
404	Experts emphasised that conveying a message of hope could help people remain
405	motivated despite of social distancing. The medical experts emphasised supportive messages
406	and acknowledging people's compliance. A communication expert described tailoring
407	different messages to different populations in the municipality by showing them activities
408	they could do during lockdown:
409 410 411 412 413	We identified that people wanted the municipality to stop telling them what they could not do, but rather what they could do. People can be outdoors even if we are on red level, and we have to think about measures to support, that involves being outdoor (No. 6).
414	3.3 Pandemic health risk for the individual and the society
415	The theme "pandemic health risk for the individual and the society" was related to three

416 sub-themes: solidarity, control of the spread, and risk-tradeoffs (Figure 4).



418 *Figure 4* Bubble chart representation of information from interviews relating to the theme 'Pandemic health risk for the individual and the society'

420

417

421 3.3.1 Solidarity

422 Experts explained the importance of informing the public of the collective moral

423 responsibility to protect others, appealing to Norwegians' sense of solidarity. As the

424 communication expert at the national level explained:

425	Initially we needed to bring the entire population to the same problem
426	definition, because we need to share the problem. We explained that it was
427	most dangerous for the oldest and groups with chronic diseases. You could
428	have chosen to be a society that said: you have to protect yourself in a
429	bubble until we are all infected. But that's ethically completely
430	unjustifiable. This means that approximately 70 percent of the population
431	should be careful to protect the remaining population, and that as an
432	individual you get some tasks to do to protect someone else than yourself
433	(No. 10).

434	However, appealing to solidarity was considered challenging, because the messages
435	need to be understood by the whole population, including people who are not in a life or work
436	situation where they can participate in the collective effort by keeping physical distance, and
437	people who come from cultural backgrounds that have a different understanding and valuation
438	of collective moral responsibility.
439	3.3.2 Control of the spread
440	Experts from different fields stressed the need to communicate the risk of losing
441	control of the spread. They stated that "while the pandemic risk is low for many individuals, it
442	is high for society". To express the risk to society, the experts reiterated the potential
443	consequences and future scenarios, conveyed as: "The virus is highly contagious. Losing
444	control of the spread of the virus, will lead to healthcare services becoming overloaded which
445	makes the pandemic deadlier".
446	The media often used the reproduction number (R-number) to describe the status of
447	the spread. According to the biostatistician:
448 449 450 451 452 453 454	The R, the reproduction number, is important because it says something about the growth of the disease, so it is the speed with which the disease will grow, the number of infected people will grow. And if this number is above 1, then the epidemic is exploding, if it is below 1, it is under control and will disappear. So, it is a fundamental number that describes the current status, where are we now, today we are 0.8, that is good, last week we were 1.3, not so good, and so on right (No. 11)
455	However, many of the experts decided against mentioning the R-number in their
456	public messaging, because of the risk of misunderstanding it. The R-number was considered
457	too complicated to explain to the public due to the need to understand uncertainties,
458	simplified assumptions, and exponential growth. Many of the experts believed that the R-
459	number could result in a misleading picture of the pandemic.

460	The experts stated that explanations of the risk of losing control should be balanced
461	with information empowering the population to control the spread. Descriptions of realistic
462	risk scenarios needed to be balanced with messages of hope. It was just as important to
463	convey the severity of the risk as it was to avoid creating panic. As the psychologist stated:
464 465 466 467 468 469	If too much fear is triggered, I think some people will defend themselves and refuse to deal with it. You should neither make the message so dangerous that it becomes overwhelming, neither so harmless that people think it does not matter what they do. I think you cannot under- communicate the consequences. You have to get a balance there (No. 5).
470	Lastly, messages implying the risk of losing control needed to be balanced with
471	messages strengthening people's sense of control. As such, the infectious diseases experts did
472	not merely reveal the status of infected and hospitalised people, but also that "we have
473	control" to ensure the public's trust in healthcare delivery. The experts believed in creating
474	situational awareness while offering specific advice on how to minimise their risk.
475	3.3.3 Risk tradeoffs
476	Some experts believed that giving people insight into their dilemmas could improve
477	the public's understanding of management strategies for the pandemic.
478	The experts emphasised the significance of informing the public of the risk tradeoffs
479	between primary and secondary consequences of the pandemic". However, their perspectives
480	on risk differed. The medical experts had low risk acceptance for the negative consequences
481	of lockdown and isolation (e.g., increased mental illness among children and adolescents).
482	Thus, they wanted all measures to be proportionate to the infection risk. Nevertheless, the
483	emergency management expert had low risk acceptance for primary consequences of the
484	pandemic, and thus communicated the importance of preventing the spread of the virus by
485	implementing measures as early as possible:

486	It's very difficult to use strict and invasive measures to prevent infection.
487	You have to wait until the infection has spread. Different professional
488	experts, health and emergency preparedness disagree. We [emergency
489	preparedness specialists] believe it is not about spreading fear. It is about
490	having a proactive approach around a potential hazardous situation in the
491	future. It is difficult to communicate well, because our message is often shot
492	down by the central health authorities (No. 3).
493	
494	The science and infectious diseases expert contended that the risk tradeoff between
495	being infected versus being vaccinated needed to be communicated. This tradeoff was
496	difficult to express, especially at the individual level:
497	To communicate that vaccines are used when a virus is so dangerous that it
498	is worth taking the risk of side-effects to avoid being exposed to the
499	infection is challenging. Where is that threshold? There is no final decision
500	on that When is the intersection where your individual risk should
501	actually take precedence over the best interest of society and herd
502	immunity? It's twisted. I try to communicate the dilemma. I do not have a
502	definitive answer, because there is no definitive answer (No. 13).

504

505 4 Discussion

506 This study uses COVID-19 as a case to explore medicine, science (medical/health research, 507 public health and communication expert' opinions and knowledge of topics to communicate 508 to the public in relation to pandemic health risks. The discussion of the findings is related to 509 the interdisciplinarity of the emerging key topics, communicating uncertainty and messages 510 which may amplify risk, and the role of public engagement along with expertise knowledge in 511 designing messages.

512 4.1 Interdisciplinarity

513 This study has identified three key topics of scientific knowledge in pandemic health

514 risk communication. The experts focused on viral replication and modes of virus

515 transmission, protection from the disease, and the health risk for individuals and the society.

These key topics relate to concepts which originate from different fields and disciplines,
including infectious diseases (e.g., virus transmission, immunity), psychology (e.g., fear,
hope, motivation), philosophy (e.g., collective moral responsibly), statistics (e.g., R-number),
risk communication (e.g., situational awareness, risk tradeoffs), health communication (e.g.,
health literacy), pedagogy (e.g., visual analogies), and marketing (e.g., targeted
communication).
Furthermore, the different experts had different priorities in terms of what messages to

523 convey. This poses a risk for contradictory communication. The multi-disciplinarity of the

524 key topics in this study, and the different views on key topics to communicate, demonstrate

525 that messages should not be developed from one scientific field only, but should rather rely on

526 interdisciplinarity to ensure coherence, clarity, and engagement.

527 4.2 Communicating uncertainty and messages which may amplify risk

528 In this study, the omission of talking about the R-number and airborne transmission in

529 public communication were related to the experts' belief that raising these topics

530 characterised by scientific uncertainty could amplify the risk. However, under those

531 circumstances people may consult other sources of information, which may lead to

532 misinformation and consequently risk amplification (42).

533 Nevertheless, evidence does not support the communication of uncertainty on all

534 occasions. The communication of uncertainty can have a range of outcomes. According to

535 Gustafson and Rices' (15) review of the experimental literature on different types of trust,

- 536 explaining the confidence interval related to the R-number and explaining probabilities
- 537 related to infection prevention measures may increase public trust, as these are related to
- 538 communicating technical uncertainty (15). Nonetheless, communicating uncertainty about

539 airborne transmission at the early phase of the pandemic may undermine public trust, as this 540 type of uncertainty is related to scientific uncertainty, and lack of experts' consensus (15). 541 The experts emphasised being open and transparent and communicating tradeoffs in crisis 542 management as vital to the creation of trust and strengthening people's comprehension of the 543 rationale behind infection prevention measures. Trust is essential during rapidly evolving 544 events characterised by scientific uncertainty (9, 13, 14). The experts reflect that the pandemic 545 measures have both intended and unintended consequences, forcing decision makers to trade 546 short-term policy goals for unintended long-term effects (43). Recent studies have shown that 547 quarantine, staying home and closing schools have implications for unemployment (44), 548 mental health issues (45, 46), interrupted learning (47) and domestic violence (48). Countries 549 strive to balance safety and health with economic security and personal freedom (49). 550 Norheim et al. (21) argue that by explaining these dilemmas, people learn about the 551 complexity of the decision making process, and strengthen social trust. Gregory et al. (20) 552 argue that experts and authorities should be transparent and acknowledge value-based 553 judgements in environmental risk management, rather than pretend that the decision making 554 is objective and non-value-based. Consistent with the literature, this study identifies risk 555 tradeoffs in pandemic health risk communication as a topic to communicate. Tradeoffs can be 556 conveyed by the messenger in their messages through transparency about what is at stake and 557 how the interests of different parties are assessed and weighted. 558 Furthermore, in this study, medical experts emphasised the balance between providing 559 people with facts and instilling fear. The experts protected this balance by suggesting future 560 scenarios and the consequences of uncontrolled virus spread with the aim of creating a 561 realistic picture of the threat while avoiding war-framing and exaggerations. Both social 562 amplification and attenuation of risk undermine the effectiveness of risk communication 563 (10). A moderate level of fear is needed to take the risk seriously and act on it (22). Consistent

with the literature, this study identifies the balance between fear and efficacy as vital when
communicating control of the spread (10, 50). By doing so, health risk communicators may
provide the public with necessary information, without attenuation or amplification of the risk
in correspondence with the SARF (42).

568 4.3 Public engagement

569 This study indicates that experts across professional fields are taking the audience's

570 knowledge (or lack of) and concerns into account. The medical and science experts are

571 mindful of relying clear messages, and of simplifying complex scientific topics and

572 principles. However, medical/health research, public health and medical experts have

573 different preferences from the communication experts in terms of selecting the content of their

574 messages. The medical and research experts, although acknowledging their audience, adopt a

575 one-way approach that consists of identifying topics based on what they consider most

576 important. The communication experts adopt a two-way approach that is intended to fill gaps

577 in the public's knowledge.

578 There are limitations to both approaches. The recipients of communication must regard 579 the message as relevant; this is a drawback of the one-way approach because without knowing 580 their audience experts' communication may be too detailed and detached from public's values 581 (9). The drawback of communication interventions informed by audience analysis only is the 582 omission of topics not emphasised by the target group. Pandemic health risk communication 583 also needs to provide people with information that they do not realise they need to protect 584 themselves and others. Communication created based on audience analysis, without expert 585 opinions may fail to fill knowledge gaps (4).

There is no one-size-fits-all to the communication of health risk during a pandemic.
There will always be unintended effects on some audience members. Both audience analysis

and interdisciplinary collaboration among experts in medicine, public health, science, and

589 communication may facilitate the creation of engaging messages with intended outcomes (4).

590 4.4 Limitations

591 This study uses a normative expert study approach to collect multi-professional domain-592 specific knowledge (25). Analytical generalisations made on a theoretical level can inform 593 frameworks for inter-professional pandemic health risk communication and public health 594 emergencies (51). This study holds strong information power due to the quality of the 595 interview dialogue (52), and the inclusion of participants with relevant experiences with the 596 phenomenon under study (53). However, the phenomenon of pandemic communication is 597 ambiguous and complex, and a larger sample could have revealed an even larger breadth of 598 pandemic communication topics. Future studies are needed to explore pandemic 599 communication in other settings and phases of a pandemic with other fast spreading infectious 600 diseases.

601 5 Conclusion

602 In this study we have explored experts' opinions and knowledge about health-related 603 information topics that need to be shared with the public in a pandemic situation. By looking 604 into the COVID-19 as a case, the experts in our sample revealed diverse key topics 605 representing several disciplinary fields. The experts found it is fundamental to communicate 606 the ways a virus enters the body and generates disease; the measures needed to protect oneself 607 and others; and enabling a wider perspective on the individual and societal risks caused by the 608 pandemic. This study demonstrates that pandemic health risk communication relies on 609 interdisciplinary expertise across medicine, public health, medical/health research, and 610 communication. Experts emphasise not merely what to communicate but understand the 611 message as meaning produced in the intersection of the messenger, the message attributes,

612	and its audience. Thus, what to communicate is highly related to the complexity of
613	communicating messages due to scientific uncertainty, fear of risk amplification,
614	heterogeneity in public health and scientific literacy, among others. Therefore, the study
615	contributes to the knowledge of complex communication processes of pandemic health risk
616	communication.
617	The results from this study can be used to create communication interventions related t
618	pandemic health risk, first of all by emphasising the need for a genuinely interdisciplinary
619	approach to what needs to be communicated. The study contributes to the field of pandemic
620	preparedness and informs effective communication interventions in future pandemic events.
621	Studies of intended audience members are required to identify what the public does and does
622	not know, as advised by the mental models' approach to risk communication (4).

623

624 Acknowledgements

We want to acknowledge the COVCOM user panel for providing feedback on our work inprogress.

627 Research team

628 This study was conducted by an interdisciplinary team of researchers in the Covid

629 Communication project (COVCOM) lead by JR. SHB (PhD, Licenced Clinical Psychologist)

- 630 is an associate professor with a background from psychology and safety science. She is an
- 631 experienced interviewer. MT (PhD) is a postdoctoral researcher with a background from
- 632 visual communication and media studies. JR (PhD) is a professor in medical statistics with a
- 633 background from mass media communication and mathematics. DAL (PhD) is an associate
- 634 professor with a background from experimental research and management studies. HT (PhD)

to

- 635 is dean of research with a background from aesthetics, gender and intermedial studies. SW
- 636 (PhD) is a professor with a background from safety science and risk communication.

637 References

638 Porta M. A dictionary of epidemiology. New York: Oxford university press; 2014. 1. 639 2. Salajan A, Tsolova S, Ciotti M, Suk JE. To what extent does evidence support decision making 640 during infectious disease outbreaks? A scoping literature review. Evidence & Policy: A Journal of 641 Research, Debate and Practice. 2020;16(3):453-75. 642 Paakkari L, Okan O. COVID-19: health literacy is an underestimated problem. The Lancet 3. 643 Public Health. 2020;5(5):e249-e50. 644 de Bruin WB, Bostrom A. Assessing what to address in science communication. Proceedings of 4. 645 the National Academy of Sciences. 2013;110(Supplement 3):14062-8. 646 Antiochou K. Science communication: challenges and dilemmas in the age of COVID-19. 5 647 History and Philosophy of the Life Sciences. 2021;43(3):87. 648 Wong-Parodi G, De Bruin WB. Informing public perceptions about climate change: A 'mental 6. 649 models' approach. Science and engineering ethics. 2017;23(5):1369-86. 650 World Health Organization. Communicating risk in public health emergencies: a WHO 7. 651 guideline for emergency risk communication (ERC) policy and practice. Switzerland: World Health 652 Organization; 2017. 653 8. Berg SH, O'Hara JK, Shortt MT, Thune H, Brønnick KK, Lungu DA, et al. Health authorities' 654 health risk communication with the public during pandemics: a rapid scoping review. BMC Public 655 Health. 2021;21(1):1401. 656 Balog-Way D, McComas K, Besley J. The Evolving Field of Risk Communication. Risk 9. 657 Analysis. 2020;40(S1):2240-62. 658 Kasperson RE, Renn O, Slovic P, Brown HS, Emel J, Goble R, et al. The social amplification of 10. 659 risk: A conceptual framework. Risk analysis. 1988;8(2):177-87. Lungu DA, Røislien J, Wiig S, Shortt MT, Ferrè F, Berg SH, et al. The Role of Recipient 660 11. 661 Characteristics in Health Video Communication Outcomes: Scoping Review. J Med Internet Res. 662 2021;23(12):e30962. 663 Van der Bles AM, Van Der Linden S, Freeman AL, Mitchell J, Galvao AB, Zaval L, et al. 12. 664 Communicating uncertainty about facts, numbers and science. Royal Society open science. 665 2019;6(5):181870. 666 Fancourt D, Steptoe A, Wright L. The Cummings effect: politics, trust, and behaviours during 13. 667 the COVID-19 pandemic. The Lancet. 2020;396(10249):464-5. 668 Plohl N, Musil B. Modeling compliance with COVID-19 prevention guidelines: The critical role 14. 669 of trust in science. Psychology, Health & Medicine. 2021;26(1):1-12. 670 Gustafson A, Rice RE. A review of the effects of uncertainty in public science communication. 15. 671 Public Understanding of Science. 2020;29(6):614-33. 672 16. Hollnagel E. The ETTO principle: efficiency-thoroughness trade-off: why things that go right 673 sometimes go wrong. London: CRC press; 2009. 674 Rasmussen J. Risk management in a dynamic society: a modelling problem. Safety Science. 17. 675 1997;27(2):183-213. 676 18. Reason J. Managing the Risks of organizational accidents. England: Ashgate; 1997. 677 Glette MK, Wiig S. The role of organizational factors in how efficiency-thoroughness trade-offs 19. 678 potentially affect clinical quality dimensions-a review of the literature. International Journal of Health 679 Governance. 2021;26(3):250-65. 680 Gregory R, Failing L, Ohlson D, McDaniels TL. Some pitfalls of an overemphasis on science in 20. 681 environmental risk management decisions. Journal of Risk Research. 2006;9(7):717-35. 682 Norheim OF, Abi-Rached JM, Bright LK, Bærøe K, Ferraz OL, Gloppen S, et al. Difficult trade-21. 683 offs in response to COVID-19: the case for open and inclusive decision making. Nature Medicine. 684 2021;27(1):10-3. 685 Harper CA, Satchell LP, Fido D, Latzman RD. Functional fear predicts public health 22. 686 compliance in the COVID-19 pandemic. International journal of mental health and addiction. 687 2020;19(5):1875-88. 688 Garfin DR, Silver RC, Holman EA. The novel coronavirus (COVID-2019) outbreak: 23. 689 Amplification of public health consequences by media exposure. Health Psychology. 2020;39(5):355-7.

690 Ocloo J, Garfield S, Franklin BD, Dawson S. Exploring the theory, barriers and enablers for 24. 691 patient and public involvement across health, social care and patient safety: a systematic review of 692 reviews. Health Research Policy and Systems. 2021;19(1):8. 693 Morgan MG, Fischhoff B, Bostrom A, Atman CJ. Risk communication: A mental models 25.694 approach: Cambridge University Press; 2002. 695 Austin LC, Fischhoff B. Injury prevention and risk communication: a mental models approach. 26. 696 Injury prevention. 2012;18(2):124-9. 697 Berg SH., MT. S, Thune H., Røislien J., JK. OH, Lungu DA., et al. Comprehending and Acting 27. 698 on Pandemic Health Risk Information Differently: A Qualitative Study Using the Mental Models' 699 Approach. BMC Public Health. 2022 Forthcoming. 700 Røislien J., O'Hara JK., Smeets I., Brønnick K., Berg SH., Shortt MT., et al. Creating Effective, 28 701 Evidence-Based Video Communication of Public Health Science (COVCOM Study): Protocol for a 702 Sequential Mixed Methods Effect Study. JMIR Res Protoc 2022;11(3):e34275. 703 Bruine de Bruin W, Fischhoff B, Brilliant L, Caruso D. Expert judgments of pandemic 20. 704 influenza risks. Global Public Health. 2006:1(2):179-94. 705 Dhar-Chowdhury P, Haque CE, Driedger SM. Dengue Disease Risk Mental Models in the City 30. 706 of Dhaka, Bangladesh: Juxtapositions and Gaps Between the Public and Experts. Risk Analysis. 707 2016;36(5):874-91. 708 Coronavirus Commission. The Norwegian Authorities pandemic management. Report from 31. 709 the Coronavirus Commision. . Kommunal- og moderniseringsdepartementet; 2021. 710 Worldometers. Reported Cases and Deaths by Country or Territory 2021 [Available from: 32. 711 https://www.worldometers.info/coronavirus/#countries. 712 Government.no. The Government is establishing clear quarantine and isolation rules 2020 33. 713 [Available from: https://www.regjeringen.no/en/aktuelt/the-government-is-establishing-clear-714 guarantine-and-isolation-rules/id2693647/. 715 Ministry of Health and Care Services. Emergency preparedness plan for infection mitigation 34. 716 during the COVID-19 pandemic Regieringen.no2020 [Available from: 717 https://www.regjeringen.no/contentassets/73a60433276240bb9247a00ecc8b23c7/beredskapsplan-718 covid-19-oppdatert-desember-2020-hbrj.pdf. 719 Norwegian Medicines Agency. The Norwegian health care system and pharmaceutical system 35. 720 legemiddelverket.no2021 [Available from: https://legemiddelverket.no/english/about-us/the-721 norwegian-health-care-system-and-pharmaceutical-system. 722 OECD.org. Trust in Government OECD2021 [Available from: http://www.oecd.org/gov/trust-36. 723 in-government.htm. 724 Sætrevik B. Realistic expectations and pro-social behavioural intentions to the early phase of 37725 the COVID-19 pandemic in the Norwegian population. Collabra Psychology 2021;7:18698. 726 38. Zickfeld JH, Schubert TW, Herting AK, Grahe J, Faasse K. Correlates of health-protective 727 behavior during the initial days of the COVID-19 outbreak in Norway. Frontiers in psychology. 728 2020;11:564083. 729 Nilsen ACE, Skarpenes O, Coping with COVID-19, Dugnad: a case of the moral premise of the 39. 730 Norwegian welfare state. International Journal of Sociology and Social Policy. 2020. 731 Braun V, Clarke V. Using thematic analysis in psychology. Qualitative Research in Psychology. 40. 732 2006;3(2):77-101. 733 41. Shortt MT, Berg SH, Wiig S, Lungu DA, Smeets I, Thune H, et al. Exploring health experts' 734 and creative communicators' focus in pandemic video communication: A qualitative study. Frontiers 735 in Communication. 2022 Forthcoming. 736 Bearth A, Siegrist M. The Social Amplification of Risk Framework: A Normative Perspective 42. 737 on Trust? Risk Analysis. 2021. 738 Balog-Way DH, McComas KA. COVID-19: Reflections on trust, tradeoffs, and preparedness. 43. 739 Journal of Risk Research. 2020:1-11. 740 Nicola M, Alsafi Z, Sohrabi C, Kerwan A, Al-Jabir A, Iosifidis C, et al. The socio-economic 44. 741 implications of the coronavirus and COVID-19 pandemic: a review. International Journal of Surgery. 742 2020;78:185-93. 743 Xiong J, Lipsitz O, Nasri F, Lui LM, Gill H, Phan L, et al. Impact of COVID-19 pandemic on 45. 744 mental health in the general population: A systematic review. Journal of affective disorders. 745 2020;277:55-64. de Lima CVC, Cândido EL, da Silva JA, Albuquerque LV, de Menezes Soares L, do Nascimento 746 46. 747 MM, et al. Effects of quarantine on mental health of populations affected by Covid-19. Journal of affective disorders. 2020;275:253-4. 748 749 47. Adverse consequences of school closures UNESCO2020 [Available from:

750 <u>https://en.unesco.org/covid19/educationresponse/consequences</u>.

- 751 48. Kofman YB, Garfin DR. Home is not always a haven: The domestic violence crisis amid the
- 752 COVID-19 pandemic. Psychological Trauma: Theory, Research, Practice, and Policy. 2020;12:199–201.
 753 .
- Norheim OF, Abi-Rached JM, Bright LK, Bærøe K, Ferraz OLM, Gloppen S, et al. Difficult
 trade-offs in response to COVID-19: the case for open and inclusive decision making. Nature Medicine.
 2021;27(1):10-3.
- 757 50. Zhang XA, Cozma R. Risk sharing on Twitter: Social amplification and attenuation of risk in 758 the early stages of the COVID-19 pandemic. Computers in Human Behavior. 2022;126:106983.
- Tracy SJ. Qualitative quality: Eight "big-tent" criteria for excellent qualitative research.
 Qualitative inquiry. 2010;16(10):837-51.
- 761 52. Malterud K, Siersma VD, Guassora AD. Sample Size in Qualitative Interview Studies: Guided 762 by Information Power. Qualitative Health Research. 2016;26(13):1753-60.
- Graneheim UH, Lindgren BM, Lundman B. Methodological challenges in qualitative content
 analysis: A discussion paper. Nurse Educ Today. 2017;56:29-34.
- 765
- 766
- 767
- 768 Supporting information
- 769 S1 File Interview guide expert interviews
- 770 S2 File. Key topics animated bubble chart. Animated bubble chart visualising concept
- 771 mapping of expert opinions.