

Transmission of SARS-CoV-2 in standardised First Few X cases and household transmission investigations: a systematic review and meta-analysis

Hannah C Lewis^{1,2*}, Adrian J Marcato^{3*}, Niamh Meagher^{3,4*}, Marta Valenciano^{1,5}, Juan-Pablo Villanueva-Cabezas^{3,6}, Violeta Spirkoska^{3,7}, James E Fielding^{3,4,7}, Amalia Karahalios⁴, Lorenzo Subissi¹, Anthony Nardone^{1,5}, Brianna Cheng^{1,8}, Soatiana Rajatonirina², Joseph Okeibunor², Eman A Aly⁹, Amal Barakat⁹, Pernille Jorgensen¹⁰, Tasnim Azim¹¹, Pushpa Ranjan Wijesinghe¹¹, Linh-Vi Le¹², Angel Rodriguez¹³, Andrea Vicari¹³, Maria Van Kerkhove¹, Jodie McVernon^{3,4,14}, Richard Pebody^{10**}, David J Price^{3,4**}, Isabel Bergeri^{1†**} and the Unity Studies Collaborator Group***

*contributed equally as co-first authors

**contributed equally as co-last authors

† Corresponding author: Isabel Bergeri, email: bergerii@who.int

***Full group author list is listed below

1 World Health Organization, Geneva, Switzerland

2 World Health Organization, Regional Office for Africa, Brazzaville, Congo

3 Department of Infectious Diseases, The University of Melbourne, at the Peter Doherty Institute for Infection and Immunity, Melbourne, Australia

4 Centre for Epidemiology & Biostatistics, Melbourne School of Population & Global Health, The University of Melbourne, Melbourne, Australia

5 Epiconcept, Paris, France

6 The Nossal Institute for Global Health, The University of Melbourne, Melbourne, Australia

7 Victorian Infectious Diseases Reference Laboratory, Royal Melbourne Hospital, at the Peter Doherty Institute for Infection and Immunity, Melbourne, Australia

8 School of Population and Global Health, McGill University, Montreal, Quebec, Canada

9 World Health Organization, Regional Office for the Eastern Mediterranean, Cairo, Egypt

10 World Health Organization Regional Office for Europe, Copenhagen, Denmark

11 World Health Organization, Regional Office for South-East Asia, New Delhi, India

12 World Health Organization, Regional Office for the Western Pacific, Manila, Philippines

13 World Health Organization, Regional Office for the Americas (Pan American Health Organization), Washington DC, United States of America

14 Murdoch Children's Research Institute, Melbourne, Australia

Abstract:

First Few X cases (FFX) investigations and Household transmission investigations (HHTI) are essential epidemiological tools for early characterisation of novel infectious pathogens and their variants. We aimed to estimate the household secondary infection attack rate (hSAR) of SARS-CoV-2 in investigations aligned with the WHO Unity Studies HHTI protocol. We conducted a systematic review and meta-analysis according to PRISMA 2020 guidelines (PROSPERO registration:CRD42021260065).

We searched Medline, Embase, Web of Science, Scopus and medRxiv/bioRxiv for ‘Unity-aligned’ FFX and HHTI published between 1 December 2019 and 26 July 2021. Standardised early results were shared by WHO Unity Studies Collaborators (to 1 October 2021). We used a bespoke tool to assess investigation methodological quality. Values for hSAR and 95% confidence intervals (CIs) were extracted or calculated from crude data. Heterogeneity was assessed by visually inspecting overlap of CIs on forest plots and quantified in meta-analyses.

Of 9988 records retrieved, 80 articles (64 from databases; 16 provided by WHO Unity Studies collaborators) were retained in the systematic review and 62 were included in the primary meta-analysis. hSAR point estimates ranged from 2%–90% (95% prediction interval: 3%–71%; $I^2=99.7\%$); I^2 values remained >99% in subgroup analyses, indicating high, unexplained heterogeneity and leading to a decision not to report pooled hSAR estimates.

The large, unexplained variance in hSAR estimates emphasises the need for improved standardisation in planning, conduct and analysis, and for clear and comprehensive reporting of FFX and HHTIs, to guide evidence-based pandemic preparedness and response efforts for SARS-CoV-2, influenza and future novel respiratory viruses.

Key words: systematic review, meta-analysis, COVID-19, SARS-CoV-2, influenza, pandemic, epidemics, transmission, household, epidemiology, infectious disease, epidemiological studies, investigation.

Introduction:

Coronavirus disease 2019 (COVID-19), caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection, was first reported in Wuhan, China, in December 2019. It was declared a Public Health Emergency of International Concern by the World Health Organization (WHO) on 30 January 2020 and characterised as a pandemic in March 2020.(1)

Since the early stages of the COVID-19 pandemic, households have been a major setting of SARS-CoV-2 transmission.(2) As such, household transmission investigations (HTIs) – including studies of the First Few X cases (FFX) in household settings – provide an opportunity to explore transmission dynamics and severity of SARS-CoV-2. HTIs facilitate the collection of important epidemiological, clinical, and virological data in well-defined closed settings, where household contacts can be more accurately ascertained and followed up to identify infector-infectee pairs than in the general population including both virological and serological evidence of infection. Conducting HTIs during the early stages of emergence of a novel respiratory pathogen provides an opportunity to swiftly characterise the transmissibility and severity of the pathogen. These estimates of key epidemiological parameters are crucial to understand and predict the health impacts, inform policy and public health interventions at the local and national level.

A systematic review and meta-analysis of within household transmission for 2009 pandemic influenza A(H1N1) found that estimates for household secondary infection rate/risk are highly heterogeneous.(3) Heterogeneity in included HTIs was at least in part attributed to varying household definitions, secondary case ascertainment and testing methods, and duration of follow up. This led to the recommendation for the development of a unified approach for such investigations. Following a review of the global response to the 2009 A(H1N1) pandemic to improve future preparedness in 2012 (4), these recommendations were actualised through the global Consortium for the Standardization of Influenza Seroepidemiology (CONSIDE) (5) and the development of a suite of standardised early investigation protocols by WHO's Global Influenza Programme and specifically the Influenza Pandemic Special Investigations and Studies (IPSS).(6) These protocols have been further adapted for other high threat respiratory pathogens, such as Middle East Respiratory Syndrome Coronavirus (MERS-CoV).(7)

With the emergence of SARS-CoV-2, the WHO quickly adapted and further developed the suite of standardised protocols for rapid use. These early investigations were re-branded as the Unity Studies (8), and included a FFX investigation protocol (9) and a HTI protocol.(10) The protocols included within the Unity Studies standardise methods to encourage rapid generation of local data for public health action and to facilitate comparison of key epidemiological parameters, such as pathogen transmissibility and severity, across regions and globally. They are intended to be adapted to enable countries to conduct local investigations irrespective of income status and resource level of the setting.

We aimed to systematically review, meta-analyse and summarise available data from standardised FFX and HTIs aligned with the objectives and methods of the WHO Unity Studies HTI protocol for SARS-CoV-2 in order to: 1) describe the implementation of investigations in context 2) assess methodological quality of aligned investigations; 3) calculate a pooled estimate of SARS-CoV-2 household secondary infection and clinical attack rate and; 4) explore sources of heterogeneity in the household secondary infection and clinical attack rates. The data from these analyses will enhance overall understanding of local, regional and international epidemiology of COVID-19 and inform future development of the HTI protocol and implementation.

Methods:

The protocol for the systematic review was registered on PROSPERO on 5 August 2021 (registration number: CRD42021260065) and is reported according to the PRISMA 2020 guidelines.(11)

Definitions

We defined Unity-aligned epidemiological SARS-CoV-2 HHTIs as investigations of index cases and all of their household contacts with longitudinal and prospective collection of epidemiological, virological and/or serological data for subsequent analysis.(10) Retrospective investigations were considered aligned where the original data source pertained to contact tracing investigations with active follow up of all household members of an index case.

Index cases were defined as the first case(s) of COVID-19 identified from a positive reverse transcription polymerase chain reaction (RT-PCR) result that subsequently triggered the recruitment of their household to the HHTI.

The WHO's HHTI protocol defines a household as "[...] a group of people (two or more) living in the same residence [...]".(10) In practice, the definition may vary across regions or when other aspects of livelihood are considered, such as income and the collective consumption of goods and services.(12) We classified individuals who lived with index cases as 'household contacts'. Only investigations with sufficient detail to characterise infection status of all household contacts were included in the systematic review and meta-analyses.

Our primary outcome, the household secondary infection attack rate (hereafter 'hSAR') was defined as the probability of a COVID-19 infection amongst susceptible household members exposed to the primary case detected using laboratory diagnostic tools (either RT-PCR or serology).(13) The household secondary clinical attack rate (hereafter 'hSCAR') was our secondary outcome and defined as above using only clinical criteria. As is often the case in HHTIs, it was not always possible to distinguish between the index and primary cases within households. As a result of these definitions, we henceforth refer to primary or index cases as 'index cases'.

Search strategy

Three sources of data were used to identify records in our systematic review. First, four databases – MEDLINE, Embase, Web of Science and Scopus, were explored to identify relevant investigations published between 1 December 2019 and 26 July 2021. The search strategy sought to identify all combinations of "COVID-19" and terms relevant to HHTIs: "COVID-19" AND ("household transmission" OR "secondary attack rate" OR "close contact" OR "contact transmission" OR "contact attack rate" OR "family transmission"). The full search strategy is detailed in the Appendix.

Second, using the same search terms and dates, we screened unpublished investigations made available on the medRxiv and bioRxiv preprint servers. Finally, we screened results from FFX and HHTIs not yet available in the literature but that had been shared by Unity Studies Collaborators with WHO prior to 1 October 2021. We facilitated the registration of these investigations to the data repository "Zenodo", with the permission of the Principal Investigators.(14)

Eligibility criteria

Records were eligible for this review where the investigation:

1. was aligned to the WHO Unity Studies FFX and HHT investigation protocols, with longitudinally collected data and active follow up of households with a RT-PCR confirmed index case.
2. reported at least five households recruited following the identification of an index case confirmed via RT-PCR.
3. reported the hSAR and/or hSCAR or provided sufficient data to calculate it (i.e., total number of secondary cases among household contacts and the total number of household contacts) with a measure of uncertainty.
4. was published in English, Chinese, French, Russian, Spanish, Portuguese, German or Italian.
5. represented an original investigation and the reported estimates are from the most complete dataset available.

Note that the initial search did not identify any records in languages other than those listed above.

Screening and selection of articles

Records were imported into Covidence for consolidation, de-duplication and storage.⁽¹⁵⁾ Initially, records were screened by title and abstract according to the eligibility criteria. Screening was performed by at least two independent reviewers (AJM, NM, JPV-C, VS, JEF) who were blind to the other reviewer's assessment. A third independent reviewer (DJP) assessed records for eligibility where consensus could not be reached. Records in languages other than English, or where eligibility was unclear from initial screening, were retained for translation or full text-assessment.

Using the same methods for the abstract screening, we conducted full text screening to determine the final eligibility of investigations. All published records and results shared by WHO Unity Studies collaborators retained for the systematic review are henceforth referred to as investigations. Where two or more investigations reported estimates based on the same dataset, the most complete and relevant record was selected. Investigations that: 1) did not report sufficient information (e.g., full text not available, conference abstract); 2) were secondary analyses of a previously included investigation or; 3) were investigations that included a significant proportion of vaccinated index cases or household contacts, were excluded.

Methodological quality assessment

We used a flexible, fit-for-purpose critical appraisal tool that consisted of 12 items to assess the methodological quality of investigations.⁽¹⁶⁾ Briefly, the tool was developed to assess specific aspects of HHTI design, which combines features of case series and longitudinal studies. The tool builds upon well-established approaches to perform critical appraisal and risk of bias assessment of observational studies.^(17, 18) Five team members (AJM, NM, JPV-C, VS, JEF) independently applied the methodological quality assessment checklist (two members per assessment) for the hSAR and hSCAR outcomes, and responses were recorded as Yes/No/Unclear. All questions were used for the overall assessment. In particular, questions 3 (was a definition of 'household' provided?), 5 (were subsequent cases identified and ascertained using appropriate methods to calculate the hSAR and hSCAR?) and 6A

(were households followed up for a period sufficient to measure primary outcomes?) were highly relevant in our methodological quality assessment to understand how aligned investigations were to the WHO Unity HHTI protocol and to indicate their suitability to produce aggregated estimates of our outcome measures, as defined above. Investigations were classified as having low, moderate or high risk of bias for each outcome according to their methodological quality, i.e., investigations with lower methodological quality were more likely to have higher risk of bias. Where consensus about the methodological quality assessment could not be reached, a sixth reviewer (DJP/AK) finalised the assessment.

High risk of bias was generally attributed to unsuitable study design, analysis methods or results, or unclear what design or analysis methods were used. These included, but were not limited to, unclear or broad definition of “household” and “household contacts”, unclear or unsuitable methods of secondary case ascertainment, unclear or insufficient follow up duration, reduced exposure of household contacts to primary/index cases or a combination of these.

Data extraction

We extracted data relevant to the primary and secondary objectives of our study, in addition to administrative information relating to each investigation (author, country). We further contextualised each investigation by WHO region (19), income status as reported by the World Bank in 2021 (20), and involvement in the United Nations Office for the Coordination of Humanitarian Affairs (OCHA) Humanitarian Response Plan (HRP) for COVID-19.(21)

We also collected data on: total index cases, total households, total household contacts, total secondary cases, dates of investigation, household transmission design (i.e., was the primary objective of the study to characterise household transmission?), method of secondary case ascertainment (e.g., RT-PCR or serology, routine or symptom-based testing) and data collection methods (i.e., retrospective or prospective).

Data were extracted by two reviewers independently (AJM, NM, JPV-C, VS, JEF). Discrepancies in the data extraction were resolved by discussion with all participating reviewers. Where the reporting of estimates was unclear, incomplete, or not explicit, authors were emailed at least twice in November and December 2021 to confirm details. Responses were collated until 17 January 2022. Investigations were excluded if authors did not respond and sufficient data were not available. No restriction was put on the number of index cases per household. The number of households was assumed to be equal to the number of index cases if either was not reported or available upon follow up, where the study design suggested that this was a reasonable assumption.

Statistical analysis

Values for SARS-CoV-2 hSAR, hSCAR and associated 95% confidence intervals (CIs) were extracted or calculated from crude data.

We conducted a meta-analysis to obtain a global estimate of SARS-CoV-2 hSAR. The primary meta-analysis only considered investigations assessed as having low or moderate risk of bias. Forest plots were produced to illustrate the hSAR and hSCAR estimates of included investigations. These are presented overall and by subgroups of interest as described below.

Heterogeneity was assessed by visually inspecting the overlap of the CIs on the forest plots, quantified in the meta-analysis using the I^2 and $\widehat{\tau}^2$ statistics, and assessed using the χ^2 test for heterogeneity.(22) We fit a binomial-normal model to pool the hSAR data (primary outcome) and the hSCAR data (secondary outcome). This model has a binomial within-study likelihood and a normal across-study likelihood, which has been demonstrated to produce unbiased estimates and accounts for variation between investigations.(23) The I^2 and $\widehat{\tau}^2$ measures of heterogeneity were presented for each model to indicate the percentage of variation across investigations attributable to heterogeneity and the estimated between-investigation variance, respectively. The p-value from the χ^2 test for heterogeneity was also presented.

We further explored SARS-CoV-2 hSAR and hSCAR heterogeneity in pre-specified subgroup analyses. Subgroups included: 1) income setting (high income vs. low- and middle-income status) according to the World Bank classification (20); 2) predominantly circulating variant at the time the investigation was conducted (variant of concern vs. other strains of SARS-CoV-2) according to data available from GISAID (24) accessed via covariants.org (25) where not reported in the investigations, and; 3) secondary case ascertainment methods (serological or RT-PCR testing of all household contacts vs. testing of symptomatic household contacts only). We also undertook a subgroup analysis of hSAR using the results from our methodological quality assessment (low or moderate risk of bias vs. high risk of bias).

Following review of forest plots and meta-analysis results, 1) three post-hoc subgroups were defined and assessed in subgroup analysis and; 2) a post-hoc decision was made to report the 95% prediction interval for the primary analysis to further demonstrate the investigation-level heterogeneity. The first post-hoc subgroup analysis examined duration of follow up of household contacts (14 days or less vs. greater than 14 days). The second compared investigations that did and did not comply with a stricter definition of adherence and alignment to the methods and objectives outlined in the Unity protocol. Strict methodological adherence was defined as a specifically designed household investigation which had prospective follow up and routine testing of all household contacts irrespective of symptoms. For example, investigations that only tested household members with symptoms would not comply with this stricter methodological definition. The third subgroup compared investigations that used RT-PCR testing to those that used RT-PCR in combination with serological testing to ascertain secondary cases.

We also planned to visually assess funnel plots of effect size versus standard error and to use Egger's test to evaluate bias due to small study effects.(26) Additionally, we planned to conduct a sensitivity analysis by fitting a beta-binomial model as applied in Thompson et al. to explore whether the chosen model impacted the results.(27)

Data cleaning and collation was performed using Stata version 16 and R version 4.0.(28, 29). All pooled meta-analyses were undertaken in R version 4.0 using the *metafor* package.(30)

Results:

Characteristics of included investigations

Figure 1 summarises the literature search and screening process. We identified 9,954 published records from database searches, and results from 34 FFX and HHTIs were provided directly from WHO Unity Studies Collaborators. Following removal of duplicates, 6,536 records went through title and abstract screening, and 284 subsequently underwent full-text assessment. A further 204 records were excluded at the full-text stage for various reasons (Figure 1). In total, 80 investigations (64 from database searches, and 16 provided by WHO Unity Studies Collaborators) were retained for data extraction, of which 62 (51 from database searches and 11 provided by WHO Unity Studies collaborators) were included in the primary meta-analyses.




Figure 1. PRISMA flow chart. Other reasons for exclusion during full-text screening were: the full text was not accessible; duplicate investigations with different indexing; commentaries; corrections; non-COVID related investigations, and conference abstracts, preprints and short reports with subsequent publications. Abbreviations: FFX, First Few X cases investigations; HHTI, Household Transmission Investigation; WHO, World Health Organization.

Table 1 provides summary characteristics of the 80 investigations that met the eligibility criteria for the systematic review and for the subgroup of 62 articles that were included in the meta-analyses - detailed characteristics of each investigation can be found in the Appendix (Supplementary Table 1). Included investigations for the primary outcome of hSAR described follow up of households between January 2020 and July 2021, with the majority ($n = 50$) completed before July 2020 (Supplementary Figure 1). Two investigations did not report a start date. Thirty-two countries were represented across all six WHO regions — fifteen countries contributed more than one investigation to this review, including 20 articles

from China, 8 from India, 7 from the United States of America, 4 from South Korea, and 4 from Canada. Fifteen lower-and-middle income countries (LMICs) accounted for 51% (41/80) of the investigations identified in the systematic review, with 78% (32/41) of these being included in the meta-analyses. Nine investigations (11%) were conducted in countries supported in the OCHA HRP for COVID-19. Correspondingly, 17 high-income countries (HICs) accounted for 49% (39/80) of the investigations in the systematic review, with 77% (30/39) of these included in the meta-analyses.

Of the 76 investigations that reported a hSAR, secondary infections were predominantly ascertained by either scheduled RT-PCR testing of all contacts irrespective of symptoms ($n = 43$), symptom-initiated RT-PCR testing ($n = 9$), serology testing alone ($n = 3$), or through use of RT-PCR in combination with serology testing ($n = 11$). In the remaining 10 investigations, the method of secondary case ascertainment was unclear — these 10 investigations were excluded from the corresponding subgroup analysis. Secondary symptomatic cases were determined by symptoms alone in all instances where hSCAR was reported ($n = 33$).

Forty-three investigations (54%) were specifically designed as HHTIs. The remaining were investigations of all close contacts that reported sufficient detail from which we could estimate hSAR and/or hSCAR. The majority of articles collected data prospectively ($n = 55$), although some involved retrospective collation of detailed contact-tracing data from which households could be reconstructed ($n = 23$).

Table 1. Summary statistics corresponding to all investigations included in the primary meta-analysis ($n = 62$) and in the systematic review ($n = 80$). Median and interquartile range [IQR, 25–75th percentiles] is reported for numeric quantities, and; number (%) is reported for categorical variables.

	Characteristic, Median [IQR] or N (%)	
	All investigations included in meta-analyses N = 62	All investigations included in the systematic review N = 80
Number of households included	100.5 [44.5, 213.5]	100 [38.75, 210.5]
Household of contacts included	286.5 [174.5, 768]	279 [154, 792.75]
Total index cases	103.5 [60.75, 259.75]	102 [53.5, 229.5]
Secondary cases	69.5 [37.75, 138.5]	71 [34.75, 140.25]
Source of article		
Peer-reviewed journal investigation – identified in literature search	48 (77.4%)	58 (72.5%)
Preprints – identified in medRxiv and bioRxiv literature search	3 (4.8%)	6 (7.5%)
Results provided directly from WHO Unity Studies collaborators†	11 (17.8%)	16 (20.0%)

WHO Region†		
AFR	3 (4.8%)	6 (7.5%)
EMR	1 (1.6%)	2 (2.5%)
EUR	13 (21.0%)	18 (22.5%)
AMR	9 (14.5%)	13 (16.3%)
SEAR	7 (11.3%)	10 (12.5%)
WPR	29 (46.8%)	31 (38.8%)
Income Status§		
High	30 (48.4%)	39 (48.8%)
Upper-Middle	20 (32.3%)	23 (28.8%)
Lower-Middle	9 (14.5%)	13 (16.3%)
Low	3 (4.8%)	5 (6.3%)
Humanitarian Response Plan (HRP) for COVID-19 Status¶		
No	58 (93.5%)	71 (88.8%)
Yes	4 (6.5%)	9 (11.3%)
Presumed Dominant Strain††		
Variant of Concern	4 (6.5%)	6 (7.5%)
Other Strain	58 (93.5%)	74 (92.5%)
HHTI design		
Yes	35 (56.5%)	43 (53.8%)
No (FFX, Case Series, Surveillance)	27 (43.5%)	37 (46.3%)
Method of Secondary Case Ascertainment (for hSAR)‡‡		
RT-PCR (irrespective of symptoms)	34 (54.8%)	43 (53.8%)
RT-PCR (symptom based)	8 (12.9%)	9 (11.3%)
RT-PCR or serology	12 (19.4%)	12 (15.0%)
Serology	-	3 (3.8%)
Unknown method of lab diagnosis	8 (12.9%)	9 (11.3%)
Did not report a hSAR	-	4 (5.0%)

Method of Secondary Case Ascertainment (for hSCAR)§§		
Symptoms only	24 (38.7%)	33 (41.3%)
Did not report a hSCAR	38 (61.3%)	47 (58.8%)
Follow up of households		
Prospective	44 (71.0%)	55 (68.8%)
Retrospective	17 (27.4%)	23 (28.8%)
Unclear	1 (1.6%)	2 (2.5%)
Strict adherence and alignment to the Unity Study protocol¶¶		
No	43 (69.4%)	58 (72.5%)
Yes	19 (30.6%)	22 (27.5%)
Duration of follow up		
14 days or less	47 (75.8%)	58 (72.5%)
Greater than 14 days	11 (17.7%)	14 (17.5%)
Unclear	4 (6.5%)	8 (10.0%)
Risk of bias in investigations reporting hSAR		
Low	24 (38.7%)	25 (31.3%)
Moderate	38 (61.3%)	38 (47.5%)
High	-	13 (16.3%)
N/A – hSAR not reported	-	4 (5.0%)
Risk of bias in investigations reporting hSCAR		
Low	21 (33.9%)	22 (27.5%)
Moderate	3 (4.8%)	11 (13.8%)
N/A – hSCAR not reported	38 (61.3%)	47 (58.8%)
<p>† Include peer-reviewed literature and early results now available on Zenodo.(14)</p> <p>‡ WHO Region (19): AFR (Africa), AMR (Americas), EMR (Eastern Mediterranean), EUR (Europe), SEAR (South East Asia), WPR (Western Pacific).</p> <p>§ Income status as reported by the World Bank in 2021.(20)</p> <p>¶ Involvement in the United Nations Office for the Coordination of Humanitarian Affairs (OCHA) Humanitarian Response Plan (HRP) for COVID-19.(21)</p> <p>†† CoVariants (CoVariants: SARS-CoV-2 Mutations and Variants of Interest. (GISAID https://www.gisaid.org/ (24) and https://covariants.org/ (25)).</p> <p>‡‡ hSAR (household secondary infection attack rate).</p> <p>§§ hSCAR (household secondary clinical attack rate).</p> <p>¶¶ Articles were defined based on strict adherence and alignment to the WHO Unity Studies HHTI protocol, including: (i) testing of all contacts (vs. only symptomatic), (ii) prospective data collection, (iii) specifically designed HH investigation (vs. FFX or surveillance data).</p>		

Methodological quality assessment

The assessment of the methodological quality of the included investigations is summarised in Table 1 and detailed in the Appendix (Supplementary Figure 2 and 3). In total, 76 investigations reported a hSAR; 25 were considered to have a low risk of bias, 38 a moderate risk of bias, and 13 a high risk of bias. Of the 33 investigations reporting hSCARs, 22 were considered to have a low risk of bias and 11 had a moderate risk of bias.

Household Secondary Attack Rate

Investigations that did not report a hSAR ($n = 4$), had a high risk of bias ($n = 13$) or were conducted in highly vaccinated cohorts ($n = 1$) were excluded from the main hSAR meta-analyses. In total, 18 investigations were excluded from the primary meta-analysis.

Point estimates for the hSAR ($n = 62$) ranged from 2% – 90% with a 95% prediction interval from 3% – 71% (Figure 2) and $I^2 = 99.7\%$, suggesting substantial heterogeneity between included investigations. The meta-analyses showed the heterogeneity between investigations was not reduced when examining the subgroups of interest (Table 2, Supplementary Table 2) — including by income setting; predominant circulating strain; testing protocol for household contacts, and; risk of bias. In all analyses, hSAR estimates varied substantially and I^2 values were $>99\%$. Due to this large amount of heterogeneity, pooled estimates of the hSAR are not reported. As a result of not producing any pooled estimates, evaluation of bias due to small study effect and sensitivity analysis to assess the effect of model choice were not carried out (Appendix).

Supplementary Figures 5 – 11 show the forest plots by pre-specified and post-hoc subgroups (including adherence to the Unity protocol; duration of follow up, and; use of serology to ascertain secondary cases), and Supplementary Table 2 examines the secondary meta-analysis to assess the effect of including high risk of bias investigations. All results indicate substantial heterogeneity. Supplementary Figure 4 shows a forest plot of hSAR estimates by WHO region, which was not examined in a subgroup meta-analysis due to correlation with income status.

Forest plots and meta-analysis of hSCAR are shown in Supplementary Figures 12 – 14 and Supplementary Table 3. As with the hSAR data, a high amount of heterogeneity between investigations was evident, which was not explained by any of the subgroups investigated. Consequently, pooled estimates of hSCAR were not produced.

Table 2. Results from meta-analyses of household secondary infection attack rate (hSAR). I^2 and $\widehat{\tau^2}$ are presented for each model to indicate the percentage of variation across investigations attributable to heterogeneity and the estimated between-investigation variance, respectively. The p-value from the χ^2 test for heterogeneity is also presented.

	No. investigations	I ²	$\widehat{\tau}^2$	P-value
Infection household secondary attack rate	62	99.7	1.190	<0.0001
Pre-Specified Subgroup Analyses				
Income Status†	62	99.6	1.188	<0.0001
<i>High income</i>	30			
<i>Low- and middle- income</i>	32			
Predominant circulating strain‡	62	99.6	1.130	<0.0001
<i>Other strain</i>	58			
<i>Variant of concern</i>	4			
Testing protocol for household contacts	54	99.4	1.174	<0.0001
<i>Testing of all contacts</i>	46			
<i>Testing of symptomatic contacts</i>	8			
Post-Hoc Subgroup Analyses				
Strict adherence and alignment to the WHO Unity Studies protocol	54	99.5	0.797	<0.0001
<i>Aligned</i>	43			
<i>Not aligned</i>	19			
Duration of follow up of household contacts	58	99.6	1.005	<0.0001
<i>14 days or less</i>	47			
<i>Greater than 14 days</i>	11			
Laboratory diagnosis method	54	99.3	1.030	<0.0001
<i>RT-PCR only</i>	42			
<i>RT-PCR and serology</i>	12			

† Income status as reported by the World Bank in 2021.(20)

‡ CoVariants (CoVariants: SARS-CoV-2 Mutations and Variants of Interest. (GISAID <https://www.gisaid.org/> (24) and <https://covariants.org/> (25)).




Figure 2: Forest plot of the household secondary infection attack rates (hSAR) in included investigations ($n = 62$), ordered from highest estimated hSAR (top) to lowest estimated hSAR (bottom). The hSAR and 95% confidence intervals (CI) are shown on the right margin.

Discussion

This is the first systematic review and meta-analysis of standardised FFX and HHTIs – aligned with the objectives and methods of the WHO Unity Studies HHTI protocol – for SARS-CoV-2. We identified 80 investigations from 32 countries, 51% (41/80) of which were from 15 LMICs, showing that implementation was feasible in a range of settings and in every WHO region. Although all investigations were classified as aligned with the objectives and methodology of the WHO Unity Studies protocol based on the information provided, we found substantial heterogeneity in the reported secondary attack rates (hSAR range: 2% – 90%; 95% prediction interval [PI]: 3% – 71%; $I^2 = 99.7$). As a result, reporting of pooled estimates of hSAR or hSCAR was not deemed appropriate. Further subgroup analyses were undertaken to understand this heterogeneity, yet a similar extent of heterogeneity (I^2 values were >99%) was still observed – consequently, pooled estimates are not provided.

The high degree of heterogeneity and wide range of estimates in reported hSAR is consistent with those of other recent systematic reviews and meta-analyses: one study reported an I^2 value of 99.4% with hSARs ranging from 0% – 74% (31) and another reported an I^2 of 97.4%, with hSARs ranging from 4% – 55% (32). Nevertheless, these reviews reported pooled hSAR estimates of 18.9% (95% CI: 16.2% – 22.0%) (31), 21.1% (95% CI: 17.4% – 24.8%) (27) and 18.1% (95% CI: 15.7% – 20.6%) (32).

In the context of quantifying transmission, relevant potential sources of heterogeneity include methodological differences such as study design and definition of household, as well as differences in study context, such as current public health and social measures (PHSM), and population or individual behaviours. The I^2 value estimates the proportion of the variance in reported estimates that is due to heterogeneity (33), and should not be interpreted in isolation. When conducting meta-analyses, sources of heterogeneity should be identified *a priori* and explored in subgroup analyses where sufficient data have been reported (34). For example, Thompson (27) identified a substantial difference in household attack rates when considering the reported duration of exposure between contact and case. This emphasises the need to precisely report the study design, epidemic context, and household dynamics, in order to meaningfully quantify transmission. Where available, we extracted information on these potential sources of heterogeneity, however sufficient detail on all aspects of the study design and implementation were not routinely or consistently reported. This highlights the importance of further standardising and strengthening the quality of reporting.

Visual inspection of the forest plots suggests potentially lower hSAR in the Western Pacific Region (WPR) compared to other regions (Supplementary Figure 4). The majority of investigations conducted in the WPR were conducted during the first half of 2020. Several countries in the WPR enacted stringent PHSM and behavioural responses in early 2020, which resulted in low SARS-CoV-2 circulation.(31, 35) Furthermore, household size and structure (e.g., high-rise apartments, high- vs low-density households) varies between regions and urban-rural localities, as well as socio-economic status of different subpopulations which may all contribute to lower observed hSAR.(36)

The timing of each investigation must be considered in relation to local epidemic activity and evolving PHSMs when interpreting the results reported in this meta-analysis (Supplementary Figure 1), however, these details were not sufficiently reported in included investigations. Most included investigations ($n = 48$) were finalised in the first six months of 2020, likely during circulation of the ancestral virus and early SARS-CoV-2 variants, i.e., prior to the designation of Alpha and Delta as Variants of Concern (37) (Appendix, Supplementary Figure 1), which are known to have increased transmissibility (38, 39). Only

six investigations in the systematic review were conducted during a period when Alpha and/or Delta variance of concern were the dominant (or equally dominant) circulating strains (24, 25). Among the six studies conducted after the designation of Alpha and Delta, two were excluded from the meta-analysis and three were conducted in the same country — providing insufficient representation for a subgroup analysis.

The frequency and type of specimen collection as well as the duration of follow up and laboratory methods employed can influence the hSAR estimate. In our review, we observed high variability in the laboratory methods used to ascertain secondary cases, including the use of RT-PCR, serology or both. However, we found no difference in observed variability in post-hoc subgroup analyses. We also explored heterogeneity in hSAR estimates by duration of follow up and whether all household contacts were tested to determine the influence of these study design aspects. Both of these analyses showed high variability across investigations. In addition, some investigations did not exclude non-susceptible individuals who tested positive by serology at baseline which may underestimate hSAR, although this information was inconsistently reported across investigations.

To assess the methodological quality of HHTIs, previous reviews adapted existing appraisal tools (17, 18) — however, these tools are limited as they were not designed for this purpose. We conducted a robust and thorough assessment of the methodological quality of investigations using a bespoke quality assessment tool for HHTIs which allowed for a more targeted critical appraisal and better understanding of strengths and limitations of HHTIs.(16) We only included investigations with a low or moderate risk of bias in our primary meta-analysis (n = 62) and subsequently only those aligned with the objectives of the WHO Unity Studies HHTI protocol. It also allowed us to carefully consider the appropriateness of pooling data across investigations that were conducted in different resource settings and environments, using different protocols and with differing internal and external validity.

The use of a bespoke tool for methodological quality assessment does not guarantee that the intricacies of HHTI designs are fully captured, particularly where insufficient details are reported. We acknowledge that tailoring of the HHTI protocol according to cultural norms (e.g., household definition), capacity (e.g., laboratory testing, degree of follow up) and context (e.g., PHSM, local incidence, quarantine practices) may be required, increasing the true variance observed across investigations. As a result, unclear reporting may have inflated the heterogeneity in our review due to inappropriate inclusion or exclusion of some investigations.

Of the 76 investigations that reported a hSAR, 13 were assessed to be at high risk of bias and subsequently excluded from the primary analyses. The assessment of methodological quality was strongly influenced by a range of factors, including: unclear or broad definition of household or household contacts (e.g., those that included more than residential contacts); unclear, unsuitable or incomplete laboratory and follow up methods in HHTIs (e.g., symptom based testing), and management of index cases that reduced exposure of household contacts to index cases (e.g., isolating index cases either within or outside the household). This demonstrates the necessity for high-quality, standardised investigations and detailed reporting of study design and household dynamics, in order to meaningfully quantify transmission.

Although the above methodological and contextual factors were deemed plausible sources of heterogeneity, none could explain a substantial amount of the variance in reported hSARs. Greater emphasis on clear, standardised and comprehensive reporting of HHTIs, and the context in which they are conducted, is required to facilitate more nuanced analysis of the sources of heterogeneity. Further,

in the presence of substantial contextual differences, we question the suitability of providing single pooled estimates of such pathogen characteristics. The first step in any meta-analysis is to consider whether the studies are all estimating the same quantity, and whether they should be pooled. Here, we have identified various sources of heterogeneity across settings which suggests they should not be pooled. While current reporting practices do not allow us to interrogate these heterogeneities further, future investigations based on consistent reporting guidelines may allow for more nuanced analyses where results can be pooled at an appropriate scale. In contrast, within a given setting, many of the identified sources of heterogeneity are likely to be consistent across investigations (e.g., household structure), or, of sufficient relevance to the interpretation that they should be reported alongside investigations repeated over time (e.g., prior infection or vaccination histories). In these cases, hSAR estimates within a given population, particularly over time, can provide crucial insight into key characteristics of a novel pathogen (e.g., changes in transmissibility of SARS-CoV-2 variants) to guide both national (40) and international (41) pandemic response and preparedness efforts.

While HHTIs from LMICs are underrepresented in the literature, this review included results from 41 investigations, of which 11 (27%) were shared by WHO Unity Studies collaborators at the end of February 2022 prior to peer review or pre-print publication. Such data would not typically be available for inclusion in a systematic review and meta-analysis. Enabled by the Unity Studies, the inclusion of these aggregate standardised results from LMICs reduces publication bias towards higher income countries. This highlights the importance of open data practices and sharing early aggregate results for the collation and analysis of timely data, particularly during public health emergencies (42). As vaccination coverage increases across the globe in high and upper-middle income countries, monitoring the transmission dynamics of COVID-19 in LMIC settings, where vaccination coverage is often still low, will be critical to overcome the pandemic.(43)

Our search strategy was highly sensitive and robust. We used broad search terms and language inclusion to capture as many relevant investigations as possible. However, we may have missed relevant investigations as we did not screen the reference lists of included investigations or other systematic reviews of hSAR. A further strength of this systematic review is that we attempted to clarify or confirm unclear or poorly reported items and contacted study investigators to request additional information.

In the future, we recommend focusing on the design of country-specific pre-planned (“at the ready”) standardised FFX and HHTIs with quality implementation by multi-disciplinary teams. These should be conducted in representative settings with available capacity, or, through collaborations to develop local capacity where it is otherwise limited. This would allow the greatest opportunity to rapidly enact FFX and HHTIs during the early stages of the emergence of novel pathogens, especially respiratory viruses of pandemic potential like influenza and coronaviruses, allowing early characterisation of transmissibility and infection-severity to inform public health responses.(44) Specific consideration should be given to prior ethics approval, governance, data collection methods and infrastructure, and resource requirements in advance of activation during a public health emergency.(45) In addition to common protocols, tools to assist quality implementation (e.g. standard operating procedures for sample collection and publicly available data analysis scripts) and dissemination of results (e.g. scientific writing skill development) are required. Furthermore, although short-term assistance can support ad hoc investigations when required, initiatives to build and strengthen surveillance and laboratory capacity in LMICs – such as the WHO’s Pandemic Influenza Preparedness Framework – are a much more sustainable approach.(46, 47) Through long-term investment and development, LMICs can enhance surveillance and implement the operational research required to monitor co-circulating influenza and SARS-CoV-2 viruses in near-real-time, and detect the emergence of novel respiratory viruses of pandemic potential. In-

country capacity building should be prioritised to ensure fit-for-purpose analytic methods for producing robust statistical inferences are implemented, such as those developed specifically for analysing household infection dynamics.(48) We further recommend development of HHTI-specific guidelines and checklists for reporting, such as those developed for clinical trials (49) or observational studies (50, 51), and those introduced in the HHTI critical appraisal tool (16).

Conclusion

As the COVID-19 pandemic progresses, FFX and HHTIs remain a critical tool to monitor population immunity to, and transmission dynamics and infection-severity of SARS-CoV-2, including the emergence of new genetic variants. These data are crucial to inform the ongoing response in different resource settings. Indeed, such estimates are key for regional and global modelling and forecasting to inform optimal application of PHSM and allocate pandemic resources including COVID-19 vaccines. The Unity Studies framework, strengthened through lessons learned from the COVID-19 pandemic, has the potential to form the basis for standardised HHTI investigations. The large and unexplained variance in hSAR estimates indicates the need for improved standardisation in the planning, conduct and analysis of HHTIs. Greater emphasis on clear and comprehensive reporting of HHTIs, and the context in which they are conducted, is required to facilitate more nuanced analysis of the sources of heterogeneity. High-quality FFX and HHTIs should continue to be conducted, ideally within a standardised framework such as the WHO Unity Studies initiative and be supported to guide evidence-based pandemic preparedness and response efforts for SARS-CoV-2, influenza and future novel respiratory viruses.

Unity Studies Collaborator Group

UNITY Studies Collaborator Group: Mikias A Alemu¹⁵, Yasir Alvi¹⁶, Elizabeth A Bukusi¹⁷, Pui Shan Chung¹², Davaalkham Dambadarjaa¹⁸, Ayan K Das^{19,20}, Timothée Dub²¹, Diba Dulacha²², Faiqa Ebrahim²³, Maritza A González-Duarte²⁴, Dinuka Guruge²⁵, Damaris C Heredia-Melo²⁶, Amy Herman-Roloff²⁷, Belinda L Herring², Farzana Islam²⁸, Kamal Chandima Jeewandara²⁹, Shashi Kant³⁰, Richard Lako³¹, Juliana Leite¹³, Gathsaurie Neelika Malavige³², Undram Mandakh³³, Warisha Mariam³⁴, Tsogt Mend³⁵, Merit Melin²¹, Valerie A Mize²², Sanjin Musa^{36,37}, Hanna Nohynek²¹, Olushayo O Olu²², May B Osorio-Merchán²⁴, Dmitriy Pereyaslov¹, James Ransom⁶³⁸, Lubna Al Ariqi⁹, Wasiq Khan⁹, Sonal Saxena³⁹, Pragya Sharma³⁴, Aswathy Sreedevi⁴⁰, Mini Satheesh^{41,42}, KJ Subhashini³⁰, Beth A Tippet-Barr^{26,43}, Anjua Usha⁴⁴, Joseph F Wamala²², Shambel H Watore¹⁵, Kapil Yadav³⁰, Francis Y Inbanathan¹¹

⁵deceased

¹⁵Ethiopian Public Health Institute

¹⁶Department of Community Medicine, Hamdard Institute of Medical Sciences and Research, New Delhi, India

¹⁷Kenya Medical Research Institute, Kenya

¹⁸School of Public Health, Mongolian National University of Medical Sciences, Ulaanbaatar, Mongolia

¹⁹Department of Microbiology, Hamdard Institute of Medical Science and Research, New Delhi, India

²⁰Hakeem Abdul Hameed Centenary Hospital, New Delhi, India

²¹Department of Health Security, Finnish Institute for Health and Welfare, Helsinki, Finland

²²WHO Country Office, Juba, South Sudan

²³World Health Organization Country Office, Ethiopia

²⁴National Institute of Health, Colombia

²⁵Colombo Municipality Council, Colombo, Sri Lanka

²⁶U.S. Centers for Disease Control and Prevention, Kenya

²⁷Department of Community Medicine, Hamdard Institute of Medical Sciences and Research, New Delhi, India

²⁸Hamdard Institute of Medical Sciences & Research (HIMSR), New Delhi, India

²⁹Allergy Immunology and Cell Biology Unit, Department of Immunology and Molecular Medicine, Faculty of Medical Sciences, University of Sri Jayewardenepura, Gangodawila, Nugegoda, Sri Lanka

³⁰Centre for Community Medicine, All India Institute of Medical Sciences, New Delhi, India

³¹Ministry of Health, South Sudan

³²Faculty of Medical Sciences, University of Sri Jayawardenapura, Nugegoda, Sri Lanka

³³Mongolian National University of Medical Sciences, Ulaanbaatar, Mongolia

³⁴Department of Community Medicine, Maulana Azad Medical College, New Delhi, India

³⁵National Center for Communicable Diseases, Ulaanbaatar, Mongolia

³⁶Institute for Public Health of the Federation of Bosnia and Herzegovina, Sarajevo, Bosnia and Herzegovina

³⁷Sarajevo School of Science and Technology, Sarajevo, Bosnia and Herzegovina

³⁸Centers for Disease Control and Prevention, South Sudan

³⁹Department of Microbiology, Maulana Azad Medical College, New Delhi, India

⁴⁰Department of Community Medicine, Amrita Institute of Medical Sciences, Kochi, Kerala, India

⁴¹Kerala University of Health Sciences, Kerala, India

⁴²Government Medical College, Thiruvananthapuram, Kerala, India

⁴³Nyanja Health Research Institute, Salima, Malawi

⁴⁴Regional Prevention of Epidemic and Infectious Disease cell, Government of Kerala, Kerala, India

Acknowledgements

Thank you to Miu-Ling Wong, Hong Kong University, for her assistance with translation of articles in Mandarin.

We thank colleagues: Mike Ryan, Keteven Glonti, Aisling Vaughan, Tony Wurda, Johson Muki, Yona Kenyi Manoah, Manuela Alphose Juma, Suzy Agrey Abas, Musah Kwaku Bukaru, Adamu Tayachew, Negussie Yohannes, Zewdu Assefa, Mengistu Biru, Musa Emmanuel, Margaret Mburu, Jayne Lewis-Kulzer, Pamela Murnane, Rachael Joseph, Elizabeth Oele, John Ndyahikayo, Ayesheshem A Tegegne, Arash Rashidian, Anna Solastie, Anu Haveri, Niina Ikonen, Lotta Hagberg, Oona Liedes, Mia Blazevec, Anes Joguncic, Faris Dizdar, Palo Mirza, Franklyn E Prieto-Alvarado, Ruth C Figueroa-Baez, Susanne C Ardila-Roldan, Mohammad Ahmad, Anisur Rahman, Shivani Rao, Mongjam Meghachandra Singh, Saurav Basu, Vikas Manchanda, Rohit Chawla, Faheem Ahmed, Mridu Dudeja, Farishta HD Singh, Iqbal Alam, Sushovan Roy, Ekta Gupta, Anil Kumar, Neethu Mohan, Soumya Gopakumar, Geetha Raveendranath, Deshni Jayathikala, Inoka S Aberathna, Saubhagya Danasekara, Buddini Gunathilake, Ruwan Wijayamuni, Ambaselmaa Amarjarga, Bilegt Altangerel, Temuulen Enebish, Oyunsuren Enebish, Ariuntuya Ochirpurev.

We would especially like to thank all WHO UNITY Study Collaborator Group, who are also named co-authors of this paper. These and other collaborators are recognized on a dedicated webpage on the UNITY and WHO website (52) in all the countries who embarked in this global response effort to COVID-19, as well as all individuals who supported, conducted or participated in each of the studies supported.

The authors alone are responsible for the views expressed in this article and they do not necessarily represent the views, decisions, or policies of the institutions with which they are affiliated.

Funding

This work was supported by WHO through funding from the WHO Solidarity Response Fund and the German Federal Ministry of Health COVID-19 Research and Development.

Data sharing

The data extraction template, data extraction forms, data used for analysis and the analysis code are publicly available on Zenodo in the WHO Unity Studies Global SARS-CoV-2 Seroepidemiological Investigations community.(14)

Appendix

Detailed searches

1. Medline – searched 17th August 2021

COVID-19/ or (COVID-19 or COVID19 or SARS-CoV-2 or coronavirus).tw

AND

(Household transmission OR secondary attack rate* OR close contact* OR contact transmission OR contact attack rate* OR family transmission).tw

2. EMBASE – searched 17th August 2021

coronavirus disease 2019/ or (COVID-19 or COVID19 or SARS-CoV-2 or coronavirus).tw

AND

(Household transmission OR secondary attack rate* OR close contact* OR contact transmission OR contact attack rate* OR family transmission).tw

3. Web of Science and Scopus – searched 17th August 2021

COVID-19 or COVID19 or SARS-CoV-2 or coronavirus

AND

"household transmission" OR "secondary attack rate*" OR "close contact*" OR "contact transmission" OR "contact attack rate*" OR "family transmission"

4. medRxiv and bioRxiv (through medRxiv – COVID-19 papers have been cross-indexed) – searched 30th August 2021

Advanced Search of (abstracts or titles)

Covid “household transmission”

Covid “secondary attack rate”

Covid “close contact”

Covid “contact transmission”

Covid “contact attack rate”

Covid “family transmission”

Detailed eligibility criteria

Inclusion criteria:

For screening of abstracts and full text:

- Any (published) investigation/study involving five or more discrete households with at least one laboratory confirmed index case of COVID-19 and all their household contacts or WHO supported investigations
- Investigations will need to provide information on transmission dynamics, severity and clinical spectrum of COVID-19 within the household setting. Specifically, they need to report on the household secondary attack rate (infection or clinical) with a measure of uncertainty, or contain

sufficient data to calculate the secondary attack rate (i.e., number of household contacts that become cases, total number of household contacts and household sizes) and may also report on the following: severity indicators such as the asymptomatic proportion of cases, hospitalisation and ICU admission rate, case fatality rate and transmission parameters such as the serial interval.

- Eligible papers will be restricted to the following languages: English, Chinese, French, Russian and Spanish, Portuguese, German and Italian.

For full text only:

- These studies will need to be aligned to the WHO Unity Studies FFX and HH transmission investigation protocols (available at: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/early-investigations>)
 - household recruitment is triggered by detection of a RT-PCR-confirmed index case
 - longitudinally collected (prospective or retrospective) data from all household members including epidemiological/virological or serological data (within the 28-day period from symptom onset in the primary case or household enrolment)
 - Active follow up of households (through case investigation and contact tracing)

Exclusion criteria:

- Duplicate studies will be removed. Where multiple papers are published on the same cohort (i.e., interim and final findings, subgroup analyses and secondary data analyses), we will use the most complete and relevant data from the article where summary statistics are appropriately reported.
- Studies that don't report sufficient information (i.e., conference abstracts or where full text is not available or where additional information can't be accessed upon contacting study investigators).

We investigated if the estimates varied by the following subgroups: country/region where study was conducted, pandemic course (epidemic activity), income setting, predominantly circulating variant at the time of the investigation/study was conducted, size of household, source population, implementation methods and index case symptom status. If data were available, we investigated if the estimates varied by sex, and age groups of the index case. We also plan to investigate if the estimates vary by publication status (i.e., comparing investigations in the literature to unpublished investigations).

Changes from PROSPERO registration and other details

- Adapted criteria for WHO Unity Studies investigations to be included irrespective of number of households included and follow up of all household contacts. This allowed us to consider inclusion of estimates that would otherwise not be included in line with our objectives. Note that all included WHO-supported investigations had at least five households.
- We did not screen reference lists of included articles. Unpublished articles and reports provided from WHO often did not contain a reference list, and we believe that our search strategy was very comprehensive and sensitive.

- Changes to subgroup analyses were determined prior to the conduct of analysis (Initial plans were as follows: We investigated if the estimates varied by the following subgroups: country/region where study was conducted, pandemic course (epidemic activity), income setting, predominantly circulating variant at the time of the investigation/study was conducted, size of household, source population, implementation methods and index case symptom status. If data were available, we investigated if the estimates varied by sex, and age groups of the index case. We also plan to investigate if the estimates vary by publication status)
- Authors in the collaborator group were included based on the ICJME criteria. Authorship was not offered in exchange for contribution of early data to the review.
- We initially intended to review all transmission and severity parameters. We decided to focus on the hSAR and hSCAR only to enable more detailed analysis of collected data.
- Description of the statistical analysis was modified to reflect change in meta-analysis models used in the analysis.
- Due to the significant amount of observed heterogeneity, we did not present a pooled HSAR and hSCAR estimate and did not visually assess funnel plots of effect size versus standard error and used Egger's test to evaluate bias due to small study effect.(18)
- Additionally, we planned to conduct a sensitivity analysis by fitting a beta-binomial model as applied in Thompson et al. to explore whether the chosen model impacted the results. This was not undertaken as we decided to not produce a pooled estimate.

Detailed questions used for methodological quality assessment

The questions detailed in the HHTI critical appraisal tool (13) used to assess these studies are as follows:

1. Was the study planning, recruitment and data collection appropriately timed to achieve the objectives of the investigation?
2. Was the cohort of households enrolled as a consequence of an adequate method of case ascertainment?
3. Was a definition of "household" provided?
4. Were all eligible cases and all householders enrolled into the investigation?
5.
 - a. Were subsequent cases identified and ascertained using appropriate methods to calculate the hSAR?
 - b. Were subsequent cases identified and ascertained using appropriate methods to calculate the hSCAR?
6.
 - a. Were households followed up for a period sufficient to measure primary outcomes?
 - b. Did all primary cases and contacts remain part of the "household" for the duration of the investigation?
7.
 - a. Were steps taken to ensure that householders were susceptible at the time of enrolment?

- b. Were steps taken to ensure that subsequent infections were due to exposure within the household?
- 8. Are the methods for analysis appropriate given the study context and design, participant definitions and the measurement of outcomes?
- 9. Has loss to follow up been appropriately accounted for in the estimated outcomes?
- 10. Has any missing data been appropriately accounted for in the estimated outcomes?

References:

1. World Health Organization. WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020 [cited 2021 6th December]. Available from: <https://www.who.int/director-general/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020>.
2. Leclerc QJ, Fuller NM, Knight LE, Group CC-W, Funk S, Knight GM. What settings have been linked to SARS-CoV-2 transmission clusters? Wellcome Open Res. 2020;5:83-.
3. Lau LLH, Nishiura H, Kelly H, Ip DKM, Leung GM, Cowling BJ. Household Transmission of 2009 Pandemic Influenza A (H1N1): A Systematic Review and Meta-analysis. Epidemiology. 2012;23(4).
4. World Health Organization. Final report of the IHR Review Committee (as presented to 64th WHA). 2011 [cited 2022 4th March]. Available from: <https://www.who.int/teams/ihr/ihr-review-committees/h1n1>.
5. Van Kerkhove MD, Broberg E, Engelhardt OG, Wood J, Nicoll A. The consortium for the standardization of influenza seroepidemiology (CONSISE): a global partnership to standardize influenza seroepidemiology and develop influenza investigation protocols to inform public health policy. Influenza Other Respir Viruses. 2013;7(3):231-4.
6. World Health Organization. Pandemic Influenza Special Investigations and Studies (IPSS) [cited 2022 4th March]. Available from: https://www.who.int/influenza/surveillance_monitoring/influenza_pandemic_special_investigations/en/.
7. World Health Organization. Middle East respiratory syndrome coronavirus (MERS-CoV) guidelines: investigation tools 2012 [cited 2022 24th March]. Available from: https://www.who.int/health-topics/middle-east-respiratory-syndrome-coronavirus-mers#tab=tab_1.
8. World Health Organization. Coronavirus disease (COVID-19) technical guidance: The Unity Studies: Early Investigation Protocols [cited 2021 6th December]. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/early-investigations>.
9. World Health Organization. The first few X cases and contacts (FFX) investigation protocol for coronavirus disease 2019 (COVID-19), version 2.2 [cited 2021 6th December]. Available from: [https://www.who.int/publications/i/item/the-first-few-x-cases-and-contacts-\(-ffx\)-investigation-protocol-for-coronavirus-disease-2019-\(-covid-19\)-version-2.2](https://www.who.int/publications/i/item/the-first-few-x-cases-and-contacts-(-ffx)-investigation-protocol-for-coronavirus-disease-2019-(-covid-19)-version-2.2).
10. World Health Organization. Household transmission investigation protocol for 2019-novel coronavirus (COVID-19) infection, version 2.2 [cited 2021 6th December]. Available from: [https://www.who.int/publications/i/item/the-first-few-x-cases-and-contacts-\(-ffx\)-investigation-protocol-for-https://www.who.int/publications/i/item/household-transmission-investigation-protocol-for-2019-novel-coronavirus-\(2019-ncov\)-infection](https://www.who.int/publications/i/item/the-first-few-x-cases-and-contacts-(-ffx)-investigation-protocol-for-https://www.who.int/publications/i/item/household-transmission-investigation-protocol-for-2019-novel-coronavirus-(2019-ncov)-infection).
11. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ. 2021;372:n71.
12. United Nations. UN Data glossary – household 1993 [cited 2021 6th December]. Available from: <http://data.un.org/Glossary.aspx?q=household>.
13. Liu Y, Eggo RM, Kucharski AJ. Secondary attack rate and superspreading events for SARS-CoV-2. The Lancet. 2020;395(10227):e47.
14. Zenodo. WHO Unity Global SARS-CoV-2 Seroepidemiological Investigations 2022 [cited 2022 16th February]. Available from: <https://zenodo.org/communities/unity-sero-2021/?page=1&size=20>.
15. Covidence systematic review software VHI, Melbourne, Australia,. [Available from: www.covidence.org.
16. Villanueva-Cabezas J SV, Marcato AJ, Meagher N, Fielding JE, Karahalios A, et al. Household Transmission Investigation: Design, Reporting and Critical Appraisal. MetaArXiv. 2021. Available from: osf.io/preprints/metaarxiv/mrf6d.
17. Wells G SB, O'Connell D, et al.,. The Newcastle-Ottawa Scale (NOS) for assessing the quality of 25 nonrandomised studies in meta-analyses. [cited 2022 4th March]. Available from: http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp.
18. Institute JBS. Critical Appraisal Tools [cited 2022 4th March]. Available from: <https://jbi.global/critical-appraisal-tools>.
19. World Health Organization. Countries [cited 2022 16th February]. Available from: <https://www.who.int/countries>.
20. The World Bank. World Bank Country and Lending Groups 2021 [cited 2021 6th December]. Available from: <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>.
21. United Nations Office for the Coordination of Humanitarian Affairs (OCHA). Global humanitarian response plan for COVID-19 2020 [cited 2022 23rd February]. Available from: https://www.unocha.org/sites/unocha/files/GHRP-COVID19_July_update.pdf.
22. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. Stat Med. 2002;21(11):1539-58.
23. Stijnen T, Hamza TH, Özdemir P. Random effects meta-analysis of event outcome in the framework of the generalized linear mixed model with applications in sparse data. Statistics in Medicine. 2010;29(29):3046-67.
24. GISAID. [Available from: <https://www.gisaid.org/>.
25. CoVariants. Overview of Variants in Countries [cited 2022 23rd February]. Available from: <https://covariants.org/per-country>.
26. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. BMJ. 1997;315(7109):629-34.
27. Thompson HA, Mousa A, Dighe A, Fu H, Arnedo-Pena A, Barrett P, et al. Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Setting-specific Transmission Rates: A Systematic Review and Meta-analysis. Clin Infect Dis. 2021;73(3):e754-e64.
28. StataCorp. Stata Statistical Software: Release 16. College Station, TX: StataCorp LLC. 2019.
29. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2020 [Available from: <https://www.R-project.org/>.
30. Viechtbauer W. Conducting Meta-Analyses in R with The metafor Package. Journal of Statistical Software. 2010;36.
31. Madewell ZJ, Yang Y, Longini IM, Jr, Halloran ME, Dean NE. Factors Associated With Household Transmission of SARS-CoV-2: An Updated Systematic Review and Meta-analysis. JAMA Network Open. 2021;4(8):e2122240-e.

32. Koh WC, Naing L, Chaw L, Rosledzana MA, Alikhan MF, Jamaludin SA, et al. What do we know about SARS-CoV-2 transmission? A systematic review and meta-analysis of the secondary attack rate and associated risk factors. *PloS one*. 2020;15(10):e0240205.
33. von Hippel PT. The heterogeneity statistic I2 can be biased in small meta-analyses. *BMC medical research methodology*. 2015;15(1):1-8.
34. Deeks J, J pT H, Altman D. Chapter 10: analysing data and undertaking meta-analyses | *Cochrane Training*. *Cochrane handbook for systematic reviews of interventions*. 2021;6.
35. Fitzgerald DA, Wong GWK. COVID-19: A tale of two pandemics across the Asia Pacific region. *Paediatr Respir Rev*. 2020;35:75-80.
36. Ghosh AK, Venkatraman S, Soroka O, Reshetnyak E, Rajan M, An A, et al. Association between overcrowded households, multigenerational households, and COVID-19: a cohort study. *Public Health*. 2021;198:273-9.
37. World Health Organization. Tracking SARS-CoV-2 variants 2021 [cited 2021 6th December]. Available from: <https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/>.
38. Davies NG, Abbott S, Barnard RC, Jarvis CI, Kucharski AJ, Munday JD, et al. Estimated transmissibility and impact of SARS-CoV-2 lineage B. 1.1. 7 in England. *Science*. 2021;372(6538):eabg3055.
39. Keeling MJ. Estimating the transmission advantage for b. 1.617. 2. 2021.
40. Public Health England. SARS-CoV-2 variants of concern and variants under investigation in England - Technical briefing 17. 2021.
41. Golding N, Shearer FM, Moss R, Dawson P, Liu D, Ross JV, et al. Situational assessment of COVID-19 in Australia Technical Report 15 March 2021. 2021 28th May 2021.
42. Dye C, Bartolomeos K, Moorthy V, Kienny MP. Data sharing in public health emergencies: a call to researchers. *Bull World Health Organ*. 2016;94(3):158.
43. World Health Organization. WHO Coronavirus (COVID-19) Dashboard 2022 [cited 2022 4th January].
44. Shearer FM, Moss R, Price DJ, Zarebski AE, Ballard PG, McVernon J, et al. Development of an influenza pandemic decision support tool linking situational analytics to national response policy. *Epidemics*. 2021;36:100478.
45. Marcatto AJ, Fielding JE, Crooks K, Massey PD, Le L-V, Bergeri I, et al. The ongoing value of first few X studies for COVID-19 in the Western Pacific Region. *Western Pac Surveill Response*. 2022;13(1).
46. Bergeri I, Lewis HC, Subissi L, Nardone A, Valenciano M, Cheng B, et al. Early epidemiological investigations: World Health Organization UNITY protocols provide a standardized and timely international investigation framework during the COVID-19 pandemic. *Influenza and Other Respiratory Viruses*. 2022;16(1):7-13.
47. World Health Organization. Pandemic influenza preparedness framework for the sharing of influenza viruses and access to vaccines and other benefits, 2nd edition. 2021.
48. Black AJ, Geard N, McCaw JM, McVernon J, Ross JV. Characterising pandemic severity and transmissibility from data collected during first few hundred studies. *Epidemics*. 2017;19:61-73.
49. Schulz KF, Altman DG, Moher D. CONSORT 2010 statement: updated guidelines for reporting parallel group randomized trials. *Ann Intern Med*. 2010;152(11):726-32.
50. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Ann Intern Med*. 2007;147(8):573-7.
51. World Health Organization Seroepidemiology Technical Working Group. ROSES-S: Statement from the World Health Organization on the reporting of seroepidemiologic studies for SARS-CoV-2. *Influenza and Other Respiratory Viruses*. 2021;15(5):561-8.
52. World Health Organization. Acknowledgements: The Unity Studies for sero-epidemiological investigation of COVID-19 2022 [cited 2022 30th March]. Available from: <https://www.who.int/publications/m/item/acknowledgements-the-unity-studies-for-sero-epidemiological-investigation-of-covid-19>.