Syndemic violence victimization, alcohol and drug use, and HIV transmission risk behavior among transgender women in India: A cross-sectional, population-based study

Venkatesan Chakrapani^{1,2}, P.V.M. Lakshmi², Peter A. Newman^{3*}, Jasvir Kaur², Alexander C.

Tsai⁴, P.P. Vijin², Bhawani Singh⁵, Pradeep Kumar⁵, Shobini Rajan⁵, Rajesh Kumar²

¹ Centre for Sexuality and Health Research and Policy (C-SHaRP), Chennai, India

² Postgraduate Institute of Medical Education and Research (PGIMER), School of Public Health, Chandigarh, India

³ University of Toronto, Factor-Inwentash Faculty of Social Work, Toronto, Canada

⁴ Massachusetts General Hospital and Harvard Medical School, Boston, United States

⁵ National AIDS Control Organisation, New Delhi, India

* Corresponding author:

Email: p.newman@utoronto.ca

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

1 Abstract

2 **Introduction:** Transgender women are disproportionately burdened by HIV. Co-occurring 3 epidemics of adverse psychosocial exposures accelerate HIV sexual risk, including among transgender women; however, studies using additive models fail to examine synergies among 4 5 psychosocial conditions that define a syndemic. We examined the impact of synergistic interactions among 4 psychosocial exposures on condomless anal sex (CAS) among a 6 national probability sample of transgender women in India. 7 8 Methods: A probability-based sample of 4,607 HIV-negative transgender women completed the Indian Integrated Bio-behavioral Surveillance survey, 2014-2015. We used linear 9 10 probability regression and logistic regression to assess 2-, 3-, and 4-way interactions among 4 exposures (physical and sexual violence, drug and alcohol use) on CAS. 11 **Results:** Overall, 27.3% reported physical and 22.3% sexual violence victimization (39.2%) 12 13 either physical or sexual violence), one-third (33.9%) reported frequent alcohol use and 11.5%

illicit drug use. Physical violence was associated with twofold higher odds of CAS in the main
effects model. Significant two- and three-way interactions were identified on both
multiplicative and additive scales between physical violence and drug use; physical and sexual
violence; physical violence, sexual violence, and alcohol use; and physical violence, alcohol
and drug use.

Conclusions: Physical and sexual violence victimization, and alcohol and drug use are highly prevalent and synergistically interact to increase CAS among transgender women in India. Targeted and integrated initiatives to improve assessment of psychosocial comorbidities, to combat transphobic violence, and to provide tailored, trauma-informed alcohol and substance use treatment services may reduce HIV risk among transgender women.

24

Keywords: syndemics, transfeminine people, HIV, syndemic theory, mental health, stigma

25 Introduction

Transgender women globally bear a disproportionate HIV burden [1] secondary to 26 rampant stigma and discrimination in various domains of their lives [2]. India's National AIDS 27 Control Organization (NACO) designates transgender women a 'high risk group' or 'key 28 29 population', with official HIV prevalence estimates ranging from 9.5% (in 2015 [3]) to 3.1% (in 2017 [4])—from 14 to over 40 times higher than the general population (0.22%) [5]. Until 30 2006, NACO-supported HIV preventive interventions subsumed transgender women under 31 32 "men who have sex with men"; but they are now reached through 41 interventions exclusively for transgender women and 152 'core-composite' targeted HIV interventions for transgender 33 34 women with other at-risk populations [5]. These targeted interventions focus on HIV and STI education, condom promotion, free condom distribution, and HIV testing referrals. 35 Transgender women in India have demonstrated high (89%) awareness of the usefulness of 36 37 condoms in preventing HIV infection; nevertheless, consistent condom use remains low at 45% to 54% [4,6]. 38

Syndemic theory has been applied to explain the clustering and concentration of 39 diseases such as HIV in vulnerable populations [7], including transgender women [1,8]. In their 40 classic paper [9], Stall et al. highlighted the public health impacts of co-occurring psychosocial 41 42 problems among gay men in the U.S., advancing a life-course theory of syndemic production rooted in the cultural marginalization of a population that is heavily stigmatized [10]. In the 43 Indian context, harmful social conditions such as societal stigma faced by transgender women 44 45 may lead to clustering of psychosocial health problems, such as violence victimization, problematic alcohol use, depression, and HIV risk [11,12]; these may mutually reinforce one 46 another and synergistically amplify disease burden. 47

48 Transgender women in India have a high prevalence of psychosocial health problems,
49 including lifetime physical or sexual violence victimization (84.0%), alcohol use (37.3%) and

depression (35.3%), all of which have demonstrated significant additive effects on HIV risk
[13]. One study reported 18% past-year prevalence of sexual violence victimization, with
higher risk for HIV among those who experienced sexual violence [14].

Exploring the presence of synergy among psychosocial health conditions requires the 53 use of appropriate analytical strategies. A recommended analytical technique for testing 54 synergy is by checking for statistical interaction (on the additive and/or multiplicative scale) 55 56 between co-occurring conditions [15,16]. Most analyses in western literature that purportedly explored synergy used the number of psychosocial conditions or exposures as a cumulative 57 58 count to predict HIV risk behavior, which does not address the interaction concept embedded in syndemic theory [15]. Two papers on syndemics among transgender women in India each 59 used analytical strategies that did not involve testing for interactions between psychosocial 60 61 conditions. One study used the number of psychosocial syndemic conditions as a cumulative count to predict HIV risk behavior [13]. Another study used latent class analysis to identify 62 different 'syndemic classes' (combinations of depression, alcohol use, and/or violence 63 64 victimization), which were associated with higher HIV risk compared to no syndemic class [6]. To address research gaps in the application of syndemic theory and support the design 65 of tailored, evidence informed interventions, we assessed potential synergistic interactions 66 between sexual and physical violence, alcohol use, and drug use on condomless anal sex (CAS) 67 68 among transgender women.

69

70 Materials and Methods

71 Study design and population

Data for these analyses were drawn from a population-based cross-sectional survey
 among transgender women recruited under India's National AIDS Control Organization

(NACO) Integrated Bio-Behavioral Surveillance (IBBS) study, last conducted from 2014–2015. The IBBS aimed to generate evidence on risk behaviors among groups deemed high-risk (e.g., transgender women, female sex workers, men who have sex with men) to support planning and prioritization of HIV program efforts. Eligibility criteria were as follows: persons aged \geq 15 years, whose self-identity does not conform unambiguously to conventional notions of male or female gender roles but combines or moves between them.

The IBBS study was approved by the ethics committees of NACO and participating Indian Council of Medical Research regional institutes. All participants provided written informed consent; for those 15-17 years-old, consent was also obtained from their parent/guardian. Monetary compensation of INR 200 (~3 USD) was provided.

This paper presents findings from secondary analysis of NACO population-based study data to examine the burden of adverse psychosocial exposures, co-occurrence of these exposures, and their potential independent and synergistic effects on CAS among transgender women.

88

89 Sampling and recruitment

90 The IBBS sample size was calculated with the primary objective to measure changes in consistent condom use with sexual partners and changes in HIV prevalence. The primary unit 91 92 of survey was a 'domain': a geographical unit of either a single district or a group of four socioculturally similar districts where a single district had an insufficient sample size. Sampling 93 94 including both conventional cluster sampling (CCS)—in sites (e.g., homes, brothels) where transgender women could be found at any time of the day, and time-location cluster sampling 95 (TLCS)—in sites such as parks and beaches where transgender women could be found on 96 particular days and times to meet or socialize with other transgender people or potential male 97 partners. The estimated sample size of transgender women for each domain was >700. 98

99 However, considering feasibility based on sample availability in different domains, a target 100 sample size of 400 per domain was chosen. In domains with sample sizes <400, a 'take all' 101 approach was used. Overall, the target sample size was 5,588 transgender women in 15 102 domains across 11 states. A valid sample of 4,966 transgender women was achieved, with an 103 81% response rate. NACO's online report [17] provides further details on sampling and 104 recruitment.

105

106 Measures

Information on participants' sociodemographic characteristics included age (in years), 107 vears of education, marital status, engagement in sex work (past year) and self-identified 108 gender-affirmative surgical status (ackwa: pre- or non-operative, or nirvan: post-operative). 109 Table 1 describes how the outcome, primary exposures, and other relevant covariates used in 110 the analysis were measured—survey items, response range and scoring, and creation of binary 111 and continuous measures. The binary outcome measure was CAS with any type (regular, 112 paying, paid, and casual) of male sexual partner in the past month. Binary adverse psychosocial 113 114 exposures included physical violence [PV], sexual violence [SV], and drug use [D] (non-115 injection or injection drugs) in the past year, and frequent alcohol use [A] (>3 days) in the past week (Table 1). Other covariates included engagement in sex work, forced sex experience 116 117 during adolescence (<18 years), HIV transmission knowledge (score), HIV risk perception (high vs. low), social support (score), and HIV program exposure (score). 118

119 Table 1. Summary of Key Measures: Outcome, Primary Exposures, and other Relevant Covariates

		Survey items		Types of variables	
Measures	Number of items	Questions asked	Range of responses, coding, and scores		
Outcome					
Condomless anal sex with male partners (past month)	4	In the last one month, how often have you used condoms when you had anal sex with your? a. regular non-paying male partner b. paying male partner c. paid male partner d. casual non-paying male partners	 0 for 'every time' and 1 for all other options (most of the time, sometimes, never) 'Condomless anal sex' was created by combining the condom use (=1) responses from these four types of partners. 	Binary	
Primary exposures					
Physical violence (past year)	2	In the last 12 months, how many times would you say someone has beaten (hurt, hit, slapped, pushed, kicked, punched, choked or burned) you? In the last 12 months, who was the person (or people) who have	0 for 'never' and 1 for other responses (Once, 2-5 times, 6-10 times, >10 times). Those who responded as "Do not remember" but reported the person who caused physical violence were also categorized as having experienced physical	Binary	
Somuel violence	1	In the last 12 months, were you physically forced to have sevuel	Violence.	Dinory	
(past year)	1	intercourse with someone even though you didn't want to?	105-1, 100-0	Dinary	
Violence victimization (past year)	-	Recoding of the final coding of physical and sexual violence variables.	'Violence victimization' was created by considering the codes of physical violence (from never=0, Once=1, 2-5 times=2, 6-10 times=3, >10 times=4) and sexual violence (Yes=1, No=0) as scores, and combining these scores. Score range: 0-5	Continuous	
Alcohol use (past week)	1	How many days did you consume alcohol in the last one week?	Responses dichotomized at the upper quartile of consumption, \leq 3 vs. >3 days/week.	Binary	
			Score range: 0-7 (Number of days of alcohol use in the past week)	Continuous	
Drug use (past year)	2	Have you consumed drugs such as Ganja, Heroine for pleasure in the last 12 months? Have you injected drugs for nonmedical reasons in the last 12 months?	Yes=1, No=0 Consumption of any drug (injection or not) was considered as 'drug use'. Score range: 0-3	Binary Continuous	
Relevant covariates					

<i>Engagement in sex work</i> (past year)	1	Have you received cash or gifts from men in exchange for sex in the last 12 months?	Yes=1, No=0	Binary
Knowledge of HIV transmission risk	5	Can a person get HIV/AIDS?	Yes=1, No=0	Continuous
		 b. by sharing infected needles 	For the analysis, each correct response was coded as	
		d. through mosquito bites	1, otherwise zero .	
		Can HIV be transmitted from an HIV infected mother to	Score range: 0 to 5	
		e. her unborn baby during pregnancy/delivery or to the new born child through breastfeeding?		
HIV risk perception	1	To what extent do you feel yourself at risk to being infected with	High=1, Moderate=2, Low=3, No risk=4	Binary
			1 and 2 as 'High risk', 3 and 4 as 'Low risk'	
Forced sex experience	2	How old were you when you had your first sex with a male?	Participants who reported first sex with a male when	Binary
auring adolescence		Were you forced to have sex during the first sexual encounter with	intercourse in India) and of them who reported forced	
		a male?	sex during the first sexual encounter with a male,	
			were categorised as having experienced forced sex	
	0		during adolescence.	Centimere
HIV program exposure	8	NGO/program/individual/group during the last 12 months?	Y es=1, No=0	Continuous
		a. Received information on STI/HIV/AIDS	Score range: 0 to 8	
		b. Received condoms	5	
		c. Received lubricants		
		d. Seen a demonstration on correct condom use		
		f Received free medicine for STIs		
		g. Visited drop in center		
		h. Referred to other services (STI clinic, HIV testing, etc.)		
Social support score	3	Have you received following services from any	Yes=1, No=0	Continuous
		NGO/program/individual/group during the last 12 months?	Score range: 0 to 3	
		a. Received help and support when faced with physical of sexual violence	Score range. 0 to 5	
		b. Received help and support when faced with trouble from		
		police		
		Are you a member of?		
		c. a self-help group or transgender women collective		

121 Analysis

The present analysis was restricted to HIV-negative transgender women (n=4,607). 122 HIV-positive participants (n=359; 7.2%) were excluded given the objective of identifying 123 correlates of HIV risk behavior to inform primary prevention initiatives. All analyses were 124 conducted in Stata (version 16; College Station, Texas, USA). Descriptive statistics were used 125 to summarize participant characteristics. Independent associations between adverse 126 psychosocial exposures (binary) and CAS (binary) were estimated using logistic regression. 127 128 The extent to which the adverse psychosocial exposures synergistically interacted with each other in their associations with CAS was examined by assessing interactions on the additive 129 scale using linear probability regression models, and on the multiplicative scale using logistic 130 regression models [15,18]. All possible two-way (e.g., physical violence x drug use), three-way 131 (e.g., sexual violence x drug use x alcohol use), and four-way interaction terms (e.g., physical 132 133 violence x sexual violence x drug use x alcohol use) were included in the regression models. All regression models were adjusted for age (in years), education, marital status (single vs. 134 heterosexually married), gender-affirmative surgical status (ackwa vs. nirvan), engagement in 135 136 sex work, forced sex experience during adolescence, knowledge of HIV transmission, HIV risk perception, social support, and exposure to HIV interventions. To account for the complex 137 survey design [19] involving cluster sampling, 'svyset' and 'svy' Stata commands were used 138 139 before estimating regression models [20].

On the additive scale in linear probability regression models, synergy is interpreted as statistically significant relative excess risk due to interaction (RERI) greater than zero; on the multiplicative scale in logistic regression models, synergy is interpreted as statistically significant coefficients on the product terms [18,21]. Multiplicative interaction estimated on the odds scale using logistic regression was re-estimated as 'semi-elasticities' on the probability scale using the Stata 'margins' command. The estimated semi-elasticities can be interpreted as the proportional change in the probability of CAS that is associated with a one
unit change in the covariate or with the interaction. For example, a semi-elasticity of .40 is
interpreted as a 40% relative change in the expected probability of CAS associated with a one
unit change in the covariate (e.g., age in years) or with the joint effect of two (PV x D) or more
(PV x SV x D) exposures, above and beyond their independent associations with the outcome.

152 Sensitivity analysis for the model of synergistically interacting

153 epidemics

To confirm the robustness of our findings, as part of the sensitivity analysis, we specified: 1) drug use (score range, 0–3) and alcohol use (range, 0–7) as continuous exposures; 2) physical and sexual violence (score range, 0–5) as a single continuous exposure, along with CAS (range, 0–4) as a continuous outcome in both sensitivity analyses.

158

159 Exploratory analyses to check for rival theoretical models

In addition to synergistically interacting epidemics, Tsai [22] described two other 160 models of co-occurring epidemics—the serially causal epidemics model and mutually causal 161 epidemics model [23]—that have been used, variably, in the literature on syndemics. Although 162 our analysis was focused specifically on testing a model of synergistically interacting 163 164 epidemics, we also tested these other two models as they may suggest alternate or complementary mechanisms of risk. To test the model of serially causal epidemics, we 165 conceptualized alcohol (score range, 0-7) and drug use (range, 0-3) as potential mediators of 166 the association between sexual and physical violence victimization and CAS (range, 0-4). 167 Similarly, we tested the model of mutually causal epidemics, in which we hypothesized that 168 violence victimization (range, 0-5) and drug use (range, 0-3) were mutually reinforcing of one 169

another, given that both were measured in the same time-period (past year). For assessing bothof these models, we conducted path analysis (sem command) in Stata-16.

172

173 **Results**

174 **Participant characteristics**

Table 2 describes participant characteristics. Participants' mean age was 28.5 years (SD 175 176 7.4) and mean number of years of education was 8.4 (SD 4.0). A majority (86.1%) were currently single. Nearly one-third (31.1%) reported sex work as their main occupation, while 177 178 over half (56.8%) reported engaging in sex work during the past month. About one-fifth (18.3%) reported having experienced forced sex during adolescence (<18 years of age). More 179 than half (57.2%) self-identified as ackwa, 42.7% as nirvan. About three-fifths (61.6%) 180 perceived themselves at high risk for HIV infection. Reported CAS with different types of male 181 partners in the past month ranged from 35.0% to 46.0%; CAS with any type of male partner 182 was 45.1%. 183

184

186 Table 2. Characteristics of HIV-negative Transgender Women who Participated in the

187 Integrated Bio-Behavioral Surveillance (IBBS) study, 2014-15 (N = 4,607)

Characteristics	Mean (SD)				
Age (in years)	28.5 (7.4)				
Education (years)	8.4 (4.0)				
HIV transmission knowledge score	4.3 (1.2)				
HIV program exposure score	4.7 (2.8)				
Social support score	1.7 (1.0)				
	N (%)				
Engagement in sex work (past month)	2617 (56.8)				
Gender-affirmative surgical status					
Ackwa (pre-operative/non-operative)	2636 (57.2)				
Nirvan (post-operative)	1969 (42.7)				
Current marital status					
Single (never married/widowed/divorced/separated)	3940 (86.1)				
Married	637 (13.9)				
HIV risk perception ^a					
Low	1594 (34.6)				
High	2839 (61.6)				
Forced sex experience during adolescence	837 (18.2)				
Condomless anal sex (past month), with ^b :					
Regular non-paying male partner	935/2033 (46.0)				
Paying male partners	884/2375 (37.2)				
Paid male partners	310/885 (35.0)				
Casual non-paying male partners	483/1146 (42.1)				
Any type of male partner	1502/3328 (45.1)				

^a Percentages may not add to 100 due to missing values.

^b Among those who had that type of partner(s) and reported anal sex during the past month.

190

191 **Co-occurrence of adverse psychosocial exposures**

Among the 4,607 HIV-negative transgender women, irrespective of other exposures, more than one-fourth (27.3%) reported having been exposed to physical violence and 22.3% to sexual violence. More than one-third (39.2%) had been exposed to either physical or sexual

195	violence and 10.5% to both physical and sexual violence. Over one-third (33.9%) reported
196	frequent alcohol use and 11.5% reported drug use. Table 3 reports the co-occurrence of
197	physical violence, sexual violence, frequent alcohol use, and drug use. Overall, 16.5% (n=698)
198	reported two co-occurring psychosocial exposures, ranging from 1.8% (n=76) co-occurrence
199	of alcohol and drug use to 6.1% (n=256) alcohol use and physical violence. Three exposures
200	were reported by 5.8% (n=243), and 2.7% (n=113) reported all four exposures. About 43% of
201	participants reported no adverse psychosocial exposures.

202

204 Table 3. Adverse Psychosocial Exposures among HIV-negative Transgender Women (N

205 = 4,607): Physical Violence, Sexual Violence, Frequent Alcohol Use, Drug Use, and their

206 Co-occurrence

Adverse psychosocial exposures and their co-occurrence	n (%) ^a
Each exposure irrespective of other three exposures	
Physical violence, past 12 months	1257 (27.3)
Sexual violence, past 12 months	1029 (22.3)
Drug use, past 12 months	488 (11.5)
Frequent alcohol use, past week (>3 days/week)	1563 (33.9)
Each exposure alone	
Physical violence	352 (8.3)
Sexual violence	264 (6.2)
Drug use	86 (2.0)
Frequent alcohol use	655 (15.5)
Co-occurrence of exposures	
Frequent alcohol use + Drug use	76 (1.8)
Frequent alcohol use + Physical violence	256 (6.1)
Frequent alcohol use + Sexual violence	124 (2.9)
Physical violence + Drug use	35 (0.8)
Sexual violence + Drug use	62 (1.5)
Physical violence + Sexual violence	145 (3.4)
Physical violence + Drug use + Frequent alcohol use	40 (1.0)
Sexual violence + Drug use + Frequent alcohol use	59 (1.4)
Physical violence + Sexual violence + Frequent alcohol use	127 (3.0)
Physical violence + Sexual violence + Drug use	17 (0.4)
Physical violence + Sexual violence + Frequent alcohol use + Drug use	113 (2.7)
No adverse psychosocial exposures	1814 (42.9)

^a Percentages may not add to 100 due to missing values

208

210 Perpetrators of physical or sexual violence

Among those who experienced sexual violence (n=1,029), over one-third (35.9%; n=369) reported more than one type of perpetrator. Perpetrators were identified as strangers (41.1%), thugs (23.9%), clients (23.2%), and family members (21.8%). Among those who experienced physical violence (n=1,257), 32.5% (n=409) reported more than one perpetrator type. Perpetrators were identified as strangers (33.0%), police (24.5%), thugs (21.2%), family members (20.0%), and clients (17.3%).

217

218 Synergistic interactions between adverse psychosocial exposures

Using logistic regression to model CAS, without product terms, physical violence [PV] 219 had a statistically significant association with CAS (aOR=1.94, 95% CI: 1.47–2.56; p<.001), 220 while frequent alcohol use [A] (aOR = 1.32, 95% CI: .98-1.77; p=.07), drug use [D] (aOR 221 =.86,95% CI: .60–1.22; p=.40) and sexual violence [SV] (aOR = .97, 95\% CI: .73–1.29; p=.86) 222 did not have statistically significant associations with CAS. Among the other covariates 223 included in the multivariable logistic regression model of main effects without product terms, 224 higher levels of education (aOR=.93, 95% CI: .90-.97; p<.001), being unmarried (aOR=.64, 225 95% CI: .45-.92; p=.016), and nirvan gender-affirmative surgical status (aOR=.68, 95% CI: 226 .52-.88; p=.003) were associated with lower odds of CAS, and forced sex during adolescence 227 with higher odds of CAS (aOR=1.35, 95% CI: 1.02–1.78; p=.04). 228

Out of six logistic regression models that included main effects and two-way product terms, the following product terms were statistically significant: PV x D (Model 1) and PV x SV (Model 5) (Table 4). In the additional five logistic regression models that included main effects and two- and three-way product terms, the following were statistically significant: PV x D (Model 7), PV x A x D (Models 8, 9, 10, and 11), D x PV x SV (Model 10) and A x PV x SV (Models 11 and 12). In the final logistic regression model that included all possible product

- terms, including a four-way product term, only A x PV x SV was statistically significant.
- Assessment of interactions on the additive scale using linear probability regression models
- showed similar results (Table 5).

238Table 4. Effect of Adverse Psychosocial Exposures on Condomless Anal Sex with Male Partners among HIV-negative Transgender

239 Women: Measures of Multiplicative Two-/Three-/Four-way Interactions between Physical Violence, Sexual Violence, Drug Use, and

240 Frequent Alcohol Use (N = 4,607)

	Effects on condomless anal sex											
	Model 1:	Model 2:	Model 3:	Model 4:	Model 5:	Model 6:	Model 7:	Model 8:	Model 9:	Model 10:	Model 11:	Model 12:
Adverse psychosocial exposures and product terms	Models with one two-way product term at a time						All two-way product terms PV x D PV x A SV x D SV x A PV x SV A x D	All two-way + one three- way product term (PV x A x D)	All two-way + two three- way product terms (PV x A x D) (SV x A x D)	All two-way + three three- way product terms (PV x A x D) (SV x A x D) (D x PV x SV)	All two-way + four three- way product terms (PV x A x D) (SV x A x D) (D x PV x SV) (A x PV x SV)	All two-way + All three-way + four-way product terms (PV x A x D) (SV x A x D) (SV x A x D) (D x PV x SV) (A x PV x SV) (A x D x PV x SV)
	PV x D	PV x A	SV x D	SV x A	PV x SV	A x D						
						Estimate	d semi-elasticity (9	5% CI), p value				
PV x D	.48						.47	17	14	39	25	11
	(.27, .69)***	12					(.23, .71)***	(83, .48)	(79, .52)	(-1.20, .42)	(99, .49)	(88, .65)
PV X A		.13					(-28, 31)	$(-43 \ 27)$	09	09	38	36
SV x D		(10, .42)	- 02				- 12	- 21	- 02	- 13	- 12	- 04
			(39, .35)				(53, .29)	(64, .22)	(54, .58)	(77, .52)	(75, .52)	(69, .61)
SV x A				01			10	06	01	.01	38	36
				(33, .30)			(43, .24)	(39, .26)	(35, .33)	(33, .34)	(91, .14)	(89, .18)
PV x SV					.29		.23	.21	.22	.13	.31	28
4 D					(.04, .53)*	0.8	(04, .49)	(06, .48)	(05, .48)	(18, .44)	(/6, .14)	(75, .19)
						.08	00	(-1.0, 19)	(-90, 44)	(-81 47)	11	03
PV x A x D						(.27,.13)	(,	.69	.69	.65	.61	.52
								(.38, 1.0)***	(.38, 1.01)***	(.29, 1.01)***	(.23, .99)**	(04, 1.08)
SV x A x D									39	65	53	74
									(-1.32, .54)	(-1.72, .43)	(-1.56, .49)	(-2.09, .61)
D x PV x SV										.51	.34	.06
A x PV x SV										(.00, .90)*	(24, .93)	(-1.15, 1.25)
											(.35, .97)***	(.27, .99)***
A x D x PV x SV						1					, , ,	.40
												(64, 1.46)

242 PV, physical violence in the past year; SV, sexual violence in the past year; D, drug use in the past year; A: frequent alcohol use in the past week.

All models adjusted for age, education, marital status, sexual identity, forced sex experience during adolescence, HIV risk perception, knowledge

244 HIV transmission risk, social support, and HIV program exposure.

245 The estimates of the main effects are not shown.

Interpretation of semi-elasticity: a semi-elasticity of .48, for example in Model 1, is interpreted as a 48% relative change in the probability of

247 condomless anal sex that is associated with the joint effect of physical violence and drug use, above and beyond their independent associations

with the outcome.

249 *p<.05, **p<.01, ***p<.001.

250

251

Table 5. Effect of Adverse Psychosocial Exposures on Condomless Anal Sex with Male Partners among HIV-negative Transgender

254 Women: Measures of Additive Two-/Three-/Four-way Interactions between Physical Violence, Sexual Violence, Drug Use, and Frequent

255 Alcohol Use (n = 4,607)

256

	Effects on condomless anal sex											
	Model 1:	Model 2:	Model 3:	Model 4:	Model 5:	Model 6:	Model 7:	Model 8:	Model 9:	Model 10:	Model 11:	Model 12:
	Models with one two-way product term at a time						All two-way product	All two-way +	All two-way + two three	All two-way + three three	All two-way + four three-	All two-way +
Adverse psychosocial exposures and product terms	PV x D	PV x A	SV x D	SV x A	PV x SV	A x D	PV x D PV x A SV x D SV x A PV x SV A x D	way product term (PV x A x D)	way product terms (PV x A x D) (SV x A x D)	way product terms (PV x A x D) (SV x A x D) (D x PV x SV)	terms (PV x A x D) (SV x A x D) (D x PV x SV) (A x PV x SV)	An inree-way + four-way product terms (PV x A x D) (SV x A x D) (D x PV x SV) (A x PV x SV) (A x D x PV x SV)
	b (95% CI), p value											
PV x D	.24 (.11, .37)***						.23 (.08, .37)**	07	05 (31, .19)	12 (39, .14)	07 (33, .19)	05 (34, .24)
PV x A		.05					01	03	04	04	14 (31, .02)	14 (31, .03)
SV x D			01				04	07	005	04	03	02 (27, .23)
SV x A				005 (-13,12)			04	02	004	002	14 (32, 04)	13 (- 33_05)
PV x SV				(.13		(-03, 22)	.09	.09	.05	11	11 (- 28, 06)
A x D					(.01, .20)	.03	02	13	08	06	03	02 (27, 22)
PV x A x D						(.12,.10)	(.10,.12)	$(08, 71)^*$.40	.35	.31	.28
SV x A x D								(.00, .71)	11	17	11	13
D x PV x SV									(+0, .10)	.20	.08	.02
A x PV x SV										(00, .+0)	(21,, 57) ,,,,,,,	.35
A x D x PV x SV											(.10, .05)	.07

257 PV, physical violence in the past year; SV, Sexual violence in the past year; D, drug use in the past year; A, frequent alcohol use in the past week.

- All models adjusted for age, education, marital status, sexual identity, forced sex experience during adolescence, HIV risk perception, knowledge
- 259 HIV transmission risk, social support, and HIV program exposure.
- 260 The estimates of the main effects are not shown.
- 261 *'b'* represents the estimated regression coefficient on the product term which indicates the relative excess risk due to interaction (RERI) used to
- assess additive interaction.
- 263 *p<.05, **p<.01, ***p<.001.

Sensitivity analysis for the model of synergistically interacting 264 epidemics

The sensitivity analysis specifying drug use and alcohol use as continuous exposures, 266 267 physical and sexual violence as a single continuous exposure (violence victimization), and CAS as a continuous outcome, showed a similar pattern of two- and three-way interactions (see S1 268 Table). 269

270

265

Rival theoretical models: serially causal and mutually causal 271

epidemics 272

In the model of serially causal epidemics, we estimated a statistically significant total 273 indirect effect of sexual violence (estimate=.04, 95% CI: .003–07; p=.03) on CAS through drug 274 use and alcohol use, but the indirect effect of physical violence was not statistically significant. 275 In the model of mutually causal epidemics, we found that violence victimization had a 276 statistically significant association with drug use (estimate=.18, 95% CI: .15-.22; p<.001) and 277 drug use had a statistically significant association with violence victimization (estimate=1.97, 278 279 95% CI: 1.38–2.56; p<.001). The model fit (SRMR<.02) and the stability index (.60) were good (see S2 Table). 280

281

Discussion 282

In this large, cross-sectional, probability-based analytical sample of HIV-negative 283 transgender women in India, we tested a model of synergistically interacting epidemics, based 284 on the joint effects of four psychosocial exposures-physical and sexual violence 285 victimization, drug use, and alcohol use-on CAS. The high prevalence of these adverse 286

psychosocial exposures, their frequent co-occurrence, and their synergistic interactions support 287 the presence of synergistically interacting epidemics, or a syndemic. On both multiplicative 288 and additive scales, we found evidence of two-way synergistic interactions between physical 289 290 violence victimization and sexual violence victimization, and between physical violence victimization and drug use. We found evidence of three-way synergistic interactions between 291 physical violence victimization, sexual violence, and alcohol use; and between physical 292 293 violence victimization, drug use and alcohol use. Our findings suggest that a harmful socialstructural environment for transgender women in India contributes to clustering of 294 psychosocial health problems significantly associated with increased HIV risk, with 295 ramifications for targeted HIV prevention initiatives. 296

The high levels of physical (27.3%) and sexual violence (22.3%) victimization faced 297 298 by transgender women are corroborated by other studies from India [13,14,24]. A study from 299 Karnataka reported past-year sexual violence prevalence of 18%, but did not disaggregate results among the combined sample of 543 men who have sex with men and transgender 300 301 women [14]. In a study with 300 transgender women in four states, 84% reported having ever experienced physical and/or sexual violence [13]. Notably, in the present study, about one-302 third of those who experienced violence reported multiple perpetrators: family 303 members/relatives, thugs, police, strangers, and sex work clients. Thus, not only do a 304 305 substantial proportion of transgender women face violence; they experience violence across a 306 myriad of settings in their social ecology-family, work, community settings, and in interactions with law enforcement officers, ostensibly positioned to offer protection from 307 violence. A latent class analysis of patterns of violence victimization among transgender 308 309 women in India (n=299) indicated that the type (verbal, physical, or sexual) of violence victimization differentially influenced HIV risk; those who experienced police-perpetrated 310 311 sexual assault vs. police-perpetrated physical violence had relatively higher risk of CAS [24].

However, that study did not assess synergistic effects, such as those identified in the present study between physical and sexual violence, nor their interactions with alcohol and drug use.

Studies among gender and sexual minority populations in other contexts [25,26] 314 indicate that the presence of multiple types and domains of violence victimization is likely to 315 substantially increase psychological distress. The present study further demonstrates that 316 multiple forms of violence victimization against transgender women, and in conjunction with 317 318 drug use and alcohol use, contribute to synergistically increasing HIV risk. In India, physical violence from family members has been shown to be a manifestation of non-acceptance of 319 320 gender identity, with violence reportedly perpetrated as a form of disciplining [27]. Physical and sexual violence from police and thugs in India have been identified as even more common 321 among those transgender women who engage in sex work [28]. The multiple forms and 322 323 domains of violence victimization can be seen as the manifestation of systemic stigma and discrimination undergirded by lack of acceptance of transgender people in the family and 324 society; and this is exacerbated by the lack of legal protections for those who engage in sex 325 work [29]. 326

The associations identified using models of synergistically interacting epidemics or 327 mediational analyses have been demonstrated among transgender women in other countries. 328 For example, studies among transgender women in North and South America indicate that life 329 330 stressors [30], discrimination [31,32] and trauma [33] can lead to alcohol and/or substance use, 331 possibly as maladaptive coping strategies. A longitudinal study among transgender women in the U.S. showed a causal association between sexual violence and drug use [34]. In several 332 other studies with transgender women, alcohol use [35] and drug use [36], including sexualized 333 334 drug use [37], were associated with increased HIV risk. As identified among other populations, such as gay and bisexual men, being under the influence of alcohol or drugs may increase the 335 likelihood of engaging in sex and the risk of CAS [38,39]. Thus, experiences of sexual violence 336

can precipitate drug or alcohol use as a coping mechanism, and alcohol and drug use (especially 337 before or during sex) in turn may increase the likelihood of being targeted or otherwise 338 exacerbate sexual or physical violence from male partners [37], which increases the risk of 339 engaging in CAS. Given evidence that internalized transprejudice and depression may stem in 340 part from discrimination and violence victimization [40,41], future studies should explore the 341 potential mediating roles of internalized transprejudice and depression in the association 342 343 between physical and sexual violence victimization and discrimination, and HIV transmission risk behavior. 344

345

346 Implications for practice and policy

Given that larger societal forces, including negative attitudes and systemic stigma and 347 discrimination against transgender persons, and criminalization of sex work, contribute to 348 349 violence victimization—as well as indirectly leading to alcohol and drug use through internalized stigma and depression [11,29,42]—addressing these social and structural factors 350 is crucial. Currently, 'crisis response' teams established in NACO-supported targeted HIV 351 352 interventions act only *after* incidents of violence are reported; and we are aware of no societallevel campaigns in India designed to prevent discrimination and violence against transgender 353 women in relation to their intersecting marginalized identities (e.g., gender minority and sex 354 355 worker and/or HIV-positive status). Important progress is evidenced in the passage of The Transgender Persons (Protection of Rights) Act, 2019 [43] and Rules, 2020 [44], in India, 356 357 which explicitly states that transgender persons should not be discriminated against and prescribes punishment for those who discriminate or perpetrate violence. Nevertheless, 358 disparities such as punishments for sexual violence against transgender women being less 359 severe than that for cisgender women highlight ongoing challenges with protective policies and 360

the devaluation of transgender women's rights and lives. It is also vital to ensure the implementation of these policies on the ground.

Our findings further indicate that physical violence victimization independently, and 363 together with sexual violence, drug use and alcohol use synergistically increase CAS. That is, 364 in the presence of physical violence victimization, relatively higher levels of CAS were 365 observed with one or more combinations of sexual violence victimization, drug use and/or 366 367 alcohol use. Thus, addressing all four exposures through integrated, multicomponent interventions may substantially reduce HIV risk [45]; however, in the context of synergistic 368 369 interactions, single-component interventions (e.g., post-care support for survivors of sexual violence or treatment of alcohol or drug dependence) may still exert an incremental impact on 370 reducing HIV transmission risk behavior. 371

Partial support for the model of serially causal epidemics-i.e., sexual violence 372 victimization leading to alcohol and drug use, and possible mutually causal associations 373 between violence victimization and drug use-suggest the utility of multilevel interventions: 374 at the individual level, screening for and managing alcohol and drug use; and at the societal 375 level, strengthening stigma and violence reduction or elimination programs. Although alcohol 376 use and drug use were not independently statistically significant predictors of CAS (main 377 effects model), in the presence of sexual and/or physical violence victimization, alcohol use 378 379 and/or drug use substantially increased CAS. Nongovernmental (NGOs) and community-based 380 organizations (CBOs) that implement targeted HIV preventive interventions for transgender 381 women should establish or strengthen referral systems with public hospitals offering treatment for alcohol and drug dependence, post-violence (physical or sexual) support services, and post-382 383 sexual exposure prophylaxis. To this end, training and monitoring healthcare providers to ensure nondiscriminatory and gender-affirmative care is crucial. Similarly, support is required 384 from police in duly filing complaints of sexual or physical violence, given that transgender 385

persons are not covered in current sexual assault/rape laws (Indian Penal Code, Section 354) a pernicious lapse in legal policy. Further, in addition to HIV and condom education and condom distribution, NGOs and CBOs should educate transgender women about the negative effects of alcohol and drug use on sexual decision-making and negotiation, counselling and supporting them to reduce or avoid drug or alcohol use before or during sex.

391

392 Limitations and strengths

393 In addition to its strengths, this study has several limitations. First, the cross-sectional data preclude causal interpretations. However, theoretical frameworks and compatible recall 394 periods for the adverse psychosocial exposures (physical and sexual violence victimization, 395 one year; drug use, one year), support the plausibility of the three models tested. Second, our 396 analysis was restricted to co-occurrence and interactions between exposures at the individual 397 398 level, while the concept of a syndemic is a population-level phenomenon; multilevel analyses at the population level should be undertaken in future research. Nevertheless, this is among the 399 400 largest probability-based studies of syndemics among transgender women [46], and had a high 401 response rate (81%). Our use of appropriate statistical analyses (interactions on both additive and multiplicative scales) to assess synergy is another strength, given several studies that 402 purportedly assessed syndemics and HIV risk among transgender women did not use such 403 404 analytic techniques [46], among very few that did [47]. In line with current practice, we dichotomized the adverse psychosocial exposures in additive and multiplicative interactions 405 [18]; however, sensitivity analyses revealed that similar findings were obtained if we used 406 different ways of specifying exposures, strengthening the evidence for synergy among 407 408 exposures.

409

411 Conclusion

This study found evidence for synergistic (additive and multiplicative) interactions between sexual and physical violence victimization, drug use, and alcohol use on HIV transmission risk behavior among transgender women. Our findings support the need to focus on preventing and eliminating violence victimization, as well as the importance of preventing or mitigating multiple co-occurring adverse psychosocial exposures, in order to reduce HIV risk among transgender women in India.

418 Acknowledgements

- 419 We thank Dr. Arvind Pandey and Ms. Abhina Aher for their helpful comments on an earlier
- 420 version of the manuscript.

422 **References**

- Poteat T, Scheim A, Xavier J, Reisner S, Baral S. Global epidemiology of HIV infection
 and related syndemics affecting transgender people. J Acquir Immune Defic Syndr.
 2016;72(Suppl 3):S210-219.
- 426 2. Hatzenbuehler ML, Pachankis JE. Stigma and minority stress as social determinants of
 427 health among lesbian, gay, bisexual, and transgender youth: Research evidence and
 428 clinical implications. Pediatr Clin North Am. 2016;63(6):985-997.
- Rajan S, Kumar P, Sangal B, Kumar A, Ramanathan S, Ammassari S. HIV/AIDS-Related
 risk behaviors, HIV prevalence, and determinants for HIV prevalence among
 hijra/transgender people in India: Findings from the 2014-2015 integrated biological and
 behavioural surveillance. Indian J Public Health. 2020;64(Supplement):S53-S60.
- 4. NACO. HIV Sentinel Surveillance: Technical Brief, India 2016-17. New Delhi: National
 AIDS Control Organisation, Ministry of Health and Family Welfare, Government of
 India.; 2017.
- 436 5. National AIDS Control Organization (NACO). Sankalak: Status of National AIDS
 437 Response. NACO. 2020. Available from: <u>http://naco.gov.in/nacoevents/release-</u>
 438 <u>sankalak-status-national-aids-response</u>
- 6. Chakrapani V, Willie TC, Shunmugam M, Kershaw TS. Syndemic classes, stigma, and
 sexual risk among transgender women in India. AIDS Behav. 2019;23(6):1518-1529.
- Singer M. A dose of drugs, a touch of violence, a case of AIDS: Conceptualizing the
 SAVA syndemic. Free Inquiry Creative Sociol. 1996;24:99-110.
- Parsons JT, Antebi-Gruszka N, Millar BM, Cain D, Gurung S. Syndemic conditions, HIV
 transmission risk behavior, and transactional sex among transgender women. AIDS
 Behav. 2018;22(7):2056-2067.

- 9. Stall R, Mills TC, Williamson J, Hart T, Greenwood G, Paul J, et al. Association of co-446 occurring psychosocial health problems and increased vulnerability to HIV/AIDS among 447 urban men who have sex with men. Am J Public Health. 2003;93(6):939-942. 448
- 10. Stall R, Friedman M, Catania JA. Interacting epidemics and gay men's health: A theory 449 of syndemic production among urban gay men. In: Wolitski RJ, Stall R, Valdiserri RO,
- editors. Unequal opportunity: health disparities affecting gay and bisexual men in the 451

United States. New York: Oxford University Press; 2008. pp. 251-274. 452

450

- Chakrapani V, Newman PA, Shunmugam M. Stigma toward and mental health of 453 11. 454 hijras/trans women and self-identified men who have sex with men in India. In: Nakamura N, Logie CH, editors. LGBTO mental health: International perspectives and 455 experiences. Washington: American Psychological Association; 2020. pp. 103-119. 456
- 12. Chakrapani V, Vijin PP, Logie CH, Newman PA, Shunmugam M, Sivasubramanian M, 457 et al. Understanding how sexual and gender minority stigmas influence depression among 458 trans women and men who have sex with men in India. LGBT Health. 2017;4(3):217-459 226. 460
- Chakrapani V, Newman PA, Shunmugam M, Logie CH, Samuel M. Syndemics of 13. 461 depression, alcohol use, and victimisation, and their association with HIV-related sexual 462 risk among men who have sex with men and transgender women in India. Glob Public 463 Health. 2017;12(2):250-265. 464
- 14. Shaw SY, Lorway RR, Deering KN, Avery L, Mohan HL, Bhattacharjee P, et al. Factors 465 associated with sexual violence against men who have sex with men and transgendered 466 individuals in Karnataka, India. PLoS One. 2012;7(3):e31705. 467
- 15. Tsai AC, Venkataramani AS. Syndemics and health disparities: A methodological note. 468 AIDS Behav. 2016;20(2):423-30. 469
- 16. VanderWeele TJ, Knol MJ. A tutorial on interaction. Epidemiol Methods. 2014;3:33-72. 470

- 471 17. National AIDS Control Organization (NACO). National IBBS 2014-15:
- 472 Hijras/Transgender people. NACO. 2016. Available from:
 473 http://naco.gov.in/sites/default/files/TG-IBBS%20ReportPrint%20text Edited.pdf
- 474 18. VanderWeele TJ. Explanation in causal inference: Methods for mediation and
 475 interaction. New York: Oxford University Press; 2015.
- 476 19. Solon G, Haider SJ, Wooldridge JM. What are we weighting for? J Hum Resour.
 477 2015;50(2):301-316.
- 478 20. Heeringa SG, West BT, Berglund PA. Applied survey data analysis. New York: Chapman
 479 and Hall/CRC; 2010.
- 480 21. Knol MJ, VanderWeele TJ. Recommendations for presenting analyses of effect
 481 modification and interaction. Int J Epidemiol. 2012;41(2):514-520.
- 482 22. Tsai AC. Syndemics: A theory in search of data or data in search of a theory? Soc Sci
 483 Med. 2018;206:117-122.
- Kuh D, Ben-Shlomo Y, Lynch J, Hallqvist J, Power C. Life course epidemiology. J
 Epidemiol Community Health. 2003;57(10):778-783.
- Willie T, Chakrapani V, White Hughto J, S. Kershaw T. Victimization and human
 immunodeficiency virus-related risk among transgender women in India: A latent profile
 analysis. Violence Gend. 2017;4(4):121-9.
- 489 25. Newman PA, Fantus S, Woodford MR, Rwigema M-J. "Pray that God will change you":
- 490 The religious social ecology of bias-based bullying targeting sexual and gender minority
- 491 youth—A qualitative study of service providers and educators. J Adolesc Res.
 492 2017;33(5):523-548.
- 26. Sterzing PR, Ratliff GA, Gartner RE, McGeough BL, Johnson KC. Social ecological
 correlates of polyvictimization among a national sample of transgender, genderqueer, and
 cisgender sexual minority adolescents. Child Abuse Negl. 2017;67:1-12.

- Chakrapani V, Dhall P. Family acceptance among self-identified men who have sex with
 men (MSM) and transgender people in India. Mumbai: Family Planning Association of
 India; 2011.
- 499 28. Ganju D, Saggurti N. Stigma, violence and HIV vulnerability among transgender persons
 500 in sex work in Maharashtra, India. Cult Health Sex. 2017;19(8):903-917.
- 29. Chakrapani V, Newman AP, Noronha E. Hijras/Transgender women and sex work in
 India: From marginalization to social protection. In: Nuttbrock LA, editor. Transgender
 sex work & society. New York: Harrington Park Press; 2018. pp. 214-235.
- 30. Hotton AL, Garofalo R, Kuhns LM, Johnson AK. Substance use as a mediator of the
 relationship between life stress and sexual risk among young transgender women. AIDS
 Educ Prev. 2013;25(1):62-71.
- 507 31. Ehlinger PP, Folger A, Cronce JM. A qualitative analysis of transgender and gender
 508 nonconforming college students' experiences of gender-based discrimination and
 509 intersections with alcohol use. Psychol Addict Behav. 2021.
- 510 32. Kcomt L, Evans-Polce RJ, Boyd CJ, McCabe SE. Association of transphobic
 511 discrimination and alcohol misuse among transgender adults: Results from the U.S.
 512 Transgender Survey. Drug Alcohol Depend. 2020;215:108223.
- Johnson EEH, Wilder SMJ, Andersen CVS, Horvath SA, Kolp HM, Gidycz CA, et al.
 Trauma and alcohol use among transgender and gender diverse women: An examination
 of the stress-buffering hypothesis of social support. J Prim Prev. 2021;42(6):567-581.
- Smith LR, Yore J, Triplett DP, Urada L, Nemoto T, Raj A, et al. Impact of sexual violence
 across the lifespan on HIV risk behaviors among transgender women and cisgender
 people living with HIV. J Acquir Immune Defic Syndr. 2017;75(4):408-416.
- 519 35. Hearld KR, Milner AN, Budhwani H, Abreau N, Rodriguez-Lauzurique RM, Charow R,
- 520 et al. Alcohol use, high risk behaviors, and experiences of discrimination among

transgender women in the Dominican Republic. Subst Use Misuse. 2019;54(10):1725-

522 1733.

- 36. Avila MM, Dos Ramos Farias MS, Fazzi L, Romero M, Reynaga E, Marone R, et al.
 High Frequency of illegal drug use influences condom use among female transgender sex
 workers in Argentina: Impact on HIV and syphilis infections. AIDS Behav.
 2017;21(7):2059-2068.
- 527 37. Fan X, Lau JTF, Cai Y, Li J, Ma T, Gu J, et al. Prevalence and associated factors of
 528 sexualized drug use in sex work among transgender women sex workers in China. AIDS
 529 Care. 2021;33(8):1098-1106.
- 38. Redina HJ, Moody RL, Ventuneac A, Grov C, Parsons JT. Aggregate and event-level
 associations between substance use and sexual behavior among gay and bisexual men:
 Comparing retrospective and prospective data. J Alcohol Drug Depend. 2015;154:199-

533 207.

- Theodore PS, Duran RE, Antoni MH. Drug use and sexual risk among gay and bisexual
 men who frequent party venues. AIDS Behav. 2014;18(11):2178-86.
- Kaighobadi F, Collier KL, Reddy V, Lane T, Sandfort TGM. Sexual violence experiences
 among black gay, bisexual, and other men who have sex with men and transgender
 women in South African townships: Contributing factors and implications for health. S
 Afr J Psychol. 2020;50(2):170-182.
- Logie CH, Lacombe-Duncan A, Poteat T, Wagner AC. Syndemic factors mediate the
 relationship between sexual stigma and depression among sexual minority women and
 gender minorities. Womens Health Issues. 2017;27(5):592-599.
- 42. Meyer IH. Prejudice, social stress, and mental health in lesbian, gay, and bisexual
 populations: Conceptual issues and research evidence. Psychol Bull. 2003;129(5):674697.

546	43.	Government of India.	The Transgender	Persons (Protection	of Rights) A	ct, 2019. No. 40
-----	-----	----------------------	-----------------	---------------------	--------------	------------------

- 547 December 5, 2019. <u>https://www.indiacode.nic.in/bitstream/123456789/13091/1/a2019-</u>
 548 40.pdf
- 549 44. Gove Government of India. The Transgender Persons (Protection of Rights) Rules, 2020.
- 550 <u>https://thc.nic.in/Central%20Governmental%20Rules/Transgender%20Persons%20(Pro</u>
 551 tection%20of%20Rights)%20Rules,%202020.pdf
- 45. Operario D, Nemoto T. HIV in transgender communities: Syndemic dynamics and a need
- for multicomponent interventions. J Acquir Immune Defic Syndr. 2010;55(Suppl 2):S91-
- 554 S93.
- 46. Tsai AC, Burns BF. Syndemics of psychosocial problems and HIV risk: A systematic
- review of empirical tests of the disease interaction concept. Soc Sci Med. 2015;139:26-
- 557 35.