

1 Title Page

2 Title: **Effect of fasting therapy on vitamin D, vitality and quality of life. A randomized**
3 **control trial**

4 **Category:** Original Article

5 **Running Title:** Prolonged fasting and Vitamin D

6 **Authors:** Gulab Rai Tewani¹, Karishma Silwal², Gita Sharma², Dinesh Yadav², Aarfa Siddiqui²,
7 Sucheta Kriplani², Varsha Vijay Nathani², Neha Sharma³, Jyoti Keswani⁴, Himanshu Sharma⁵,
8 Pradeep M.K. Nair^{6*}

- 9 1- Chief Medical Officer- Sant Hirdaram Yoga and Nature Cure Hospital, Bhopal
10 2- BNYS[#], Medical Officer, Sant Hirdaram Yoga and Nature Cure Hospital, Bhopal
11 3- BPT, Physiotherapist, Sant Hirdaram Yoga and Nature Cure Hospital, Bhopal
12 4- BNYS[#], Ph.D^α – Professor & Head, Yoga, Sant Hirdaram Medical College of
13 Naturopathy & Yogic Sciences, Bhopal
14 5- BNYS[#], MPH[§], Principal, Sant Hirdaram Medical College of Naturopathy & Yogic
15 Sciences, Bhopal
16 6- BNYS[#], MSc^Ω, Ph.D^α Scholar, Professor & Head, Research, Sant Hirdaram Medical
17 College of Naturopathy & Yogic Sciences, Bhopal

18 *Corresponding author

19 [#] Bachelor of Naturopathy and Yogic Sciences

20 ^Ω Masters in Science

21 ^α Doctorate in Philosophy

22 [§] Masters in Public Health

23

24

25 **Corresponding authors Address:**

26 Dr. Pradeep MK Nair,

27 Professor & Head, Research,

28 Sant Hirdaram Medical College of Naturopathy & Yogic Sciences,

29 Sant Hirdaram Nagar, Bhopal-462020

30 drpradeep18bnys@gmail.com

31 **Author contribution statement:** Conceptualization: PMK, HS,GRT, KS, VVN, DY,; Data
32 curation: PMK, SK, KS,KG, AS Formal analysis: PMK, KS,HS, GRT; Investigation:
33 PMK,GS,MJ,SK,AS,DY,JK; Methodology: PMK, GRT, HS, KS,VVN,DY; Project
34 administration: PMK, KS,KG, NS; Resources: GRT, HS; Supervision: PMK,HS, GRT;
35 Visualization: PMK,GRT, HS; Writing - original draft: PMK; Writing - review & editing: PMK,
36 KS, HS, GRT, SK,GS,JK,NS.

37 **Funding:** No funds were received for this study

38 **Conflicts of interest:** None declared

39 **Ethics approval :** The study was approved by the institutional ethics committee of Sant
40 Hirdaram Medical College for Naturopathy & Yogic Sciences, Bhopal, India.

41 **Consent to participate :** All the participants signed a written consent to express their consent to
42 participate in the study.

43 **Consent to Publish:** All the participants consented to publish their de-identified data on a
44 medical journal.

45 **Availability of data and material:** The data will be made available on request to the
46 corresponding author.

48 **Effect of fasting therapy on vitamin D, vitality and quality of life. A randomized control**
49 **trial**

50 **Abstract**

51 **Background**

52 The aim of the present study was to determine the effects of prolonged fasting (10 days) in the
53 vitamin D, B12 levels, body mass index (BMI), weight, hemoglobin, vitality and quality of life
54 (QoL) compared to normal diet.

55 **Methods**

56 This randomized control trial included 52 participants (aged 19-74 years) randomized in to a
57 fasting group (FG) or a normal diet group (NDG) with 26 participants in each group. The study
58 was conducted at an in-patient setting where the FG were on a fasting diet (500 kCal/day) which
59 included holy basil herbal tea, lemon honey juice and water (3 L). The NDG (1500 kCal/day)
60 consumed routine diet that included Indian breads, pulses, steamed rice, vegetable salads and
61 beverages.

62 **Results**

63 The FG has shown significant increase in the Vitamin D levels ($p=0.003$, $d=0.475$), vitality
64 ($p=0.006$, $d=0.425$), physical QoL ($p<0.001$, $d=0.549$), psychological QoL ($p=0.002$, $d=0.488$),
65 environmental QoL ($p=0.004$, $d=0.457$) compared to NDG. No significant changes were
66 observed in Vitamin B12, weight, BMI, hemoglobin and social QoL. A weak to moderate ($p=$
67 $0.330-0.483$) positive correlation was observed between vitality scores and QoL domains,
68 whereas BMI scores showed an inverse correlation ($p=-0.280$) with vitamin D levels.

69 **Discussion**

70 The results suggest that prolonged fasting can improve the vitamin D levels, vitality and promote
71 quality life compared to normal diet. Unlike previous studies FG does not differ from NDG with
72 respect to weight and BMI. Nevertheless, fasting may be utilized as an effective tool to tackle
73 vitamin d deficiency and associated health insufficiencies.

74 **Keywords:** Vitamin D deficiency; Fasting; Diet; Therapeutic fasting; Vitality; Quality of life

75 **Trial Registry:** Clinical Trial Registry of India CTRI/2022/02/040446.

77 **Introduction**

78 Therapeutic fasting is increasingly becoming popular among scientists, physicians and patients
79 owing to its health benefits. The role of fasting in the prevention and management of various
80 cardio-metabolic, musculoskeletal disorders are widely reckoned.(1)Majority of the reports on
81 fasting have reported the efficacy of intermittent fasting (lasting for 16-48 hours) or calorie
82 restriction but not on prolonged fasting (more than 4 days) barring few studies.(2,3) Prolonged
83 fasting is defined as a medically supervised fasting regimen that lasts for a minimum of 7 days to
84 21 days with an average calorie intake of (200–500 kcal nutritional intake/day).(4) Fasting has
85 potential therapeutic benefits as it has shown to reduce body weight,(5) body mass index (BMI),
86 adiposity,(6) inflammation,(7) blood pressure, cholesterol, regulate blood sugar, insulin, glycated
87 hemoglobin(8) etc. Hence fasting is viewed as a therapy with homoeostatic potential.

88 Vitamin D deficiency is reckoned as one of the growing public health issue worldwide reportedly
89 affecting 30-80% of global population.(9)Similarly, the role of vitamin D from a mere
90 micronutrient to an essential element in preserving health is been substantiated by numerous
91 observational studies, which links vitamin D deficiency as one of prime precursor in cancers,
92 immune disorders, metabolic disorders and neurodegenerative studies.(10)While Vitamin D
93 supplementation is considered as a probable measure in alleviating Vitamin D deficiency,
94 previous data suggests that a hypocaloric diet induced weight loss can improve the vitamin D
95 status among patients.(11)Recently Arankale et al and Żychowska et al has reported that fasting
96 can improve the circulating levels of vitamin D and has an impact on its metabolites as
97 well.(3,12)However, both these studies were conducted on healthy volunteers without any
98 controls.

99 It will be interesting to study the impact of prolonged fasting on the vitamin D levels of
100 participants with different diseases, as fasting is widely practiced both as a therapeutic, cultural
101 and religious practice. Further, the association of prolonged fasting with vitality and quality of
102 life is an unexplored domain. Fewer studies have reported fasting to act as a mood enhancer in
103 pain,(8) moderately relieve stress, anxiety, depression(13) and offers rewarding psychological
104 experiences.(14) However, all the previous reports necessitate the need for future randomized
105 controlled trials. Further barring Ramadan fasting and calorie restriction, the impact of prolonged
106 fasting on health indices like weight, BMI etc were not studied on individuals with various
107 medical co-morbidities.

108 The study hypothesized that a 10 days prolonged fasting program will increase the vitamin D
109 levels, Vitamin B12 levels, enhance the vitality, quality of life, reduce body weight and BMI.

110 **Methods**

111 *Study settings and design:* This study was conducted at a private yoga and naturopathy medical
112 college hospital in India. The study approved by institutional ethics committee via
113 (F.No:12/SHMCNYS-IEC/P45/2021-2022) and was registered in Clinical Trial Registry of India
114 CTRI/2022/02/040446. Written informed consent was obtained from all participants prior to
115 participation in the study. This was a pragmatic, parallel group, randomized controlled trial
116 (n=52). Eligible participants were randomized by the study coordinator at a ratio of 1:1 using
117 computer random software in to fasting arm or normal diet arm. Figure 1 depicts the trial profile.

118 **Figure 1: Trial Profile**

119 *Study Participants:* The study participants were those volunteers with diverse medical conditions
120 enrolled for a 10 days lifestyle modification program at the in-patient department of a private

121 yoga and naturopathy medical college hospital. Both men and women aged between 20 to 75
122 years participated in this program. All the patients who were underweight, with known
123 nutritional deficiency, psychiatric disorders were excluded from participating in this study. Any
124 participants who have a previous history of fasting for more than 10 days in the last 6 months
125 were also excluded from the study. Participants were also excluded if they are a part of any other
126 clinical trial or following any dietary regimens/food supplements.

127 **Sample size:** Based on a previous pilot study,(3) for an effect size of 0.8, level of significance
128 (α)=0.05 and power 80% the sample size was calculated to be n=40 by using G Power software.
129 Considering a drop out of 30% the final sample size was n=52 (n=26 in each arm).

130 **Interventions:** The fasting arm received 10 days of fasting regimen which includes two days of
131 feeding (preparation day 1 and re-feeding day 10) that included herbal tea (100 mL), lemon
132 honey juice(100 mL), cucumber juice (200 mL), boiled green grams (75gms), Rice and lentil
133 porridge (100 g), Plain vegetable soup (150 mL), Water (3 L) at different intervals during the
134 day, which amounts for 1000 kCal/day approximately. For the rest of 8 days (day 2 to day 9) the
135 participants were on a fasting diet (500 kCal/day) which included holy basil herbal tea (100 mL),
136 lemon honey water (1,000 mL), and water (3 L). The normal diet arm consumed routine diet that
137 included Indian breads, pulses, steamed rice, vegetable salads and beverages that amounts to
138 1500 kCal/day. In addition to the dietary regimen both the study group and the control group
139 received yoga and naturopathy therapies like hip bath, cold water enema, and mud packs to the
140 abdomen and eyes which are intended to promote elimination.(3)

141 **Assessments:** Blood samples were collected from both the study group at baseline and at the 10th
142 day after intervention to evaluate the changes in the serum levels of vitamin D, vitamin B12 and

143 hemoglobin. Additionally, we evaluated the changes in weight, body mass index, quality of life
144 and vitality. World Health Organization Quality of Life (WHO-QoL BREF)(15) instrument was
145 used to assess the quality of life, whereas vitality (defined as the innate power to heal and to be
146 resilient)(16) was measured on a scale of 1 to 10, using a visual analogue scale where ‘0’
147 represents poor vitality and ‘10’ represents higher vitality.

148 **Statistical analysis:** All the data were analyzed using Jeffreys's Amazing Statistics Program
149 (JASP) version 0.16. Shapiro’s Wilk’s test was used to test the normality of the data. Wilcoxon
150 sign-rank test (within the group testing) and Mann-Whitney test (between the group testing) were
151 used to analyze the non-normally distributed data, where as paired T test and Independent T test
152 were used for analyzing the normally distributed samples. A P-value of 0.05 is considered as a
153 significant change.

154 **Results**

155 Of the 103 individuals screened for the trial, n=52 were randomized to the fasting (n=26) or
156 control groups (n=26). All the 52 participants completed the study without any adverse events.
157 The baseline characteristics are tabulated in table 1. The changes in the body weight (p= 0.51),
158 BMI (P=0.99), vitamin B12 (P=0.40) and hemoglobin (P=0.77) were not significant in the
159 fasting group compared to the control group. However, a within the group analysis has shown
160 that the changes in weight and BMI were significant (P < .001) before and after the intervention
161 in both the fasting and control group. However, the changes in vitamin B12 and hemoglobin
162 remained non-significant in both the groups. The mean changes in these parameters from the
163 baseline to end point are tabulated in table 2.

164

165 **Table 1: Baseline demographic characteristics of the participants**

Parameters assessed	Fasting Group (n=26)	Control group(n=26)
Age	53.57±10.21	49.96±15.62
Sex	Males-11 Females-15	Males-11 Females-15
Height	159.65±9.68	158.92±9.55
Weight	67.9±10.53	70.88±28.46
Body Mass Index	26.86±4.71	27.8±9.11
Socioeconomic status	Upper Class-9 Middle class-17	Upper class-1 Upper Middle class-2 Middle class- 23
Habits	Smoking-1	Smoking-1
Medications	Under Medications-9 No Medication-17	Under Medications-14 No Medication-12
Conditions	Diabetes-2 Dyslipidemia-1 Gastrointestinal disorders-2 Hypertension-5 Hyperthyroidism-1 Hypothyroidism-1 Insomnia-1 Lumbago-3 Osteoarthritis-2 Obesity-5 Skin disorders-3	Asthma-2 Diabetes-6 Fatty Liver-1 Gastrointestinal disorder-3 Hypertension-5 Lumbago-1 Migraine-1 Osteoarthritis-3 Obesity-4

167 **Table 2: Comparison of variables between the fasting and control group**

Variables assessed	Fasting group		Within group analysis p value	Normal diet group		Within group analysis p value	Between group analysis p value	Effect size
	Pre (Mean±SD)	Post (Mean±SD)		Pre (Mean±SD)	Post (Mean±SD)			
Weight (kg)	67.9±10.53	65.38±10.64	<0.001	70.88±28.46	68.68±28.16	<0.001	0.57 [^]	0.155
Body mass index (kg/m²)	26.86±4.71	25.99±4.53	<0.001	27.8±9.11	26.92±8.99	<0.001	0.63 [^]	0.131
Vitamin D (ng/ml)	25.69±10.66	31.5±11.13	<0.001	20.92±8.89	22.4±11.29	0.23	0.003 [^]	0.475
Vitamin B12 (pg/ml)	384.42±336.43	365.5±207.92	0.75	355.91±224.9	358.26±140.51	0.25	0.40 [^]	0.136
Hemoglobin (gm/dl)	13.12±1.14	13.04±1.09	0.91	12.9±1.39	12.95±1.34	0.72	0.77 [*]	0.07
Vitality	6.73±1.18	7.61±1.02	<0.001	5.42±1.39	8.46±0.9	<0.001	0.006 [^]	0.425
WHO-Quality of life (QoL)								
Physical QoL	70.38±14.45	82.5±11.85	<0.001	55.61±17.23	68.8±13.64	<0.001	<0.001 [^]	0.549
Psychological QoL	71.03±15.01	83.26±14.09	<0.001	59.46±15.17	70.73±15.66	0.001	0.002 [^]	0.488
Social QoL	78.46±13.6	82.88±13.56	0.02	72.15±18.7	73.34±19.53	0.42	0.06 [^]	0.302
Environment QoL	84.53±11.34	88.26±11.88	0.01	71.53±16.05	77.92±13.69	0.006	0.004 [^]	0.457

168 *Students T test; [^]Mann-Whitney U test, QoL-Quality of Life

169

170 Significant increase in the levels of vitamin D (P=0.003) was observed in fasting group compared to the control group. Similarly,
 171 statistically significant increase was observed in vitality scores (P= 0.003) of the fasting group when compared to the controls. A
 172 within group analysis between the mean values before and after the intervention has shown a significant rise in the vitamin D levels (P

173 < .001) and vitality ($P < .001$) in the fasting group. However, the control group has shown
174 significant increase only in the vitality scores ($P < .001$) whereas the changes in vitamin D in this
175 group remained insignificant ($P=0.23$). The changes are depicted in figure 2.

176 **Figure 2: Changes in the mean values of outcome measures between the fasting and control**
177 **group**

178 All the participants in the fasting group has shown significant improvement in the quality of life
179 domains of WHOQOL namely physical domain ($P < .001$), psychological domain ($P=0.002$),
180 environmental domain ($P=0.004$) except social domain ($P=0.06$) compared to their controls.
181 Nevertheless, a within group analysis has shown significant improvement in all the QOL
182 domains in the fasting group and control group, except for psychological domain ($P=0.09$) in the
183 control group. The detailed results are tabulated in table 2.

184 Spearman's correlation has shown a weak negative correlation between BMI and Vitamin D
185 levels ($\rho=-0.280, P=0.04$). Vitality status was reported to have a moderate positive correlation
186 with physical QoL ($\rho= 0.442, P=0.001$) and environmental QoL ($\rho= 0.483, P< .001$). Further
187 vitality status also demonstrated to have a weak positive correlation with psychological QoL ($\rho=$
188 $0.348, P=0.001$) and social QoL ($\rho= 0.330, P=0.01$). The correlations between these variables
189 are plotted in figure 3 and 4.

190 **Figure3: Relationship between Vitamin D level and Body mass index**

191 **Figure 4: Correlation between Vitality and Quality of life**

192 **Discussion**

193 This study was conducted to validate the findings of our previous pilot study which reported 10
194 days fasting significantly improve the levels of vitamin D and B12, weight reduction, reduces

195 BMI and prime the other physiological functions among female healthy volunteers.(3) This
196 randomized controlled trial indicates a positive role of fasting therapy over the vitamin D levels,
197 quality of life and vitality of the patients. The present study demonstrated an increase in the
198 vitamin D levels after fasting compared to normal diet. This is strengthening the observations
199 from earlier studies which reported 8-10 days of fasting can increase the levels of serum vitamin
200 D.(3,12)

201 Earlier studies postulate that a reduction in body weight during fasting and/or exercise as the rate
202 limiting factor that stimulates the redistribution of vitamin D stored in the adipose tissue and
203 skeletal muscles.(3,11,12,17) In the present study, participants from both the groups had reported
204 to have significant reduction in their weight and BMI. Compared to the baseline, fasting group
205 had an average of 3.7% reduction in weight and 0.25% reduction in the BMI scores, whereas the
206 control group had 3.1% and 0.23% reduction in weight and BMI respectively.

207 Statistically fasting group did not differ from that of the control group with respect to weight and
208 BMI scores. This is in contrary to the results reported by a recent systematic review that has
209 shown intermittent fasting to reduce body weight.(18) A previous observational study reported
210 significant reduction in weight on a 20 day fasting program.(2) However, the absence of a
211 control group in this study limits its generalizability. To our knowledge this is the first
212 randomized control trial to compare the impact of prolonged fasting (10 days) with normal diet
213 on body weight in a controlled residential setting.

214 The present data suggests that the changes in vitamin D level post fasting may be independent of
215 reduction in weight or BMI. However, a correlation analysis suggests that reduction in BMI was
216 associated with an increase in the vitamin D levels. This is strengthening the earlier reports that

217 vitamin D levels are inversely proportional to the body weight and BMI.(19) Adequate vitamin
218 D levels are warranted in humans, as vitamin D is necessitated in all major functions of the body
219 and mind.(20) Reduced levels of vitamin D predisposes to obesity, insulin resistance and other
220 metabolic disorders.(21) Further optimum levels of vitamin D is considered as a pre-requisite in
221 treating and preventing diseases ranging from diabetes to cancer.(22)As on date vitamin D
222 deficiency and insufficiency is a global concern. In this scenario fasting therapy may be a
223 promising solution for not only mitigating vitamin D deficiency but also can prevent its
224 ramifications.

225 We observed 7% reduction in the vitamin B12 levels in the fasting group, whereas the control
226 group had an increase of 1% in their vitamin B12 levels. However, these changes were not
227 significant. The present study contradict the results of the previous pilot study which reported
228 fasting to increase the vitamin B12 levels on a group of healthy volunteers.(3)Vitamin B12
229 shares an inverse relationship with weight,(23) the insignificant changes in the weight of the
230 study participants may be a reason for the insignificant changes in vitamin B12 levels.
231 Nevertheless, this needs to be revalidated by future studies.

232 Another significant change observed in the present study was the improvement in the vitality
233 status and the quality of life in the fasting group. Vitality is reckoned as one of the important
234 component in measuring one's abilities irrespective of his/her disabilities.(24)There is a growing
235 interests in measuring the vitality as one of the indicator of well-being among various medical
236 disciplines.(25–27)Similarly, improving the quality of life remains as one the primarily sought
237 endpoint in health care delivery and research.(28)

238 The present data suggests a compelling positive relationship between the vitality status and the
239 quality of life. Therefore, understanding and prodding the vitality status among the patients may
240 be a key factor to be considered in future research and clinical practice, as the paradigm of health
241 care and research is shifting towards a patient-centric approach. However, there is a need for a
242 better scale to measure vitality as VAS scoring may not return an accurate measure.

243 ***Limitations of the study***

244 The status of vitamins D, B12, weight, BMI, vitality and quality of life were not measured post
245 fasting program. This remains as one of the limitations of this study as the changes in these
246 parameters during re-feeding phase in an uncontrolled atmosphere will add more information
247 about the sustainability of the results. Further, due to operational constraints at the clinical
248 setting, the authors could not control for the difference in disease status and medication use of
249 the participants, which may be viewed as a potential confounder in this study. The authors have
250 not assessed the specific disease related effects of fasting in these participants which may be
251 included in future studies. Irrespective of the limitations, this is the first randomized controlled
252 trial to report an association between therapeutic fasting and vitamin D.

253 **Conclusion**

254 Prolonged fasting up to 10 days has demonstrated to be safe and helps in improving the vitamin
255 D levels, vitality and quality of life of patients. This may have considerable preventive and
256 prophylactic benefits. Future studies should evaluate the long-term effects and cost effectiveness
257 of fasting as a public health tool.

258 **Conflicts of interest**

259 The authors declare that there are no conflicts of interest.

260 **References:**

- 261 1. Mattson MP, Longo VD, Harvie M. Impact of intermittent fasting on health and disease
262 processes. *Ageing Res Rev* [Internet]. 2017 Oct 1 [cited 2022 Mar 12];39:46–58.
263 Available from: <https://pubmed.ncbi.nlm.nih.gov/27810402/>
- 264 2. De Toledo FW, Grundler F, Bergouignan A, Drinda S, Michalsen A. Safety, health
265 improvement and well-being during a 4 to 21-day fasting period in an observational study
266 including 1422 subjects. *PLoS One* [Internet]. 2019 Jan 1 [cited 2022 Mar 12];14(1).
267 Available from: </pmc/articles/PMC6314618/>
- 268 3. Arankalle D, Tewani GR, Nair PM, Wardle J. Outcomes of a Medically Supervised
269 Fasting Module on Healthy Females in a Controlled Residential Environment: A Brief
270 Report. *Perm J* [Internet]. 2021 Dec 6 [cited 2022 Mar 3];25(4). Available from:
271 </pmc/articles/PMC8784085/>
- 272 4. Michalsen A. Prolonged Fasting as a Method of Mood Enhancement in Chronic Pain
273 Syndromes: A Review of Clinical Evidence and Mechanisms. *Curr Pain Headache*
274 *Reports* 2010 142 [Internet]. 2010 Mar 10 [cited 2022 Mar 14];14(2):80–7. Available
275 from: <https://link.springer.com/article/10.1007/s11916-010-0104-z>
- 276 5. Varady KA, Bhutani S, Church EC, Klempel MC. Short-term modified alternate-day
277 fasting: a novel dietary strategy for weight loss and cardioprotection in obese adults. *Am J*
278 *Clin Nutr* [Internet]. 2009 Nov 1 [cited 2022 Mar 12];90(5):1138–43. Available from:
279 <https://pubmed.ncbi.nlm.nih.gov/19793855/>

- 280 6. Klempel MC, Kroeger CM, Bhutani S, Trepanowski JF, Varady KA. Intermittent fasting
281 combined with calorie restriction is effective for weight loss and cardio-protection in
282 obese women. *Nutr J* [Internet]. 2012 [cited 2022 Mar 12];11(1):98. Available from:
283 </pmc/articles/PMC3511220/>
- 284 7. Faris MAIE, Kacimi S, Al-Kurd RA, Fararjeh MA, Bustanji YK, Mohammad MK, et al.
285 Intermittent fasting during Ramadan attenuates proinflammatory cytokines and immune
286 cells in healthy subjects. *Nutr Res* [Internet]. 2012 Dec [cited 2022 Mar 12];32(12):947–
287 55. Available from: <https://pubmed.ncbi.nlm.nih.gov/23244540/>
- 288 8. Harvie MN, Pegington M, Mattson MP, Frystyk J, Dillon B, Evans G, et al. The effects of
289 intermittent or continuous energy restriction on weight loss and metabolic disease risk
290 markers: a randomised trial in young overweight women. *Int J Obes (Lond)* [Internet].
291 2011 May [cited 2022 Mar 12];35(5):714. Available from: </pmc/articles/PMC3017674/>
- 292 9. Wakayo T, Belachew T, Vatanparast H, Whiting SJ. Vitamin D Deficiency and Its
293 Predictors in a Country with Thirteen Months of Sunshine: The Case of School Children
294 in Central Ethiopia. *PLoS One* [Internet]. 2015 Mar 30 [cited 2022 Apr
295 5];10(3):e0120963. Available from:
296 <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0120963>
- 297 10. Lockau L, Atkinson SA. Vitamin D's role in health and disease: How does the present
298 inform our understanding of the past? *Int J Paleopathol*. 2018 Dec 1;23:6–14.
- 299 11. Tzotzas T, Papadopoulou FG, Tziomalos K, Karras S, Gastaris K, Perros P, et al. Rising
300 serum 25-hydroxy-vitamin D levels after weight loss in obese women correlate with
301 improvement in insulin resistance. *J Clin Endocrinol Metab* [Internet]. 2010 [cited 2022

- 302 Mar 9];95(9):4251–7. Available from: <https://pubmed.ncbi.nlm.nih.gov/20534751/>
- 303 12. Żychowska M, Rola R, Borkowska A, Tomczyk M, Kortas J, Anczykowska K, et al.
304 Fasting and Exercise Induce Changes in Serum Vitamin D Metabolites in Healthy Men.
305 Nutr 2021, Vol 13, Page 1963 [Internet]. 2021 Jun 8 [cited 2022 Mar 9];13(6):1963.
306 Available from: <https://www.mdpi.com/2072-6643/13/6/1963/htm>
- 307 13. Berthelot E, Etchecopar-Etchart D, Thellier D, Lancon C, Boyer L, Fond G. Fasting
308 interventions for stress, anxiety and depressive symptoms: A systematic review and meta-
309 analysis. Nutrients [Internet]. 2021 Nov 1 [cited 2022 Mar 14];13(11). Available from:
310 </pmc/articles/PMC8624477/>
- 311 14. Watkins E, Serpell L. The Psychological Effects of Short-Term Fasting in Healthy
312 Women. Front Nutr [Internet]. 2016 Aug 22;3:27. Available from:
313 <https://pubmed.ncbi.nlm.nih.gov/27597946>
- 314 15. World Health Organization's Quality of Life group: Measuring Quality of Life.
315 Development of the World Health Organization Quality of Life Instrument (WHOQOL).
316 1992.
- 317 16. Pradeep MK Nair. The Scientific Naturopath: A leap into the evidence behind naturopathy
318 phislophies. 1 Edition. Noida: Bluerose Publishers; 2022.
- 319 17. Mieszkowski J, Stankiewicz B, Kochanowicz A, Niespodziński B, Kowalik T, Żmijewski
320 M, et al. Ultra-Marathon-Induced Increase in Serum Levels of Vitamin D Metabolites: A
321 Double-Blind Randomized Controlled Trial. Nutr 2020, Vol 12, Page 3629 [Internet].
322 2020 Nov 25 [cited 2022 Mar 9];12(12):3629. Available from:

- 323 <https://www.mdpi.com/2072-6643/12/12/3629/htm>
- 324 18. Welton S, Minty R, O’Driscoll T, Willms H, Poirier D, Madden S, et al. Intermittent
325 fasting and weight loss: Systematic review. *Can Fam Physician* [Internet]. 2020 Feb 1
326 [cited 2022 Mar 12];66(2):117. Available from: [/pmc/articles/PMC7021351/](https://pubmed.ncbi.nlm.nih.gov/327021351/)
- 327 19. Zakharova I, Klimov L, Kuryaninova V, Nikitina I, Malyavskaya S, Dolbnya S, et al.
328 Vitamin D insufficiency in overweight and obese children and adolescents. *Front*
329 *Endocrinol (Lausanne)*. 2019;10(MAR):103.
- 330 20. Wierzbicka J, ... AP-AB, 2014 undefined. The renaissance of vitamin D.
331 ojs.ptbioch.edu.pl [Internet]. [cited 2022 Mar 9]; Available from:
332 <http://ojs.ptbioch.edu.pl/index.php/abp/article/view/1830>
- 333 21. Chacko SA, Song Y, Manson JAE, Van Horn L, Eaton C, Martin LW, et al. Serum 25-
334 hydroxyvitamin D concentrations in relation to cardiometabolic risk factors and metabolic
335 syndrome in postmenopausal women. *Am J Clin Nutr* [Internet]. 2011 Jul 1 [cited 2022
336 Mar 9];94(1):209–17. Available from: <https://pubmed.ncbi.nlm.nih.gov/21613558/>
- 337 22. Herman RH, Holick MF. Vitamin D: importance in the prevention of cancers, type 1
338 diabetes, heart disease, and osteoporosis. *Am J Clin Nutr* [Internet]. 2004 Mar 1 [cited
339 2022 Mar 9];79(3):362–71. Available from:
340 <https://academic.oup.com/ajcn/article/79/3/362/4690120>
- 341 23. Sun Y, Sun M, Liu B, Du Y, Rong S, Xu G, et al. Inverse Association Between Serum
342 Vitamin B12 Concentration and Obesity Among Adults in the United States. *Front*
343 *Endocrinol (Lausanne)* [Internet]. 2019 [cited 2022 Mar 12];10(JUN):414. Available

344 from: /pmc/articles/PMC6610317/

345 24. Van Steenbergen E, Van Dongen JM, Wendel-Vos GCW, Hildebrandt VH, Strijk JE.
346 Insights into the concept of vitality: associations with participation and societal costs. Eur
347 J Public Health [Internet]. 2016 Apr 1 [cited 2022 Mar 12];26(2):354–9. Available from:
348 <https://academic.oup.com/eurpub/article/26/2/354/2570374>

349 25. Kurita N, Akizawa T, Fukuhara S. Vitality Measured as Self-reported Energy Level and
350 Clinical Outcomes in Hemodialysis Patients: The Japanese Dialysis Outcomes and
351 Practice Pattern Study (J-DOPPS). Am J Kidney Dis [Internet]. 2019 Apr 1;73(4):486–95.
352 Available from: <https://doi.org/10.1053/j.ajkd.2018.10.001>

353 26. Kubzansky LD, Thurston RC. Emotional Vitality and Incident Coronary Heart Disease:
354 Benefits of Healthy Psychological Functioning. Arch Gen Psychiatry [Internet]. 2007 Dec
355 1;64(12):1393–401. Available from: <https://doi.org/10.1001/archpsyc.64.12.1393>

356 27. Ganz PA, Moinpour CM, McCoy S, Pauler DK, Press OW, Fisher RI. Predictors of
357 vitality (energy/fatigue) in early stage Hodgkin's disease (HD): Results from Southwest
358 Oncology Group (SWOG) Study 9133. J Clin Oncol [Internet]. 2004 Jul
359 15;22(14_suppl):6546. Available from: <https://doi.org/10.1200/jco.2004.22.90140.6546>

360 28. Haraldstad K, Wahl A, Andenæs R, Andersen JR, Andersen MH, Beisland E, et al. A
361 systematic review of quality of life research in medicine and health sciences. Qual Life
362 Res [Internet]. 2019 Oct 1 [cited 2022 Mar 12];28(10):2641. Available from:
363 /pmc/articles/PMC6761255/







Figure 1: Trial Profile




Changes in Vitamin D levels between the fasting and control group




Changes in weight between the fasting and control group




Changes in BMI Scores between the fasting and control group



Changes in Vitamin B12 between the fasting and control group




Changes in Vitality Scores between the fasting and control group







Changes in hemoglobin between the fasting and control group

Figure 2: Changes in mean values of outcome measures between the fasting and control group



Figure

medRxiv preprint doi: <https://doi.org/10.1101/2022.04.08.22273614>; this version posted April 16, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).



WHO - World Health Organisation
QoL-Quality of Life

Figure 3: Association between vitality scores and quality of life domain