

Neurodevelopmental, cognitive, behavioural and mental health impairments following childhood malnutrition: a systematic review

Amir Kirolos 1,2, Magdalena Goyheneix 3, Mike Kalmus Eliazs 1, Mphatso Chisala 4, Samantha Lissauer 2,5, Melissa Gladstone* 1, Marko Kerac* 6

1. Institute of Life Course and Medical Sciences, Department of Women's and Children's Health, University of Liverpool, Liverpool, UK
2. Malawi-Liverpool-Wellcome Trust Clinical Research Programme, Blantyre, Malawi
3. ACNUR (La Agencia de la ONU para los Refugiados), United Nations High Commissioner for Refugees, Buenos Aires, Argentina
4. Malawi Epidemiology and Intervention Research Unit, Lilongwe/Karonga, Malawi
5. Institute of Infection, Veterinary & Ecological Sciences, University of Liverpool, Liverpool, UK
6. Centre for MARCH (Maternal, Adolescent, Reproductive & Child Health), London School of Hygiene & Tropical Medicine, London, UK

* Indicates joint senior authors

Corresponding Author: Dr Amir Kirolos (a.kirolos@liverpool.ac.uk)

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

ABSTRACT

Background: Severe childhood malnutrition impairs growth and development short-term, but current understanding of long-term outcomes is limited. We aimed to identify studies assessing neurodevelopmental, cognitive, behavioural and mental health outcomes following childhood malnutrition.

Methods: We systematically searched MEDLINE, EMBASE, Global Health and PsychINFO for studies assessing these outcomes in those exposed to childhood malnutrition in low- and middle-income settings. We included studies assessing undernutrition measured by low mid-upper arm circumference, weight-for-height, weight-for-age or nutritional oedema. We used guidelines for synthesis of results without meta-analysis to analyse three outcome areas: neurodevelopment, cognition/academic achievement, behaviour/mental health.

Results: We identified 30 studies, including some long-term cohorts reporting outcomes through to adulthood. There is strong evidence that malnutrition in childhood negatively impacts neurodevelopment based on high-quality studies using validated neurodevelopmental assessment tools. There is also strong evidence that malnutrition impairs academic achievement with agreement across seven studies investigating this outcome. 8 of 11 studies showed association between childhood malnutrition and impaired cognition. This moderate evidence is limited by some studies failing to measure important confounders such as socioeconomic status. 5 of 7 studies found a difference in behavioural assessment scores in those exposed to childhood malnutrition compared to controls but this moderate evidence is similarly limited by unmeasured confounders. Mental health impacts were difficult to ascertain due to few studies with mixed results.

Conclusions: Childhood malnutrition is associated with impaired neurodevelopment, academic achievement, cognition and behavioural problems but evidence regarding possible mental health impacts is inconclusive. Future research should explore the interplay of childhood and later-life adversities on these outcomes. Whilst evidence on improving nutritional and clinical therapies to reduce long-term risks is also needed, preventing and eliminating child malnutrition is likely to be the best way of preventing long-term neurocognitive harms.

PROSPERO registration number: CRD42021260498

57

58 **KEY QUESTIONS**

59

60 **What is already known?**

61

62 - High mortality risk and impaired growth are well-recognised short-term risks of
63 childhood malnutrition

64

65 - Whilst there is increasing appreciation of longer-term risks for survivors, notably
66 adult cardiometabolic non-communicable disease, other longer-term risks have
67 been poorly described.

68

69 **What are the new findings?**

70 - There is strong evidence that malnutrition impairs neurodevelopment and
71 academic achievement in childhood which has significant implications for future
72 wellbeing and prospects of those affected

73

74 - Childhood malnutrition is associated with impaired cognition and behavioural
75 problems with evidence of effects through to adolescence and adulthood but the
76 effect of nutritional treatment and interplay with childhood adversity, co-existing
77 illness such as HIV and environmental factors in influencing these outcomes is
78 unclear

79

80 **What do the new findings imply?**

81 - Study findings imply that there are likely to be long-term effects of childhood
82 malnutrition on cognition and wellbeing lasting through adolescence and
83 adulthood

84

- Long-term needs of malnutrition survivors need to be carefully considered in treatment programmes. Further research is needed on the effects of nutritional therapy, adversity and environmental factors to tailor future interventions, particularly with regards to mental health which has been little researched

INTRODUCTION

Severe childhood malnutrition is widespread and has a high disease burden concentrated in low- and middle-income settings¹. To date, most malnutrition policies and treatment programmes have focused on short-term risks, notably infections and death^{2,3}. There is however growing evidence of long-term risks for malnutrition survivors, including that of later-life non-communicable disease⁴. The prevalence of malnutrition has decreased in recent years due to concerted global efforts¹. However, climate change, conflict and food insecurity in many settings with fragile food supply chains risk resurgence and perpetuation of childhood malnutrition⁵.

Understanding long-term outcomes following child malnutrition is especially important because improved treatment has thankfully resulted in more children with malnutrition surviving into adolescence and adulthood^{3,6}. Most previous research and programmatic investment have focused on child mortality rather than thriving and long-term outcomes. Fewer studies explore long-term impacts of malnutrition on cognitive, behavioural and mental health outcomes in survivors. Improved understanding of these outcomes can inform disease burden estimates, support ongoing investment and inform follow-up care for children with malnutrition.

Causal pathways linking malnutrition with neurodevelopment, cognition, behaviour and mental health are complex. Previous studies have found children admitted to hospital with severe malnutrition often have severe developmental delays and children with neurodisability are inherently at higher risk of becoming malnourished^{7,8}. The relationship between malnutrition and development is also influenced by other associated factors such as childhood adversity, HIV-exposure, risk of infectious disease, parental engagement, school attendance and impaired growth⁹. These mediating factors also explain in part the difficulty in predicting developmental trajectories after an episode of severe malnutrition and why

studies investigating outcomes are significantly influenced by potential confounders affecting the internal validity of results ¹⁰.

A 1995 review found school-age children who suffered from early childhood malnutrition generally had poorer IQ levels, cognitive function, school achievement and greater behavioural problems than matched controls and, to a lesser extent, siblings ¹¹. Despite these associations, previous evidence of direct causal relationships is limited due to a lack of long-term follow-up studies, retrospective study designs and few studies having investigated behaviour and mental health. With this lack of up-to-date evidence on an increasingly important topic we aimed to identify studies reporting neurodevelopmental, cognitive, behavioural and mental health outcomes for children exposed to malnutrition in childhood.

METHODS

Our systematic review protocol was registered on PROSPERO (CRD42021260498). We searched MEDLINE, EMBASE, Global Health and PsycINFO for studies published between 1st January 1995 and 6th January 2022 using PRISMA guidelines ¹². Detailed search strategies are included in Supplementary Appendix 1.

Selection criteria

We included studies from low- and middle-income countries reporting neurodevelopmental, cognitive, school/academic achievement, mental health or behavioural outcomes in children under five exposed to malnutrition (we did not exclude studies that included small numbers of children between the ages of five and six years). We defined childhood malnutrition as undernutrition using standard definitions (definitions which are commonly measured in research and often related to severe adverse outcomes ¹³): moderate or severe wasting defined by low weight for height or low mid-upper arm circumference; presence of nutritional oedema; low weight-for-age as per older definitions of severe malnutrition. We did not include studies focusing on stunting (low height-for-age), chronic malnutrition or micronutrient deficiencies.

We included cross-sectional, cohort, case control and controlled trial study designs which had a comparator group not exposed to childhood malnutrition. We excluded conference papers

and reviews which did not include original data. We excluded studies which failed to define how malnutrition, child neurodevelopment, cognition, school/academic achievement, mental health or behaviour were measured. We also excluded studies looking at specific sub-populations of children (e.g. only children with a specific medical condition). No language restrictions were placed on studies.

Literature search

Titles and abstracts were screened by two independent reviewers (two of AK, MGo, MKE, MC). Selected full texts by any reviewer were then independently reviewed by two reviewers (two of AK, MGo, MKE, MC) against our selection criteria. We also screened reference lists of included studies. Data extraction of included studies was done by one of AK or MGo into a standard template (including study characteristics, assessment tool, malnutrition definition, sample size, results in cases and controls, reported effect sizes, results of statistical significance tests, results of analyses adjusted for confounding variables). Data extraction was subsequently checked by a second separate reviewer (one of AK, ME, MC). Disagreements regarding inclusion of full texts and data extraction were resolved through mutual discussion.

Quality assessment

We assessed study quality using the National Institute of Health and Care Excellence (NICE) quality appraisal checklist¹⁴. Studies were scored independently by two reviewers (two of AK, MGo, MKE, MC) with scoring disagreements resolved through mutual discussion. Internal and external validity were rated as poor, acceptable and very good based on NICE quality appraisal checklist results. We considered overall study quality to be high-quality when both internal and external validity scores were rated as very good, adequate-quality when either score was rated as acceptable and poor-quality where either internal or external validity was rated as poor.

Synthesis of study results

We grouped studies into three outcome areas: neurodevelopmental, cognition/academic achievement and mental health/behavioural outcomes (with some studies reporting outcomes from more than one category). We undertook a narrative synthesis of results within each of these areas as diverse outcomes and measurement tools precluded meta-analysis.

We followed the Synthesis without meta-analysis (SWiM) reporting guidelines for analysing and reporting results¹⁵.

For neurodevelopmental studies we grouped studies and compared results by neurodevelopmental assessment tool used. For studies investigating cognition/academic achievement, we grouped those that measured IQ or executive function and those that measured either school or language performance/assessment results. For mental health/behaviour studies, we grouped studies that used behavioural assessment tools and then into groups by mental health condition or domain assessed.

For each study outcome where available, we recorded the unadjusted and adjusted results in cases and controls, reported effect sizes, and results from any statistical significance tests. When synthesizing results by the groupings described above, we used vote counting by direction of effect to assess the number of studies which recorded differences between cases and controls. We summarised results in tables indicating the number of studies with results indicating an effect of malnutrition on the study outcome within each outcome area. We also recorded the age group (childhood 0-10 years, adolescence 10-18 years, adulthood 18+ years) at which outcomes were measured to compare results between studies. When drawing conclusions from our synthesis we accounted for study quality scoring, whether results adjusted for confounding variables, sample size and the use of validated outcome tools/assessments. Where several high-quality studies in one grouped outcome area reported results of an association with malnutrition, we graded conclusions as strong evidence. Where most adequate and high-quality studies reported an association with malnutrition, but there were uncertainties or limitations of studies identified, we graded conclusions as moderate evidence. Where there were few studies identified or studies with mixed results or poor study quality, we graded associations as inconclusive.

RESULTS

Study Characteristics

Thirty studies met our selection criteria (Supplementary Figure 1)¹⁶⁻⁴⁵. Full study characteristics are included in Supplementary Table 1. Studies were conducted between

1967-2019 across several countries in Africa, Asia and South America and included some long-term cohorts following participants through to adolescence and adulthood. Nine studies were part of the same Barbados Nutrition Study (BNS), a prospective lifelong cohort study assessing multiple outcomes at different follow-up points for children who suffered from malnutrition in the first year of life³⁷⁻⁴⁵. Nine studies assessed neurodevelopmental outcomes, 12 studies assessed cognition or academic achievement (three from BNS) and 14 studies assessed mental health conditions or behaviour (eight from BNS). Study designs included cross-sectional, retrospective cohorts, prospective cohorts and randomised controlled trial studies. Neurodevelopmental studies were either cross-sectional or had a short follow-up time during childhood. The age at follow-up following exposure to malnutrition in other studies varied from childhood through to adulthood.

Study quality scoring is included in Supplementary Table 2. Twenty-three studies had very good external validity, with the seven others apart from one with acceptable external validity. This was due to most providing good descriptions of study settings, populations and selection criteria. Fourteen studies had very good internal validity, 12 had acceptable internal validity and four studies had poor internal validity. Studies often scored poorly where important confounding variables such as socioeconomic status weren't accounted for in analyses, sample size was small or study designs were poor or poorly described, leading to potential biases. Overall, 14 studies were rated as high-quality, 12 were adequate-quality and four were poor-quality.

Malnutrition and neurodevelopment

We identified nine studies assessing neurodevelopmental outcomes of malnutrition (Table 1). A summary of results from these studies is included in Supplementary Table 3. Five studies used the Bayley Scales of Infant Development (BSID). Three of these were adequate-quality studies which found impaired neurodevelopment in those with malnutrition unadjusted for confounding variables such as socioeconomic status and family characteristics^{19,30,36}. Two high-quality studies using BSID found impaired neurodevelopment in those with malnutrition but one found that differences were no longer significant when adjusting for differences in current weight during follow-up at two years old^{23,33}. A further high-quality study using the Denver-II tool found children with malnutrition had lower scores across fine motor, gross motor, language and personal-social domains compared to controls²². One adequate-quality

study using a neurodevelopment tool developed in India (Indian Council Medical Research Psychosocial Developmental Screening Test) found poorer development outcomes for those with malnutrition³⁴. Another adequate-quality study using UNICEF Multiple Indicator Cluster Surveys (MICS) Early Childhood Development Indicators (ECDI) did not find an effect of malnutrition on neurodevelopment²⁰. This was carried out on a large population sample from five country surveys in Asia, but the MICS ECDI tool consists of few basic developmental items which span a large age range. This is notably different to the detailed neurodevelopmental assessments by trained assessors used in most other studies. A poor-quality study using an Indian specific development tool (Indian Development Inventory), found poorer neurodevelopment scores in cases, but this was no longer significant when adjusted for confounding variables. The results from high- and adequate-quality studies provides strong evidence of an association between malnutrition in childhood and impaired neurodevelopment across multiple developmental domains.

Malnutrition and cognition/academic achievement

We identified 12 studies (three from BNS) assessing cognitive outcomes of malnutrition (Table 2). A summary of results from these studies is included in Supplementary Table 4. Seven studies assessed academic performance, with three based on self-report of school performance or schooling years achieved, and four based on tests of school skill such as mathematics or country specific language tests (Supplementary Table 4)^{17,18,21,28,32,38,43}. Three studies were high-quality, three were adequate-quality and one was poor-quality. All studies (two from BNS) found worse school/academic performance in those exposed to malnutrition in childhood compared to controls. The results from high- and adequate-quality studies identified provides strong evidence of an association between exposure to malnutrition and impaired academic performance/achievement in childhood and adolescence.

Eleven studies investigated cognition using several tools assessing intelligence and executive function (Table 2, Supplementary Table 4). Eight of these studies (three from BNS) found impaired intelligence/executive function in those exposed to malnutrition in childhood compared to controls^{17,26,28,32,35,38,39,43}. Two high-quality studies found no significant association between malnutrition and cognition^{16,21}. One study from China found no difference between cases and controls using Wechsler Pre-school and Primary Scale of Intelligence (WPPSI-IV), and the other from Malawi found the differences seen between cases

and controls using the Cambridge Neuropsychological Test Automated Battery (CANTAB) were no longer significant when adjusting for confounding variables including HIV, stunting, socioeconomic status and household characteristics. Of the studies which found a significant difference in cognition between cases and controls, there were four high quality studies (two from BNS) ^{26,32,38,39}. These four high-quality studies all used different cognitive assessment tools, including Kauffman-ABC, Bohem Test, Trail Making Test and Wechsler Scale of Intelligence (Table 2), and found poorer scores in those exposed to malnutrition compared to controls. Three adequate-quality studies (one from BNS) using different cognitive assessment tools, including Mini-Mental State Exam, Wechsler Scale of Intelligence and Anstey Domino test (Table 2) also found those exposed to childhood malnutrition had impaired cognition compared to controls, however only one of these studies adjusted for any confounding variables (BNS study which adjusted for childhood standard of living) ^{17,35,43}. The results from high- and adequate-quality studies provide moderate evidence of an association between exposure to malnutrition in childhood and impaired cognition, but definitive conclusions are limited by mixed results from two high-quality studies showing no association after adjusting for important confounders such as HIV and socioeconomic status and results from several adequate-quality studies which failed to adjust for any confounding variables.

Malnutrition and mental health/behaviour

We identified 14 studies (eight from BNS) assessing mental health and behavioural outcomes of malnutrition (Table 3). A summary of results from these studies is included in Supplementary Table 5. Seven studies (three from BNS) assessed behaviour using different behavioural rating tools including the Strengths and Difficulties (SDQ), Child Behaviour Checklist, Ages and Stages (socio-emotional questions) and other adapted behavioural questionnaires (Table 3) ^{16,22,29,31,39,42,44}. Five of these studies, four of which were rated as high-quality, found significantly higher behavioural problems in those exposed to malnutrition in childhood ^{22,29,31,39,44}. One high-quality study found no difference in behavioural scores between cases and controls using SDQ and one adequate-quality study looking at conduct problems using found differences in behaviour were no longer significant when adjusting for confounding variables including childhood standard of living. The results from adequate- and high-quality studies provide moderate evidence of an association between exposure to malnutrition in childhood and increased behavioural problems but is

limited by some mixed results in high-quality studies and potential confounding in some studies reporting an association.

Seven studies (five from BNS) investigated different mental health outcomes (Table 3) ^{17,24,37,40,41,43,45}. Studies indicated possible associations with self-esteem, quality of life, personality disorders, attention deficits and low morale. There were small numbers of studies investigating specific mental health domains and there were significant study limitations in several studies included. The results from the studies identified provide inconclusive evidence regarding possible associations between exposure to malnutrition in childhood and poorer mental health outcomes.

DISCUSSION

Our review finds strong evidence that exposure to malnutrition in childhood impairs neurodevelopment and academic achievement. There is moderate evidence that childhood malnutrition is associated with impaired cognition and is associated with more behavioural problems throughout childhood and adolescence. However, there is uncertainty around the relative contributions of malnutrition and other associated factors (such as childhood adversity, HIV-exposure, socioeconomic status and household characteristics) to these outcomes. Research investigating mental health outcomes in children with malnutrition is inconclusive and there are few studies investigating specific mental health domains such as depression. These results have implications for policy makers surrounding the long-term care needs of those treated for childhood malnutrition and future research such as exploring how nutritional therapies affect these outcomes.

Due to study heterogeneity, we were unable to perform a meta-analysis of any results. We therefore used published guidelines to synthesize results and determine the strength of evidence for each outcome area. Despite this, there are still limitations given potential publication bias of positive associations between malnutrition and the outcomes investigated. Our review however builds on previously published evidence which has suggested links between malnutrition and poorer IQ levels, cognitive function, school achievement and greater behavioural problems ¹¹. Our findings strengthen the evidence-base regarding these associations as previous findings were limited by difficulties in interpreting

retrospective case control studies, but our review includes evidence from prospective studies which have sought to minimise these biases. There are however several poor- or adequate-quality studies we identified which failed to account for important confounders such as socioeconomic status and family characteristics. Of the 30 studies we identified, nine were from the same prospective cohort study in Barbados, with many of these publications from the same cohort testing multiple cognitive, behavioural and mental health outcomes. This may have increased the likelihood of type one error. For example, different studies from the same cohort found mixed results regarding morale and personality disorder scores when different assessment tools were used (Table 3) ^{41,43,45}.

Impaired neurodevelopment in childhood is likely linked to subsequently poorer academic achievement in childhood and adolescence and may also be linked to increased behavioural problems seen in some studies ⁴⁶. The studies assessing IQ/executive function outcomes spanned childhood through to adulthood and there was moderate evidence of a link between malnutrition and cognition. Two high-quality studies showed no effect ^{16,21} and other studies which showed an effect did not adjust for any confounding variables ^{17,28,35}. There are therefore outstanding questions over how much early impairment in neurodevelopment due to malnutrition affects future cognition and functioning, and to what extent other related environmental factors such as prenatal nutrition, family characteristics and infections contribute to these outcomes ^{47,48}. Possible sex-specific differences are another area which data here does not allow us to comment on but is important for future work to explore ^{49,50}. Mental health outcomes such as depression and inattention are likely to be influenced by similar confounding variables, and whether early insults from malnutrition independently contribute to poor mental health outcomes in later life is yet to be established. There is also uncertainty around the effect of nutritional therapy on long-term cognition and functioning ⁵¹. However, even if some of the effects we report are due to confounding, evidence is clear from our review that children who experience an episode of malnutrition in childhood are at high risk of poorer development, behaviour and cognition. Policy makers should therefore prioritise targeted support for these vulnerable children and programmes to optimise life chances after recovery.

While there is still uncertainty around the relative contributions of associated medical and social factors, there is increasing evidence from this review that the early impacts of

malnutrition are related to worse academic, cognitive and behavioural outcomes compared to well-nourished peers. Preventing and decreasing childhood malnutrition is therefore of key importance to prevent serious long-term neurocognitive issues in affected children, particularly given the ongoing high malnutrition prevalence in many low- and middle-income settings. Further research is needed on how to optimise treatment and to best support ongoing care for survivors to improve outcomes in the long-term.

Contributions

AK, MGo, MGI, MK conceived the study design. **AK** wrote the main draft of the paper. **AK, MGo, MKE, MC** screened studies against selection criteria, extracted/checked data from studies and quality scored studies. **AK, MGo** analysed and synthesized data. All authors reviewed and contributed to the manuscript.

Funding

AK is supported by a Wellcome Trust Clinical PhD Programme Fellowship (203919/Z/16/Z).

Ethics Statement

This study did not receive nor require ethics approval, as it does not involve human & animal participants

Acknowledgements

Patients and public were not involved in the design and conduct of this study

Competing Interests

None declared

REFERENCES

- 1 Global Nutrition Report: The state of global nutrition. (Development Initiatives, Bristol, UK, 2021).
- 2 Black, R. E. *et al.* Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet* **382**, 427-451, doi:10.1016/s0140-6736(13)60937-x (2013).

- 3 Bhutta, Z. A. *et al.* Severe childhood malnutrition. *Nat Rev Dis Primers* **3**, 17067, doi:10.1038/nrdp.2017.67 (2017).
- 4 Grey, K. *et al.* Severe malnutrition or famine exposure in childhood and cardiometabolic non-communicable disease later in life: a systematic review. *BMJ Global Health* **6**, e003161, doi:10.1136/bmjgh-2020-003161 (2021).
- 5 United Nations Children's Fund (UNICEF) WHO, W. B. Levels and trends in child malnutrition: key findings of the 2021 edition of the joint child malnutrition estimates. (World Health Organization, Geneva, 2021).
- 6 Fergusson, P., Tomkins, A. & Kerac, M. Improving survival of children with severe acute malnutrition in HIV-prevalent settings. *Int Health* **1**, 10-16, doi:10.1016/j.inhe.2009.03.001 (2009).
- 7 van den Heuvel, M. *et al.* Developmental and behavioural problems in children with severe acute malnutrition in Malawi: A cross-sectional study. *J Glob Health* **7**, 020416, doi:10.7189/jogh.07.020416 (2017).
- 8 Gladstone, M. *et al.* Assessment of neurodisability and malnutrition in children in Africa. *Semin Pediatr Neurol* **21**, 50-57, doi:10.1016/j.spen.2014.01.002 (2014).
- 9 Martins, V. J. *et al.* Long-lasting effects of undernutrition. *Int J Environ Res Public Health* **8**, 1817-1846, doi:10.3390/ijerph8061817 (2011).
- 10 Pollitt, E. Developmental sequel from early nutritional deficiencies: conclusive and probability judgements. *J Nutr* **130**, 350s-353s, doi:10.1093/jn/130.2.350S (2000).
- 11 Grantham-McGregor, S. A review of studies of the effect of severe malnutrition on mental development. *J Nutr* **125**, 2233s-2238s, doi:10.1093/jn/125.suppl_8.2233S (1995).
- 12 Page, M. J. *et al.* PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *Bmj* **372**, n160, doi:10.1136/bmj.n160 (2021).
- 13 Kerac, M. *et al.* 'Severe malnutrition': thinking deeply, communicating simply. *BMJ Global Health* **5**, e003023, doi:10.1136/bmjgh-2020-003023 (2020).
- 14 NICE. *Quality appraisal checklist - quantitative studies reporting correlations and associations*, <<https://www.nice.org.uk/process/pmg4/chapter/appendix-g-quality-appraisal-checklist-quantitative-studies-reporting-correlations-and>> (2012).
- 15 Campbell, M. *et al.* Synthesis without meta-analysis (SWiM) in systematic reviews: reporting guideline. *Bmj* **368**, l6890, doi:10.1136/bmj.l6890 (2020).
- 16 Chen, K. *et al.* Nutrition, Cognition, and Social Emotion among Preschoolers in Poor, Rural Areas of South Central China: Status and Correlates. *Nutrients* **13**, doi:10.3390/nu13041322 (2021).
- 17 Mwene-Batu, P. *et al.* Long-term effects of severe acute malnutrition during childhood on adult cognitive, academic and behavioural development in African fragile countries: The Lwiro cohort study in Democratic Republic of the Congo. *PLoS One* **15**, e0244486, doi:10.1371/journal.pone.0244486 (2020).
- 18 Asiki, G., Newton, R., Marions, L., Kamali, A. & Smedman, L. The effect of childhood stunting and wasting on adolescent cardiovascular diseases risk and educational achievement in rural Uganda: a retrospective cohort study. *Glob Health Action* **12**, 1626184, doi:10.1080/16549716.2019.1626184 (2019).
- 19 Dwivedi, D., Singh, S., Singh, J., Bajaj, N. & Singh, H. P. Neurodevelopmental Status of Children aged 6-30 months with Severe Acute Malnutrition. *Indian Pediatr* **55**, 131-133 (2018).
- 20 Kang, Y. *et al.* Nutritional status and risk factors for stunting in preschool children in Bhutan. *Matern Child Nutr* **14 Suppl 4**, e12653, doi:10.1111/mcn.12653 (2018).
- 21 Lelijveld, N. *et al.* Brain MRI and cognitive function seven years after surviving an episode of severe acute malnutrition in a cohort of Malawian children. *Public Health Nutr* **22**, 1406-1414, doi:10.1017/s1368980018003282 (2018).

- 433 22 Abessa, T. G., Bruckers, L., Kolsteren, P. & Granitzer, M. Developmental performance of
434 hospitalized severely acutely malnourished under-six children in low- income setting. *BMC*
435 *Pediatr* **17**, 197, doi:10.1186/s12887-017-0950-5 (2017).
- 436 23 Sudfeld, C. R. *et al.* Malnutrition and Its Determinants Are Associated with Suboptimal
437 Cognitive, Communication, and Motor Development in Tanzanian Children. *J Nutr* **145**, 2705-
438 2714, doi:10.3945/jn.115.215996 (2015).
- 439 24 De Grandis, E. S., Armelini, P. A. & Cuestas, E. [Evaluation of quality of life in school children
440 with a history of early severe malnutrition]. *An Pediatr (Barc)* **81**, 368-373,
441 doi:10.1016/j.anpedi.2013.11.021 (2014).
- 442 25 Malhi, P., Bharti, B. & Sidhu, M. Developmental Functioning of Young Indian
443 Children with Malnutrition. *Psychol Study* **58**, 259-264, doi:10.1007/s12646-013-0194-x (2013).
- 444 26 Bogale, A. *et al.* Nutritional status and cognitive performance of mother-child pairs in
445 Sidama, Southern Ethiopia. *Matern Child Nutr* **9**, 274-284, doi:10.1111/j.1740-
446 8709.2011.00345.x (2013).
- 447 27 Warsito, O., Khomsan, A., Hernawati, N. & Anwar, F. Relationship between nutritional status,
448 psychosocial stimulation, and cognitive development in preschool children in Indonesia. *Nutr*
449 *Res Pract* **6**, 451-457, doi:10.4162/nrp.2012.6.5.451 (2012).
- 450 28 Nassar, M. F., Shaaban, S. Y., Nassar, J. F., Younis, N. T. & Abdel-Mobdy, A. E. Language skills
451 and intelligence quotient of protein energy malnutrition survivors. *J Trop Pediatr* **58**, 226-
452 230, doi:10.1093/tropej/fmr081 (2012).
- 453 29 Baker-Henningham, H., Hamadani, J. D., Huda, S. N. & Grantham-McGregor, S. M.
454 Undernourished children have different temperaments than better-nourished children in
455 rural Bangladesh. *J Nutr* **139**, 1765-1771, doi:10.3945/jn.109.106294 (2009).
- 456 30 al, H. E.-k. e. Cognitive Functions in Protein-Energy Malnutrition: In Relation to Long Chain-
457 polyunsaturated Fatty Acids. *Pakistan Journal of Biological Sciences* **10**, 1773-1781 (2007).
- 458 31 Liu, J., Raine, A., Venables, P. H. & Mednick, S. A. Malnutrition at age 3 years and
459 externalizing behavior problems at ages 8, 11, and 17 years. *Am J Psychiatry* **161**, 2005-2013,
460 doi:10.1176/appi.ajp.161.11.2005 (2004).
- 461 32 Liu, J., Raine, A., Venables, P. H., Dalais, C. & Mednick, S. A. Malnutrition at age 3 years and
462 lower cognitive ability at age 11 years: independence from psychosocial adversity. *Arch*
463 *Pediatr Adolesc Med* **157**, 593-600, doi:10.1001/archpedi.157.6.593 (2003).
- 464 33 Drewett, R., Wolke, D., Asefa, M., Kaba, M. & Tessema, F. Malnutrition and mental
465 development: is there a sensitive period? A nested case-control study. *J Child Psychol*
466 *Psychiatry* **42**, 181-187, doi:10.1111/1469-7610.00709 (2001).
- 467 34 Vazir, S., Naidu, A. N. & Vidyasagar, P. Nutritional status, psychosocial development and the
468 home environment of Indian rural children. *Indian Pediatr* **35**, 959-966 (1998).
- 469 35 Perales, C. G., Heresi, E., Pizarro, F. & Colombo, M. [Cognitive functions of school children
470 with normal IQ and histories of severe and early malnutrition]. *Arch Latinoam Nutr* **46**, 282-
471 286 (1996).
- 472 36 Meeta, K. & I.J.S, J. Malnutrition and Mental Development during Infancy. *J. Dairying, Foods*
473 *& Home Sci.*, **14**, 159-165 (1995).
- 474 37 Hock, R. S. *et al.* Childhood malnutrition and maltreatment are linked with personality
475 disorder symptoms in adulthood: Results from a Barbados lifespan cohort. *Psychiatry Res*
476 **269**, 301-308, doi:10.1016/j.psychres.2018.05.085 (2018).
- 477 38 Waber, D. P. *et al.* Impaired IQ and academic skills in adults who experienced moderate to
478 severe infantile malnutrition: a 40-year study. *Nutr Neurosci* **17**, 58-64,
479 doi:10.1179/1476830513y.0000000061 (2014).
- 480 39 Waber, D. P. *et al.* Neuropsychological outcomes at midlife following moderate to severe
481 malnutrition in infancy. *Neuropsychology* **28**, 530-540, doi:10.1037/neu0000058 (2014).
- 482 40 Galler, J. R. *et al.* Malnutrition in the first year of life and personality at age 40. *J Child*
483 *Psychol Psychiatry* **54**, 911-919, doi:10.1111/jcpp.12066 (2013).

- 41 Galler, J. R. *et al.* Infant malnutrition is associated with persisting attention deficits in middle adulthood. *J Nutr* **142**, 788-794, doi:10.3945/jn.111.145441 (2012).
- 42 Galler, J. R. *et al.* Infant malnutrition predicts conduct problems in adolescents. *Nutr Neurosci* **15**, 186-192, doi:10.1179/1476830512y.0000000012 (2012).
- 43 Waber, D. P. *et al.* Cognitive impairment as a mediator in the developmental pathway from infant malnutrition to adolescent depressive symptoms in Barbadian youth. *J Dev Behav Pediatr* **32**, 225-232, doi:10.1097/DBP.0b013e31820b7707 (2011).
- 44 Galler, J. R. *et al.* Early malnutrition predicts parent reports of externalizing behaviors at ages 9-17. *Nutr Neurosci* **14**, 138-144, doi:10.1179/147683011x13009738172521 (2011).
- 45 Galler, J. R. *et al.* Early childhood malnutrition predicts depressive symptoms at ages 11-17. *J Child Psychol Psychiatry* **51**, 789-798, doi:10.1111/j.1469-7610.2010.02208.x (2010).
- 46 Krebs, N. F., Lozoff, B. & Georgieff, M. K. Neurodevelopment: The Impact of Nutrition and Inflammation During Infancy in Low-Resource Settings. *Pediatrics* **139**, S50-S58, doi:10.1542/peds.2016-2828G (2017).
- 47 Germine, L., Dunn, E. C., McLaughlin, K. A. & Smoller, J. W. Childhood Adversity Is Associated with Adult Theory of Mind and Social Affiliation, but Not Face Processing. *PLoS One* **10**, e0129612, doi:10.1371/journal.pone.0129612 (2015).
- 48 McLaughlin, K. A., Weissman, D. & Bitrán, D. Childhood Adversity and Neural Development: A Systematic Review. *Annu Rev Dev Psychol* **1**, 277-312, doi:10.1146/annurev-devpsych-121318-084950 (2019).
- 49 Shapiro, J. R., Klein, S. L. & Morgan, R. Stop 'controlling' for sex and gender in global health research. *BMJ Global Health* **6**, e005714, doi:10.1136/bmjgh-2021-005714 (2021).
- 50 Thurstans, S. *et al.* Boys are more likely to be undernourished than girls: a systematic review and meta-analysis of sex differences in undernutrition. *BMJ Global Health* **5**, e004030, doi:10.1136/bmjgh-2020-004030 (2020).
- 51 Stephenson, K. *et al.* Low linoleic acid foods with added DHA given to Malawian children with severe acute malnutrition improve cognition: a randomized, triple blinded, controlled clinical trial. *Am J Clin Nutr*, doi:10.1093/ajcn/nqab363 (2021).

Table 1: Summary of results from studies assessing the effect of malnutrition on neurodevelopment

Study (quality) Country	Neurodevelopmental assessment tool				
	BSID	MICS ECDI	Denver	IDI	ICMR
Dwivedi et al 2018 (+) India	Effect (unadjusted)*				
Kang et al 2018 (+) multiple in Asia		No effect (adjusted)			
Abessa et al 2017 (++) Ethiopia			Effect (adjusted)		
Sudfeld et al 2015 (++) Tanzania	Effect (adjusted)				
Malhi et al 2013 (-) India				No effect (adjusted)	
El-Khayat et al 2007 (+) Egypt	Effect (unadjusted)				
Drewett et al 2001 (++) Ethiopia	No effect (adjusted)				
Vazir et al 1998 (+) India					Effect (unadjusted)
Kaul et al 1995 (+) India	Effect (unadjusted)				

* Indian modification of BSID

Effect - statistically significant difference in neurodevelopment between cases and controls

No effect – no difference, or statistically insignificant difference, or statistically insignificant difference after adjusting for confounding variables, in neurodevelopment between cases and controls

Adjusted – results adjusted for confounding variables

Unadjusted – results unadjusted for confounding variables

++ = high-quality

+ = adequate-quality

- = poor-quality

BSID – Bayley Scales of Infant Development, IDI – Indian Developmental Inventory, ICMR – Indian Council Medical Research Psychosocial Developmental Screening Test, MICS ECDI – Multiple Indicator Cluster Survey Early Child Development Index

Table 2: Summary of results from studies assessing the effect of malnutrition on cognition and academic achievement

Study (quality) [outcome age] Country	School, academic or language performance	Executive function/ Intelligence [assessment tool]
Chen <i>et al</i> 2021 (++) [childhood] China		No effect (adjusted) [WPPSI-IV]
Mwene-Batu <i>et al</i> 2020 (+) [adulthood] DRC	Effect (unadjusted)	Effect (unadjusted) [MMSE]
Asiki <i>et al</i> 2018 (++) [adolescence] Uganda	Effect (adjusted)	
Lelijveld <i>et al</i> 2018 (++) [childhood] Malawi	Effect (adjusted)	No effect (adjusted) [CANTAB]
Bogale <i>et al</i> 2013 (++) [childhood] Ethiopia		Effect (adjusted) [Raven's CPM, KABC-II]
Warsito <i>et al</i> 2012 (-) [childhood] Indonesia		No effect (unadjusted) [Indonesian Department of National Education – child development instrument score]
Nassar <i>et al</i> 2012 (-) [childhood] Egypt	Effect (unadjusted)	Effect (unadjusted) [Stanford-Binet-Intelligence-Scale]
Liu <i>et al</i> 2003 (++) [adolescence] Mauritius	Effect (adjusted)	Effect (adjusted) [Bohem Test, Trail Making Test, WISC]
Perales <i>et al</i> 1996 (+) [childhood] Chile		Effect (unadjusted) [Continuous Performance Task, Anstey Domino test]
Barbados Nutrition Study		
Waber (1) <i>et al</i> 2014* (++) [adulthood] Barbados	Effect (adjusted)	Effect (adjusted) [WASI]
Waber (2) <i>et al</i> 2014* (++) [adulthood] Barbados		Effect (adjusted) [WAIS-III, D-KEFS, WRAML-2, Wisconsin card sorting, Metacognitive index]
Waber <i>et al</i> 2011* (+) [adolescence] Barbados	Effect (adjusted)	Effect (adjusted) [WISC]

*Studies are drawn from the same Barbados Nutrition Study

Effect - statistically significant difference in academic achievement/cognition between cases and controls

No effect –no difference, or statistically insignificant difference, or statistically insignificant difference after adjusting for confounding variables, in academic achievement/cognition between cases and controls

Adjusted – results adjusted for confounding variables

Unadjusted – results unadjusted for confounding variables

++ = high-quality

+ = adequate-quality

- = poor-quality

CANTAB – Cambridge Neuropsychological Test Automated Battery, D-KEFS – Delis-Kaplan Executive Function System, KABC-II – Kaufman Assessment Battery for Children-II, MMSE – Mini-Mental State Exam, Raven’s CPM – Raven’s Colored Progressive Matrices, WAIS-III – Wechsler Adult Intelligence Scale-III, WASI – Adult IQ: Wechsler Abbreviated Scale of Intelligence – Vocabulary and Matrix Reasoning subsets, WISC – Weschler Intelligence Scale for Children, WPPSI-IV – Wechsler Preschool and Primary Scale of Intelligence Fourth Edition, WRAML-II – Wide Range Assessment of Memory and Learning-II

Table 3: Summary of results from studies assessing the effect of malnutrition on mental health and behaviour

Study (quality) [outcome age] Country	Behavioural problems [tool]	Self-esteem [tool]	Social-related disability [tool]	Quality of life [tool]	Personality [tool]	Attention [tool]	Morale [tool]
Chen <i>et al</i> 2021 (++) [childhood] China	No effect (adjusted) [SDQ]						
Mwene-Batu et al 2020 (+) [adulthood] DRC		Effect (unadjusted) [Rosenberg Self- Esteem Scale]	No effect (unadjusted) [WHODAS]				
Abessa et al 2017 (++) [childhood] Ethiopia	Effect (adjusted) [ASQ:SE]						
De Grandis et al 2014 (-) [childhood] Argentina				Effect (unadjusted) [Paediatric quality of life inventory]			
Baker- Henningham et al 2009 (++) [childhood] Bangladesh	Effect (adjusted) [Wachs temperament questionnaire]						
Liu et al 2004 (+) [childhood, adolescence] Mauritius	Effect (adjusted) [Child Behaviour Checklist]						
Barbados Nutrition Study							

Hock et al 2018* (+) [adulthood] Barbados					Mixed (adjusted) [SCID-II-PQ, NEO PI-R FFM]		
Waber (2) et al 2014* (++) [adulthood] Barbados	Effect (adjusted) [Behavioural regulation index]						
Galler et al 2013* (+) [adulthood] Barbados					Effect (adjusted) [NEO-PI-R]		
Galler (1) et al 2012* (+) [adulthood] Barbados						Mixed (adjusted) [CAARS, CPT]	
Galler (2) et al 2012* (+) [adolescence] Barbados	No effect (adjusted) [Conduct: CBQ, Teacher behaviour questionnaire]						
Waber et al 2011* (+) [adolescence] Barbados						No effect (adjusted) [Attention: CBQ]	No effect (adjusted) [Minnesota Adjustment and Morale Scale]
Galler et al 2011* (++) [adolescence] Barbados	Effect (adjusted) [BCBS]						
Galler et al 2010* (++)							Effect

[adolescence] Barbados							(adjusted) [Minnesota Adjustment and Morale Scale]
----------------------------------	--	--	--	--	--	--	---

*Studies are drawn from the same Barbados Nutrition Study

Effect - statistically significant difference in mental health/behavior between cases and controls

No effect –no difference, or statistically insignificant difference, or statistically insignificant difference after adjusting for confounding variables, in mental health/behaviour between cases and controls

Adjusted – results adjusted for confounding variables

Unadjusted – results unadjusted for confounding variables

++ = high-quality

+ = adequate-quality

- = poor-quality

ASQ:SE – Ages and Stages Questionnaire Socio emotional, BCBS – Barbados Child Behaviour Scale, CAARS - Attention-domain: Conners-ADHD-Rating-Scales, CBQ – Child Behaviour Questionnaire, CPT – Conners-Continuous-Performance Test, NEO PI-R – NEO Personality Inventory-Revised, NEO PI-R FFM – NEO Personality Inventory-Revised derived Five-Factor Model, SCID-II-PQ – Structured Clinical Interview for DSM-IV Axis II Personality Disorders Personality Questionnaire, SDQ – Strengths and Difficulties Questionnaire, WHODAS – World Health Organization Disability Assessment Schedule

Supplementary Appendix 1 – Search strategies

EMBASE

1. severe acute malnutrition.ti,ab.
2. acute severe malnutrition.ti,ab.
3. acute malnutrition.ti,ab.
4. (acute adj2 malnutrition).ti,ab.
5. "sever* acute* maln*".ti,ab.
6. protein-energy malnutrition.ti,ab.
7. protein calorie malnutrition.ti,ab.
8. (protein malnutrition or energy malnutrition or energy-protein malnutrition).ti,ab.
9. "acute* maln*".ti,ab.
10. "kwas?io?kor*".ti,ab.
11. "marasm*".ti,ab.
12. "marasm* kwas?io?kor*".ti,ab.
13. wasting.ti,ab.
14. "acute wast*".ti,ab.
15. wasted.ti,ab.
16. (acute adj2 wasting).ti,ab.
17. (severe adj2 wasting).ti,ab.
18. acute undernutrition.ti,ab.
19. (acute adj2 undernutrition).ti,ab.
20. severe undernutrition.ti,ab.
21. (severe adj2 undernutrition).ti,ab.
22. "mid* upper arm circumference".ti,ab.
23. (mid* upper arm circumference adj "115").ti,ab.
24. (mid* upper arm circumference adj "110").ti,ab.

25. MUAC.ti,ab.
26. "acute* undernourish*".ti,ab.
27. "acut* emaciat*".ti,ab.
28. "acut* maln*".ti,ab.
29. *malnutrition/
30. exp kwashiorkor/
31. exp marasmus/
32. exp wasting syndrome/
33. exp protein calorie malnutrition/
34. *nutritional disorder/
35. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23 or 24 or 25 or 26 or 27 or 28 or 29 or 30 or 31 or 32 or 33 or 34
36. infant/
37. child/
38. exp toddler/
39. "child*".ti,ab.
40. "infan*".ti,ab.
41. "pre-school*".ti,ab.
42. "preschool*".ti,ab.
43. "toddler*".ti,ab.
44. 36 or 37 or 38 or 39 or 40 or 41 or 42 or 43
45. 35 and 44
46. "child* developm* ".ti,ab.
47. "child* neurodevelopm*".ti,ab.
48. "neurodevelopm*".ti,ab.
49. (socio-emotion* develop* or socioemotion* develop*).ti,ab.

50. "social developm*".ti,ab.
51. "emotion* developm*".ti,ab.
52. (sensorymotor developm* or sensory-motor developm*).ti,ab.
53. "motor developm*".ti,ab.
54. "motor neurodevelopm*".ti,ab.
55. exp child development/ or exp postnatal development/
56. 46 or 47 or 48 or 49 or 50 or 51 or 52 or 53 or 54 or 55
57. exp cognition assessment/ or exp cognition/
58. exp learning/ or exp learning disorder/
59. exp intelligence quotient/ or exp intelligence/
60. exp language development/ or exp human development/ or exp mental development/
61. *brain function/ or *central nervous system function/
62. cognition.ti,ab.
63. "cogniti* performance".ti,ab.
64. intelligence.ti,ab.
65. IQ.ti,ab.
66. "executive function*".ti,ab.
67. reasoning.ti,ab.
68. language.ti,ab.
69. attention.ti,ab.
70. memory.ti,ab.
71. memory.ti,ab.
72. learning.ti,ab.
73. early learning.ti,ab.
74. information processing.ti,ab.
75. literacy.ti,ab.

76. reading.ti,ab.
77. math.ti,ab.
78. 57 or 58 or 59 or 60 or 61 or 62 or 63 or 64 or 65 or 66 or 67 or 69 or 70 or 71 or 72 or 73 or 74 or 75 or 76 or 77
79. education.ti,ab.
80. "school* achievement".ti,ab.
81. "school* performance".ti,ab.
82. school retention.ti,ab.
83. academic.ti,ab.
84. pre-academic.ti,ab.
85. schooling.ti,ab.
86. exp academic achievement/
87. 79 or 80 or 81 or 82 or 83 or 84 or 85 or 86
88. exp behavior/
89. exp mental health/
90. exp psychological aspect/ or exp neuropsychology/ or exp attention/ or exp cognitive defect/ or exp mental disease/
91. exp depression/
92. *hyperactivity/ or *psychomotor disorder/
93. exp attention deficit disorder/
94. "behavi?r* problem*".ti,ab.
95. "emotion* problem*".ti,ab.
96. "temperament*".ti,ab.
97. self regulation.ti,ab.
98. attachment.ti,ab.
99. self esteem.ti,ab.
100. self efficacy.ti,ab.
101. "social competen*".ti,ab.

102. "peer relationship*".ti,ab.
103. pro-social behavi?r.ti,ab.
104. hyperactivity.ti,ab.
105. impulsivity.ti,ab.
106. attention* deficit hyperactivity disorder*.ti,ab.
107. agresion.ti,ab.
108. 88 or 89 or 90 or 91 or 92 or 93 or 94 or 95 or 96 or 97 or 98 or 99 or 100 or 101 or 102 or 103 or 104 or 105 or 106 or 107
109. 56 or 78 or 90 or 108
110. 45 and 109
111. limit 110 to (human and yr="1995 -Current")

MEDLINE

1. "acute severe malnutrition".ti,ab.
2. severe acute malnutrition.ti,ab.
3. acute malnutrition.ti,ab.
4. (acute adj2 malnutrition).ti,ab.
5. "sever* acute* maln*".ti,ab.
6. protein-energy malnutrition.ti,ab.
7. acute severe malnutrition.ti,ab.
8. (protein malnutrition or energy malnutrition or energy-protein malnutrition).ti,ab.
9. "acute* maln*".ti,ab.
10. "kwas?io?kor*".ti,ab.
11. "marasm*".ti,ab.
12. "marasm* kwas?io?kor*".ti,ab.
13. wasting.ti,ab.

14. "acute wast*".ti,ab.
15. wasted.ti,ab.
16. (acute adj2 wasting).ti,ab.
17. (severe adj2 wasting).ti,ab.
18. acute undernutrition.ti,ab.
19. (acute adj2 undernutrition).ti,ab.
20. severe undernutrition.ti,ab.
21. (severe adj2 undernutrition).ti,ab.
22. "mid* upper arm circumference".ti,ab.
23. (mid* upper arm circumference adj "115").ti,ab.
24. (mid* upper arm circumference adj "110").ti,ab.
25. MUAC.ti,ab.
26. "acute* undernourish*".ti,ab.
27. "acut* maln*".ti,ab.
28. exp child nutrition disorders/ or exp deficiency diseases/ or exp starvation/ or *wasting syndrome/
29. exp Protein-Energy Malnutrition/
30. exp Infant Nutrition Disorders/
31. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23 or 24 or 25 or 26 or 27 or 28 or 29 or 30
32. "child*".ti,ab.
33. "infan*".ti,ab.
34. "pre-school*".ti,ab.
35. "preschool*".ti,ab.
36. "toddler*".ti,ab.
37. exp Infant/
38. exp Child/

39. Child, Preschool/
40. 32 or 33 or 34 or 35 or 36 or 37 or 38 or 39
41. 31 and 40
42. "child* developm* ".ti,ab.
43. "child* neurodevelopm*".ti,ab.
44. "neurodevelopm*".ti,ab.
45. (socio-emotion* develop* or socioemotion* develop*).ti,ab.
46. "social developm*".ti,ab.
47. "emotion* developm*".ti,ab.
48. (sensorymotor developm* or sensory-motor developm*).ti,ab.
49. "motor developm*".ti,ab.
50. "motor neurodevelopm*".ti,ab.
51. exp Child Development/
52. 42 or 43 or 44 or 45 or 46 or 47 or 48 or 49 or 50 or 51
53. cognition.ti,ab.
54. "cogniti* performance".ti,ab.
55. intelligence.ti,ab.
56. IQ.ti,ab.
57. "executive function*".ti,ab.
58. reasoning.ti,ab.
59. language.ti,ab.
60. attention.ti,ab.
61. memory.ti,ab.
62. learning.ti,ab.
63. early learning.ti,ab.
64. information processing.ti,ab.

- 65. literacy.ti,ab.
- 66. reading.ti,ab.
- 67. math.ti,ab.
- 68. *cognition/ or exp cognitive reserve/ or exp comprehension/ or exp executive function/ or exp learning/
- 69. exp Mild Cognitive Impairment/
- 70. *Intelligence/
- 71. *language development/ or *child language/
- 72. 53 or 54 or 55 or 56 or 57 or 58 or 59 or 60 or 61 or 62 or 63 or 64 or 65 or 66 or 67 or 68 or 69 or 70 or 71
- 73. education.ti,ab.
- 74. "school* achievement".ti,ab.
- 75. "school* performance".ti,ab.
- 76. school retention.ti,ab.
- 77. academic.ti,ab.
- 78. pre-academic.ti,ab.
- 79. schooling.ti,ab.
- 80. exp Educational Status/
- 81. 73 or 74 or 75 or 76 or 77 or 78 or 79 or 80
- 82. "behavi?r* problem*".ti,ab.
- 83. "emotion* problem*".ti,ab.
- 84. "temperament*".ti,ab.
- 85. self regulation.ti,ab.
- 86. attachment.ti,ab.
- 87. self esteem.ti,ab.
- 88. self efficacy.ti,ab.
- 89. "social competen*".ti,ab.
- 90. "peer relationship*".ti,ab.

91. pro-social behavi?r.ti,ab.
92. hyperactivity.ti,ab.
93. impulsivity.ti,ab.
94. attention* deficit hyperactivity disorder*.ti,ab.
95. aggression.ti,ab.
96. exp Child Behavior Disorders/
97. exp Mental Health/
98. *psychology/ or exp psychology, adolescent/ or exp psychology, child/
99. exp Depression/
100. exp Attention Deficit Disorder with Hyperactivity/
101. *Anxiety Disorders/
102. 82 or 83 or 84 or 85 or 86 or 87 or 88 or 89 or 90 or 91 or 92 or 93 or 94 or 95 or 96 or 97 or 98 or 99 or 100 or 101
103. 52 or 72 or 81 or 102
104. 41 and 103
105. limit 104 to (humans and yr="1995 -Current")

GLOBAL HEALTH

1. severe acute malnutrition.ti,ab.
2. acute severe malnutrition.ti,ab.
3. acute malnutrition.ti,ab.
4. (acute adj2 malnutrition).ti,ab.
5. "sever* acute* maln*".ti,ab.
6. protein-energy malnutrition.ti,ab.
7. protein calorie malnutrition.ti,ab.
8. (protein malnutrition or energy malnutrition or energy-protein malnutrition).ti,ab.

9. "acute* maln*".ti,ab.
10. "kwas?io?kor*".ti,ab.
11. "marasm*".ti,ab.
12. "marasm* kwas?io?kor*".ti,ab.
13. wasting.ti,ab.
14. "acute wast*".ti,ab.
15. wasted.ti,ab.
16. (acute adj2 wasting).ti,ab.
17. (severe adj2 wasting).ti,ab.
18. acute undernutrition.ti,ab.
19. (acute adj2 undernutrition).ti,ab.
20. severe undernutrition.ti,ab.
21. (severe adj2 undernutrition).ti,ab.
22. "mid* upper arm circumference".ti,ab.
23. (mid* upper arm circumference adj "115").ti,ab.
24. (mid* upper arm circumference adj "110").ti,ab.
25. MUAC.ti,ab.
26. "acute* undernourish*".ti,ab.
27. "acut* maln*".ti,ab.
28. emaciat*.ti,ab.
29. nutritional disorders/ or malnutrition/ or exp protein energy malnutrition/ or exp undernutrition/ or exp nutritional oedema/
30. wasting disease/ or exp emaciation/
31. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23 or 24 or 25 or 26 or 27 or 28 or 29 or 30
32. "child*".ti,ab.
33. "infan*".ti,ab.

34. "pre-school*".ti,ab.
35. "preschool*".ti,ab.
36. "toddler*".ti,ab.
37. exp infants/
38. exp preschool children/ or children/
39. 32 or 33 or 34 or 35 or 36 or 37 or 38
40. 31 and 39
41. "child* developm* ".ti,ab.
42. "child* neurodevelopm*".ti,ab.
43. "neurodevelopm*".ti,ab.
44. (socio-emotion* develop* or socioemotion* develop*).ti,ab.
45. "social developm*".ti,ab.
46. "emotion* developm*".ti,ab.
47. (sensorymotor developm* or sensory-motor developm*).ti,ab.
48. "motor developm*".ti,ab.
49. child development/ or exp early childhood development/ or exp infant development/ or exp psychomotor development/
50. 41 or 42 or 43 or 44 or 45 or 46 or 47 or 48 or 49
51. cognition.ti,ab.
52. "cogniti* performance".ti,ab.
53. intelligence.ti,ab.
54. IQ.ti,ab.
55. "executive function*".ti,ab.
56. reasoning.ti,ab.
57. language.ti,ab.
58. attention.ti,ab.
59. memory.ti,ab.

- 60. learning.ti,ab.
- 61. early learning.ti,ab.
- 62. information processing.ti,ab.
- 63. literacy.ti,ab.
- 64. reading.ti,ab.
- 65. math.ti,ab.
- 66. exp mental ability/ or exp cognitive development/ or exp memory/
- 67. exp learning/
- 68. 51 or 52 or 53 or 54 or 55 or 56 or 57 or 58 or 59 or 60 or 61 or 62 or 63 or 64 or 65 or 66 or 67
- 69. education.ti,ab.
- 70. "school* achievement".ti,ab.
- 71. "school* performance".ti,ab.
- 72. school retention.ti,ab.
- 73. academic.ti,ab.
- 74. schooling.ti,ab.
- 75. preacademic.ti,ab.
- 76. education/ or exp early childhood education/ or exp elementary education/ or exp extension education/ or exp higher education/ or exp primary education/ or exp professional education/ or exp secondary education/ or exp academic achievement/ or exp educational performance/
- 77. 69 or 70 or 71 or 72 or 73 or 74 or 75 or 76
- 78. "behavi?r* problem*".ti,ab.
- 79. "emotion* problem*".ti,ab.
- 80. "temperament*".ti,ab.
- 81. self regulation.ti,ab.
- 82. attachment.ti,ab.
- 83. self esteem.ti,ab.

84. self efficacy.ti,ab.
85. "social competen*".ti,ab.
86. "peer relationship*".ti,ab.
87. pro-social behavi?r.ti,ab.
88. hyperactivity.ti,ab.
89. impulsivity.ti,ab.
90. attention* deficit hyperactivity disorder*.ti,ab.
91. aggression.ti,ab.
92. exp mental health/ or exp mental disorders/
93. depression/
94. psychology/ or exp adolescent development/ or exp adult development/ or exp emotional development/ or exp self esteem/ or exp self reliance/
95. exp attention deficit hyperactivity disorder/
96. 78 or 79 or 80 or 81 or 82 or 83 or 84 or 85 or 86 or 87 or 88 or 89 or 90 or 91 or 92 or 93 or 94 or 95
97. 50 or 68 or 77 or 96
98. 40 and 97
99. limit 98 to yr="1995 -Current"

PsyclINFO

1. severe acute malnutrition.ti,ab.
2. acute malnutrition.ti,ab.
3. (acute adj2 malnutrition).ti,ab.
4. "sever* acute* maln*".ti,ab.
5. protein-energy malnutrition.ti,ab.
6. protein calorie malnutrition.ti,ab.


7. (protein malnutrition or energy malnutrition or energy-protein malnutrition).ti,ab.
8. "acute* maln*".ti,ab.
9. "kwas?io?kor*".ti,ab.
10. "marasm*".ti,ab.
11. "marasm* kwas?io?kor*".ti,ab.
12. wasting.ti,ab.
13. wasted.ti,ab.
14. (acute adj2 wasting).ti,ab.
15. (severe adj2 wasting).ti,ab.
16. acute undernutrition.ti,ab.
17. (acute adj2 undernutrition).ti,ab.
18. severe undernutrition.ti,ab.
19. (severe adj2 undernutrition).ti,ab.
20. "mid* upper arm circumference".ti,ab.
21. MUAC.ti,ab.
22. "acute* undernourish*".ti,ab.
23. "emaciat*".ti,ab.
24. "acut* maln*".ti,ab.
25. nutritional deficiencies/ or protein deficiency disorders/ or exp failure to thrive/ or exp underweight/
26. exp Kwashiorkor/
27. *protein deficiency disorders/
28. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23 or 24 or 25 or 26 or 27
29. "child*".ti,ab.
30. "infan*".ti,ab.
31. "pre-school*".ti,ab.

32. "preschool*".ti,ab.
33. "toddler*".ti,ab.
34. 29 or 30 or 31 or 32 or 33
35. 28 and 34
36. "child* developm* ".ti,ab.
37. "child* neurodevelopm*".ti,ab.
38. "neurodevelopm*".ti,ab.
39. (socio-emotion* develop* or socioemotion* develop*).ti,ab.
40. "social developm*".ti,ab.
41. "emotion* developm*".ti,ab.
42. (sensorymotor developm* or sensory-motor developm*).ti,ab.
43. "motor developm*".ti,ab.
44. "motor neurodevelopm*".ti,ab.
45. psychomotor developm*.ti,ab.
46. exp infant development/ or exp early childhood development/
47. *childhood development/ or exp early childhood development/ or exp motor development/ or exp psychological development/ or exp psychomotor development/
48. exp emotional development/ or exp psychological development/
49. 36 or 37 or 38 or 39 or 40 or 41 or 42 or 43 or 44 or 45 or 46 or 47 or 48
50. cognition.ti,ab.
51. "cogniti* performance".ti,ab.
52. intelligence.ti,ab.
53. IQ.ti,ab.
54. "executive function*".ti,ab.
55. reasoning.ti,ab.
56. language.ti,ab.

- 57. attention.ti,ab.
- 58. memory.ti,ab.
- 59. learning.ti,ab.
- 60. early learning.ti,ab.
- 61. information processing.ti,ab.
- 62. literacy.ti,ab.
- 63. reading.ti,ab.
- 64. math.ti,ab.
- 65. *cognitive processes/ or exp cognitive assessment/ or exp learning/ or exp memory/ or exp neurocognition/
- 66. *learning/ or adult learning/ or problem based learning/ or school learning/ or skill learning/ or verbal learning/
- 67. exp Fine Motor Skill Learning/ or exp Learning Disabilities/ or exp Learning Disorders/ or exp Gross Motor Skill Learning/
- 68. *language disorders/ or *specific language impairment/ or exp language delay/ or exp language development/
- 69. *cognitive development/ or exp brain development/ or exp speech development/
- 70. *cognition/ or exp cognitive impairment/
- 71. *intelligence/ or exp intellectual development/ or exp intelligence quotient/
- 72. 50 or 51 or 52 or 53 or 54 or 55 or 56 or 57 or 58 or 59 or 60 or 61 or 62 or 63 or 64 or 65 or 66 or 67 or 68 or 69 or 70 or 71
- 73. education.ti,ab.
- 74. "school* achievement".ti,ab.
- 75. "school* performance".ti,ab.
- 76. school retention.ti,ab.
- 77. academic.ti,ab.
- 78. pre-academic.ti,ab.
- 79. schooling.ti,ab.
- 80. *education/
- 81. education/ or exp academic achievement/ or exp school attendance/ or exp school dropouts/ or exp school learning/
- 82. 73 or 74 or 75 or 76 or 77 or 78 or 79 or 80 or 81

83. "behavi?r* problem*".ti,ab.
84. "emotion* problem*".ti,ab.
85. "temperament*".ti,ab.
86. self regulation.ti,ab.
87. attachment.ti,ab.
88. self esteem.ti,ab.
89. self efficacy.ti,ab.
90. "social competen*".ti,ab.
91. "peer relationship*".ti,ab.
92. pro-social behavi?r.ti,ab.
93. hyperactivity.ti,ab.
94. impulsivity.ti,ab.
95. attention* deficit hyperactivity disorder*.ti,ab.
96. aggression.ti,ab.
97. *mental health/
98. exp behavior problems/ or exp behavior disorders/ or exp conduct disorder/
99. exp anxiety disorders/
100. exp hyperkinesis/ or exp attention deficit disorder with hyperactivity/
101. *deprivation/ or exp psychological stress/ or exp stress/
102. *affective disorders/ or *major depression/
103. 83 or 84 or 85 or 86 or 87 or 88 or 89 or 90 or 91 or 92 or 93 or 94 or 95 or 96 or 97 or 98 or 99 or 100 or 101 or 102
104. 49 or 72 or 82 or 103
105. 35 and 104
106. limit 105 to (human and yr="1995 –Current)

Supplementary Figure 1 – Flow chart of literature search



Supplementary Table 1 – Study Characteristics

Study (Year published)	Country (Years conducted)	Study Design (Sample Size)	Age exposed to malnutrition (Age at follow- up if different)	Malnutrition definition (reference population)	Outcome	Outcome Measurement Tool
Chen <i>et al</i> (2021)	China (2018)	Cross-sectional study (23 cases, 1270 controls)	3-5 years	w/h z score <-2 (WHO)	Cognition/ School achievement, Behaviour/ mental health	WPPSI-IV SDQ
Mwene-Batu <i>et al</i> (2020)	DRC (1988 – 2019)	Retrospective cohort (524 cases, 407 controls)	median 41 months (median 22 years)	w/h z score <-3 (WHO)	Cognition/ School achievement, Behaviour/ mental health	MMSE, Rosenberg self-esteem scale, WHODAS
Asiki <i>et al</i> (2019)	Uganda (1999-2011)	Retrospective cohort (12 cases with wasting, 145 cases recovered from wasting, 464 controls)	2-5 years (13-16 years)	w/h z score <-2 (WHO)	Cognition/ School achievement	Parental report of school years achieved
Dwivedi <i>et al</i> (2018)	India (2014-15)	Cross-sectional study (102 cases, 101 controls)	6-30 months	WHO definition of severe acute malnutrition	Neurodevelopment	DASII (Indian modification of BSID)

Study (Year published)	Country (Years conducted)	Study Design (Sample Size)	Age exposed to malnutrition (Age at follow- up if different)	Malnutrition definition (reference population)	Outcome	Outcome Measurement Tool
Kang <i>et al</i> (2018)	Bangladesh (2014), Bhutan (2011), Nepal (2015), Punjab, Pakistan (2016), Sindh, Pakistan (2015)	Cross-sectional study (31037 participants)	36-59 months	w/h z score <-2 (WHO)	Neurodevelopment	MICS ECDI
Lelijveld <i>et al</i> (2018)	Malawi (2006-2013)	Prospective cohort (171 cases, 155 controls)	median 21.5 months [IQR 15-32] (9 years)	w/h <70% of the median (NCHS)	Cognition/ School achievement	CANTAB, Parental report of school grade
Abessa <i>et al</i> (2017)	Ethiopia (2011-13)	Cross-sectional study (310 cases, 310 controls)	cases - mean 30.7 months (SD 15.2) controls – mean 29.6 months (SD 15.4)	w/h <70% of the median (NCHS), MUAC <110mm, or bilateral pitting oedema due to malnutrition	Neurodevelopment, Behaviour/ Mental Health	Denver II, ASQ
Sudfeld <i>et al</i> (2015)	Tanzania (2010-14)	Randomised Controlled Trial	18-36 months	w/h z score <-2 (WHO)	Neurodevelopment	BSID-III

Study (Year published)	Country (Years conducted)	Study Design (Sample Size)	Age exposed to malnutrition (Age at follow- up if different)	Malnutrition definition (reference population)	Outcome	Outcome Measurement Tool
		(47 cases, 989 controls)				
De Grandis <i>et al</i> (2014)	Argentina (not stated)	Retrospective cohort (25 cases, 28 controls)	0-2 years (5-12 years)	w/h z score <-3 (or <70% expected) (WHO)	Behaviour/ Mental Health	Paediatric quality of life inventory
Malhi <i>et al</i> (2013)	India (not stated)	Cross-sectional study (20 cases, 44 controls)	0-6 years	w/a z-score < -2 (WHO)	Neurodevelopment	Indian development inventory
Bogale <i>et al</i> (2013)	Ethiopia (2007)	Cross-sectional study (12 cases, 88 controls)	61.0 months +- 3.0	w/a z-score < -2 (WHO)	Cognition / School achievement	Raven's CPM, KABC-II
Warsito <i>et al</i> (2012)	Indonesia (not stated)	Cross-sectional study (5 cases, 48 controls)	3-5 years old	w/h z-score < -2 (WHO)	Cognition/ School achievement	Indonesian Department of National Education – child development instrument
Nassar <i>et al</i>	Egypt	Cross-sectional study	3-6 years old	Wellcome classification	Cognition/ School achievement	Stanford-Binet-Intelligence-Scale (Arabic-translation)

Study (Year published)	Country (Years conducted)	Study Design (Sample Size)	Age exposed to malnutrition (Age at follow- up if different)	Malnutrition definition (reference population)	Outcome	Outcome Measurement Tool
(2012)	(not stated)	(33 cases, 30 controls)		(WHO)		Receptive/Expressive/-Total- Language (Arabic-Language-Test)
Baker- Henningham <i>et al</i> (2009)	Bangladesh (not stated)	Cross-sectional study (212 cases, 108 controls)	6-24 months old	w/a z-score < -2 (NCHS)	Behaviour/ Mental Health	Temperament Questionnaire derived from existing validated instruments (modified by Wachs, Purdue University)
El-Khayat <i>et al</i> (2007)	Egypt (2006)	Cross-sectional study (42 cases, 15 controls)	6-25 months	w/h z-score <-2 (WHO)	Neurodevelopment	BSID-II
Liu <i>et al</i> (2004)	Mauritius (1972-)	Prospective Cohort (235 cases, 807 controls)	3 years (8,11, 17 years)	Malnutrition and one of angular stomatitis, kwashiorkor, sparse thin hair, haemoglobin <8.5gr/dl	Behaviour/ Mental Health	Child behaviour questionnaire at age 8, Child Behaviour Checklist at age 11, Revised Behaviour Checklist at Age 17
Liu <i>et al</i> (2003)	Mauritius (1972-)	Prospective Cohort (253 cases, 837 controls)	3 years (11 years)	Malnutrition and one of angular stomatitis, kwashiorkor, sparse thin hair,	Cognition/ School achievement	Bohem Test WISC Academic Tests Holborn-Reading Scale Trail Making Test

Study (Year published)	Country (Years conducted)	Study Design (Sample Size)	Age exposed to malnutrition (Age at follow- up if different)	Malnutrition definition (reference population)	Outcome	Outcome Measurement Tool
				haemoglobin <8.5gr/dl		
Drewett <i>et al</i> (2001)	Ethiopia (not stated)	Prospective Cohort (97 cases, 100 controls)	2 years	w/a z-score < - 1.88 (NCHS)	Neurodevelopment	BSID
Vazir <i>et al</i> (1998)	India (not stated)	Cross-sectional study (1456 cases, 2212 controls)	0-6 years	<75% expected w/a (NCHS)	Neurodevelopment	ICMR Psychosocial Developmental Screening Test
Perales <i>et al</i> (1996)	Chile (1987 – not stated)	Retrospective Cohort (40 cases, 40 controls)	0-2 years (8-10 years)	Protein energy malnutrition (defined by Sempe (1979) <i>et al</i>)	Cognition/ School achievement	Continuous Performance Task Anstey Domino Test
Kaul <i>et al</i> (1995)	India (not stated)	Cross-sectional study (50 cases, 102 controls)	0-12 months	Gomez classification – moderate malnutrition (NCHS)	Neurodevelopment	BSID
Barbados Nutrition Study						

Study (Year published)	Country (Years conducted)	Study Design (Sample Size)	Age exposed to malnutrition (Age at follow- up if different)	Malnutrition definition (reference population)	Outcome	Outcome Measurement Tool
Hock <i>et al</i>* (2018)	Barbados (1977-not stated)	Prospective cohort (77 cases, 62 controls)	0-1 years (40-45 years)	Gomez classification	Behaviour/mental health	SCID-II-PQ, NEO PI-R FFM
Waber (1) <i>et al</i>* (2014)	Barbados (1977-2010)	Prospective cohort (77 cases, 59 controls)	0-1 years (38 years)	Gomez classification	Cognition/ School achievement	WASI, WRAT-III
Waber (2) <i>et al</i>* (2014)	Barbados (1977-2010)	Prospective cohort (77 cases, 59 controls)	0-1 years (38 years)	Gomez classification	Cognition / School achievement, Behaviour/ Mental Health	WAIS-III, D-KEFS, WRAML-II, Wisconsin card sorting, Metacognitive index, Behavioural regulation index
Galler <i>et al</i>* (2013)	Barbados (2006-2010)	Prospective cohort (77 cases, 57 controls)	0-1 years (37 - 43 years)	Gomez classification	Behaviour/ Mental Health	NEO-PI-R personality inventory
Galler (1) <i>et al</i>* (2012)	Barbados (1967-2010)	Prospective cohort (80 cases, 65 controls)	0-1 years (40 years)	Gomez classification	Behaviour/ Mental Health	CAARS, CPT
Galler (2) <i>et al</i>* (2012)	Barbados (1967-1985)	Prospective cohort	0-1 years (15 years old)	Gomez classification	Behaviour/ Mental Health	Child-Behaviour-Questionnaire, Teacher-Behaviour- Questionnaire

Study (Year published)	Country (Years conducted)	Study Design (Sample Size)	Age exposed to malnutrition (Age at follow- up if different)	Malnutrition definition (reference population)	Outcome	Outcome Measurement Tool
(2012)		(56 cases, 60 controls)				
Waber <i>et al</i>* (2011)	Barbados (1967-1985)	Prospective cohort (57 cases, 60 controls)	0-1 years (11-18 years)	Gomez classification	Cognition/ School achievement and Behaviour/ Mental Health	WISC, Common entrance examination (local school test), Teacher-Behaviour Questionnaire, School functioning scale Minnesota-General-Adjustment- and Morale-Scale
Galler <i>et al</i>* (2011)	Barbados (1967-1985)	Prospective cohort (109 cases, 107 controls)	0-1 years (9-17 years)	Gomez classification	Behaviour/ Mental Health	Barbados-Child-Behaviour- Scale (BCBS)
Galler <i>et al</i>* (2010)	Barbados (1967-1985)	Prospective cohort (116 cases, 61 controls)	0-1 years (11-17 years old)	Gomez classification	Behaviour/ Mental Health	Minnesota-General- Adjustment-and Morale-Scale

*Studies are drawn from the same Barbados Nutrition Study Cohort

ASQ – Ages & Stages Questionnaire, BSID – Bayley scales of infant development, CANTAB – Cambridge Neuropsychological Test Automated Battery, CAARS - Attention-domain: Conners-ADHD-Rating-Scales, CPT – Conners-Continuous-Performance Test, DASII – Developmental Assessment Scale of Indian Infants, DRC – Democratic Republic of Congo, D-KEFS – Delis-Kaplan Executive Function System, ICMR – Indian Council Medical Research Psychosocial Developmental Screening Test, KABC-II – Kaufman Assessment Battery for Children-II, MDAT – Malawi Developmental Assessment Tool, MICS ECDI – Multiple Indicator Cluster Survey Early Child Development Index, MMSE – Mini Mental State Exam, MUAC – Mean Upper Arm Circumference, NEO PI-R FFM – NEO Personality Inventory-Revised derived Five-Factor Model, NCHS – National Centre for Health Statistics reference population, Raven’s CPM – Raven’s

Colored Progressive Matrices, SCID-II-PQ – Structured Clinical Interview for DSM-IV Axis II Personality Disorders Personality Questionnaire, SDQ – Strengths and Difficulties Questionnaire, w/h – Weight-for-Height, WAIS-III – Wechsler Adult Intelligence Scale-III, WASI – Adult IQ: Weschler Abbreviated Scale of Intelligence – Vocabulary and Matrix Reasoning subsets, WHO – World Health Organization reference population, WHODAS – World Health Organization Disability Assessment Schedule, WISC – Weschler Intelligence Scale, WPPSI-IV – Wechsler Preschool and Primary Scale of Intelligence Fourth Edition, WRAML-II – Wide Range Assessment of Memory and Learning-II, WRAT-III – Academic achievement: Wide Range Achievement Test-III – Reading Spelling and Calculation subsets

Supplementary Table 2 – Study quality scoring

Study (Year published)	External Validity	Internal Validity	Overall Study Quality
Chen <i>et al</i> (2021)	++	++	++
Mwene-Batu <i>et al</i> (2020)	++	+	+
Asiki <i>et al</i> (2019)	++	++	++
Dwivedi <i>et al</i> (2018)	+	+	+
Hock <i>et al</i> (2018)	++	+	+
Kang <i>et al</i> (2018)	++	+	+
Lelijveld <i>et al</i> (2018)	++	++	++
Abessa <i>et al</i> (2017)	++	++	++
Sudfeld <i>et al</i> (2015)	++	++	++
De Grandis <i>et al</i> (2014)	+	-	-
Waber (1) <i>et al</i> (2014)	++	++	++
Waber (2) <i>et al</i> (2014)	++	++	++
Malhi <i>et al</i> (2013)	+	-	-
Galler <i>et al</i> (2013)	++	+	+
Bogale <i>et al</i> (2013)	++	++	++
Galler (1) <i>et al</i> (2012)	++	+	+
Galler (2) <i>et al</i> (2012)	++	+	+
Warsito <i>et al</i> (2012)	++	-	-
Nassar <i>et al</i> (2012)	-	-	-
Waber <i>et al</i> (2011)	++	+	+
Galler <i>et al</i> (2011)	++	++	++
Galler <i>et al</i> (2010)	++	++	++
Baker-Henningham <i>et al</i> (2009)	++	++	++
El-Khayat <i>et al</i> (2007)	+	+	+
Liu <i>et al</i> (2004)	++	++	++
Liu <i>et al</i> (2003)	++	++	++
Drewett <i>et al</i> (2001)	++	++	++
Vazir <i>et al</i> (1998)	++	+	+
Perales <i>et al</i> (1996)	+	+	+
Kaul <i>et al</i> (1995)	+	+	+

++ = very good

+ = adequate

- = poor

Supplementary Table 3 – Results from studies assessing neurodevelopment in children with malnutrition compared to controls

Study (quality)	Neurodevelopment tool/measurement	Summary of results		
Dwivedi <i>et al</i> (2018) (+)	DASII (Indian modification of BSID)	<hr/>		
		Controls n=101	Cases n=102	
		<u>Mental age:</u>		
		10.4 (SE 0.5)	8.2 (SE 0.4)	
		<u>Mental development quotient:</u>		
		83.5 (SE 1.0)	62.1 (SE 0.6)	
		<u>Motor age:</u>		
		10.1 (SE 0.5)	7.9 (SE 0.4)	
		<u>Motor development quotient:</u>		
		80.0 (SE 1.0)	59.0 (SE 0.7)	
<u>Adjusted analysis:</u> (unadjusted analysis)				
<u>Overall findings:</u> Significant developmental delay in children with severe malnutrition compared to controls unadjusted for confounding variables.				
Kang <i>et al</i> 2018 (+)	MICS ECDI	<u>Learning-cognition development (odds of cases being on track compared to controls):</u>		
		Country	Unadjusted Odds Ratio	Adjusted Odds Ratio
		Bangladesh	1.27 [95% CI 0.92, 1.75]	1.27 [95% CI 0.90, 1.79]
		Bhutan	1.60 [95% CI 0.32, 7.85]	1.93 [95% CI 0.39, 9.48]
		Nepal	1.09 [95% CI 0.66, 1.79]	1.52 [95% CI 0.90, 2.58]

		<p>Punjab 0.63 [95% CI 0.50, 0.79] 0.67 [95% CI 0.53, 0.86]</p> <p>Sindh 0.69 [95% CI 0.53, 0.89] 0.76 [95% CI 0.59, 0.97]</p> <p>All 0.98 [95% CI 0.70, 1.37]</p> <p>Socio-emotional development (odds of cases being on track compared to controls):</p> <table> <tr> <th>Country</th><th>Unadjusted Odds Ratio</th><th>Adjusted Odds Ratio</th></tr> <tr> <td>Bangladesh</td><td>1.14 [95% CI 0.92, 1.42]</td><td>1.10 [95% CI 0.40, 1.88]</td></tr> <tr> <td>Bhutan</td><td>1.08 [95% CI 0.58, 2.01]</td><td>1.08 [95% CI 0.58, 2.03]</td></tr> <tr> <td>Nepal</td><td>2.40 [95% CI 1.25, 4.58]</td><td>2.54 [95% CI 1.33, 4.85]</td></tr> <tr> <td>Punjab</td><td>0.86 [95% CI 0.73, 1.02]</td><td>0.88 [95% CI 0.74, 1.04]</td></tr> <tr> <td>Sindh</td><td>0.92 [95% CI 0.74, 1.14]</td><td>0.94 [95% CI 0.76, 1.15]</td></tr> <tr> <td>All</td><td></td><td>1.07 [95% CI 0.86, 1.33]</td></tr> </table> <p>Adjusted analysis: (results adjusted for rural/urban, wealth quintile, household size, household head, water source, toilet facility, maternal education, child age and sex, diarrhoea/cough in past 2 weeks, study design effect, early childhood education, support for learning, availability of children's books and playthings, inadequate care)</p> <p>Sample size: n = 31037, wasting prevalence = between 4.1-12.8% across study sites</p> <p>Overall findings: Mixed results between study sites with no significant difference between cases and controls including when adjusted for confounding variables</p>	Country	Unadjusted Odds Ratio	Adjusted Odds Ratio	Bangladesh	1.14 [95% CI 0.92, 1.42]	1.10 [95% CI 0.40, 1.88]	Bhutan	1.08 [95% CI 0.58, 2.01]	1.08 [95% CI 0.58, 2.03]	Nepal	2.40 [95% CI 1.25, 4.58]	2.54 [95% CI 1.33, 4.85]	Punjab	0.86 [95% CI 0.73, 1.02]	0.88 [95% CI 0.74, 1.04]	Sindh	0.92 [95% CI 0.74, 1.14]	0.94 [95% CI 0.76, 1.15]	All		1.07 [95% CI 0.86, 1.33]
Country	Unadjusted Odds Ratio	Adjusted Odds Ratio																					
Bangladesh	1.14 [95% CI 0.92, 1.42]	1.10 [95% CI 0.40, 1.88]																					
Bhutan	1.08 [95% CI 0.58, 2.01]	1.08 [95% CI 0.58, 2.03]																					
Nepal	2.40 [95% CI 1.25, 4.58]	2.54 [95% CI 1.33, 4.85]																					
Punjab	0.86 [95% CI 0.73, 1.02]	0.88 [95% CI 0.74, 1.04]																					
Sindh	0.92 [95% CI 0.74, 1.14]	0.94 [95% CI 0.76, 1.15]																					
All		1.07 [95% CI 0.86, 1.33]																					
Abessa <i>et al</i> (2017) (++)	<p>Denver II</p> <p>ASQ: social-emotional score</p>	<p>Denver II score (incidence rate ratio in cases compared to controls):</p> <p>Fine motor IRR 0.76 [95% CI 0.71, 0.8]</p> <p>Gross motor IRR 0.73 [95% CI 0.71, 0.74]</p> <p>Language IRR 0.98 [95% CI 0.86, 1.11]</p> <p>Personal-social IRR 0.7 [95% CI 0.65, 0.75]</p> <p>Adjusted analysis: (results adjusted for significant variables from age, sex, maternal religion)</p>																					

		<p><u>Sample size:</u> Cases n=310, Controls n=310</p> <p><u>Overall Findings:</u> Children with malnutrition performed worse on personal-social, fine motor, and gross motor skills compared to age and sex-matched controls adjusted for confounding variables. Motor skills are the most and personal social the least affected domains assessed on the Denver II Jimma.</p>		
<p>Sudfeld <i>et al</i> (2015)</p> <p>(++)</p>	BSID-III	<p><u>BSID-II score (association between score and malnutrition):</u> Cognitive -0.63 [95% CI -0.97, -0.29] Communication -0.32 [95% CI -0.64, -0.01] Motor -0.54 [95% CI -0.86, -0.23]</p> <p><u>Adjusted analysis:</u> (results adjusted for height-for-age, stimulation category, sex, age, assessor, maternal education, wealth quintile, randomized regimen)</p> <p><u>Sample size:</u> Cases n=47, Controls n=989</p> <p><u>Overall Findings:</u> Children experiencing wasting had across domain deficits compared to controls adjusted for confounding variables.</p>		
<p>Malhi <i>et al</i> 2013</p> <p>(-)</p>	Indian Development Inventory	<p>Controls n=20</p> <p><u>Inventory domain:</u> Social score 104.64 (SD 17.11) Adaptive score 105.36 (SD 9.71) Motor score 112 (SD 14.5) Communication score 97.82 (SD 14.34) Cognitive score 92.82 (SD 13.92)</p>	<p>Cases n=44</p> <p>101.30 (SD 17.31) 105.25 (SD 8.33) 108.85 (SD 10.93) 94.50 (SD 8.61) 90.1 (SD 9.79)</p>	<p><u>p value</u></p> <p>0.036 0.239 0.074 0.047 0.080</p>

		<div>Total score100.86 (SD 18.15)95 (SD 13.51)0.007</div> <div>Adjusted analysis: Multivariate regression revealed that 26.7% of variance was accounted by household income and stunting, indicating results were no longer significant</div> <div>Overall Findings: Children with wasting compared to controls did not have significantly lower domain scores adjusted for confounding variables.</div>																
<div>El-Khayat et al 2007</div> <div>(+)</div>	BSID-II	<table><tr><td></td><td>Controls n=15</td><td>Cases n=42 (before rehab)</td><td>Cases (after rehab)</td></tr><tr><td>Mental Development Index</td><td>97 (SD 10.92)</td><td>76.3 (SD 8.78)</td><td>88.19 (SD 8.93)</td></tr><tr><td>Psychomotor Development Index</td><td>97 (SD 12.38)</td><td>72.76 (SD 11.27)</td><td>87.39 (SD 8.85)</td></tr></table> <div>Adjusted analysis: (unadjusted analysis)</div> <div>Overall Findings: Cases had lower scores at the start of the study compared to controls unadjusted for confounding variables. Scores were improved after 8 weeks of nutritional rehabilitation but were still significantly lower when compared to controls unadjusted for confounding variables.</div>		Controls n=15	Cases n=42 (before rehab)	Cases (after rehab)	Mental Development Index	97 (SD 10.92)	76.3 (SD 8.78)	88.19 (SD 8.93)	Psychomotor Development Index	97 (SD 12.38)	72.76 (SD 11.27)	87.39 (SD 8.85)				
	Controls n=15	Cases n=42 (before rehab)	Cases (after rehab)															
Mental Development Index	97 (SD 10.92)	76.3 (SD 8.78)	88.19 (SD 8.93)															
Psychomotor Development Index	97 (SD 12.38)	72.76 (SD 11.27)	87.39 (SD 8.85)															
<div>Drewett et al 2001</div> <div>(++)</div>	BSID	<table><tr><td></td><td>Controls n=100</td><td>Cases n=27</td><td>Cases n=70</td></tr><tr><td></td><td></td><td>Early Malnutrition at 4 months</td><td>Late Malnutrition at 12 months</td></tr></table> <div>BSID (score at 2 years):</div> <table><tr><td>Psychomotor score</td><td>10.2 (SD3.7)</td><td>6.6 (SD 4.2)</td><td>8.5 (SD 4.3)</td></tr><tr><td>Mental score</td><td>28.9 (SD5.8)</td><td>22.6 (SD 6.2)</td><td>26.6 (SD 6.1)</td></tr></table> <div>Adjusted analysis: Weight at 2 years associated with BSID outcome, but early and late growth faltering before 1 year not independently associated with BSID scores after adjusting for weight at 2 years and other covariates (sex, mother’s education, possessions).</div>		Controls n=100	Cases n=27	Cases n=70			Early Malnutrition at 4 months	Late Malnutrition at 12 months	Psychomotor score	10.2 (SD3.7)	6.6 (SD 4.2)	8.5 (SD 4.3)	Mental score	28.9 (SD5.8)	22.6 (SD 6.2)	26.6 (SD 6.1)
	Controls n=100	Cases n=27	Cases n=70															
		Early Malnutrition at 4 months	Late Malnutrition at 12 months															
Psychomotor score	10.2 (SD3.7)	6.6 (SD 4.2)	8.5 (SD 4.3)															
Mental score	28.9 (SD5.8)	22.6 (SD 6.2)	26.6 (SD 6.1)															

		<p><u>Overall Findings:</u> Significantly lower psychomotor and mental scores at 2 years in early growth falterers and late growth falterers compared to controls. Difference is accounted for by differences in weight at 2 years when adjusted for in analysis.</p>																		
<p>Vazir <i>et al</i> 1998</p> <p>(+)</p>	ICMR Psychosocial Developmental Screening Test	<p><u>ICMR Psychosocial Development Screening Test Results:</u> Well nourished children attained developmental milestones at a younger age in the five areas of development compared to undernourished children.</p> <p><u>Adjusted analysis:</u> (unadjusted analysis)</p> <p><u>Sample size:</u> Cases n=1456, Controls n=2212</p> <p><u>Overall Findings:</u> Malnourished children attained developmental milestones at a later age unadjusted for confounders.</p>																		
<p>Kaul <i>et al</i> 1995</p> <p>(+)</p>	BSID	<table> <tr> <th></th><th>Controls n=102</th><th>Cases n=50</th></tr> <tr> <td colspan="3"><u>BSID (mental quotient):</u></td></tr> <tr> <td>Male (0-6 months)</td><td>98.80 (SD 11.94)</td><td>57.60 (SD 6.77)</td></tr> <tr> <td>Female (0-6 months)</td><td>97.28 (SD 10.22)</td><td>59.85 (SD 11.94)</td></tr> <tr> <td>Male (6-12 months)</td><td>100.86 (SD 4.39)</td><td>79.93 (SD 5.57)</td></tr> <tr> <td>Female (6-12 months)</td><td>103.68 (SD 4.31)</td><td>75.15 (SD 3.39)</td></tr> </table> <p><u>Adjusted analysis:</u> (unadjusted analysis)</p> <p><u>Overall Findings:</u> Children with moderate malnutrition had a lower mental quotient than controls unadjusted for confounding variables.</p>		Controls n=102	Cases n=50	<u>BSID (mental quotient):</u>			Male (0-6 months)	98.80 (SD 11.94)	57.60 (SD 6.77)	Female (0-6 months)	97.28 (SD 10.22)	59.85 (SD 11.94)	Male (6-12 months)	100.86 (SD 4.39)	79.93 (SD 5.57)	Female (6-12 months)	103.68 (SD 4.31)	75.15 (SD 3.39)
	Controls n=102	Cases n=50																		
<u>BSID (mental quotient):</u>																				
Male (0-6 months)	98.80 (SD 11.94)	57.60 (SD 6.77)																		
Female (0-6 months)	97.28 (SD 10.22)	59.85 (SD 11.94)																		
Male (6-12 months)	100.86 (SD 4.39)	79.93 (SD 5.57)																		
Female (6-12 months)	103.68 (SD 4.31)	75.15 (SD 3.39)																		

95% CI – 95% confidence intervals, BSID – Bayley Scales of Infant Development, DASII – Developmental Assessment Scale of Indian Infants, ICMR – Indian Council Medical Research Psychosocial Developmental Screening Test, IDI – Indian Developmental Inventory, IRR – incidence rate ratio, MDAT – Malawi Developmental Assessment Tool, MICS ECDI – Multiple Indicator Cluster Survey Early Child Development Index, SD – Standard Deviation, SE – standard error

Supplementary Table 4 - Results from studies assessing cognitive outcomes in those exposed to childhood malnutrition compared to controls

Study (quality)	Cognition tool/measurement (outcome age)	Summary of results		
Chen <i>et al</i> 2021 (++)	WPPSI-IV (childhood)	Controls n=1270		Cases n=23
		<u>WPPSI-IV:</u>		
		Working Memory Index	90.35 (SD 0.38) [95% CI 89.60, 91.09]	95.09 (SD 2.67) [95% CI 89.86, 100.32]
		Verbal Comprehension Index	86.00 (SD 0.36) [95% CI 85.30, 86.70]	87.26 (SD 3.29) [95% CI 80.80, 93.72]
		<u>Adjusted analysis:</u> Wasting was not statistically associated with any developmental outcome variables after adjusting for vector of child, parent, caregiver and household-level characteristics.		
Mwene-Batu <i>et al</i> 2020 (+)	MMSE Academic performance (adulthood)	Controls n=407		Cases n=524
		<u>MMSE:</u>		p value
		Mean score	25.6 (SD 2.6)	27.8 (SD 2.2)
		Normal score (%)	78.0	90.1
		<u>Academic performance (based on self-report):</u>		0.001 <0.001

		<div>Low (%)15.223.8</div> <div>Average (%)49.045.1</div> <div>High (%)35.831.00.014</div> <div>Adjusted analysis: (unadjusted analysis)</div> <div>Overall findings: Malnutrition survivors had a significantly lower probability of attaining a high education level, reported worse academic performance and a significantly lower mean cognition score unadjusted for confounding variables.</div>																					
Asiki <i>et al</i> 2019 (++)	Academic achievement (adolescence)	<table><tr><td>Controls n=464</td><td>Children with ongoing wasting n=12</td><td>Children recovered from wasting n=145</td></tr><tr><td colspan="3">School years achieved (parental report):</td></tr><tr><td>9.2 years (SD 11)</td><td>6.2 (SD 2.5)</td><td>7.8 (SD 3.8)</td></tr><tr><td colspan="3">Adjusted correlation coefficient:</td></tr><tr><td></td><td>-1.91 [95% CI -4.51, 0.68]</td><td>- 2.05 [95% CI -3.30, -0.79]</td></tr><tr><td colspan="3">Adjusted analysis: (adjusted for age group, sex, education, tribe, wall of house type, reported alcohol consumption)</td></tr><tr><td colspan="3">Overall findings: Wasting was associated with fewer schooling years regardless of recovery from undernutrition adjusted for confounding variables.</td></tr></table>	Controls n=464	Children with ongoing wasting n=12	Children recovered from wasting n=145	School years achieved (parental report):			9.2 years (SD 11)	6.2 (SD 2.5)	7.8 (SD 3.8)	Adjusted correlation coefficient:				-1.91 [95% CI -4.51, 0.68]	- 2.05 [95% CI -3.30, -0.79]	Adjusted analysis: (adjusted for age group, sex, education, tribe, wall of house type, reported alcohol consumption)			Overall findings: Wasting was associated with fewer schooling years regardless of recovery from undernutrition adjusted for confounding variables.		
Controls n=464	Children with ongoing wasting n=12	Children recovered from wasting n=145																					
School years achieved (parental report):																							
9.2 years (SD 11)	6.2 (SD 2.5)	7.8 (SD 3.8)																					
Adjusted correlation coefficient:																							
	-1.91 [95% CI -4.51, 0.68]	- 2.05 [95% CI -3.30, -0.79]																					
Adjusted analysis: (adjusted for age group, sex, education, tribe, wall of house type, reported alcohol consumption)																							
Overall findings: Wasting was associated with fewer schooling years regardless of recovery from undernutrition adjusted for confounding variables.																							
Lelijveld <i>et al</i> 2018 (++)	Academic achievement CANTAB (adolescence)	<table><tr><td>Controls n=178</td><td>Cases n=315</td></tr><tr><td colspan="2">School grade achieved (parental report):</td></tr><tr><td>3.1 (SD 1.6)</td><td>2.5 (SD 1.3)</td></tr><tr><td colspan="2">Logistic regression comparing school grade in groups:</td></tr><tr><td>OR 0.5 [95% CI 0.4, 0.7]</td><td></td></tr><tr><td>Adjusted OR 0.54 [95% CI 0.35, 0.81]</td><td></td></tr></table>	Controls n=178	Cases n=315	School grade achieved (parental report):		3.1 (SD 1.6)	2.5 (SD 1.3)	Logistic regression comparing school grade in groups:		OR 0.5 [95% CI 0.4, 0.7]		Adjusted OR 0.54 [95% CI 0.35, 0.81]										
Controls n=178	Cases n=315																						
School grade achieved (parental report):																							
3.1 (SD 1.6)	2.5 (SD 1.3)																						
Logistic regression comparing school grade in groups:																							
OR 0.5 [95% CI 0.4, 0.7]																							
Adjusted OR 0.54 [95% CI 0.35, 0.81]																							

Warsito et al 2012 (-)	Indonesian Department of National Education – child development instrument (childhood)	<div> <div>Controls n=48</div> <div>Wasting n=3</div> <div>Severe Wasting n=2</div> </div>		
		<p>Child development score:</p> <p>Percentage score 68.7 73.9 73.8</p> <p>Adjusted analysis: (results unadjusted)</p> <p>Overall findings: Those with wasting in this study had better development scores than those without wasting. The study is limited by small case numbers.</p>		
Nassar et al 2012 (-)	Stanford-Binet-Intelligence-Scale Arabic-Language-Test (childhood)	<div> <div>Controls n=30</div> <div>Cases n=33</div> <div>p value</div> </div>		
		<p>Stanford-Binet-Intelligence-Scale (mean):</p> <p>IQ score 92.07 (SD 11.52) 80.52 (SD 13.65) 0.001</p> <p>Mental age (months) 50.50 (SD 13.40) 42.58 (SD 8.66) 0.017</p> <p>Arabic Language Test (mean):</p> <p>Receptive language age (months) 51.67 (SD 13.98) 40.09 (SD 17.36) 0.005</p> <p>Expressive language age (months) 51.67 (SD 14.23) 40.76 (SD 16.74) 0.003</p> <p>Total language age (months) 51.40 (SD 14.15) 40.00 (SD 17.66) 0.007</p> <p>Adjusted analysis: (results unadjusted)</p> <p>Overall findings: Those with malnutrition had poorer scores for both tests than age- and sex-matched controls unadjusted for confounding variables.</p>		
Liu et al 2003 (++)	Bohem Test WISC Academic Tests Holborn-Reading Scale Trail Making Test	<div> <div>Control n=837</div> <div>Cases n=253</div> <div>p value</div> </div>		
		<p>Age 3y:</p> <p>Verbal IQ 100.92 (SD 15.04) 96.93 (SD 14.36) 0.001</p> <p>Spatial IQ 100.34 (SD 15.21) 98.95 (SD 13.52) 0.18</p> <p>Full-Scale IQ 100.92 (SD 15.04) 96.93 (SD 14.36) 0.001</p>		

	(adolescence)	<p><u>Age 11y:</u></p> <table><tr><td>Verbal IQ</td><td>100.90 (SD 14.41)</td><td>96.16 (SD 15.40)</td><td>0.001</td></tr><tr><td>Spatial IQ</td><td>100.75 (SD 14.55)</td><td>96.23 (SD 15.41)</td><td>0.001</td></tr><tr><td>Full-Scale IQ</td><td>100.88 (SD 14.43)</td><td>95.83 (SD 15.49)</td><td>0.001</td></tr><tr><td>Reading</td><td>95.11 (SD 55.30)</td><td>76.81 (SD 56.58)</td><td>0.001</td></tr><tr><td>School performance</td><td>10.33 (SD 7.07)</td><td>7.63 (SD 6.48)</td><td>0.001</td></tr><tr><td>Trails A</td><td>-0.049 (SD 0.95)</td><td>0.24 (SD 1.10)</td><td>0.001</td></tr><tr><td>Trails B</td><td>-0.023 (SD 0.98)</td><td>0.20 (SD 1.07)</td><td>0.004</td></tr></table> <p><u>Adjusted analysis:</u> Differences remain after controlling for psychosocial adversity.</p> <p><u>Overall Findings:</u> Malnourished children had poorer cognition and school/academic performance at both ages after controlling for psychosocial adversity.</p>	Verbal IQ	100.90 (SD 14.41)	96.16 (SD 15.40)	0.001	Spatial IQ	100.75 (SD 14.55)	96.23 (SD 15.41)	0.001	Full-Scale IQ	100.88 (SD 14.43)	95.83 (SD 15.49)	0.001	Reading	95.11 (SD 55.30)	76.81 (SD 56.58)	0.001	School performance	10.33 (SD 7.07)	7.63 (SD 6.48)	0.001	Trails A	-0.049 (SD 0.95)	0.24 (SD 1.10)	0.001	Trails B	-0.023 (SD 0.98)	0.20 (SD 1.07)	0.004				
Verbal IQ	100.90 (SD 14.41)	96.16 (SD 15.40)	0.001																															
Spatial IQ	100.75 (SD 14.55)	96.23 (SD 15.41)	0.001																															
Full-Scale IQ	100.88 (SD 14.43)	95.83 (SD 15.49)	0.001																															
Reading	95.11 (SD 55.30)	76.81 (SD 56.58)	0.001																															
School performance	10.33 (SD 7.07)	7.63 (SD 6.48)	0.001																															
Trails A	-0.049 (SD 0.95)	0.24 (SD 1.10)	0.001																															
Trails B	-0.023 (SD 0.98)	0.20 (SD 1.07)	0.004																															
<p>Perales <i>et al</i> 1996</p> <p>(+)</p>	<p>Continuous Performance Task</p> <p>Anstey Domino Test</p> <p>(childhood)</p>	<table><tr><td></td><td>Controls n=40</td><td>Cases n=40</td><td>p value</td></tr><tr><td colspan="4"><u>Short-term Memory:</u></td></tr><tr><td>Number-of-digits score</td><td>4.5 (SD 0.87)</td><td>3.8 (SD 0.88)</td><td><0.05</td></tr><tr><td colspan="4"><u>Problem-Solving:</u></td></tr><tr><td>Correct score</td><td>4.7 (SD 2.1)</td><td>3.4 (SD 1.4)</td><td><0.05</td></tr><tr><td colspan="4"><u>Attention:</u> No differences in correct answers between groups. Lower performance in cases response time for correct answers (p<0.02).</td></tr><tr><td colspan="4"><u>Adjusted analysis:</u> (unadjusted analysis)</td></tr><tr><td colspan="4"><u>Overall Findings:</u></td></tr></table>		Controls n=40	Cases n=40	p value	<u>Short-term Memory:</u>				Number-of-digits score	4.5 (SD 0.87)	3.8 (SD 0.88)	<0.05	<u>Problem-Solving:</u>				Correct score	4.7 (SD 2.1)	3.4 (SD 1.4)	<0.05	<u>Attention:</u> No differences in correct answers between groups. Lower performance in cases response time for correct answers (p<0.02).				<u>Adjusted analysis:</u> (unadjusted analysis)				<u>Overall Findings:</u>			
	Controls n=40	Cases n=40	p value																															
<u>Short-term Memory:</u>																																		
Number-of-digits score	4.5 (SD 0.87)	3.8 (SD 0.88)	<0.05																															
<u>Problem-Solving:</u>																																		
Correct score	4.7 (SD 2.1)	3.4 (SD 1.4)	<0.05																															
<u>Attention:</u> No differences in correct answers between groups. Lower performance in cases response time for correct answers (p<0.02).																																		
<u>Adjusted analysis:</u> (unadjusted analysis)																																		
<u>Overall Findings:</u>																																		

		Cases had lower memory scores and problem solving scores unadjusted for confounding variables. Cases had worse performance than controls for response times in the attention test unadjusted for confounding variables.																		
Barbados Nutrition Study																				
Waber (1) <i>et al 2014*</i> (++)	Academic achievement, WASI (adulthood)	<p><u>WRAT-III academic achievement score (adjusted odds ratio):</u> Reading OR 3.44 [95% CI 1.88–6.30] Spelling OR 6.10 [95% CI 3.34–11.16] Calculation OR 6.23 [95% CI 3.15–12.32]</p> <p><u>WASI score (adjusted odds ratio):</u> IQ OR 9.18 [95% CI 3.50, 24.13]</p> <p><u>Adjusted analysis:</u> (Results adjusted for childhood standard of living)</p> <p><u>Sample size:</u> Cases n=77, Controls n=59</p> <p><u>Overall findings:</u> Academic skills and IQ were lower in the malnourished group compared to controls after adjusting for childhood standard of living.</p>																		
Waber (2) <i>et al 2014*</i> (++)	WAIS-III, D-KEFS, WRAML-II, Wisconsin card sorting, Metacognitive index (adulthood)	<table border="1"> <thead> <tr> <th></th><th>Controls n=59</th><th>Cases n=77</th></tr> </thead> <tbody> <tr> <td><u>Metacognitive index (General executive composite):</u></td><td></td><td></td></tr> <tr> <td>Mean score</td><td>41.81 (SD 6.05)</td><td>47.36 (SD 9.33)</td></tr> <tr> <td>Adjusted Mean difference</td><td colspan="2">3.70 (p<0.05)</td></tr> <tr> <td><u>WRAML-2:</u></td><td></td><td></td></tr> <tr> <td>Scaled score</td><td>8.49 (SD 1.81)</td><td>7.45 (SD 2.28)</td></tr> </tbody> </table>		Controls n=59	Cases n=77	<u>Metacognitive index (General executive composite):</u>			Mean score	41.81 (SD 6.05)	47.36 (SD 9.33)	Adjusted Mean difference	3.70 (p<0.05)		<u>WRAML-2:</u>			Scaled score	8.49 (SD 1.81)	7.45 (SD 2.28)
	Controls n=59	Cases n=77																		
<u>Metacognitive index (General executive composite):</u>																				
Mean score	41.81 (SD 6.05)	47.36 (SD 9.33)																		
Adjusted Mean difference	3.70 (p<0.05)																			
<u>WRAML-2:</u>																				
Scaled score	8.49 (SD 1.81)	7.45 (SD 2.28)																		

		<p>Adjusted Mean difference -0.42 (p>0.05)</p> <p><u>Wisconsin card sorting:</u> Total errors score 94.56 (SD 12.70) 80.67 (SD 12.53)</p> <p>Adjusted Mean difference -9.74 (p<0.001)</p> <p><u>Adjusted analysis</u> (results adjusted for childhood standard of living and IQ)</p> <p>Significant differences persisted for the WAIS Digit Symbol and Letter-Number, D-KEFS Trails (Numbers) , Verbal Fluency (Letters) and Rey-Osterrieth Complex Figure after adjusting for covariates</p> <p><u>Overall findings:</u> Malnutrition during infancy is associated with impaired IQ in adulthood. In models adjusted only for childhood standard of living, the malnutrition group performed more poorly on nearly all measures. In the IQ-adjusted models, significant differences persisted for the WAIS Digit Symbol and Letter-Number, D-KEFS Trails (Numbers) and Verbal Fluency (Letters), Rey-Osterrieth Complex Figure, and Wisconsin Card Sorting Test.</p>																								
Waber <i>et al</i> 2011 (+)	WISC, Common entrance examination (local school test) (adolescence)	<table><tr><th></th><th>Controls n=60</th><th>Malnutrition n=56</th><th>p value</th></tr><tr><td colspan="4"><u>WISC:</u></td></tr><tr><td>IQ</td><td>104.92 (SD 10.54)</td><td>92.54 (SD 13.86)</td><td><0.0001</td></tr><tr><td colspan="4"><u>Common Entrance Exam:</u></td></tr><tr><td>Score</td><td>210.42 (SD 22.13)</td><td>184.10 (29.27)</td><td><0.0001</td></tr><tr><td colspan="4"><u>Adjusted analysis:</u></td></tr></table>		Controls n=60	Malnutrition n=56	p value	<u>WISC:</u>				IQ	104.92 (SD 10.54)	92.54 (SD 13.86)	<0.0001	<u>Common Entrance Exam:</u>				Score	210.42 (SD 22.13)	184.10 (29.27)	<0.0001	<u>Adjusted analysis:</u>			
	Controls n=60	Malnutrition n=56	p value																							
<u>WISC:</u>																										
IQ	104.92 (SD 10.54)	92.54 (SD 13.86)	<0.0001																							
<u>Common Entrance Exam:</u>																										
Score	210.42 (SD 22.13)	184.10 (29.27)	<0.0001																							
<u>Adjusted analysis:</u>																										

		<p>Early malnutrition predicted cognitive function and exam score in childhood after adjusting for childhood standard of living.</p> <p>Overall findings: Malnutrition predicted common entrance exam score and cognitive functioning after adjusting for childhood standard of living.</p>
--	--	--

*Studies are drawn from the same Barbados Nutrition Study

95% CI – 95% Confidence Intervals, CANTAB – Cambridge Neuropsychological Test Automated Battery, D-KEFS – Delis-Kaplan Executive Function System, MMSE – Mini-Mental State Exam, OR – Odds Ratio, SD – Standard Deviation, WAIS-III – Wechsler Adult Intelligence Scale-III, WASI – Adult IQ: Weschler Abbreviated Scale of Intelligence – Vocabulary and Matrix Reasoning subsets, WISC – Wechsler Intelligence Scale, WPPSI-IV – Wechsler Preschool and Primary Scale of Intelligence Fourth Edition, WRAML-II – Wide Range Assessment of Memory and Learning-II, WRAT-III – Academic achievement: Wide Range Achievement Test-III – Reading Spelling and Calculation subsets

Supplementary Table 5 - Results from studies assessing mental health outcomes in those exposed to childhood malnutrition compared to controls

Study (quality)	Mental health tool/measurement (outcome age)	Summary of results		
Chen <i>et al</i> 2021 (++)	SDQ (childhood)		Controls n=1270	Wasting n=23
		SDQ:		
		Prosocial score	6.83 (SD 0.06) [95% CI 6.71, 6.95]	6.70 (SD 0.42) [95% CI 5.86, 7.53]
		Total difficulties score	12.35 (SD 0.13)	12.09 (SD 1.09)

		[95% CI 12.10, 12.61]	[95% CI 9.95, 14.22]																																				
		<p><u>Adjusted analysis:</u> Wasting was not statistically associated with any developmental outcome variables after adjusting for vector of child, parent, caregiver and household-level characteristics.</p> <p><u>Overall findings:</u> Wasting was not associated with a difference in behaviour after adjusting for confounding variables.</p>																																					
Mwene-Batu et al 2020	Rosenberg self-esteem scale, WHODAS	<table> <tr> <th></th><th>Controls n=407</th><th>Cases n=524</th><th>p value</th></tr> <tr> <td colspan="4"><u>Self-esteem:</u></td></tr> <tr> <td>Low (%)</td><td>12.1</td><td>20.5</td><td></td></tr> <tr> <td>Average (%)</td><td>78.5</td><td>72.6</td><td></td></tr> <tr> <td>High (%)</td><td>9.4</td><td>6.9</td><td>0.003</td></tr> <tr> <td colspan="4"><u>Social related disability:</u></td></tr> <tr> <td>No disability (%)</td><td>91.0</td><td>94.2</td><td></td></tr> <tr> <td>Moderate disability (%)</td><td>8.8</td><td>5.0</td><td></td></tr> <tr> <td>Severe disability (%)</td><td>0.3</td><td>0.8</td><td>0.03</td></tr> </table>			Controls n=407	Cases n=524	p value	<u>Self-esteem:</u>				Low (%)	12.1	20.5		Average (%)	78.5	72.6		High (%)	9.4	6.9	0.003	<u>Social related disability:</u>				No disability (%)	91.0	94.2		Moderate disability (%)	8.8	5.0		Severe disability (%)	0.3	0.8	0.03
	Controls n=407	Cases n=524	p value																																				
<u>Self-esteem:</u>																																							
Low (%)	12.1	20.5																																					
Average (%)	78.5	72.6																																					
High (%)	9.4	6.9	0.003																																				
<u>Social related disability:</u>																																							
No disability (%)	91.0	94.2																																					
Moderate disability (%)	8.8	5.0																																					
Severe disability (%)	0.3	0.8	0.03																																				
(+)	(adulthood)	<p><u>Adjusted analysis:</u> (unadjusted results)</p> <p><u>Overall findings:</u> Cases had significantly less disability in terms of social relationships than the community controls but had statistically significant lower self-esteem than the controls unadjusted for confounding variables.</p>																																					
Abessa et al (2017)	ASQ: social-emotional score	<p><u>ASQ: Socio-emotional score (incidence rate ratio in cases compared to controls):</u> Socioemotional IRR 3.1 [95% CI 2.24, 4.26]</p>																																					
(++)	(childhood)	<p><u>Adjusted analysis:</u> (results adjusted for significant variables from age, sex, maternal religion)</p> <p><u>Sample size:</u></p>																																					

		Cases n=310, Controls n=310			
		<u>Overall findings:</u> Children with malnutrition performed worse on socio-emotional competences compared to age and sex-matched non-malnourished children adjusted for confounding variables.			
De Grandis et al 2014	Paediatric quality of life inventory		Controls n=28	Malnutrition n=25	p value
(-)	(childhood)	<u>Peds Q:</u>			
		Global score	89.18 (SD 1.84)	80.82 (SD 1.94)	p<0.0001
		Health/physical dimension	94.75 (SD 1.87)	87.75 (SD 3.37)	p<0.0001
		Psychosocial health	86.57 (SD 1.42)	77.77 (SD 2.90)	p<0.0001
		Emotional dimension	78.75 (SD 2.96)	67.80 (SD 4.40)	p<0.0001
		Social dimension	95.71 (SD 1.52)	88.80 (SD 3.05)	p<0.0001
		School dimension	74.58 (SD 3.80)	85.00 (SD 3.51)	p<0.0001
		<u>Adjusted analysis:</u> (unadjusted analysis)			
		<u>Overall findings:</u> Those with a history of early severe malnutrition, showed significantly lower quality of life scores compared to controls unadjusted for confounding variables.			
Baker-Henningham et al 2009	Temperament Questionnaire derived from existing validated instruments		Controls n=108	Undernourished n=212	p value
(++)	(modified by Wachs, Purdue University)	<u>Temperament questionnaire:</u>			
		Manageability	8.67 (SD 3.51)	9.56 (SD 3.26)	p<0.05
		Activity	10.40 (SD 3.00)	10.59 (SD 2.73)	p>0.05
		Emotionality	13.66 (SD 4.74)	14.80 (SD 4.24)	p<0.05
		Sociability	8.15 (SD 3.87)	9.29 (SD 3.98)	p<0.05
		Attention	12.58 (SD 3.31)	13.53 (SD 3.06)	p<0.05
		Soothability	8.23 (SD 2.77)	8.42 (SD 2.57)	p<0.01
	(childhood)	<u>Adjusted analysis:</u>			

		<p>After adjusting for significant covariates (parental schooling and occupation, sanitation index, asset index, housing index, mother's BMI and home), the undernourished children were less sociable (regression coefficient (B) = -0.96 [95% CI -0.04, -1.88]), less attentive (B = -0.94 [95% CI -0.19, -1.69]), more fearful (B = 1.43 [95% CI 2.44, 0.42]), and had more negative emotionality (B = -1.08 [95% CI -2.16, 0.006]).</p> <p><u>Overall findings:</u> There were significant differences in temperament traits between undernourished and better-nourished groups after adjusting for confounding variables.</p>
Liu <i>et al</i> (2004)	<p>Child behaviour questionnaire at age 8, Child Behaviour Checklist at age 11, Revised Behaviour Checklist at Age 17</p> <p>(childhood, adolescence)</p>	<p><u>Externalizing behaviour problems at age 8:</u> Malnourished children had higher overall externalizing behaviour scores (both hyperactivity and aggression) than controls.</p> <p><u>Adjusted analysis:</u> Main effect of malnutrition remained significant after controlling for psychosocial adversity but was abolished after controlling for cognitive ability.</p> <p><u>Externalizing behaviour problems at age 11:</u> Malnourished children had higher externalising behaviour scores (hyperactivity only) at age 11 than non-malnourished children.</p> <p><u>Adjusted analysis:</u> Main effect of malnutrition remained significant after controlling for psychosocial adversity but was abolished after controlling for cognitive ability.</p> <p><u>Externalizing behaviour problems at age 17:</u> Malnourished group had significantly higher scores on conduct disorder and motor excess but not attention problems or socialized aggression.</p> <p><u>Adjusted analysis:</u> Main effect of malnutrition remained significant after adjusting for psychosocial adversity and cognitive ability.</p>

		<p><u>Sample size:</u> Malnutrition group n=235, Control group n=807</p> <p><u>Overall Findings:</u> Children with malnutrition were more aggressive or hyperactive at age 8 years, had more externalizing problems at age 11, and had greater conduct disorder and excessive motor activity at age 17 after adjusting for confounding variables.</p>
Barbados Nutrition Study		
<p>Hock et al 2018*</p> <p>(+)</p>	<p>SCID-II-PQ, NEO PI-R FFM</p> <p>(adulthood)</p>	<p><u>SCID-II-PQ (linear regression with malnutrition as predictor of score):</u> Paranoid personality score 0.6 [95% CI 0.0, 1.2] (p=0.0499)</p> <p><u>Adjusted analysis:</u> PD paranoid scale score had unique association with malnutrition after adjusting for maltreatment and childhood standard of living</p> <p><u>NEO PI-R FFM (linear regression with malnutrition as predictor of score):</u> Avoidant personality score 1.0 [95% CI 0.2, 1.7] (p=0.009), schizoid personality score 0.7 [95% CI 0.1, 1.4] (p=0.03) and dependant personality score 0.7 [95% CI 0.1, 1.4] (p=0.04)</p> <p><u>Adjusted analysis:</u> PD avoidant, schizoid and dependant scale scores had unique associations with malnutrition after adjusting for maltreatment and childhood standard of living</p> <p><u>Sample size:</u> Malnutrition group n=77, control group n=62</p> <p><u>Overall findings:</u> Adults with history of infant malnutrition showed an increase in most SCID-II and NEO FFM PD personality scores compared to those with little or no exposure to either adversity. Few personality disorder score differences were statistically significant however and results differed depending on the assessment tool used.</p>

Waber (2) et al 2014*	Behavioural Regulation Index score	Controls n=59	Malnutrition n=77
(++)	(adulthood)	<u>Behavioural regulation index:</u>	
		Mean score	43.34 (SD 7.52) 49.51 (SD 11.21)
		Mean difference (adjusted) 3.40 (p<0.05)	
		<u>Adjusted analysis</u> (results adjusted for childhood standard of living and IQ)	
		<u>Overall findings:</u> Adults with a history of infant malnutrition showed a significantly higher behavioural regulation index score which is indicative of more behavioural problems, with a significant difference in score after adjusting for confounding variables.	
Galler et al 2013*	NEO-PI-R personality inventory	<u>NEO-PI-R score:</u>	
		At the broad domain or factor level, previously malnourished participants had higher scores on Neuroticism and lower scores on Extraversion, Openness, Agreeableness and Conscientiousness than healthy controls. At the sub-domain or facet level, previously malnourished participants reported more anxiety, vulnerability, shyness and lowered sociability, less intellectual curiosity, greater suspiciousness of others, a more egocentric than altruistic orientation, and a lowered sense of efficacy or competence.	
(+)	(adulthood)	<u>Adjusted analysis:</u> (results adjusted for childhood standard of living)	
		<u>Sample size</u> Malnutrition group n=77, Control group n=57	
		<u>Overall findings:</u> Childhood malnutrition was associated with higher chance of personality trait scores outside of the average range after adjusting for childhood standard of living.	

Galler (1) et al 2012* (+)	CAARS, CPT (adulthood)	<hr/>			
		Controls n=63		Malnutrition n=80	p value
		<u>CAARS (mean score):</u>			
		Inattentive symptoms	3.6 (SD 0.1)	3.7 (SD 0.2)	<0.05
		Hyperactive symptoms	3.7 (SD 0.1)	3.7 (SD 0.2)	>0.05
		DSM-IV ADHD symptoms	3.6 (SD 0.1)	3.7 (SD 0.1)	<0.01
		ADHD index	3.8 (SD 0.2)	3.7 (SD 0.1)	>0.05
		<hr/>			
		Controls n=59		Malnutrition n=77	p value
		<u>CPT (mean score):</u>			
		Commission errors	3.8 (SD 0.2)	3.9 (SD 0.2)	>0.05
		Omission errors	3.9 (SD 0.2)	4.1 (SD 0.3)	>0.05
		Reaction time	4.0 (SD 0.2)	4.0 (SD 0.2)	>0.05
<u>Adjusted analysis:</u> (results adjusted for childhood standard of living and IQ)					
<u>Overall findings:</u> CAARS scores were higher (indicating more attention problems) in the previously malnourished group. There were significant differences in inattention and DSM-IV ADHD symptom scores after adjusting for confounding variables. Differences in CPT scores were attenuated after adjusting for confounding variables.					
Galler (2) et al 2012* (+)	Conduct-problems: Child-Behaviour-Questionnaire, Teacher-Behaviour-Questionnaire (adolescence)	Controls n=60		Malnutrition n=56	p value
		<u>Child-Behaviour-Questionnaire (coefficient):</u>			
		Conduct problems	-0.27 (SD 0.84)	0.29 (SD 1.18)	<0.01
		<u>Teacher-behaviour-questionnaire:</u>			
		All four items showed a higher level prevalence of aberrant behavior in the previously malnourished children as compared with the comparison group.			
		<u>Adjusted analysis:</u>			
		The difference was no-longer-significant with IQ and childhood standard of living included in the model			

		Overall findings: Self-reported conduct problems were more prevalent among previously malnourished youth. The associated was no longer significant when adjusted for confounding variables.																																											
Waber <i>et al</i> 2011* (+)	Teacher-Behaviour Questionnaire, School functioning scale Minnesota-General-Adjustment-and Morale-Scale (adolescence)	<table><tr><td></td><td>Controls n=60</td><td>Malnutrition n=56</td><td>p value</td></tr><tr><td colspan="4">Teacher questionnaire:</td></tr><tr><td>Attention symptoms</td><td>-0.36 (SD 0.81)</td><td>0.33 (SD 0.96)</td><td><0.0001</td></tr><tr><td colspan="4">Morale scale:</td></tr><tr><td>Depressive symptoms</td><td>-0.33 (SD 0.93)</td><td>0.30 (SD 1.01)</td><td>0.001</td></tr></table>		Controls n=60	Malnutrition n=56	p value	Teacher questionnaire:				Attention symptoms	-0.36 (SD 0.81)	0.33 (SD 0.96)	<0.0001	Morale scale:				Depressive symptoms	-0.33 (SD 0.93)	0.30 (SD 1.01)	0.001																							
	Controls n=60	Malnutrition n=56	p value																																										
Teacher questionnaire:																																													
Attention symptoms	-0.36 (SD 0.81)	0.33 (SD 0.96)	<0.0001																																										
Morale scale:																																													
Depressive symptoms	-0.33 (SD 0.93)	0.30 (SD 1.01)	0.001																																										
		Adjusted analysis: Association attenuated when IQ and attention problems included in model. Difference in attention symptoms not significant when adjusting childhood standard of living.																																											
		Overall findings: Depressive symptoms and inattention were higher in those with malnutrition but there were no significant differences when adjusting for confounding variables.																																											
Galler <i>et al</i> 2011* (++)	Barbados-Child-Behaviour-Scale adolescence	<table><tr><td></td><td>Controls n=109</td><td>Malnutrition n=107</td></tr><tr><td colspan="3">(correlations with mean score)</td></tr><tr><td colspan="3">Peer aggression:</td></tr><tr><td>Time point 1 (9-15y)</td><td>-0.05 (SD 0.96)</td><td>0.39 (SD 1.16)</td></tr><tr><td>Time point 2 (11-17y)</td><td>-0.34 (SD 0.79)</td><td>-0.34 (SD 0.80)</td></tr><tr><td colspan="3">Cooperative with adults:</td></tr><tr><td>Time point 1 (9-15y)</td><td>0.19 (SD 1.11)</td><td>-0.01 (SD 1.15)</td></tr><tr><td>Time point 2 (11-17y)</td><td>0.15 (SD 0.73)</td><td>-0.28 (SD 1.00)</td></tr><tr><td colspan="3">Activity level:</td></tr><tr><td>Time point 1 (9-15y)</td><td>0.32 (SD 1.09)</td><td>0.18 (SD 1.03)</td></tr><tr><td>Time point 2 (11-17y)</td><td>-0.35 (SD 0.72)</td><td>-0.29 (SD 0.78)</td></tr><tr><td colspan="3">Executive function deficits:</td></tr><tr><td>Time point 1 (9-15y)</td><td>0.0 (SD 0.94)</td><td>0.24 (SD 1.00)</td></tr><tr><td>Time point 2 (11-17y)</td><td>-0.38 (SD 0.82)</td><td>-0.09 (SD 0.94)</td></tr></table>		Controls n=109	Malnutrition n=107	(correlations with mean score)			Peer aggression:			Time point 1 (9-15y)	-0.05 (SD 0.96)	0.39 (SD 1.16)	Time point 2 (11-17y)	-0.34 (SD 0.79)	-0.34 (SD 0.80)	Cooperative with adults:			Time point 1 (9-15y)	0.19 (SD 1.11)	-0.01 (SD 1.15)	Time point 2 (11-17y)	0.15 (SD 0.73)	-0.28 (SD 1.00)	Activity level:			Time point 1 (9-15y)	0.32 (SD 1.09)	0.18 (SD 1.03)	Time point 2 (11-17y)	-0.35 (SD 0.72)	-0.29 (SD 0.78)	Executive function deficits:			Time point 1 (9-15y)	0.0 (SD 0.94)	0.24 (SD 1.00)	Time point 2 (11-17y)	-0.38 (SD 0.82)	-0.09 (SD 0.94)	
	Controls n=109	Malnutrition n=107																																											
(correlations with mean score)																																													
Peer aggression:																																													
Time point 1 (9-15y)	-0.05 (SD 0.96)	0.39 (SD 1.16)																																											
Time point 2 (11-17y)	-0.34 (SD 0.79)	-0.34 (SD 0.80)																																											
Cooperative with adults:																																													
Time point 1 (9-15y)	0.19 (SD 1.11)	-0.01 (SD 1.15)																																											
Time point 2 (11-17y)	0.15 (SD 0.73)	-0.28 (SD 1.00)																																											
Activity level:																																													
Time point 1 (9-15y)	0.32 (SD 1.09)	0.18 (SD 1.03)																																											
Time point 2 (11-17y)	-0.35 (SD 0.72)	-0.29 (SD 0.78)																																											
Executive function deficits:																																													
Time point 1 (9-15y)	0.0 (SD 0.94)	0.24 (SD 1.00)																																											
Time point 2 (11-17y)	-0.38 (SD 0.82)	-0.09 (SD 0.94)																																											

		<p><u>Sleep problems:</u></p> <p>Time point 1 (9-15y) -0.45 (SD 1.17) -0.24 (SD 1.08)</p> <p>Time point 2 (11-17y) 0.21 (SD 0.61) 0.43 (SD 0.68)</p> <p><u>Adjusted analysis:</u></p> <p>Cases had increased prevalence of executive function deficits at both time points after adjusting for age, sex, childhood standard of living and maternal depressive symptoms. No significant association with aggression, cooperation, activity and sleep after adjusting for covariates at second time point. Cases had more problem aggression at the first time point than at the second.</p> <p><u>Overall findings:</u></p> <p>Persisting parent-reported executive functioning deficits through adolescence at two time points after adjusting for confounding variables.</p>		
<p>Galler <i>et al</i> 2010*</p> <p>(++)</p>	<p>Minnesota-General-Adjustment-and Morale-Scale</p> <p>(adolescence)</p>	<p>Controls n=60</p> <p>Cases n=56</p> <p>p value</p> <p><u>Morale Scale:</u></p> <p>Youth depression score -0.33 (SD 0.98) 0.30 (SD 1.01) <0.001</p> <p><u>Adjusted analysis:</u></p> <p>Adjusting for age, maternal depression symptoms and household standard of living the effect of malnutrition is similar to unadjusted estimates and remains significant.</p> <p><u>Overall findings:</u></p> <p>Early childhood malnutrition contributed independently to depressive symptoms after adjusting for confounding variables.</p>		

*Studies are drawn from the same Barbados Nutrition Study

95% CI – 95% Confidence Intervals, CAARS - Attention-domain: Conners-ADHD-Rating-Scales, CPT – Conners-Continuous-Performance Test, NEO PI-R FFM – NEO Personality Inventory-Revised derived Five-Factor Model, SD – Standard Deviation, SDQ – Strengths and Difficulties Questionnaire, SCID-II-PQ – Structured Clinical Interview for DSM-IV Axis II Personality Disorders Personality Questionnaire, WHODAS – World Health Organization Disability Assessment Schedule