

1 **Title**

2 **Effects of multiple-dose intranasal oxytocin treatment on social responsiveness in**
3 **children with autism: A randomized, placebo-controlled trial**

4 **Authors and affiliations**

5 *Daniels Nicky*^{a,c,*}, *Moerkerke Matthijs*^{b,c,*}, *Steyaert Jean*^c, *Bamps Annelies*^c, *Debbaut Edward*
6 *b,c*, *Prinsen Jellina*^{a,c}, *Tang Tiffany*^{b,c}, *Van der Donck Stephanie*^{b,c}, *Boets Bart*^{b,c,†}, *Alaerts Kaat*
7 *a,c,†,**

8 *Joined first-authorship, † senior authors

9 ^a KU Leuven, Department of Rehabilitation Sciences, Research Group for
10 Neurorehabilitation, Tervuursevest 101 box 1501, 3001 Leuven, Belgium.

11 ^b KU Leuven, Department of Neurosciences, Center for Developmental Psychiatry, O&N5b
12 Herestraat 49 box 7003, 3000 Leuven

13 ^c Leuven Autism Research consortium, KU Leuven, Leuven, Belgium

14

15 ✖ *Corresponding author.*

16 Kaat Alaerts

17 KU Leuven, Department of Rehabilitation Sciences, Research Group for Neurorehabilitation,
18 Tervuursevest 101 box 1501, 3001 Leuven, Belgium.

19 E-mail: kaat.alaerts@kuleuven.be

20 Tel.: +32 16 37 64 46, Fax: +32 16 32 91 97

21 **Abstract**

22 In the past decade, intranasal administration of the neuropeptide oxytocin is increasingly
23 explored as a new treatment for reducing the core symptoms of autism spectrum disorder
24 (ASD). The efficacy of continual oxytocin treatment in school-aged children with ASD is,
25 however, not well established. Using a double-blind, randomized, placebo-controlled, parallel
26 design, the current trial explored the effects of four weeks of intranasal oxytocin treatment
27 (12 IU, twice daily) on social functioning in pre-pubertal school-aged children (aged 8-12
28 years, 61 boys, 16 girls). The double-blind phase was followed by a four-week single-blind
29 extension phase during which all participants received intranasal oxytocin. In the double-
30 blind phase, no treatment-specific effects were identified in the primary outcome assessing
31 social functioning (parent-rated Social Responsiveness Scale), as well as on secondary
32 outcomes assessing repetitive behaviors, anxiety, and attachment. Exploratory moderator
33 analyses revealed that children who received the oxytocin treatment in combination with
34 concomitant psychosocial treatment displayed a greater benefit than those who received
35 psychosocial treatment or oxytocin alone. A modulating effect of parents' beliefs about
36 allocated treatment was also identified, indicating that parents who believed their child
37 assigned to the active treatment reported greater benefit than those who believed their child
38 received placebo, particularly in the actual oxytocin group. Finally, participants who were
39 allocated to receive the placebo treatment during the double-blind phase of the trial and later
40 crossed-over to receive the active treatment during the single-blind extension phase,
41 displayed a significant within-group improvement in social responsiveness, over and above
42 the placebo-induced improvements noted in the first phase. While no overall treatment-
43 specific improvements were identified, our results provide important indications that clinical
44 efficacy can be augmented when oxytocin administration is paired with targeted psychosocial
45 interventions that similarly stimulate socio-communicative behaviors. Future trials are urged
46 to further elucidate the potential of embedding oxytocin treatment within a socially stimulating
47 context.

48 Introduction

49 Autism spectrum disorder (ASD) is a neurodevelopmental condition characterized by
50 impairments in social communication and interaction, combined with restricted and repetitive
51 behaviors and interests (American Psychiatric Association, 2013). Thus far, the means of
52 treatment of ASD's core symptoms are primarily based on behavioral interventions (e.g.,
53 stimulation of social communication, lessening the impairment due to restricted and repetitive
54 behaviors), since biomedical or pharmacological therapies targeting social impairment or
55 repetitive behaviors are largely unproven.

56 In the past decade, intranasal administration of the neuropeptide oxytocin (OT) has been
57 increasingly explored as a new treatment option for reducing ASD's social symptoms
58 (recently summarized in Huang et al., 2021). OT is an endogenous neuropeptide that is
59 mainly produced in paraventricular nuclei of the hypothalamus. In the brain, OT acts as an
60 important neuromodulator for a broad range of affiliative and prosocial behaviors, including
61 interpersonal bonding, social attunement and attachment (Bakermans-Kranenburg & van
62 IJzendoorn, 2013; Bartz et al., 2011; Jurek & Neumann, 2018), presumably mediated
63 through its postulated top-down enhancing effect on 'social salience' and bottom-up effect on
64 regulating (social) stress and anxiety (Guastella & Hickie, 2016; Shamay-Tsoory & Abu-Akel,
65 2016).

66 Following a myriad of single-dose proof-of-principle studies (Alvares et al., 2017; Huang et
67 al., 2021), an initial multiple-dose pilot study assessed the safety and efficacy of six weeks of
68 chronic intranasal OT treatment on core autism symptoms in 19 adults with ASD (10
69 receiving OT, 9 receiving placebo), and showed improved emotion recognition and quality of
70 life, and tentative improvements in repetitive behaviors after OT treatment (Anagnostou et
71 al., 2012). Later, Kosaka et al. (2016) showed significant improvements on the Clinical
72 Global Impression-Improvement scale after twelve weeks of OT treatment in adult men with
73 ASD, albeit only in the subgroup of participants receiving the high-dose treatment (32 IU/day;
74 $n = 13$), and not in the low-dose (16 IU; $n = 15$) or placebo groups ($n = 16$). In an exploratory
75 cross-over study by Watanabe et al. (2015), the effects of six weeks of daily intranasal OT
76 administration on core autism characteristics were studied in 20 adult men with ASD, and
77 significant improvements in social reciprocity and social functioning (social-judgement task)
78 were identified. Yamasue et al. (2020) conducted a confirmatory trial with an identical
79 protocol as Watanabe et al. (2015) in 106 adult men with ASD (53 OT / 53 placebo). While
80 these authors identified significant improvements in terms of repetitive behaviors, the effects
81 on social reciprocity and social functioning could not be replicated. Bernaerts et al. (2020)
82 extended these observations in an exploratory sample of 40 young adult men with ASD (22

83 OT / 18 placebo), demonstrating long-term improvements in repetitive behaviors and feelings
84 of attachment after a four-week course which outlasted the period of administration till one
85 year post-treatment.

86 Given that ASD is an early-onset neurodevelopmental condition, it is important to extend
87 these insights to pediatric populations, allowing evaluations of OT treatment efficacy within
88 an early developmental window and whether it can be facilitatory for enriching social
89 behaviors and experiences from an early age onwards. To date, a handful of trials explored
90 the effects of multiple-dose OT administration in children with ASD. Two initial trials reported
91 a consistent pattern of results, indicating improvements in the social domain (parent-reported
92 social responsiveness) after five weeks of intranasal OT treatment in 3-to-6-year-old children
93 ($n = 31$; cross-over; Yatawara et al., 2016) and after four weeks of treatment in 6-to-12-year-
94 old children with ASD (14 OT / 18 placebo; Parker et al., 2017). No significant improvements
95 on core autism symptoms were demonstrated, however, after an eight-week OT treatment in
96 adolescent boys with ASD (26 OT / 24 placebo; 12-18 years; Guastella et al., 2015) or in a
97 preliminary 12-week administration trial encompassing a broad age range of 5-to-17-year-old
98 children (8 OT / 10 placebo) with Phelan-McDermid syndrome (characterized by ASD
99 symptoms; Fastman et al., 2021). Also, in a recent confirmatory trial including 3-to-17-year-
100 old children with ASD (139 OT / 138 placebo) and an age-adjusted dosing scheme ranging
101 from 8-80 IU, no improvements on outcomes of social functioning were evident after 24
102 weeks of OT treatment (Sikich et al., 2021).

103 Several factors have been put forward to understand these inconsistent results, ranging from
104 heterogeneity in trial design (e.g., parallel versus cross-over design, adopted outcomes,
105 dosing schema) to variation in participant characteristics. For instance, the well-powered
106 confirmatory trial by Sikich et al. (2021) covered a broad age range (3-17 years),
107 encompassing a critical period of pubertal development, which could have rendered
108 heterogeneity due to differential physiologic effects of OT during different developmental
109 stages (Geschwind, 2021).

110 Here, results are presented from a single-center, randomized, double-blind, placebo-
111 controlled clinical trial (RCT with parallel design), testing efficacy on improvement in social
112 functioning and safety of multiple-dose OT treatment (four weeks of twice daily intranasal
113 administration of 12 IU) in a representative sample of 8-to-12-year-old children with ASD (40
114 OT / 40 placebo). Accordingly, this is the largest trial to date, examining OT treatment effects
115 in a relatively strict age range of pre-pubertal, school-aged children, aged 8 to 12 years,
116 thereby allowing to overcome some of the raised issues regarding sample heterogeneity.
117 Further, following prior observations of long-lasting retention effects of OT treatment in adults

118 with ASD (Bernaerts et al., 2020), the current trial also included a follow-up session four
119 weeks after cessation of the daily OT administrations, testing the possibility of crucial
120 retention effects in the current pediatric sample.

121 **Methods**

122 **2.1. General study design**

123 A single-center, two-arm, double-blind, randomized, placebo-controlled parallel study was
124 performed at the Leuven University Hospital (Leuven, Belgium) to assess effects of four
125 weeks of twice daily intranasal administration of OT on core autism characteristics, social
126 anxiety, and experience of attachment in school-aged children with ASD. The double-blind
127 phase (phase I) was followed by a four-week single-blind extension phase (phase II) during
128 which all participants received intranasal OT. In both phases, treatment effects were
129 assessed immediately after the four-week treatment (post) and at a follow-up session, four
130 weeks after cessation of the daily administrations (follow-up). See [Figure 1](#), CONSORT Flow
131 diagram for number of participants randomized and analyzed.

132 Written informed consent from the parents and assent from the child were obtained prior to
133 the study. Consent forms and study design were approved by the local Ethics Committee for
134 Biomedical Research at the University of Leuven, KU Leuven (S61358) in accordance with
135 The Code of Ethics of the World Medical Association (Declaration of Helsinki). The trial was
136 registered at the European Clinical Trial Registry (Eudract 2018-000769-35) and the Belgian
137 Federal Agency for Medicines and Health products.

138 **2.2. Participants**

139 Children with a formal diagnosis of ASD were recruited through the Autism Expertise Centre
140 at the Leuven University Hospital between July 2019 and January 2021. The diagnosis was
141 established by a multidisciplinary neuropsychiatric team based on the strict criteria of the DSM-
142 5 (Diagnostic and Statistical Manual of Mental Disorders; American Psychiatric Association,
143 2013). Prior to randomization, the Autism Diagnostic Observation Schedule (ADOS-2; Lord
144 et al., 2012) and estimates of intelligence (four subtests of the Wechsler Intelligence Scale
145 for Children, Fifth Edition, Dutch version; Wechsler, 2018) were acquired ([Table 1](#)). Principal
146 inclusion criteria comprised a clinical diagnosis of ASD, age (8-12 years old), intelligence
147 quotient (IQ) above 70, native Dutch speaker, a stable background treatment for at least four
148 weeks prior to the screening and no anticipated changes during the trial. Only premenstrual
149 girls were included. Principal criteria for exclusion comprised any neurological (e.g., stroke,
150 epilepsy, concussion) or significant physical disorder (liver, renal, cardiac pathology) or prior

151 treatment with OT (**Supplementary Table 1**). The presence of comorbid psychiatric
152 disorders, current psychoactive medication use, and concomitant participation in
153 psychosocial therapies were screened for and logged (see **Table 1 and Supplementary**
154 **Table 2**).

155 A sample size of 40 participants in each treatment group was determined to be able to detect
156 a medium effect size ($d = 0.60$) with $\alpha = 0.05$ and 80% power, corresponding to effect sizes
157 previously reported in the four-week oxytocin trial with school-aged children by Parker et al.,
158 2017.

159 **2.3. Intervention**

160 **Study medication.** Participants were randomized to receive OT (Syntocinon®, Sigma-tau)
161 or placebo nasal sprays, administered in identical blinded amber 10 ml glass bottles with
162 metered pump. The placebo spray consisted of all the ingredients used in the active solution
163 except the OT compound. Nasal spray preparation, packaging, blinding and randomization
164 (permuted-block randomization, RITA software; Pahlke et al., 2004) was performed by the
165 pharmacy of Heidelberg University Hospital (Germany). Participants were randomly assigned
166 in a 1:1 ratio, with stratification according to gender. During the initial double-blind phase
167 (phase I), all research staff conducting the trial, participants and their parents were blinded to
168 treatment allocation. During the subsequent single-blind extension phase (phase II),
169 experimenters were aware that all participants received intranasal OT, but participants and
170 parents were still fully blinded regarding treatment allocation.

171 **Dosing.** Children (assisted by their parents) were asked to self-administer a daily dose of 2 x
172 12 IU nasal spray or placebo equivalent (3 puffs of 2 IU in each nostril), 12 IU in the morning
173 and 12 IU in the afternoon, during 28 consecutive days during the initial double-blind phase
174 (phase I), and for another 28 days during the single-blind extension phase (phase II).
175 Participants received clear instructions about use of the nasal sprays (based on Guastella et
176 al., 2013) through a demonstration together with the experimenter.

177 **Compliance monitoring.** Compliance was assured using a daily medication diary that
178 recorded date and time of administration (phase I percentage compliance; OT: $96.75 \pm$
179 5.26% ; placebo: $96.11 \pm 5.29\%$; $t(74) = .52$, $p = .603$; phase II percentage compliance; OT-
180 first: $94.55 \pm 11.69\%$; placebo-first: $92.98 \pm 13.92\%$; $t(74) = .53$, $p = .597$). The total amount
181 of administered fluid was also monitored (phase I: OT: 14.86 ± 2.37 ml; PL: 13.79 ± 2.35 ml;
182 $t(75) = 2.00$, $p = .050$; phase II: OT-first: 13.72 ± 3.47 ml; placebo-first: 12.83 ± 3.52 ml; $t(74)$
183 $= 1.10$, $p = .275$).

184 **Side effects.** During the course of the treatment, participants were screened for potential
185 adverse events (weekly parent report) or changes in affect and arousal (daily diary by child
186 and parent). Overall, reports of side effects were minimal and not treatment-specific (see
187 [Supplementary Tables 3 and 4](#)).

188 **Parent reported treatment beliefs.** At the end of each trial phase (I and II), parents reported
189 beliefs about treatment allocation (see **Results** and [Figure 3](#)). In the double-blind phase
190 (phase I), the proportion of parents that believed their child had received the OT treatment
191 was similar in both treatment arms: 39.5% in the OT group, 35.9% in the placebo group ($p =$
192 $.75$). In the OT group 18.4% of parents indicated to 'have no explicit belief' about treatment
193 allocation versus 10.3% in the placebo group. In the single blind phase (phase II), during
194 which all participants received the actual OT treatment, 51.9% of the parents believed their
195 child received the OT treatment, 35.1% believed their child received the placebo treatment
196 and 13.0% indicated to 'have no explicit belief'.

197 **2.4. Outcome Measures**

198 The primary outcome measure was change from baseline in parent-rated social
199 responsiveness on the Social Responsiveness Scale-Children, second edition (SRS-2;
200 Constantino & Gruber, 2012; Roeyers et al., 2015). which comprises four subscales
201 examining social communication, social awareness, social motivation, and
202 rigidity/repetitiveness, using a four-point Likert-scale (65 items). Higher scores indicate
203 greater deficit.

204 Secondary outcome measures included changes from baseline in parent-rated repetitive
205 behaviors (Repetitive Behavior Scale-Revised (RBS-R; Bodfish et al., 2000), self and parent-
206 rated presence of anxiety symptoms (Screen for Child Anxiety Related Emotional Disorders
207 (SCARED-NL: Muris et al., 2007), and changes from baseline in constructs of self-rated
208 attachment towards their mother (Attachment Questionnaire child-report; Bosmans et al.,
209 2014) and peers (Attachment Style Classification Questionnaire child-report; Finzi et al.,
210 2000) (see also [Supplementary Table 5](#)).

211 All outcomes were assessed five times: (i) at baseline, (ii) immediately after the four-week
212 double-blind treatment (phase I - post); (iii) at a follow-up session, four weeks after cessation
213 of the double-blind treatment (phase I - follow-up); (iv) immediately after the four-week
214 single-blind treatment (phase II - post); and (v) at a follow-up session four weeks after
215 cessation of the single-blind treatment (phase II - follow-up). Post sessions were scheduled
216 approximately 24h after the last administration, follow-up sessions within 28 ± 7 days.

217 2.5. Data Analysis

218 Analyses were performed using a modified intention-to-treat approach that included all
219 randomized participants who completed the baseline session and at least one post or follow-
220 up session (see **Figure 1**, **CONSORT flow-chart**). All statistics were executed with Statistica
221 14 (Tibco Software Inc.).

222 First, as outlined in **Table 1**, possible baseline differences between treatment groups on the
223 primary or secondary outcomes were assessed using independent sample t-tests, but no
224 statistically significant differences were identified. Next, between-group differences in
225 treatment responses of *phase I (double-blind)* on the primary and secondary outcome
226 measures were assessed, by subjecting change from baseline scores of the post and follow-
227 up session to independent sample t-tests. Additionally, single-sample *t*-tests were adopted to
228 assess within-group changes (compared to baseline) in the OT and placebo group
229 separately.

230 Subsequent exploratory analyses of the primary outcome were performed to investigate the
231 potential influence of possible moderator variables on phase I treatment outcome. To do so,
232 change from baseline scores of the primary outcome (SRS-2) were subjected to general
233 linear models with the within-subject factor ‘assessment session’ (post, follow-up) and the
234 between-subject factors ‘treatment’ (OT, placebo) and specific moderator variables. Separate
235 models were constructed to assess the modulating effect of concomitant psychosocial
236 treatments (present, not present); medication use (present, not present; as listed in **Table 1**);
237 and parent reported beliefs (OT, placebo).

238 To evaluate the effect of phase, within-subject changes from *phase I (double-blind)* to *phase*
239 *II (single-blind extension)* were assessed, across groups and separately within the OT-first
240 and placebo-first groups. To do so, change from baseline scores of the primary outcome
241 (SRS-2) were subjected to general linear models with the within-subject factors ‘assessment
242 session’ (post, follow-up) and ‘phase’ (phase I, phase II).

243 Finally, to assess whether the magnitude of the observed change from baseline scores at the
244 last session of the trial were reliable for individual participants (more than can be expected by
245 measurement error), the Reliable Change Index (RCI) as proposed by Jacobson et al. (1999)
246 was calculated, based on the test-retest reliability of the adopted Dutch parent-reported SRS
247 scale (Cronbach’s alpha = .94; Roeyers et al., 2015). Change scores higher than the RCI-
248 value (14.8) were considered reliable.

249 Results

250 Double-blind phase (phase I)

251 Between-group analyses revealed no significant effect of treatment on the primary outcome
252 parent-reported social responsiveness (SRS-2), either immediately post-treatment ($p = .839$),
253 or at the follow-up session, four weeks after cessation of the daily nasal spray
254 administrations ($p = .626$) (see [Table 2](#) and [Supplementary Table 6](#) for the raw scores).
255 Both groups displayed similar significant pre-to-post-treatment improvements in social
256 responsiveness (reduced SRS-2 scores) immediately after treatment (OT: $p = .017$; placebo:
257 $p = .009$) and at the follow-up session (OT: $p = .001$; placebo: $p = .017$). A similar pattern of
258 non-treatment-specific improvements was evident for the secondary outcomes ([Table 2](#)).

259 Interestingly, exploratory moderator analyses showed a significant interaction between
260 treatment and the presence of **concomitant psychosocial treatment** ($F(1,72) = 6.87$; $p =$
261 $.011$; [Figure 2](#)), indicating that, across assessment sessions (post, follow-up), participants
262 who received the OT treatment combined with psychosocial treatment displayed greater
263 benefits compared to children receiving only psychosocial treatment (combined with placebo)
264 ($t(26) = 2.40$; $p = .012$, one-tailed) or only the OT treatment ($t(36) = 1.90$; $p = .033$, one-
265 tailed). In children without psychosocial treatment, OT treatment responses were not
266 significantly different from those in the placebo group ($t(46) = 1.40$; $p = .084$, one-tailed).
267 None of the other main or interaction effects (e.g., with assessment session) were significant
268 (all, $p > .05$).

269 In terms of modulating effects of **parent reported beliefs**, a trend-level interaction with
270 'assessment session' was evident ($F(1,61) = 3.22$; $p = .077$), indicating that the parents' own
271 belief about allocated treatment differentially modulated treatment responses at the post and
272 follow-up assessment session ([Figure 3](#)). Direct exploration of this effect, separately for each
273 treatment group, showed that for participants receiving the actual OT treatment, parents' own
274 belief moderated treatment immediately post-treatment ($t(29) = -3.18$; $p = .001$, one-tailed),
275 but no longer at the follow-up session ($t(29) = -1.02$; $p = .158$, one-tailed). Specifically,
276 parents who believed their child had received OT, reported significantly greater
277 improvements in social responsiveness immediately post-treatment, compared to those who
278 believed their child had received placebo. In the placebo group, no significant modulations of
279 treatment responses were evident either immediately post-treatment ($t(32) = -.32$; $p = .374$,
280 one-tailed) or at the follow-up session ($t(32) = .37$; $p = .358$, one-tailed). None of the other
281 main or interaction effects were significant (all, $p > .05$). Further, for **concomitant**
282 **medication use**, no significant modulating effects were evident (all, $p > .148$).

283 **Single-blind extension phase (phase II)**

284 Examination of within-subject changes from phase I to phase II yielded a significant effect of
285 'phase' ($F(1,70) = 12.94$; $p < .001$), but no 'phase x treatment interaction' ($F(1,70) = 2.24$; $p =$
286 $.14$), indicating a further improvement in social responsiveness across treatment groups from
287 phase I to phase II, (**Figure 4** and **Supplementary Table 6** for the raw scores). The main
288 effects of treatment and assessment session were not significant ($p > .05$).

289 When examined separately for each treatment group, the effect of phase was particularly
290 strong in the placebo-first group ($F(1,35) = 14.54$; $p < .001$), indicating that for children who
291 crossed over from placebo (in phase I) to OT treatment (in phase II), improvements in social
292 responsiveness were significantly more pronounced in phase II, during which the child
293 received the actual OT treatment (**Figure 4, right panel**).

294 Within the OT-first group, only non-significant within-subject changes from phase I to phase II
295 were noted ($F(1,35) = 1.99$; $p = .17$), indicating that OT treatment effects of phase I were not
296 significantly augmented by receiving the additional four-week OT treatment of phase II.
297 Analysis of the OT-first group did reveal a significant effect of 'session', indicating that
298 irrespective of phase, treatment-related improvements were more pronounced at the follow-
299 up session, compared to the post session ($F(1,35) = 6.11$; $p = .018$), with maximal treatment
300 responses at the last assessment session of the trial (four-week follow-up of phase II; **Figure**
301 **4, left panel**).

302 Accordingly, at the last session of the trial, both the OT-first (receiving a total of eight weeks
303 of OT treatment) and the placebo-first group (receiving a total of four weeks of OT treatment)
304 displayed significant pre-to-post improvements in social responsiveness (OT-first; pre-post
305 change: -9.61 ± 12.18 ; $t(35) = -4.74$; $p < .001$; (placebo-first; pre-post change: -9.81 ± 14.83 ;
306 $t(35) = -3.97$; $p < .001$). In the OT-first group, 27 (out of 36: 75%) participants displayed a pre-
307 to-post improvement, and this change was identified to be reliable for 12 participants (higher
308 than the Reliable Change Index: >14.8). Similarly, also in the PL-first group, 27 (out of 36:
309 75%) participants displayed a pre-to-post improvement at the last session of the trial, which
310 was reliable for 11 participants.

311 Discussion

312 The current pediatric trial with pre-pubertal school-aged children with ASD demonstrated no
313 significant treatment-specific effects of four weeks of intranasal OT administration on the
314 primary outcome, assessing parent-rated social responsiveness (SRS-2), nor on the
315 secondary outcomes assessing parent and child self-reports of repetitive behaviors, anxiety,
316 and attachment. Both the OT and the placebo group displayed similar improvements, both
317 immediately after the multiple-dose treatment and at the four-week follow-up session.
318 Notably, exploratory analyses showed that children who received the OT treatment in
319 combination with concomitant psychosocial treatment displayed a greater improvement in
320 social responsiveness than those who received psychosocial treatment or OT alone. A
321 modulating effect of parents' belief about allocated treatment was also identified, indicating
322 that parents who believed their child had been assigned the active treatment reported greater
323 benefit than those who believed their child received placebo, particularly in the experimental
324 group receiving actual OT. Finally, participants who were allocated to receive the placebo
325 treatment during the first double-blind phase of the trial and later crossed-over to receive the
326 active treatment during the second (single-blind) phase, displayed a significant improvement
327 in social responsiveness, over and above the 'placebo-induced' improvement noted in the
328 first phase. The next sections will address these observations in more detail.

329 The results of earlier continual OT trials in children with ASD have been equivocal: some with
330 beneficial outcomes (Parker et al., 2017; Yatawara et al., 2016), while others without
331 significant effect (Fastman et al., 2021; Guastella et al., 2015; Sikich et al., 2021). While it is
332 difficult to pinpoint the different factors contributing to variability in study results, several key
333 differences in adopted dosing schema, trial design, and participant demographics have been
334 put forward as important moderators. Furthermore, the particular context in which the OT
335 treatment is administered is also increasingly put forward as a vital factor for understanding
336 variability in treatment responses within and across studies. Initial single-dose administration
337 studies already noted that acute effects of OT can be modulated by contextual factors,
338 indicating for instance that OT-induced facilitation of cooperation and trust is most
339 pronounced towards in-group members, but is diminished, absent or even reversed towards
340 out-group members, particularly within threat-emanating contexts (de Dreu et al., 2010;
341 Mikolajczak et al., 2010). Also, in an early study by Heinrichs et al (2003), stress-reducing
342 effects of OT were significantly augmented when accompanied by a supportive context (i.e.,
343 social support from a friend).

344 Against this background, it has been theorized that OT may indeed open a 'window of
345 opportunity' to enhance prosocial behavior, but its potential can only be fully realized when

346 OT treatment is paired with a supportive context, such as effective concomitant behavioral
347 interventions that can support social skill development and improve prosocial behavior (Ford
348 & Young, 2021; Geschwind, 2021). In line with this notion, exploratory assessments within
349 our current trial revealed a significant synergetic modulation of treatment response related to
350 the presence of **concomitant psychosocial treatment** during the course of the OT trial
351 (screened through parent report, see **Table 1**), indicating maximal treatment effects in
352 children receiving the OT treatment in combination with ongoing psychosocial treatment.
353 Administration of OT as an adjunct to other therapeutic approaches has been explored
354 before. For example, in a study with schizophrenic patients, a six-week (12 session) social
355 cognition training was combined with OT administration (shortly before the start of each
356 session), and a significant improvement in empathic accuracy was observed (Davis et al.,
357 2014). Also, in patients with social anxiety disorders, OT treatment administered as an
358 adjunct to 4 sessions of public speaking-exposure therapy induced significant improvements
359 in mental representations of the self (Guastella et al., 2009). While preliminary, a pilot 6-week
360 OT administration study in which parents were stimulated to systematically engage with their
361 child in a positive social interaction or play session in the first hour after spray administration,
362 yielded unanimously positive treatment outcomes in 3-to-8-year-old children with ASD (n =
363 46), both in terms of social improvements and repetitive behaviors (Le et al., 2022).
364 Together, these and our study highlight the relevance of context and urge future clinical trials
365 to further elucidate whether clinical efficacy can be augmented when OT administration is
366 paired with targeted behavioral interventions that support similar states and (social)
367 behaviors.

368 Another notable observation was the identification that the **parent's belief about allocated**
369 **treatment** constituted a potentially important moderator of treatment response, indicating
370 that – within the OT group - parents who believed their child had been assigned the active
371 treatment reported greater benefit than those who believed their child received placebo.
372 Notably, the modulation was only significant in the group receiving the actual OT treatment,
373 not in the placebo group, and only for the immediate post-treatment outcome assessment,
374 not for the four-week follow-up assessment. These results therefore only partly concur with a
375 prior negative OT trial in which moderator effects by parent belief were evident, both in the
376 actual OT group, as well as in the placebo group (Guastella et al., 2015). On the one hand,
377 the modulation by parents' belief may reflect an expectancy bias, as noted in many prior
378 studies, especially in pediatric trials (King et al., 2009). Particularly in relation to OT
379 intervention research, increased biases can be expected, considering the large media
380 coverage and hype about purported prosocial benefits of the OT “love hormone” that may
381 eventually impact parents' expectancies (Guastella et al., 2015). However, a sole effect of

382 expectancy bias may be unlikely, since in that case, one would expect response modulations
383 to be present both in the OT and placebo groups. Since the modulating effect was specific to
384 the OT group, the possibility cannot be ruled out that parents may have actually correctly
385 identified real treatment responders, yielding maximal treatment responses in a particular
386 subgroup of children that displayed actual beneficial effects. Further, in line with the notion
387 that context may constitute an important moderator of treatment, one could also envisage
388 that parents who believed their child to receive the active treatment, may have provided their
389 children with more active socio-interactive family contexts during the four-week treatment
390 period, i.e., prompting them to increasingly engage in social experiences and learning,
391 thereby effectively boosting treatment responses.

392 Another important result relates to the observation that children who **crossed over** from
393 placebo (in phase I) to the actual OT treatment (in phase II), showed a significant further
394 improvement in social responsiveness over and above the substantial placebo-induced
395 improvement noted in phase I. Trial designs in which a phase of blinded placebo intervention
396 is administered before actual treatment allocation have been put forward before as an
397 efficient method to control for placebo effects and to improve detection of 'real' therapeutic
398 responses (Yatawara et al., 2016). The current observation of a significant further
399 improvement from a blinded placebo phase (double-blinded) to the active treatment (single-
400 blinded) provides support to this notion. While not explicitly addressed in the current study, it
401 is also noteworthy to stipulate that the oxytocinergic system itself has been suggested to
402 form a key mediator for facilitating placebo-induced improvements. Indeed, as postulated in
403 the recent oxytocin-placebo account of Itskovich et al. (2022); OT is suggested to mediate
404 social facilitation of placebo effects, an effect that is thought to be facilitated by increased
405 social connectedness between patient and clinician during trial participation. In line with this
406 notion, a prior OT administration trial showed that placebo-induced improvements in social
407 functioning coincided with endogenous increases in oxytocin secretion (Parker et al., 2017)

408 Further, in our trial, children who received the actual OT treatment in the first phase and
409 crossed over to a second phase of active treatment, showed only non-significant within-
410 subject improvements from phase I to phase II, particularly at the four-week follow-up
411 session of phase II - supporting prior observations of a retention of OT's beneficial effects,
412 also after cessation of the daily nasal spray administrations (Alaerts et al., 2020; Bernaerts et
413 al., 2020). It is noted indeed, that at the last follow-up session of the trial, the majority of
414 children of both the OT-first group and the placebo-first group displayed (reliable) beneficial
415 effects in social responsiveness, indicating that both an eight-week (with a four-week break
416 in the middle) or a continual four-week OT treatment were similarly able to induce a
417 significant beneficial outcome on a core ASD symptom domain. This observation adds to the

418 field's uncertainty regarding to-be-administered dosing schemas and durations. In multiple-
419 dose OT trials with individuals with ASD, daily dosing ranged from 8-80 IU and durations
420 from 4 continual days to 24 weeks, but strong empirical support for favoring one dosing
421 scheme over another is currently lacking. Some earlier single-dose trials suggested dose-
422 response curves to exhibit U-shaped forms (Lieberz et al., 2020; Spengler et al., 2017), a
423 notion that is supported by a recent chronic four-week OT administration trial in ASD,
424 identifying a daily total dose of 6 IU of TTA-121 (a new formulation of intranasal OT spray
425 with an enhanced bioavailability) to be the most efficacious one, compared to a lower (3 IU)
426 or higher (10 IU) daily dose (Yamasue et al., 2022). Furthermore, in terms of dosing scheme,
427 recent work showed that intermittent (every other day) administration may be therapeutically
428 more efficient than continual administration to obtain anxiolytic effects and reduce amygdala
429 reactivity (Kou et al., 2020). These observations were attributed to reflect a desensitization of
430 the endogenous oxytocinergic system upon too high concentrations and/or too high
431 frequencies of exogenous OT administration. The current observation that a single four-week
432 course can yield the same beneficial effects as a twice four-week course therefore reinforces
433 the notion that longer treatment durations do not necessarily facilitate higher treatment
434 responses. Similarly, in a recent large-scale trial administering OT over a 24-week period, it
435 was noted that the long duration might have attenuated initial early responses to OT (Sikich
436 et al., 2021). In light of these observations, future trials should be directed at identifying the
437 optimal dosing, administration length, and intervals of intranasal OT administration.

438 To conclude, while the current study showed no overall treatment-specific improvements,
439 important moderator effects were identified, providing initial indications that clinical efficacy
440 can be augmented when OT administration is paired with targeted behavioral interventions
441 that support similar states and (social) behaviors. Future trials are urged to further elucidate
442 the potential of embedding OT treatment within a (socially) stimulating context. Also, the role
443 of parental belief in modulating OT treatment responses needs further attention in
444 subsequent clinical trials. Finally, the observation that (reliable) improvements were
445 established in a large subset of children at the last session of the trial, either after a single
446 four-week course or after two four-week courses, should stimulate future trials to further
447 identify optimal dosing schemas, not only in terms of (daily) concentration, but also in terms
448 of administration length and the possibility of intermittent dosing.

449 **Funding and Disclosure**

450 This research was supported by an internal C1 fund of the KU Leuven [ELG-D2857-
451 C14/17/102], a Doctor Gustave Delpont fund of the King Baudouin Foundation and the
452 Branco Weiss fellowship of the Society in Science - ETH Zurich granted to KA. JP is
453 supported by the Marguerite-Marie Delacroix foundation and a postdoctoral fellowship of the
454 Flanders Fund for Scientific Research (FWO; 1257621N).

455 The funding sources had no further role in study design, data collection, analysis and
456 interpretation of data, writing of the report or in the decision to submit the paper for
457 publication.

458 **Conflict of Interest**

459 All authors declare that they have no conflicts of interest.

460 **Acknowledgements**

461 We would like to thank all the participants of the study and our colleagues of the Leuven
462 Autism Research Consortium (LAuRes).

463 References

- 464 Alaerts, K., Bernaerts, S., Prinsen, J., Dillen, C., Steyaert, J., & Wenderoth, N. (2020).
465 Oxytocin induces long-lasting adaptations within amygdala circuitry in autism: a
466 treatment-mechanism study with randomized placebo-controlled design.
467 *Neuropsychopharmacology*, *45*(7), 1141–1149. [https://doi.org/10.1038/s41386-020-](https://doi.org/10.1038/s41386-020-0653-8)
468 [0653-8](https://doi.org/10.1038/s41386-020-0653-8)
- 469 Alvares, G. A., Quintana, D. S., & Whitehouse, A. J. O. (2017). Beyond the hype and hope:
470 Critical considerations for intranasal oxytocin research in autism spectrum disorder.
471 *Autism Research*, *10*(1), 25–41. <https://doi.org/10.1002/aur.1692>
- 472 American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental*
473 *Disorders (DSM-5®)* (5th ed.). American Psychiatric Association.
474 <https://doi.org/10.1176/appi.books.9780890425596>
- 475 Anagnostou, E., Soorya, L., Chaplin, W., Bartz, J., Halpern, D., Wasserman, S., Wang, A. T.,
476 Pepa, L., Tanel, N., Kushki, A., & Hollander, E. (2012). Intranasal oxytocin versus
477 placebo in the treatment of adults with autism spectrum disorders: a randomized
478 controlled trial. *Molecular Autism*, *3*(1). <https://doi.org/10.1186/2040-2392-3-16>
- 479 Bakermans-Kranenburg, M. J., & van IJzendoorn, M. H. (2013). Sniffing around oxytocin:
480 review and meta-analyses of trials in healthy and clinical groups with implications for
481 pharmacotherapy. *Translational Psychiatry*, *3*(5), 1–14.
482 <https://doi.org/10.1038/TP.2013.34>
- 483 Bartz, J. A., Zaki, J., Bolger, N., & Ochsner, K. N. (2011). Social effects of oxytocin in
484 humans: context and person matter. *Trends in Cognitive Sciences*, *15*(7), 301–309.
485 <https://doi.org/10.1016/J.TICS.2011.05.002>
- 486 Bernaerts, S., Boets, B., Bosmans, G., Steyaert, J., & Alaerts, K. (2020). Behavioral effects
487 of multiple-dose oxytocin treatment in autism: a randomized, placebo-controlled trial
488 with long-term follow-up. *Molecular Autism*, *11*(1). [https://doi.org/10.1186/S13229-020-](https://doi.org/10.1186/S13229-020-0313-1)
489 [0313-1](https://doi.org/10.1186/S13229-020-0313-1)
- 490 Bodfish, J. W., Symons, F. J., Parker, D. E., & Lewis, M. H. (2000). Varieties of repetitive
491 behavior in autism: comparisons to mental retardation. *Journal of Autism and*
492 *Developmental Disorders*, *30*(3), 237–243. <https://doi.org/10.1023/A:1005596502855>
- 493 Bosmans, G., van de Walle, M., Goossens, L., & Ceulemans, E. (2014). (In)variability of
494 attachment in middle childhood: Secure base script evidence in diary data. *Behaviour*
495 *Change*, *31*(4), 225–242. <https://doi.org/10.1017/BEC.2014.18>
- 496 Constantino, J. N., & Gruber, C. P. (2012). *Social responsiveness scale (2nd. ed.). Manual.*
497 Western Psychological Services.
- 498 Davis, M. C., Green, M. F., Lee, J., Horan, W. P., Senturk, D., Clarke, A. D., & Marder, S. R.
499 (2014). Oxytocin-Augmented Social Cognitive Skills Training in Schizophrenia.
500 *Neuropsychopharmacology*, *39*(9), 2070. <https://doi.org/10.1038/NPP.2014.68>
- 501 de Dreu, C. K. W., Greer, L. L., Handgraaf, M. J. J., Shalvi, S., van Kleef, G. A., Baas, M.,
502 ten Velden, F. S., van Dijk, E., & Feith, S. W. W. (2010). The neuropeptide oxytocin
503 regulates parochial altruism in intergroup conflict among humans. *Science (New York,*
504 *N.Y.)*, *328*(5984), 1408–1411. <https://doi.org/10.1126/SCIENCE.1189047>

- 505 Fastman, J., Foss-Feig, J., Frank, Y., Halpern, D., Harony-Nicolas, H., Layton, C., Sandin,
506 S., Siper, P., Tang, L., Trelles, P., Zweifach, J., Buxbaum, J. D., & Kolevzon, A. (2021).
507 A randomized controlled trial of intranasal oxytocin in Phelan-McDermid syndrome.
508 *Molecular Autism*, 12(1). <https://doi.org/10.1186/s13229-021-00459-1>
- 509 Finzi, R., Cohen, O., Sapir, Y., & Weizman, A. (2000). Attachment styles in maltreated
510 children: a comparative study. *Child Psychiatry and Human Development*, 31(2), 113–
511 128. <https://doi.org/10.1023/A:1001944509409>
- 512 Ford, C. L., & Young, L. J. (2021). Refining oxytocin therapy for autism: context is key.
513 *Nature Reviews Neurology* 2021 18:2, 18(2), 67–68. [https://doi.org/10.1038/S41582-](https://doi.org/10.1038/S41582-021-00602-9)
514 [021-00602-9](https://doi.org/10.1038/S41582-021-00602-9)
- 515 Geschwind, D. H. (2021). Oxytocin for Autism Spectrum Disorder — Down, but Not Out. *New*
516 *England Journal of Medicine*, 385(16), 1524–1525.
517 https://doi.org/10.1056/NEJME2110158/SUPPL_FILE/NEJME2110158_DISCLOSURES
518 .PDF
- 519 Guastella, A. J., Gray, K. M., Rinehart, N. J., Alvares, G. A., Tonge, B. J., Hickie, I. B.,
520 Keating, C. M., Cacciotti-Saija, C., & Einfeld, S. L. (2015). The effects of a course of
521 intranasal oxytocin on social behaviors in youth diagnosed with autism spectrum
522 disorders: a randomized controlled trial. *Journal of Child Psychology and Psychiatry,*
523 *and Allied Disciplines*, 56(4), 444–452. <https://doi.org/10.1111/JCPP.12305>
- 524 Guastella, A. J., & Hickie, I. B. (2016). Oxytocin treatment, circuitry, and autism: A critical
525 review of the literature placing oxytocin into the autism context. *Biological Psychiatry,*
526 79(3), 234–242. <https://doi.org/10.1016/j.biopsych.2015.06.028>
- 527 Guastella, A. J., Hickie, I. B., McGuinness, M. M., Otis, M., Woods, E. A., Disinger, H. M.,
528 Chan, H. K., Chen, T. F., & Banati, R. B. (2013). Recommendations for the
529 standardisation of oxytocin nasal administration and guidelines for its reporting in
530 human research. *Psychoneuroendocrinology*, 38(5), 612–625.
531 <https://doi.org/10.1016/J.PSYNEUEN.2012.11.019>
- 532 Guastella, A. J., Howard, A. L., Dadds, M. R., Mitchell, P., & Carson, D. S. (2009). A
533 randomized controlled trial of intranasal oxytocin as an adjunct to exposure therapy for
534 social anxiety disorder. *Psychoneuroendocrinology*, 34(6), 917–923.
535 <https://doi.org/10.1016/J.PSYNEUEN.2009.01.005>
- 536 Heinrichs, M., Baumgartner, T., Kirschbaum, C., & Ehlert, U. (2003). Social support and
537 oxytocin interact to suppress cortisol and subjective responses to psychosocial stress.
538 *Biological Psychiatry*, 54(12), 1389–1398. [https://doi.org/10.1016/S0006-](https://doi.org/10.1016/S0006-3223(03)00465-7)
539 [3223\(03\)00465-7](https://doi.org/10.1016/S0006-3223(03)00465-7)
- 540 Huang, Y., Huang, X., Ebstein, R. P., & Yu, R. (2021). Intranasal oxytocin in the treatment of
541 autism spectrum disorders: A multilevel meta-analysis. *Neuroscience and Biobehavioral*
542 *Reviews*, 122, 18–27. <https://doi.org/10.1016/j.neubiorev.2020.12.028>
- 543 Itskovich, E., Bowling, D. L., Garner, J. P., & Parker, K. J. (2022). Oxytocin and the social
544 facilitation of placebo effects. *Molecular Psychiatry* 2022, 1–10.
545 <https://doi.org/10.1038/S41380-022-01515-9>
- 546 Jacobson, N. S., Roberts, L. J., Berns, S. B., & McGlinchey, J. B. (1999). Methods for
547 defining and determining the clinical significance of treatment effects: Description,

- 548 application, and alternatives. *Journal of Consulting and Clinical Psychology*, 67(3), 300–
549 307. <https://doi.org/10.1037/0022-006X.67.3.300>
- 550 Jurek, B., & Neumann, I. D. (2018). The Oxytocin Receptor: From Intracellular Signaling to
551 Behavior. *Physiological Reviews*, 98(3), 1805–1908.
552 <https://doi.org/10.1152/PHYSREV.00031.2017>
- 553 King, B. H., Hollander, E., Sikich, L., McCracken, J. T., Scahill, L., Bregman, J. D., Donnelly,
554 C. L., Anagnostou, E., Dukes, K., Sullivan, L., Hirtz, D., Wagner, A., & Ritz, L. (2009).
555 Lack of efficacy of citalopram in children with autism spectrum disorders and high levels
556 of repetitive behavior: citalopram ineffective in children with autism. *Archives of General
557 Psychiatry*, 66(6), 583–590. <https://doi.org/10.1001/ARCHGENPSYCHIATRY.2009.30>
- 558 Kosaka, H., Okamoto, Y., Munesue, T., Yamasue, H., Inohara, K., Fujioka, T., Anme, T.,
559 Orisaka, M., Ishitobi, M., Jung, M., Fujisawa, T. X., Tanaka, S., Arai, S., Asano, M.,
560 Saito, D. N., Sadato, N., Tomoda, A., Omori, M., Sato, M., ... Wada, Y. (2016). Oxytocin
561 efficacy is modulated by dosage and oxytocin receptor genotype in young adults with
562 high-functioning autism: a 24-week randomized clinical trial. *Translational Psychiatry*,
563 6(8), e872. <https://doi.org/10.1038/TP.2016.152>
- 564 Kou, J., Zhang, Y., Zhou, F., Sindermann, C., Montag, C., Becker, B., & Kendrick, K. M.
565 (2020). A randomized trial shows dose-frequency and genotype may determine the
566 therapeutic efficacy of intranasal oxytocin. *Psychological Medicine*, 1–10.
567 <https://doi.org/10.1017/S0033291720003803>
- 568 Le, J., Zhang, L., Zhao, W., Zhu, S., Lan, C., Kou, J., Zhang, Q., Zhang, Y., Li, Q., Chen, Z.,
569 Fu, M., Montag, C., Zhang, R., Yang, W., Becker, B., & Kendrick, K. M. (2022).
570 Infrequent intranasal oxytocin followed by positive social interaction improves symptoms
571 in autistic children: a randomized clinical trial. *MedRxiv*, 2022.01.03.22268708.
572 <https://doi.org/10.1101/2022.01.03.22268708>
- 573 Lieberz, J., Scheele, D., Spengler, F. B., Matheisen, T., Schneider, L., Stoffel-Wagner, B.,
574 Kiefe, T. M., & Hurlmann, R. (2020). Kinetics of oxytocin effects on amygdala and
575 striatal reactivity vary between women and men. *Neuropsychopharmacology*, 45(7),
576 1134. <https://doi.org/10.1038/S41386-019-0582-6>
- 577 Lord, C., Rutter, M., DiLavore, P. C., Risi, S., Gotham, K., & Bishop, S. L. (2012). *Autism
578 Diagnostic Observation Schedule: ADOS-2*. Western Psychological Services.
- 579 Mikolajczak, M., Gross, J. J., Lane, A., Corneille, O., de Timary, P., & Luminet, O. (2010).
580 Oxytocin Makes People Trusting, Not Gullible. *Psychological Science*, 21(8), 1072–
581 1074. <https://doi.org/10.1177/0956797610377343>
- 582 Muris P., Bodden D., Hale W, Birmaher B, & Mayer B. (2007). *SCARED-NL. Vragenlijst over
583 angst en bang-zijn bij kinderen en adolescenten. Handleiding bij de gereviseerde
584 Nederlandse versie van de Screen for Child Anxiety Related Emotional Disorders.*
585 Boom test uitgevers.
- 586 Pahlke, F., König, I., & Ziegler, A. (2004). Randomization In Treatment Arms (RITA): Ein
587 Randomisierungs-Programm für klinische Studien. *Inform Biom Epidemiol Med Biol*, 35,
588 1–22.
- 589 Parker, K. J., Oztan, O., Libove, R. A., Sumiyoshi, R. D., Jackson, L. P., Karhson, D. S.,
590 Summers, J. E., Hinman, K. E., Motonaga, K. S., Phillips, J. M., Carson, D. S., Garner,
591 J. P., & Hardan, A. Y. (2017). Intranasal oxytocin treatment for social deficits and

- 592 biomarkers of response in children with autism. *Proceedings of the National Academy of*
593 *Sciences of the United States of America*, 114(30), 8119–8124.
594 <https://doi.org/10.1073/PNAS.1705521114>
- 595 Roeyers, H., Thys, M., Druart, C., de Schryver, M., & Schittekatte, M. (2015). *SRS-2:*
596 *Screeningslijst voor autismespectrumstoornissen*. Hogrefe Uitgevers.
- 597 Shamay-Tsoory, S. G., & Abu-Akel, A. (2016). The Social Salience Hypothesis of Oxytocin.
598 *Biological Psychiatry*, 79(3), 194–202.
599 <https://doi.org/10.1016/J.BIOPSYCH.2015.07.020>
- 600 Sikich, L., Kolevzon, A., King, B. H., McDougle, C. J., Sanders, K. B., Kim, S.-J., Spanos, M.,
601 Chandrasekhar, T., Trelles, M. D. P., Rockhill, C. M., Palumbo, M. L., Witters Cundiff,
602 A., Montgomery, A., Siper, P., Minjarez, M., Nowinski, L. A., Marler, S., Shuffrey, L. C.,
603 Alderman, C., ... Veenstra-VanderWeele, J. (2021). Intranasal Oxytocin in Children and
604 Adolescents with Autism Spectrum Disorder. *New England Journal of Medicine*,
605 385(16), 1462–1473.
606 [https://doi.org/10.1056/NEJMOA2103583/SUPPL_FILE/NEJMOA2103583_DATA-](https://doi.org/10.1056/NEJMOA2103583/SUPPL_FILE/NEJMOA2103583_DATA-SHARING.PDF)
607 SHARING.PDF
- 608 Spengler, F. B., Schultz, J., Scheele, D., Essel, M., Maier, W., Heinrichs, M., & Hurlmann,
609 R. (2017). Kinetics and Dose Dependency of Intranasal Oxytocin Effects on Amygdala
610 Reactivity. *Biological Psychiatry*, 82(12), 885–894.
611 <https://doi.org/10.1016/J.BIOPSYCH.2017.04.015>
- 612 Watanabe, T., Kuroda, M., Kuwabara, H., Aoki, Y., Iwashiro, N., Tatsunobu, N., Takao, H.,
613 Nippashi, Y., Kawakubo, Y., Kunimatsu, A., Kasai, K., & Yamasue, H. (2015). Clinical
614 and neural effects of six-week administration of oxytocin on core symptoms of autism.
615 *Brain : A Journal of Neurology*, 138(Pt 11), 3400–3412.
616 <https://doi.org/10.1093/BRAIN/AWV249>
- 617 Wechsler, D. (2018). *WISC-V-NL. Wechsler Intelligence Scale for Children, Fifth Edition,*
618 *Dutch version*. Amsterdam: Pearson Benelux B.V.
- 619 Yamasue, H., Kojima, M., Kuwabara, H., Kuroda, M., Matsumoto, K., Kanai, C., Inada, N.,
620 Owada, K., Ochi, K., Ono, N., Benner, S., Wakuda, T., Kamen, Y., Inoue, J., Harada,
621 T., Tsuchiya, K., Umemura, K., Yamauchi, A., Ogawa, N., ... Yamasue, H. (2022).
622 Effect of a novel nasal oxytocin spray with enhanced bioavailability on autism: a
623 randomized trial. *Brain*. <https://doi.org/10.1093/BRAIN/AWAB291>
- 624 Yamasue, H., Okada, T., Munesue, T., Kuroda, M., Fujioka, T., Uno, Y., Matsumoto, K.,
625 Kuwabara, H., Mori, D., Okamoto, Y., Yoshimura, Y., Kawakubo, Y., Arioka, Y., Kojima,
626 M., Yuhi, T., Owada, K., Yassin, W., Kushima, I., Benner, S., ... Kosaka, H. (2020).
627 Effect of intranasal oxytocin on the core social symptoms of autism spectrum disorder: a
628 randomized clinical trial. *Molecular Psychiatry*, 25(8), 1849–1858.
629 <https://doi.org/10.1038/S41380-018-0097-2>
- 630 Yatawara, C. J., Einfeld, S. L., Hickie, I. B., Davenport, T. A., & Guastella, A. J. (2016). The
631 effect of oxytocin nasal spray on social interaction deficits observed in young children
632 with autism: a randomized clinical crossover trial. *Molecular Psychiatry*, 21(9), 1225–
633 1231. <https://doi.org/10.1038/MP.2015.162>

Table 1.**Demographic characteristics of the trial participants at baseline.**

Mean baseline scores listed separately for the oxytocin and placebo treatment groups. *T*- and *p*-values correspond to independent sample *t*-tests assessing between-group differences in baseline scores.

| | Oxytocin | | | Placebo | | | <i>t</i> -value | <i>p</i> -value |
|---|------------|--------|---------|------------|--------|---------|-----------------|-----------------|
| | N | mean | ± SD | N | mean | ± SD | | |
| Age | 38 | 10.48 | ± 1.32 | 39 | 10.39 | ± 1.23 | 0.28 | 0.779 |
| Gender | 30 M / 8 F | | | 31 M / 8 F | | | | |
| Handedness | 35 R / 3 L | | | 33 R / 6 L | | | | |
| WISC-V* | | | | | | | | |
| Verbal IQ | 37 | 105.84 | ± 14.41 | 38 | 109.42 | ± 15.77 | -1.03 | 0.308 |
| Performance IQ | 38 | 104.05 | ± 15.36 | 38 | 101.66 | ± 12.75 | 0.74 | 0.462 |
| ADOS-2 | | | | | | | | |
| Total | 33 | 9.48 | ± 3.78 | 32 | 9.16 | ± 4.15 | 0.33 | 0.740 |
| Social Affect | 31 | 7.13 | ± 3.55 | 32 | 7.47 | ± 3.72 | -0.37 | 0.712 |
| Restricted and Repetitive Behavior | 31 | 2.10 | ± 1.19 | 31 | 1.71 | ± 1.30 | 1.22 | 0.226 |
| Primary Outcome | | | | | | | | |
| SRS-2 | 38 | 89.26 | ± 21.66 | 39 | 87.87 | ± 20.03 | 0.29 | 0.771 |
| Secondary outcomes - parent report | | | | | | | | |
| RBS-R | 38 | 27.29 | ± 15.24 | 39 | 26.64 | ± 16.43 | 0.18 | 0.858 |
| SCARED Parent | 38 | 39.74 | ± 21.74 | 39 | 45.15 | ± 18.31 | -1.18 | 0.240 |
| Secondary outcomes - self report | | | | | | | | |
| SCARED child | 38 | 38.29 | ± 20.99 | 39 | 39.05 | ± 20.21 | -0.16 | 0.872 |
| ASCQ Anxious | 38 | 13.45 | ± 5.19 | 39 | 12.85 | ± 4.15 | 0.56 | 0.575 |
| ASCQ Avoidant | 38 | 13.79 | ± 4.00 | 39 | 14.08 | ± 3.86 | -0.32 | 0.749 |

| | | | | | | |
|---|----|--------------|----|--------------|-------------------------------|----------------|
| ASCQ Secure | 38 | 19.97 ± 3.50 | 39 | 19.23 ± 2.78 | 1.03 | 0.305 |
| Attachment Mother Anxiety | 38 | 4.74 ± 2.89 | 39 | 4.82 ± 2.78 | -0.13 | 0.897 |
| Attachment Mother Avoidance | 38 | 9.05 ± 4.76 | 39 | 7.97 ± 4.03 | 1.07 | 0.286 |
| Attachment Mother Secure | 38 | 16.76 ± 4.24 | 39 | 17.87 ± 3.13 | -1.31 | 0.195 |
| | | | | | Pearson chi square | p-value |
| Comorbidity** | 15 | | 15 | | 0.01 | 0.927 |
| Psychoactive Medication** | 22 | | 23 | | 0.01 | 0.923 |
| Psychosocial Therapy*** (N sessions/month) | 15 | 3.34 ± 0.97 | 14 | 3.31 ± 1.08 | 0.10 | 0.746 |

Data are shown as mean ± standard deviation. M male, F female, R right, L left, ADOS autism diagnostic observation schedule, SRS Social Responsiveness Scale, RBS-R Repetitive Behavior Scale-Revised, SCARED Screen for Child Anxiety Related Disorders, ASCQ Attachment Style Classification Questionnaire.

*WISC-V Wechsler Intelligence Scale for Children. The verbal intelligence quotient (IQ) was derived from the subtests Block design and Figure puzzles. The performance IQ was derived from the subtests Similarities and Vocabulary. **Detailed information on comorbidities and medication use is provided in [Supplementary Table 2](#). ***Concomitant participation in psychosocial therapies were screened for and if present (minimum of one session/month), frequency was determined as number of sessions/month. Reported psychosocial therapies include: Theory of mind training; emotion recognition training; social skills training; cognitive behavioral therapy; psychotherapy; self-esteem training; mood regulation; music therapy; hippotherapy; autism coach.

Table 2.

Effects of oxytocin treatment on primary and secondary outcome measures of the double-blind phase I.

Mean change from baseline scores are listed separately for the oxytocin and placebo treatment groups, and separately for the post assessment session (immediately after the four-week treatment) and the follow-up assessment (four weeks after cessation of the treatment).

| Outcome Measure | Within-group | | | | | | | | | | Between-group | | | | |
|---|--------------|-------|---|-------|-----------------|-----------------|----|-------|---|-------|------------------|-----------------|-----------------|-----------------|-----------------|
| | Oxytocin | | | | | Placebo | | | | | Cohen's <i>d</i> | <i>t</i> -value | <i>p</i> -value | | |
| | N | Mean | ± | SD | <i>t</i> -value | <i>p</i> -value | N | Mean | ± | SD | | | | <i>t</i> -value | <i>p</i> -value |
| Post assessment | | | | | | | | | | | | | | | |
| Primary Outcome | | | | | | | | | | | | | | | |
| SRS-2 | 38 | -4.08 | ± | 10.05 | -2.50 | 0.017 | 38 | -4.55 | ± | 10.23 | -2.74 | 0.009 | 0.05 | 0.20 | 0.839 |
| Secondary Outcomes - parent report | | | | | | | | | | | | | | | |
| RBS-R | 38 | -6.53 | ± | 11.26 | -3.57 | 0.001 | 38 | -6.76 | ± | 9.92 | -4.20 | 0.000 | 0.02 | 0.10 | 0.923 |
| SCARED Parent | 38 | -0.47 | ± | 11.61 | -0.25 | 0.803 | 38 | -4.95 | ± | 11.93 | -2.56 | 0.015 | 0.38 | 1.66 | 0.102 |
| Secondary Outcomes - self report | | | | | | | | | | | | | | | |
| SCARED Child | 38 | -4.74 | ± | 11.47 | -2.55 | 0.015 | 39 | -3.38 | ± | 11.32 | -1.87 | 0.070 | -0.12 | -0.52 | 0.604 |
| ASCQ Secure | 38 | -0.89 | ± | 2.42 | -2.27 | 0.029 | 39 | -0.92 | ± | 2.93 | -1.97 | 0.057 | 0.01 | 0.05 | 0.963 |
| ASCQ Anxious | 38 | -0.92 | ± | 3.24 | -1.75 | 0.088 | 39 | -1.00 | ± | 2.50 | -2.50 | 0.017 | 0.03 | 0.12 | 0.905 |
| ASCQ Avoidant | 38 | -0.29 | ± | 3.73 | -0.48 | 0.636 | 39 | -1.08 | ± | 3.30 | -2.04 | 0.049 | 0.22 | 0.98 | 0.330 |
| Attachment Mother Anxiety | 38 | 0.76 | ± | 2.95 | 1.59 | 0.120 | 39 | -0.33 | ± | 2.57 | -0.81 | 0.423 | 0.40 | 1.74 | 0.086 |
| Attachment Mother Avoidance | 38 | -0.21 | ± | 3.18 | -0.41 | 0.686 | 39 | -0.82 | ± | 3.94 | -1.30 | 0.201 | 0.17 | 0.75 | 0.458 |
| Attachment Mother Secure | 38 | 0.24 | ± | 4.18 | 0.35 | 0.729 | 39 | -0.18 | ± | 2.55 | -0.44 | 0.663 | 0.12 | 0.53 | 0.598 |
| Follow-up assessment | | | | | | | | | | | | | | | |
| Primary Outcome | | | | | | | | | | | | | | | |
| SRS-2 | 38 | -6.76 | ± | 11.19 | -3.73 | 0.001 | 39 | -5.38 | ± | 13.42 | -2.51 | 0.017 | -0.11 | -0.49 | 0.626 |
| Secondary Outcomes - parent report | | | | | | | | | | | | | | | |
| RBS-R | 38 | -4.55 | ± | 10.76 | -2.61 | 0.013 | 39 | -4.41 | ± | 8.28 | -3.33 | 0.002 | -0.01 | -0.07 | 0.948 |
| SCARED Parent | 38 | -2.92 | ± | 11.53 | -1.56 | 0.127 | 39 | -5.38 | ± | 9.52 | -3.53 | 0.001 | 0.23 | 1.02 | 0.309 |
| Secondary Outcomes - self report | | | | | | | | | | | | | | | |
| SCARED Child | 38 | -5.97 | ± | 9.84 | -3.74 | 0.001 | 39 | -6.36 | ± | 13.38 | -2.97 | 0.005 | 0.03 | 0.14 | 0.886 |
| ASCQ Anxious | 38 | -1.00 | ± | 3.92 | -1.57 | 0.125 | 39 | -2.38 | ± | 3.75 | -3.98 | 0.000 | 0.36 | 1.58 | 0.117 |

| | | | | | | | | | | | |
|-----------------------------|----|--------------|-------|--------------|----|--------------|-------|--------------|-------|-------|-------|
| ASCQ Avoidant | 38 | -0.84 ± 4.04 | -1.28 | 0.207 | 39 | -1.46 ± 3.09 | -2.95 | 0.005 | 0.17 | 0.76 | 0.452 |
| ASCQ Secure | 38 | -1.37 ± 3.34 | -2.53 | 0.016 | 39 | -0.82 ± 3.78 | -1.35 | 0.184 | -0.15 | -0.67 | 0.503 |
| Attachment Mother Anxiety | 38 | 0.84 ± 3.11 | 1.67 | 0.103 | 39 | 0.08 ± 3.07 | 0.16 | 0.877 | 0.25 | 1.09 | 0.281 |
| Attachment Mother Avoidance | 38 | -0.61 ± 3.89 | -0.96 | 0.343 | 39 | -0.67 ± 3.50 | -1.19 | 0.241 | 0.02 | 0.07 | 0.942 |
| Attachment Mother Secure | 38 | 0.66 ± 3.93 | 1.03 | 0.308 | 39 | 0.13 ± 2.83 | 0.28 | 0.779 | 0.16 | 0.68 | 0.498 |

Within-group *t*- and *p*-values correspond to single-sample *t*-tests assessing within-group change from baseline separately for the oxytocin and placebo group. Between-group *t*- and *p*-values correspond to independent sample *t*-tests assessing between-group differences in change from baseline scores. Cohen's *d* effect sizes (change from baseline_{OT}–change from baseline_{PL})/pooled SD) are reported where 0.2 is indicative of a small effect, 0.5 a medium effect and 0.8 a large effect. Data printed in bold show *p*-values or less than 0.05.

SRS Social Responsiveness Scale, RBS-R Repetitive Behavior Scale-Revised, SCARED Screen for Child Anxiety Related Disorders, ASCQ Attachment Style Classification Questionnaire. For all outcomes, except ASCQ secure and Attachment mother secure, negative change from baseline scores indicate pre-to-post improvement.

Figure 1

CONSORT flow diagram of participants in the trial.

Participants first underwent a double-blind phase (phase I) during which they were allocated to receive either oxytocin or placebo (four weeks of twice daily intranasal administration), followed by a four-week single-blind extension phase (phase II) during which all participants received four weeks of intranasal oxytocin. In both phases, treatment effects were assessed immediately after the four-week treatment (post) and at a follow-up session four weeks after cessation of the daily administrations (follow-up). For each assessment session, completed assessments are indicated separately for parent informant- and child self-reports.




Figure 2

Change in treatment responses according to the presence of concomitant psychosocial treatment.

Visualization of changes from baseline in parent-reported social responsiveness (SRS) of the double-blind phase (phase I), separately for children receiving only the oxytocin (n = 23) or placebo (n = 25) treatment and children receiving oxytocin (n = 15) or placebo (n = 13) treatment in combination with concomitant psychosocial treatment (pooled across the immediate post and four-week follow-up session). Lower scores indicate improvement. Vertical bars denote \pm standard errors.




Figure 3

Change in treatment responses according to parent reported beliefs about the allocated treatment.

Visualization of changes from baseline in parent-reported social responsiveness (SRS) of the double-blind phase (phase I) at the post (immediately after the four week treatment) and four-week follow-up session, separately for each treatment group (actual spray: oxytocin or placebo) and according to parent reported beliefs about the allocated treatment (oxytocin or placebo): oxytocin_{spray}/oxytocin_{belief}: n = 15; oxytocin_{spray}/placebo_{belief}: n = 16; placebo_{spray}/oxytocin_{belief}: n = 13; placebo_{spray}/placebo_{belief}: n = 21. Lower scores indicate improvement. Vertical bars denote \pm standard errors.




Figure 4

Treatment responses of the single-blind extension phase (phase II).

Visualization of changes from baseline in caregiver-reported social responsiveness (SRS) of the double-blind phase (phase I) and the single-blind extension phase (phase II), separately for each original treatment group (oxytocin-first, placebo-first) and assessment session (immediate post and four-week follow-up). Lower scores indicate improvement. Vertical bars denote \pm standard errors.

