

1 **Postural control telerehabilitation with a low-cost virtual reality protocol for children**
2 **with cerebral palsy: Protocol for a clinical trial.**

3

4 Valeska Gatica-Rojas^{a*}, Ricardo Cartes-Velásquez^b, Alex Soto-Poblete^c and Luis Eduardo
5 Cofré Lizama^d

6

7 a*. Tele-rehabilitation technology center and Neurosciences in Human Movement, Faculty
8 of Health Sciences, Universidad de Talca. Av. Lircay S/N, Talca, Chile. Email address:
9 valeskagaticar@gmail.com

10 b. Fundación Kimntrum, Concepción, Chile. Email address: cartesvelasquez@gmail.com

11 c. Institute of Mathematics and Physics, Universidad de Talca. Av. Lircay S/N, Talca,
12 Chile. Email address: alsoto@utalca.cl

13 d. School of Allied Health, Human Services and Sports, La Trobe University, Victoria,
14 Australia. Email address: e.cofrelizama@latrobe.edu.au

15

16 Corresponding author: Dr Valeska Gatica-Rojas at Tele-rehabilitation technology center
17 and Neurosciences in Human Movement, Faculty of Health Sciences, Universidad de
18 Talca, Av. Lircay S/N, Talca, Chile. Tel: +56 71 2418859. E-mail:
19 valeskagaticar@gmail.com

20

21

22

23

24

25

26

27

28

29

30

31

32 **Abstract**

33 **Objective:** To establish the feasibility and effectiveness of a rehabilitation programme
34 using low-cost virtual reality aimed at improving postural control in children with cerebral
35 type palsy spastic hemiplegia. It also seeks to compare low-cost virtual reality under two
36 delivery modalities, telerehabilitation (TR) and face-to-face (FtF).

37 **Methods:** Randomized controlled clinical trial from the ACTRN platform
38 ACTRN12621000117819. Eighteen sessions of low-cost virtual reality therapy will be
39 provided through both, FtF and TR modalities using a Nintendo Wii balance board. Each
40 programme will last for 6 weeks and will consist of 3 sessions per week of 25 minutes
41 each. The participants will include 40 patients diagnosed with cerebral palsy type spastic
42 hemiplegia. Twenty participants for each group. Participants will be assessed at baseline,
43 by the end of weeks 2, 4, and 6, and at weeks 8 and 10 (post-intervention follow-ups).
44 Clinical measures include the Modified-Modified Ashworth Scale for lower limbs,
45 Modified Ashworth Scale for upper limbs, timed up-and-go tests, the timed one-leg
46 standing and 6-minute walk test. Posturographic measures, including sway area and
47 velocity, under six conditions will be used: 2 statics and 4 dynamic conditions, which
48 include voluntary sway in the mediolateral direction following a metronome set at 30Hz
49 and 60Hz, and sway while playing 2 different videogames.

50 **Results:** This study provides an assessment of the feasibility and effectiveness of an
51 affordable rehabilitation programme using low-cost virtual reality aimed at improving
52 postural control in children with cerebral palsy.

53 **Conclusion:** Rehabilitation programme using low-cost virtual reality will improve postural
54 control in children with cerebral palsy type spastic hemiplegia and this programme
55 delivered using TR will be as effective as a FtF modality. The TR programme has be
56 designed to expand the coverage of physiotherapy services for children with cerebral palsy
57 in low-resource settings and in remote areas.

58

59 **Keywords:** telerehabilitation; cerebral palsy; balance, virtual reality; covid-19 pandemic.

60

61 **Introduction**

62 Cerebral palsy (CP) is the most common condition treated by therapists in infant
63 neurological units, and it is the most common cause of motor disability in children and
64 adolescents,^{1,2} with an increasing prevalence in developing countries.³ According to the
65 topographical classification, spastic hemiplegia (SHE) is one of the most prevalent types of
66 CP.^{1,4} In SHE, the mobility of one side of the body is more affected, generating poor
67 postural control during standing, leading to impaired performance in of daily functional
68 activities.^{5,6} To date, most studies have focused in developing interventions to improve
69 upper-limb function in SHE, for example, virtual reality interventions, which further allow
70 undertaking therapeutic training at home.⁷⁻⁹ The development of therapeutic interventions
71 aimed at improving postural control, on the other hand, has received less attention.¹⁰⁻¹²

72 Efficient, inexpensive, accessible, and engaging are some of the highlighted characteristics
73 sought after in the development of novel physiotherapy interventions, particularly in
74 children.¹³⁻¹⁶ Since the recent COVID-19 pandemic has interrupted non-essential
75 rehabilitation services, including those focusing on patients with CP, an urgent need for
76 remotely deliverable and less supervised interventions has also been highlighted to provide
77 health access to underserved populations.¹⁷ In this regard, telerehabilitation (TR) with the
78 use low-cost technological tools and exergames provide with an opportunity to meet the
79 before mentioned needs and which can further expand qualitatively and quantitatively
80 neurorehabilitation practice. This TR approach is not excluding of at-clinic interventions,
81 which combined may allow for a flexible dual-care model that integrates hospital and
82 home-based rehabilitation.¹³ To date, two earlier studies using TR in children with CP have
83 shown promising results in terms of feasibility and effectiveness compared with standard
84 face-to-face (FtF) interventions, however, these studies have not reported objective
85 outcomes.^{14,15}

86 Low-cost virtual reality exergames are well-accepted by children and adolescents due to
87 their ludic, motivating and engaging features, which facilitate a nearly seamless
88 incorporation into TR.¹⁶ However, research on TR using low-cost virtual reality is yet in its
89 infancy as high-level evidence comparing this modality with standard FtF interventions is
90 still lacking. It is noteworthy that low-cost virtual reality TR also brings the possibility to
91 standardize rehabilitation interventions, which is a pending task in neurorehabilitation.¹⁸

92 This clinical trial protocol seeks to establish the feasibility and effectiveness of a
93 rehabilitation programme using low-cost virtual reality aimed at improving postural control
94 in children with CP. It also seeks to compare low-cost virtual reality under two delivery
95 modalities, telerehabilitation (TR) and face-to-face (FtF, control group). We hypothesize
96 that a rehabilitation programme using low-cost virtual reality will improve postural control
97 in children with CP and that this programme delivered using TR will be as effective as a
98 FtF modality. The TR programme is designed to expand the coverage of physiotherapy
99 services for children with CP in low-resource settings and in remote areas, but also to
100 provide access to physiotherapy during restricted mobility as in the case of recent COVID-
101 19-related lockdowns.

102

103 **Materials and methods**

104 **Funding**

105 This study was supported by the Innovation Fund for Competitiveness, Maule Regional
106 Government – Chile No. 30.481.923-0.

107

108 **Design**

109 This is a double blinded, randomized, controlled trial designed following the Standard
110 Protocol Items: Recommendations for Interventional Trials (SPIRIT), (figure 1).¹⁹

111 The trial adheres to the Declaration of Helsinki and the Chilean laws of rights and duties of
112 the patient and research in humans. Ethical approval was obtained from the Ethics
113 Committee of the University of Talca (Ref. No. 24-2018). This study was registered with
114 the Australian and New Zealand Clinical Trials Registry (ACTRN12621000117819).
115 Written informed consent will be obtained from parents and participants over 7 years old;
116 verbal assent will be obtained from younger children.

117

118 **Study setting**

119 This study compares the effects of low-cost virtual reality in two distinct settings. The first
120 setting is clinical (Telerehabilitation Technology Centre and Neurosciences in Human
121 Movement, Universidad de Talca, Talca, Chile) and in which low-cost virtual reality is
122 delivered in a FtF modality (control group) with therapist interacting directly with the

123 patient. The second setting is a remote facility with no therapist on-site for the delivery of
124 low-cost virtual reality through TR, and could be an at-home, school, or rehabilitation
125 centre. The choice of setting will depend on each participant's adequate access to space and
126 equipment. All assessments will be conducted at the facility in which therapy was
127 conducted.

128

129 **Sample size**

130 The sample size was calculated to detect clinically relevant changes in postural control after
131 low-cost virtual reality in both groups by the end of week 6. Using data from earlier
132 studies,^{11,28} we considered a standard deviation of 4.974 cm² for the CoP sway area
133 (CoP_{Sway}), an alpha level of 0.05, 80% statistical power, and an allowance for 5% attrition;
134 we required a minimum of 20 participants in each group (total sample of 40).

135

136 **Eligibility criteria**

137 The inclusion criteria are as follows: (1) levels I or II on the Gross Motor Function
138 Classification System (GMFCS)²⁰ and on the Expanded and Revised Gross Motor Function
139 Classification System (GMFCS-ER)²¹; (2) aged 7 to 14 years (both sexes) with CP type
140 spastic hemiplegia; and (3) having mild or no cognitive impairment. Exclusion criteria
141 includes patients with other than CP neurological disorders, epilepsy, and uncorrected
142 vision or hearing. Participants with access to a Nintendo Wii at home before the
143 intervention commences will be also excluded.

144

145 **Interventions**

146 This study will compare the effects of low-cost virtual reality delivered through FtT and TR
147 modalities. The low-cost virtual reality will consist of in using a Nintendo Wii balance
148 board to play 4 exergames. In the FtF group therapy will be provided by a physiotherapist
149 with experience in the use of exergames for rehabilitation. In the TR modality, the
150 intervention (low-cost virtual reality) will be delivered by the patient's parent or caregivers.
151 In total, both groups (FtF and TR) will receive 18 sessions of low-cost virtual reality
152 divided in 3 sessions per week (Monday, Wednesday, and Friday) over a period of 6 weeks.
153 Each session will last for 25 minutes in which each participant will perform three series of

154 exergames resting 2 minutes in between. The first two series will include Snowboard,
155 Penguin Slide, and Super Hula Hoop. The first series is played with arms on the side and
156 the second one with hands on the waist to avoid the potential effect of training under one or
157 the other strategy. The third series is deep breathing in the Yoga game. During the first 2
158 weeks of training (first 6 sessions), patients will receive manual guidance and verbal
159 instructions after which (7th session onwards) only verbal instructions will be provided by
160 the parents, caregiver, or physiotherapist.

161 In the Snowboard exergame patients displace their centre of pressure (CoP) in the anterior-
162 posterior direction to swerve around flags whereas in the Penguin Slide movement occur on
163 the medial-lateral direction as patients need to score by catching fishes while maintaining a
164 penguin inside a continuously tilting platform. Finally, the Hula Hoop consists in
165 maintaining hoops on the waist while moving the CoP in a clockwise and anticlockwise
166 circular pattern. In the Yoga game the aim is to stand as quite as possible first with eyes
167 open and then with the eyes closed

168 This low-cost virtual reality programme is based on previous studies that have
169 demonstrated similar interventions can effectively train overall postural control.¹¹ For all
170 participants in both groups (FtF and TR), adherence will be encouraged through telephone
171 reminders. No modifications of the intervention protocols throughout the study are
172 considered at this stage.

173

174 **Outcomes**

175 Differences between the effects of FtF and TR using low-cost virtual reality on postural
176 control will be assessed using posturographic measures. In brief, posturography assesses
177 postural control through measures of the centre-of-pressure sway, of which the most
178 commonly used are sway area and trajectories and velocity; total and in the medial-lateral
179 and anterior-posterior directions.²² These CoP variables have been shown to be sensitive in
180 determining changes in postural control due to training in young adults,²³ patients with
181 post-stroke hemiparesis,²⁴ patients with Parkinson's disease,²⁵ and children with CP.²⁶⁻³⁰

182 Primary outcome in this study will be the CoP sway area (CoP_{area}), which has been shown
183 to be a reliable and valid measure of postural control in different clinical and nonclinical
184 populations.²²⁻³⁰ The CoP_{area} is an overall measure of the ability of the balance control

185 system to maintain a stable upright posture, to note, higher CoP_{area} values indicate poorer
186 balance control.

187 Secondaries outcome measures from the posturographic assessments to be used in this
188 study will be: standard deviation and velocity of the CoP in the medial-lateral and anterior-
189 posterior directions. The CoP standard deviation is a measure of the variability of CoP
190 displacements whereas the CoP velocity is a measure of the ability to react to changes in
191 posture with larger values indicating poorer balance control. The latter is due to late
192 postural adjustments lead to large CoP excursion that demand rapid corrections to maintain
193 stability.²²

194 Tertiary outcomes of this study will be: the Modified Modified Ashworth Scale (MMAS)
195 for lower limbs (ankle and knee flexor-extensor musculature), Modified Ashworth Scale
196 (MAS) for upper limbs (elbow and wrist flexor-extensor musculature), timed up-and-go
197 (TUG) tests, the timed one-leg standing (TOLS) and 6-minute walk test (6MWT). These
198 are widely used clinical measures of spasticity (MMAS and MAS) and overall physical
199 performance, which include some aspects of postural control (TOLS) and dynamic balance
200 (TUG and 6MWT).

201

202 **Intervention timeline**

203 Participants will be assessed at baseline, by the end of weeks 2, 4, and 6, and at weeks 8
204 and 10 (post-intervention follow-ups) using posturography and clinical measures.

205

206 **Recruitment**

207 Children and adolescents with congenital SHE-CP will be recruited from the Rehabilitation
208 Institute, Talca, Chile. This is an outpatient clinic that provides rehabilitation services to
209 patients with neurological disorders including CP children. It is noteworthy, that patients
210 from this centre will have little or no experience using a Nintendo Wii balance board. An
211 experienced staff of health professionals, comprised of a physician and two physical
212 therapists, will recruit participants according to the inclusion/exclusion criteria. Screening
213 for potential participants will continue until the calculated sample size is achieved.

214

215 **Allocation**

216 Participants will be randomly assigned to either control (FtF) or experimental group (TR)
217 with a 1:1 allocation as per a computer-generated randomization schedule stratified by site
218 and the baseline score of the Action Arm Research Test (ARAT; ≤ 21 versus >21) using
219 permuted blocks of random sizes. The block sizes will not be disclosed, to ensure
220 concealment.

221

222 **Management**

223 Participants will be informed of the group they will be randomly assigned to at the
224 beginning of the study. Participants in each group will be systematically asked about the
225 content of the interventions received to maintain engagement and compliance. Due to the
226 nature of the intervention neither participants nor staff can be blinded to allocation, but are
227 strongly inculcated not to disclose the allocation status of the participant at the follow up
228 assessments. An employee outside the research team will feed data into the computer in
229 separate datasheets so that the researchers can analyse data without having access to
230 information about the allocation.

231 All raw data will be electronically saved in coded text files whereas calculated and clinical
232 outcomes measures will be saved in a coded Excel spreadsheet. Data backups will be made
233 every week.

234

235 **Data collection**

236 All measurements will be performed at the same time of the day, under the same conditions
237 and by the same assessor at each time point. Posturographic measures (CoP displacements)
238 for both groups will be collected using an AMTI OR67 force platform (Watertown, MA,
239 USA). The AMTI NetForce software (Watertown, MA, USA) will be used for the
240 acquisition of moments and forces at 200Hz Data will be recorded, coded, and stored on a
241 personal computer. CoP variables will be calculated in Matlab R2012 (Mathworks Inc.,
242 Natick, MA, USA).

243 All primary and secondary posturographic measures will be assessed under 2 static and 4
244 dynamic conditions for 30 seconds each: (i) standing still with eyes open, (ii) standing still
245 with eyes closed, voluntary sway in the mediolateral direction following a metronome set at
246 (iii) 30Hz and (iv) 60Hz, and multidirectional sway while playing 2 different videogames

247 that challenge (v) anterior-posterior postural control (snowboard) and (vi) medial-lateral
248 postural control (penguin). It is expected that all assessments, including clinical ones, will
249 be completed within 30 minutes.

250

251 **Data analysis**

252 Statistical analyses will be performed using SPSS 20.00 for Windows. Descriptive statistics
253 will be used to calculate the demographic and clinical characteristics of participants in both,
254 FtF and TR groups (unpaired t-tests and χ^2 tests).

255 For all outcome measures, the Shapiro–Wilk and Levene tests will be used to measure the
256 normality and homogeneity of the variance, respectively. The Mann–Whitney U test will be
257 used to determine differences between the FtF and TR groups. Friedman’s one-way
258 analysis of variance (ANOVA) with post hoc pairwise comparisons (Wilcoxon signed-rank
259 test) will be used to determine the effect over time for each group (FtF and TR). For all
260 analyses, a p -value ≤ 0.05 will be considered statistically significant.

261

262 **Data monitoring**

263 A permanent monitoring will be made by the clinical team, oversights by the research
264 team and the Ethics Committee of the University of Talca. All the possible adverse events
265 will be informed immediately to the Ethics Committee of the University of Talca to
266 determine trial’s modifications or stop.

267

268 **Discussion**

269 The aims of this clinical trial protocol are to establish the feasibility and effectiveness of a
270 low-cost virtual reality to improve postural control in children with CP and to compare two
271 low-cost virtual reality delivery modalities, telerehabilitation (TR) and face-to-face (FtF).
272 Demonstrating that a TR is as efficient as a FtF intervention is crucial to adopt TR as an
273 alternative to expand the coverage of rehabilitation services for children with CP in low-
274 resource settings and in remote areas. Further, by showing the efficacy of TR as a method
275 to deliver low-cost virtual reality, physiotherapy can be potentially delivered when services
276 are inaccessible or closed as in the case of recent covid-19-related lockdowns. In this
277 protocol we hypothesize that low-cost virtual reality will improve postural control and that

278 both delivery methods will be equally effective. The latter is reasonable considering that
279 our previous research has shown that the FtF modality is effective in improving static and
280 dynamic postural control and reducing spasticity in children and adolescents with spastic
281 cerebral palsy.²⁹⁻³²

282

283 High-intensity physiotherapy is necessary for optimizing motor learning and improve
284 physical skills in patients with CP.³³ Regrettably, children and adolescents with CP
285 typically receive limited physiotherapy sessions, which implies a lost opportunity to
286 improve the functionality and quality of life of these patients during a critical period of
287 their lives. It is important to note that the neural maturity of sensory systems responsible for
288 the control of postural balance is sequential and dependent on age, reaching full maturity at
289 15-16 years.^{34, 35} Therefore, providing postural control rehabilitation during this age period
290 is essential to generate changes in the lives of these patients that will accompany them into
291 adulthood.

292 There are several factors that affect standing balance: not only the type of CP but also other
293 medical and environmental conditions. This diversity habitually demands for tailored
294 therapy that is not easily accessible, especially in low-resource and/or rural settings. Then,
295 this heterogeneity is consistent with the plethora of interventions available that are hard to
296 standardize precisely because of the diversity of patients, their needs, environments, and
297 resources, among other aspects³⁶⁻³⁹.

298 Virtual reality and low-cost virtual reality could alleviate this tension by creating a plethora
299 of programs that can be replicated in different settings in a tailored way.

300 In this regard, virtual environments created by software are characterized as artificial
301 functional spaces of multi-sensory interaction with a motivational purpose and are easy to
302 manipulate.⁴⁰ These virtual environments can be tailored for each patient, but all these
303 modifications are transferable for other patients and settings; this eventuality provides the
304 possibility to standardize most of the rehabilitation therapy. Furthermore, the
305 standardization of virtual reality-based therapies seems to be related to the adoption of
306 commercial video games,^{41,42} which are widely accessible worldwide. However, this does
307 not guarantee that these kinds of therapies are really standardized. The number of sessions,
308 duration, sequence, games, intervals, and other aspects need to be defined to obtain a

309 transferable standard. A recent systematic review on the effectiveness of virtual reality in
310 children and young adults with CP found limited confidence in effect estimation for
311 utilising virtual reality in this population; thus, further research is necessary.¹⁸

312 A low-cost virtual reality device provides greater accessibility and transferability to patients
313 with CP at neurological care centers. The Nintendo Wii system offers a simple and
314 affordable mode of virtual reality therapy.^{43,44} The Nintendo Wii and its peripheral balance
315 board have been proposed as a training tool to improve standing balance in the elderly,
316 adults with total knee replacements, post-stroke patients, and patients with Parkinson's
317 disease.^{9,45-48}

318 Another important consideration to generate relevant evidence is comprehensive
319 measurements. Thus, evaluations to build postural control construct and to determine the
320 effects of physical interventions in these populations must include clinical and laboratory
321 measurements. This clinical trial includes a set of clinical and laboratory evaluations that
322 enable knowing the effects of these therapies in the different aspects of postural control, for
323 example, the levels of the sensory systems—visual, vestibular, and mechanoreceptive
324 systems—in static and dynamic motor skills, within them include walking.

325 Home-based therapies for patients with CP seem to be feasible; however, the available
326 studies have used diverse methods and strategies, and, therefore, it is hard to determine the
327 impact of these kinds of therapies.⁴⁹ An efficient way to deliver these home-based therapies
328 is using TR, which appears to be an option to improve physical function in patients with
329 disabilities. However, most research has included low-quality evidence studies that make it
330 impossible to determine the effectiveness of TR.⁵⁰ In the case of patients with CP, two
331 studies have shown promising results,^{13,14} but the evidence is far from conclusive.

332

333 We have selected posturographic measures that have shown to be sensitive to the effects of
334 physiotherapy interventions.⁵¹⁻⁵² Velocity of CoP describes the neuromuscular response to
335 shifts in the body's centre of mass and serves as an indicator of stability. The mean velocity
336 of CoP has high reliability (ICC > 0.8) as a measurement parameter during standing
337 balance and is sensitive to changes of age-related postural impairment.⁵² However, the area
338 of the CoP presents behaviour that is more difficult to modify after physical interventions
339 in physiotherapy. For the area of the CoP (CoP_{Sway}), greater values indicate poorer balance

340 control as CoP gets closer to the limit of stability, which demands rapid stabilization
341 responses expressed in greater CoP velocities (V_{ML} and V_{AP}). Generating a modification in
342 postural control, specifically reducing the CoP area, involves working exhaustively at a
343 dose capable of achieving it. The dose in this study has been reviewed and verified both
344 with the scientific literature and tested in previous research.²⁸ There are correlation studies
345 between posturographic platform variables and clinical variables. In line with this, the area
346 of the CoP and other variables are poorly correlated with the clinical variables.^{53,54} These
347 findings suggest that the dose of physical therapies should be constantly reviewed, and
348 second, that clinical variables are more difficult to be transformed after a physical
349 intervention. This is related to the interpretation of the patients themselves in the face of
350 physical changes that their body and motor control has. There is evidence that the
351 perceptions of better postural control, balance, static, and dynamic stability by patients are
352 conditioned by the positive changes in the clinical variables investigated in these
353 evaluations. Therefore, it is essential to know the effects of this type of virtual physical
354 therapy in clinical trials and to consider pertinent clinical and gold standard laboratory
355 evaluations, such as the force platform, both of which will build the postural control
356 construct and interpret the results towards the patients.

357

358 Despite being discontinued in 2014, the Nintendo Wii continues to be a relevant tool for
359 neurorehabilitation research. Recent research using this console to treat patients with CP
360 has shown positive effects compared to other kinds of rehabilitation services.⁵⁵⁻⁵⁷

361

362 This will be the first study assessing the effects of a six-week exercise programme remote
363 (TR) using a Nintendo Wii balance board to improve standing balance in children and
364 adolescents with CP type spastic hemiplegia compared with the effects of an FtF modality.
365 This trial will provide evidence on the effectiveness of the Nintendo Wii for improving
366 standing balance in patients with CP, a matter in which the current evidence is lacking. This
367 low-cost virtual reality TR programme is designed to expand the coverage of physical
368 therapy services for patients with CP in low-resource settings using affordable products,
369 such as the Nintendo Wii balance board. The scientific contribution towards improving
370 static or dynamic standing balance will support the idea that affordable technological tools

371 can positively contribute to the usual activities, such as walking, running, and climbing
372 stairs, that children with CP perform in their environment, namely playgrounds, schools,
373 and in the cities where they live. During the pandemic, the development of new
374 technological strategies in physical therapies has contributed to the advancement in
375 rehabilitation, this should allow in the future to install new proposals in physical
376 rehabilitation for the children with CP type spastic hemiplegia.

377

378

380 **References**

- 381 [1] Xianrong L, Zhujiang T, Guojun Y, et al. Effectiveness of exercise interventions for
382 children with cerebral palsy: A systematic review and meta-analysis of randomized
383 controlled trials. *J Rehabil Med* 2021 Apr 1;53(4):jrm00176.
- 384 [2] Oskoui M, Coutinho F, Dykeman J, et al. An update on the prevalence of cerebral
385 palsy: a systematic review and meta-analysis. *Dev. Med. Child Neurol.* 2013; 55: 509-19.
- 386 [3] Shengyi Y, Jiayue X, Jing G, et al. Increasing prevalence of cerebral palsy among
387 children and adolescents in China 1988-2020: A systematic review and meta-analysis. *J*
388 *Rehabil Med* 2021 May 24;53(5):jrm00195.
- 389 [4] Saxena S, Rao BK, Kumaran S. Analysis of postural stability in children with
390 cerebral palsy and children with typical development: an observational study. *Pediatr. Phys.*
391 *Ther.* 2014; 26: 325-30.
- 392 [5] Rojas VG, Rebolledo GM, Muñoz EG, et al. Differences in standing balance
393 between patients with diplegic and hemiplegic cerebral palsy. *Neural Regen. Res.* 2013; 8:
394 2478-83.
- 395 [6] Kassee C, Hunt C, Holmes MWR, et al. Home-based Nintendo Wii training to
396 improve upper-limb function in children ages 7 to 12 with spastic hemiplegic cerebral
397 palsy. *J. Pediatr. Rehabil. Med.* 2017; 10: 145-154.
- 398 [7] El-Shamy SM. Efficacy of Armeo Robotic Therapy Versus Conventional Therapy
399 on Upper Limb Function in Children With Hemiplegic Cerebral Palsy. *Am. J. Phys. Med.*
400 *Rehabil.* 2018; 97: 164-169.
- 401 [8] Shum LC, Valdes BA, Hodges NJ, et al. Error Augmentation in Immersive Virtual
402 Reality for Bimanual Upper-Limb Rehabilitation in Individuals With and Without
403 Hemiplegic Cerebral Palsy. *IEEE Trans. Neural Syst. Rehabil. Eng.* 2020; 28: 541-549.
- 404 [9] Jelsma J, Pronk M, Ferguson G, et al. The effect of the Nintendo Wii Fit on balance
405 control and gross motor function of children with spastic hemiplegic cerebral palsy. *Dev.*
406 *Neurorehabil.* 2013; 16: 27–37.
- 407 [10] Gatica-Rojas V, Méndez-Rebolledo G, Guzman-Muñoz E, et al. Does Nintendo Wii
408 Balance Board improve standing balance? A randomized controlled trial in children with
409 cerebral palsy. *Eur. J. Phys. Rehabil. Med.* 2017; 53: 535-544.


- 410 [11] Do JH, Yoo EY, Jung MY, et al. The effects of virtual reality-based bilateral arm
411 training on hemiplegic children's upper limb motor skills. *NeuroRehabilitation*. 2016; 38:
412 115-27.
- 413 [12] Nuara A, Fabbri-Destro M, Scalona E, et al. Telerehabilitation in response to
414 constrained physical distance: an opportunity to rethink neurorehabilitative routines. *J.*
415 *Neurol*. 2021; 2021: 1-12.
- 416 [13] Beani E, Menici V, Ferrari A, et al. Feasibility of a Home-Based Action
417 Observation Training for Children With Unilateral Cerebral Palsy: An Explorative Study.
418 *Front. Neurol*. 2020; 11: 16.
- 419 [14] Molinaro A, Micheletti S, Pagani F, et al. Action Observation Treatment in a tele-
420 rehabilitation setting: a pilot study in children with cerebral palsy. *Disabil. Rehabil*. 2020;
421 17: 1-6.
- 422 [15] Demers M, Martinie O, Winstein C, et al. Active Video Games and Low-Cost
423 Virtual Reality: An Ideal Therapeutic Modality for Children With Physical Disabilities
424 During a Global Pandemic. *Front. Neurol*. 2020;14: 601898.
- 425 [16] Westby A, Nissly T, Giesecker R, et al. Achieving Equity in Telehealth: "Centering
426 at the Margins" in Access, Provision, and Reimbursement. *J. Am. Board Fam. Med*. 2021;
427 34: S29-S32.
- 428 [17] Ayala L, Winter S, Byrne R, et al. Assessments and Interventions for Spasticity in
429 Infants With or at High Risk for Cerebral Palsy: A Systematic Review. *Pediatr. Neurol*.
430 2020; 20: S0887-8994(20)30349-0.
- 431 [18] Fandim JV, Saragiotto BT, Porfirio GJM, et al. Effectiveness of virtual reality in
432 children and young adults with cerebral palsy: a systematic review of randomized
433 controlled trial. *Braz. J. Phys. Ther*. 2020; 20: S1413-3555(20)31132-1.
- 434 [19] Chan A-W, Tetzlaff JM, Altman DG, et al. SPIRIT 2013 Statement: Defining
435 standard protocol items for clinical trials. *Ann Intern Med* 2013;158:200-207.
- 436 [20] Palisano R, Rosenbaum P, Walter S, et al. Development and reliability of a system
437 to classify gross motor function in children with cerebral palsy. *Dev. Med. Child Neurol*.
438 1997; 39: 214-23.

- 439 [21] Fowler EG, Staudt LA, Greenberg MB, et al. Selective Control Assessment of the
440 Lower Extremity (SCALE): development, validation, and interrater reliability of a clinical
441 tool for patients with cerebral palsy. *Dev. Med. Child Neurol.* 2009; 51: 607-14.
- 442 [22] Duarte M, Freitas SM. Revision of posturography based on force plate for balance
443 evaluation. *Rev. Bras. Fisioter.* 2010; 14: 183-92.
- 444 [23] Kenny RPW, Atkinson G, Eaves DL, et al. The effects of textured materials on
445 static balance in healthy young and older adults: A systematic review with meta-analysis.
446 *Gait Posture.* 2019; 71: 79-86.
- 447 [24] Ioffe ME, Ustinova KI, Chernikova LA, et al. Supervised learning of postural tasks
448 in patients with poststroke hemiparesis, Parkinson's disease or cerebellar ataxia. *Exp. Brain*
449 *Res.* 2006; 168: 384–94.
- 450 [25] Rocchi L, Chiari L, Cappello A, et al. Identification of distinct characteristics of
451 postural sway in Parkinson's disease: a feature selection procedure based on principal
452 component analysis. *Neurosci. Lett.* 2006; 394: 140–5.
- 453 [26] Donker SF, Ledebt A, Roerdink M, et al. Children with cerebral palsy exhibit
454 greater and more regular postural sway than typically developing children. *Exp. Brain Res.*
455 2008; 184: 363-70.
- 456 [27] Cherng RJ, Su FC, Chen JJ, et al. Performance of static standing balance in children
457 with spastic diplegic cerebral palsy under altered sensory environments. *Am. J. Phys. Med.*
458 *Rehabil.* 2007; 78: 336–43.
- 459 [28] Gatica VF, Velásquez IS, Méndez GA, et al. Differences in standing balance in
460 patients with cerebral palsy and typically developing children. *Biomedica.* 2014; 34: 102-9.
- 461 [29] Gatica-Rojas V, Cartes-Velásquez R, Méndez-Rebolledo G, et al. Change in
462 functional balance after an exercise program with Nintendo Wii in Latino patients with
463 cerebral palsy: a case series. *J Phys Ther Sci.* 2016 Aug;28(8):2414-7.
- 464 [30] Gatica-Rojas V, Cartes-Velásquez R, Guzmán-Muñoz E, et al. Effectiveness of a
465 Nintendo Wii balance board exercise programme on standing balance of children with
466 cerebral palsy: A randomised clinical trial protocol. *Contemp Clin Trials Commun.*
467 2017;6:17-21

- 468 [31] Gatica-Rojas V, Méndez-Rebolledo G, Guzman-Muñoz E, et al. Does Nintendo Wii
469 Balance Board improve standing balance? A randomised controlled trial in children with
470 cerebral palsy. *Eur J Phys Rehabil Med.* 2017;53(4):535-544.
- 471 [32] Gatica-Rojas V, Cartes-Velásquez R, Méndez-Rebolledo G, et al. Effects of a
472 Nintendo Wii exercise program on spasticity and static standing balance in spastic cerebral
473 palsy. *Dev Neurorehabil.* 2017;20(6):388-391.
- 474 [33] Sakzewski L, Provan K, Ziviani J, et al. Comparison of dosage of intensive upper
475 limb therapy for children with unilateral cerebral palsy: how big should the therapy pill be?
476 *Res. Dev. Disabil.* 2015; 37: 9-16.
- 477 [34] Kyvelidou A, Harbourne RT, Shostrom VK, et al. Reliability of center of pressure
478 measures for assessing the development of sitting postural control in infants with or at risk
479 of cerebral palsy. *Arch. Phys. Med. Rehabil.* 2010; 91: 1593–601.
- 480 [35] Mitteregger E, Marsico P, Balzer J, et al. Translation and construct validity of the
481 Trunk Control Measurement Scale in children and youths with brain lesions. *Res. Dev.*
482 *Disabil.* 2015; 45: 343-52.
- 483 [36] Fahr A, Keller JW, van Hedel HWA. A systematic review of training methods that
484 may improve selective voluntary motor control in children with spastic cerebral palsy.
485 *Front. Neurol.* 2020; 11: 572038.
- 486 [37] O'Brien SM, Lichtwark GA, Carroll TJ, et al. Impact of Lower Limb Active
487 Movement Training in Individuals With Spastic Type Cerebral Palsy on Neuromuscular
488 Control Outcomes: A Systematic Review. *Front. Neurol.* 2020; 11: 581892.
- 489 [38] Inamdar K, Molinini RM, Panibatla ST, et al. Physical therapy interventions to
490 improve sitting ability in children with or at-risk for cerebral palsy: a systematic review and
491 meta-analysis. *Dev. Med. Child Neurol.* 2021; 63: 396-406.
- 492 [39] Liang X, Tan Z, Yun G, et al. Chen. Effectiveness of exercise interventions for
493 children with cerebral palsy: A systematic review and meta-analysis of randomized
494 controlled trials. *J. Rehabil, Med.* 2021; 53: jrm00176.
- 495 [40] Levin M and Demers M. Motor learning in neurological rehabilitation. *Disabil*
496 *Rehabil.* 2021 Dec;43(24):3445-3453
- 497 [41] Bonnechère B, Jansen B, Omelina L, et al. The use of commercial video games in
498 rehabilitation: a systematic review. *Int. J. Rehabil. Res.* 2016; 39: 277-290.

- 499 [42] Johansen T, Strøm V, Simic J, et al. Effectiveness of training with motion-
500 controlled commercial video games on hand and arm function in young people with
501 cerebral palsy: A systematic review and meta-analysis. *J. Rehabil. Med.* 2019; 52:
502 jrm00012.
- 503 [43] Gatica-Rojas V, Méndez-Rebolledo G. Virtual reality interface devices in the
504 reorganization of neural networks in the brain of patients with neurological diseases. *Neural*
505 *Regen. Res.* 2014; 9: 888-96.
- 506 [44] AlSaif AA, Alsenany S. Effects of interactive games on motor performance in
507 children with spastic cerebral palsy. *J. Phys. Ther. Sci.* 2015; 27: 2001-3.
- 508 [45] Gatica RV, Elgueta CE, Vidal SC, et al. Impacto del entrenamiento del balance a
509 través de realidad virtual en una población de adultos mayores. *Int. J. Morphol.* 2010; 28;
510 303-8.
- 511 [46] Bateni H. Changes in balance in older adults based on use of physical therapy vs the
512 Wii Fit gaming system: a preliminary study. *Physiotherapy.* 2012; 98: 211-6.
- 513 [47] Fung V, Ho A, Shaffer J, et al. Use of Nintendo Wii Fit™ in the rehabilitation of
514 outpatients following total knee replacement: a preliminary randomised controlled trial.
515 *Physiotherapy.* 2012; 98: 183-8.
- 516 [48] Pompeu JE, Mendes FA, Silva KG, et al. Effect of Nintendo Wii™-based motor
517 and cognitive training on activities of daily living in patients with Parkinson's disease: a
518 randomised clinical trial. *Physiotherapy.* 2012; 98: 196-204.
- 519 [49] Beckers LWME, Geijen MME, Kleijnen J, et al. Feasibility and effectiveness of
520 home-based therapy programmes for children with cerebral palsy: a systematic review.
521 *BMJ Open.* 2020; 10: e035454.
- 522 [50] Dias JF, Oliveira VC, Borges PRT, et al. Effectiveness of exercises by
523 telerehabilitation on pain, physical function and quality of life in people with physical
524 disabilities: a systematic review of randomised controlled trials with GRADE
525 recommendations. *Br. J. Sports Med.* 2021; 5
- 526 [51] Cofré Lizama LE, Pijnappels M, Faber GH, et al.: Age effects on mediolateral
527 balance control. *PLoS One*, 2014, 9: e110757.

- 528 [52] Rafał S, Janusz M, Wiesław O, et al.: Test-retest reliability of measurements of the
529 center of pressure displacement in quiet standing and during maximal voluntary body
530 leaning among healthy elderly men. *J Hum Kinet*, 2011, 28: 15-23.
- 531 [53] Kubicki A. Functional assessment in older adults: should we use timed up and go or
532 gait speed test?. *Neurosci Lett*, 2014, 577: 89-94.
- 533 [54] Gatica-Rojas et al. Correlation between center of pressure and functional balance in
534 non-faller elderly practitioners of Tai Chi Chuan. *J. Phys. Ther. Sci.* 28: 2350–2352, 2016
- 535 [55] Wang TN, Chen YL, Shieh JY, et al. Commercial Exergaming in Home-Based
536 Pediatric Constraint-Induced Therapy: A Randomized Trial. *OTJR (Thorofare N J)* 2021;
537 1539449220984110.
- 538 [56] Avcil E, Tarakci D, Arman N, et al. Upper extremity rehabilitation using video
539 games in cerebral palsy: a randomized clinical trial. *Acta Neurol. Belg.* 2021; 121(4):1053-
540 1060.
- 541 [57] Farr WJ, Green D, Bremner S, et al. Feasibility of a randomised controlled trial to
542 evaluate home-based virtual reality therapy in children with cerebral palsy. *Disabil.*
543 *Rehabil.* 2021; 43: 85-97.



medRxiv preprint doi: <https://doi.org/10.1101/2022.04.25.22274289>; this version posted April 28, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

1
2
3
4
5
6
7
8
9
10
11

12

13

14

15

16

17

18

19

medRxiv preprint doi: <https://doi.org/10.1101/2022.04.25.22274289>; this version posted April 28, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](#).

21

22 **Figure 1:** Flow chart according to the Standard Protocol Items: Recommendations for
23 Interventional Trials (SPIRIT). Participants, assessments and interventions. Through the
24 two groups of the trial.