

Genetic risk score for intracranial aneurysms to predict aneurysmal subarachnoid hemorrhage and identify associations with patient characteristics

Mark K. Bakker¹, Jos P. Kanning¹, Gad Abraham^{2,3}, Amy E. Martinsen^{4,5,6}, Bendik S. Winsvold^{4,6,7}, John-Anker Zwart^{4,5,6}, Romain Bourcier^{8,9}, Tomonobu Sawada¹⁰, Masaru Koido^{11,12}, Yoichiro Kamatani¹⁰, Sandrine Morel^{13,14}, Philippe Amouyel¹⁵, Stéphanie Debette^{16,17}, Philippe Bijlenga¹³, Takiy Berrandou¹⁸, Santhi K. Ganesh^{19,20}, Nabila Bouatia-Naji¹⁸, Gregory Jones²¹, Matthew Bown^{22,23}, HUNT All-In Stroke, CADISP group, International Consortium for Blood Pressure, International Headache Genetics Consortium, International Stroke Genetics Consortium (ISGC) Intracranial Aneurysm Working Group, Gabriël J.E. Rinkel¹, Jan H. Veldink¹, Ynte M. Ruigrok¹.

- 1 Department of Neurology and Neurosurgery, University Medical Center Utrecht Brain Center, Utrecht University, Utrecht, The Netherlands.
- 2 Cambridge Baker Systems Genomics Initiative, Baker Heart and Diabetes Institute, Melbourne, VIC, Australia.
- 3 Department of Clinical Pathology, University of Melbourne, Parkville, VIC, Australia.
- 4 Department of Research and Innovation, Division of Clinical Neuroscience, Oslo University Hospital, Oslo, Norway.
- 5 Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway.
- 6 K.G. Jebsen Center for Genetic Epidemiology, Department of Public Health and Nursing, NTNU, Norwegian University of Science and Technology, Trondheim, Norway.
- 7 Department of Neurology, Oslo University Hospital, Oslo, Norway
- 8 Université de Nantes, CHU Nantes, INSERM, CNRS, l'institut du thorax, Nantes, France.
- 9 CHU Nantes, Department of Neuroradiology, Nantes, France.
- 10 Graduate School of Frontier Sciences, The University of Tokyo, Tokyo, Japan.

- 11 Laboratory for Statistical and Translational Genetics, RIKEN Center for Integrative Medical Sciences, Yokohama, Japan.
- 12 Department of Cancer Biology, Institute of Medical Science, The University of Tokyo, Tokyo, Japan.
- 13 Neurosurgery Division, Department of Clinical Neurosciences, Faculty of Medicine, Geneva University Hospitals, Geneva, Switzerland.
- 14 Department of Pathology and Immunology, Faculty of Medicine, University of Geneva, Geneva, Switzerland.
- 15 LabEx DISTALZ-U1167, RID-AGE-Risk Factors and Molecular Determinants of Aging-Related Diseases, University of Lille, Lille, France; Inserm U1167, Lille, France; Centre Hospitalier Universitaire Lille, Lille, France; Institut Pasteur de Lille, Lille, France.
- 16 University of Bordeaux, INSERM, Bordeaux Population Health Center, UMR1219, F-33000 Bordeaux, France.
- 17 Bordeaux University Hospital, Department of Neurology, Institute of Neurodegenerative Diseases, F-33000 Bordeaux, France.
- 18 PARCC, INSERM, Université de Paris, F-750015, Paris, France.
- 19 Division of Cardiovascular Medicine, Department of Internal Medicine, University of Michigan Medical School, Ann Arbor, MI, USA.
- 20 Department of Human Genetics, University of Michigan Medical School, Ann Arbor, MI, USA.
- 21 Department of Surgery, University of Otago, Dunedin, New Zealand.
- 22 Department of Cardiovascular Sciences and National Institute for Health Research, University of Leicester, UK.
- 23 Leicester Biomedical Research Centre, University of Leicester, Glenfield Hospital, Leicester, UK.

Corresponding authors: Mark K. Bakker, MSc. Department of Neurology, University Medical Center Utrecht Brain Center, University Medical Center Utrecht, The Netherlands.

Heidelberglaan 100, 3584 CX Utrecht, The Netherlands. (m.k.bakker-25@umcutrecht.nl);

Ynte M. Ruigrok, MD PhD. Department of Neurology, University Medical Center Utrecht Brain Center, University Medical Center Utrecht, The Netherlands. Heidelberglaan 100, 3584 CX Utrecht, The Netherlands. (ij.m.ruigrok@umcutrecht.nl).

Word count: 3054

Reference count: 44

Abstract

Background: Rupture of an intracranial aneurysm (IA) causes aneurysmal subarachnoid hemorrhage (ASAH). There is no accurate prediction model for IA or ASAH in the general population. Recent discoveries in genetic risk for IA may allow improved risk prediction.

Methods: We constructed a genetic risk score including genetic association data for IA and 17 traits related to IA (a metaGRS) to predict ASAH incidence and IA presence. The metaGRS was trained in 1,161 IA cases and 407,392 controls in the UK Biobank and validated in combination with risk factors blood pressure, sex, and smoking in 828 IA cases and 68,568 controls from the Nordic HUNT study. We further assessed association between genetic risk load and patient characteristics in a cohort of 5,560 IA patients.

Results: The hazard ratio for ASAH incidence was 1.34 (95% confidence interval = 1.20-1.51) per SD increase of metaGRS. Concordance index increased from 0.63 [0.59-0.67] to 0.65 [0.62-0.69] upon including the metaGRS on top of clinical risk factors. The odds ratio for prediction of IA presence was 1.09 [95% confidence interval: 1.01-1.18], but did not improve area under the curve. The metaGRS was statistically significantly associated with age at ASAH ($\beta=-4.82\times 10^{-3}$ per year [-6.49×10^{-3} to -3.14×10^{-3}], $P=1.82\times 10^{-8}$), and location at the internal carotid artery (OR=0.92 [0.86 to 0.98], $P=0.0041$).

Conclusions: The metaGRS was predictive of ASAH incidence with modest added value over clinical risk factors. Genetic risk plays a role in clinical heterogeneity of IA. Additional studies are needed to identify the biological mechanisms underlying this heterogeneity.

Keywords:

Intracranial aneurysm, aneurysmal subarachnoid hemorrhage, genetics, risk prediction

KEY MESSAGES

What is already known on this topic

Recent advances in the understanding of genetic risk for IA opened an opportunity for risk prediction by combining genetic and conventional risk factors.

What this study adds

Here, we developed a genetic risk score based on genetic association information for IA and 17 related traits. This risk score improved prediction compared to a model including only conventional risk factors. Further, genetic risk was associated with age at ASAH and IA location.

How this study might affect research, practice, or policy

This study emphasizes the importance of combining conventional and genetic risk factors in prediction of IA. It provides a metric to develop an accurate risk assessment method including conventional and genetic risk factors.

Introduction

Rupture of an intracranial aneurysm (IA) leads to aneurysmal subarachnoid hemorrhage (ASAH), a severe type of stroke causing death in one third of cases, and permanent disability in another third.¹ It is one of few cardiovascular diseases in which women are at higher risk than men and is caused by a complex interplay of genetic factors and environmental risk factors,^{2,3} including smoking and hypertension.^{4,5} Aneurysmal rupture can be prevented by endovascular treatment or surgery, with relatively low risk of complications compared to the high case fatality and morbidity of ASAH.⁶ Therefore, prediction of ASAH has high potential in reducing disease burden.

Genetic risk scores (GRSs) showed potential in risk prediction of common diseases.⁷ New techniques improved prediction potential of GRSs by: 1. providing methods to include a large number of genetic variants⁸ and 2. combining GRSs for multiple traits (a so-called metaGRS), leading to improved prediction of, amongst others, coronary artery disease⁹ and ischemic stroke.¹⁰ These advances, combined with a substantial portion of heritability of IA being explained in the latest genome-wide association study (GWAS) of IA,² provide an opportunity for genetic risk prediction of IA.

A broad spectrum of clinical heterogeneity of IAs exists, including number, size and different locations of IAs.^{11,12} A GRS constructed with only seven single nucleotide polymorphisms (SNPs) was higher in patients with IAs at the middle cerebral artery (MCA) compared to those with IAs at other locations, in a cohort of 1,691 IA patients.¹³ In a cohort of 4,890 patients of whom 109 had an unruptured IA (UIA), a 10-SNP GRS was associated with

aneurysmal diameter and volume.¹⁴ These studies show a potential link between genetic predisposition and patient characteristics, but additional studies using larger populations and assessing across the genome are warranted.

We created a metaGRS for IA that incorporates GWAS summary statistics for IA together with summary statistics for other stroke subtypes and risk factors for IAs, and assessed its predictive performance for ASAH incidence and IA presence. Next, we assessed how the metaGRS associates with clinical characteristics of IA patients.

Methods

Figure 1 shows an overview of the study methods. In short, trait-level GRSs were constructed using summary statistics of the largest publicly available GWASs of IA and related traits (Supplementary Table 1). Optimal GRS model selection, and combining these GRSs into a metaGRS, was performed using the UK Biobank, a prospective cohort including 1,161 IA patients (959 with ASAH and 202 with UIA) and 407,392 controls (Figure 1A, Table 1). Predictive performance of the metaGRS was assessed in the HUNT prospective cohort study consisting of 828 IA patients (318 with ASAH and 510 with UIA) and 68,568 controls (Table 1). Associations of the metaGRS with patient characteristics were assessed in the well-phenotyped cohort of the international stroke genetics consortium (ISGC, www.strokegenetics.org) IA working group (ISGC-IA), including 5,560 IA patients of whom 3,918 with ASAH and 1,642 with UIA (Supplementary Table 2; Supplementary Data for participant selection and quality control of all cohorts).

Table 1. Baseline characteristics of the study populations

Characteristic	UK Biobank	HUNT study	ISGC-IA phenotype cohort
Age in years, mean (SD)	65.3 (8.0)	53.7 (17.5)	53.3 (13.0)**
Women, N (%)	220,835 (54.1%)	36,887 (53.0%)	3,786 (68.1%)
IA cases, N (%)	1,161 (0.28%)	828 (1.2%)	5,560 (100%)
ASAH cases, N (%)	959 (0.23%)	318 (0.46%)	3,916 (70.4%)
Controls, N	407,392	68,568	<i>not applicable</i>
SBP, mean (SD)	138.2 (18.1)	134.4 (21.1)	<i>no data</i>
Self-reported hypertension, N (%)***	<i>no data</i>	<i>no data</i>	2,150 (39.4%)
Ever smokers, N (%)	123,604 (30.3%)	35,204 (50.6%)	3,481 (69.4%)
Smoking packs/day, mean (SD)*	0.56 (0.42)	0.39 (0.30)	<i>no data</i>

UK Biobank characteristics are estimated based on the most recent assessment date.

**Packs/day are cumulative pack-years divided by age minus 16 based on ever smokers only.*

***In the ISGC-IA cohort only age at ASAH was recorded, and these numbers are shown.*

****In the ISGC-IA phenotype cohort, only self-reported hypertension was recorded.*

Constructing the metaGRS

Methods of the metaGRS construction are shown in Figure 1A. Summary statistics of large GWASs of IA and 24 traits (potentially) associated with IA were obtained (Supplementary Table 1).² These 24 traits included: 1. established risk factors for IA and/or ASAH, being diastolic and systolic blood pressure (SBP), smoking (cigarettes per day), and alcohol consumption (drinks per week);^{4, 5, 15-19} 2. suggestive risk factors including those related to female hormones (age at menarche, age at menopause and number of births),²⁰⁻²³ and cardiovascular disease risk (diabetes type II, body-mass index, waist-to-hip ratio, low- and high density lipoprotein levels, total cholesterol, and triglyceride levels),^{15, 24-29} migraine,³⁰⁻³² epilepsy (focal and generalized),^{2, 33} and years of education^{34, 35} and 3. diseases genetically

correlated with IA, being intracerebral hemorrhage, ischemic stroke, abdominal aortic aneurysms, and fibromuscular dysplasia (multifocal and any type).³⁶⁻³⁹ Only data of European-ancestry individuals were used while UK Biobank participants were excluded. Data pre-processing steps are described in the Supplementary Data.

For traits genetically correlated with IA ($P < 0.05$) trait-level GRSs were created (Supplementary Data for more details) using three methods: LD-based clumping with 9 different LD thresholds, summary statistics-based best linear unbiased predictor⁴⁰ and summary statistics-based BayesR⁸ (Figure 1A, Supplementary Data). To assess whether the trait-level GRSs captured risk of the respective trait, we assessed the association and predictive performance of optimal trait-level GRS with those traits in the UK Biobank. Samples with a missing genotype were ignored for association of the specific variant, while samples with missing phenotype were excluded (Supplementary Data for detailed methods).

For each trait, the trait-level GRS with highest Nagelkerke pseudo- R^2 in predicting IA status in the UK Biobank cohort was selected (Supplementary Table 3) and subsequently jointly analyzed in an elastic-net regression to obtain per-trait weights (Figure 1A). Trait-level SNP weights were scaled according to the per-trait weights and population standard deviation, and then summed over traits to create metaGRS SNP weights (see Supplementary Data for extensive methods). These analyses were performed for the whole cohort and for men and women separately. A separate GRS was constructed only considering the IA GRS, to assess potential added value of a metaGRS over an IA-only GRS.

Prediction of ASAH and IA by the metaGRS

Prediction by the metaGRS was evaluated in the HUNT study in two models: 1. for ASAH incidence using cox regression with age at ASAH as outcome and age at last assessment for controls as censoring time, and 2. for IA presence (including both UIA and ASAH) using a logistic regression with IA case-control status as outcome. For logistic regression, age, SBP, and average smoking packs per day since age 16 (SBP and smoking as three-knot polynomial spline), were included as covariates. For Cox regression, age was left out as covariate (further details described in the Supplementary Data). The added value of specific predictors was assessed using various models: 1. a reference model (sex and age), clinical model (only sex, age, SBP, and smoking), 2. a reference+metaGRS model, 3. a full model (clinical model + metaGRS), and 4. models leaving out a single predictor from the full model. Predictive value was determined in the HUNT study using a metaGRS created with GWAS summary statistics for IA leaving out samples from the HUNT study (Figure 1B).

Association between metaGRS and patient characteristics

The ISGC-IA phenotype cohort of 5,560 IA patients was used to determine the association of the metaGRS with the following patient characteristics: sex, smoking status (ever or never), self-reported hypertension, age at ASAH, and family history of IA (≥ 1 first degree relative with ASAH and/or UIA), IA location, number of IAs (single vs multiple), rupture status (UIA vs ASAH), and aneurysmal size at rupture. Locations of IA were grouped: 1) internal carotid artery (ICA) including the ICA, ophthalmic artery, and cavernous artery, 2) posterior communicating artery (PCOM), 3) anterior cerebral arteries (ACA) including the A1 anterior

segment, anterior communicating artery, and A2 segment, 4) MCA, and 5) posterior circulation (PC), including the vertebrobasilar system. IA at other locations were excluded from these analyses.

Since most cases of the ISGC-IA phenotype cohort were included in the IA GWAS, we created stratum-specific metaGRSs leaving out samples from the ISGC-IA phenotype cohort one GWAS stratum at a time from the IA GWAS summary statistics, resulting in nine metaGRS versions (Figure 1B, Supplementary Data). To control for differences in metaGRS versions between strata, we used the different cohorts of the ISGC-IA phenotype cohort (Supplementary Table 4) as covariate in all subsequent analyses.

We calculated associations between metaGRS and patient characteristics, correcting for sex and cohort using a generalized linear model. We tested whether statistically significantly associated phenotypes were independently associated from one another using a multivariate model. For each phenotype, samples with missing values were excluded for analysis of that specific phenotype. In analyses studying the association with IA location, we included only patients with one IA. Statistical analyses were done in R 4.1.2. Statistical significance was determined by Bonferroni correction for the number of phenotypes in the primary analyses: rupture status, sex, family history, IA multiplicity, five locations, age at ASAH, and size at rupture ($P < 0.05/11$). More details on the statistics are described in the Supplementary Data.

Results

Constructing the metaGRS

Seventeen out of 24 traits showed genetic correlation with IA ($P < 0.05$, Figure 1, Supplementary Table 5). Characteristics of the UK Biobank training cohort are shown in Table 1. We checked for the trait-level GRSs of which a representative phenotype was available in the UK Biobank, whether it was associated with that trait and an improvement of area under the curve (AUC) or R^2 compared to a reference model (for example, the trait-level GRS for SBP being associated with SBP in the UK biobank). This was the case for all phenotypes except intracerebral hemorrhage (Supplementary Table 6). Traits with an effect in the elastic-net regression were included in the metaGRS. Elastic-net regression weights for each trait-level GRS are shown in Supplementary Table 7. In total, 7,078,955 SNPs were included in the metaGRS. In separate models trained in only men or women in the UK Biobank, 6,618,190 and 6,671,269 SNPs remained, respectively.

Prediction of ASAH by the metaGRS

Characteristics of the HUNT study validation cohort are shown in Table 1. The metaGRS ranged from -0.83 to +0.50, with mean -0.22 and standard deviation 0.14. In the HUNT study, the metaGRS showed improved prediction of ASAH incidence compared to a reference model including only sex (hazard ratio [HR] = 1.34 [95% confidence interval, CI: 1.20 – 1.51], Supplementary Table 8). The C-index increased from 0.53 [0.49-0.56] to 0.58 [0.55-0.62] upon including the metaGRS to the reference model (Supplementary Table 9).

With a higher HR, the metaGRS seemed to outperform a GRS constructed using only summary statistics of IA (HR=1.25 [1.12-1.41], C-index=0.57 [0.53-0.61]). Maximum prediction was reached upon including the metaGRS on top of clinical risk factors, where the C-index increased from 0.63 [0.59–0.67] to 0.65 [0.62–0.69] (Figure 2). Given the HR per standard deviation, HR for a person at the highest 1% of metaGRS compared to median metaGRS is 1.99 [1.52-2.61], while the highest 1% versus lowest 1% had a HR of 3.96 [2.31-6.81].

In the model trained in women in the UK Biobank and validated in women in the HUNT study, the metaGRS alone had a larger effect compared to the model of both women and men and the model of men only (women: HR per SD of metaGRS=1.36 [1.18-1.60], men: 1.12 [0.93-1.34], Supplementary Figure 1, Supplementary Table 8). Similarly, clinical risk factors combined provided better prediction in women, and worse in men (women: C-index = 0.71 [0.67-0.75], men: 0.57 [0.52-0.62], Supplementary Table 9). Furthermore, metaGRS outperformed IA-only GRS in women, similar to in the whole cohort (IA-only GRS in women: HR=1.30 [1.11-1.51]).

Prediction of IA by the metaGRS

In prediction of IA presence (either UIA or ASAH) in the HUNT study, the metaGRS provided marginal added value on top of clinical risk factors (odds ratio [OR] = 1.09 [95% CI: 1.01-1.18], Supplementary Table 10). The metaGRS did not improve prediction above a model including clinical risk factors (AUC clinical model = 0.76 [0.75–0.78], AUC clinical + metaGRS = 0.76 [0.75–0.78], Supplementary Figure 2, Supplementary Table 11). Only the predictor

age showed independent added value (Δ AUC excluding age versus full model = -0.067 [-0.092 to -0.043]).

Association between metaGRS and patient characteristics

In the ISGC-IA phenotype cohort, patients with multiple IAs had a higher metaGRS than patients with a single IA, with nominal significance (OR=1.05 [1.01–1.09], $P=0.010$, Table 2, Figure 3A). Younger age at ASAH was associated with a higher metaGRS ($\beta=-4.82\times 10^{-3}$ per year [- 6.49×10^{-3} to -3.14×10^{-3}], $P=1.82\times 10^{-8}$, Figure 3B). Accordingly, the effect of one standard deviation increase of metaGRS on age at ASAH was -1.70 [-2.30 to -1.11] years. This equates to patients with a top 5% metaGRS suffering ASAH on average 2.70 [0.65–4.74] years earlier compared to patient with a mean metaGRS, while this is 3.83 [0.95–6.70] years earlier in patients with a top versus bottom 1% metaGRS. Of all aneurysmal locations, only patients with an IA at the ICA had lower genetic risk (OR=0.92 [0.86 to 0.98], $P=0.0041$, Figure 3C, Supplementary Figure 3-7). This effect reduced and was not statistically significant anymore when considering ruptured IAs only (OR=0.94 [0.86–1.03], $P=0.16$, and Supplementary Figure 8-12). No effect was observed for sex, positive family history, rupture status of an IA, or aneurysmal size at rupture (Table 2, Supplementary Figure 13-16). A higher metaGRS was associated with hypertension (OR=1.10 [95% CI=1.06–1.14], $P=3.82\times 10^{-7}$) and ever smokers (OR=1.14 [1.10–1.18], $P=9.30\times 10^{-10}$, Supplementary Table 12, Supplementary Figure 17-18), which is expected due to including summary statistics for these traits in the metaGRS.

Table 2. Associations between metaGRS and patient characteristics in the ISGC-IA phenotype cohort.

Subset	Phenotype	Comparison	P Value	Cases / controls	OR/Beta*	95% CI
All cases (N=5,560)	Ruptured	Yes / No	0.77	3918 / 1642	1.01	[0.96-1.05]
	Sex	Men / Women	0.74	1774 / 3786	1.01	[0.97-1.04]
	Positive Family History	Yes / No	0.80	1258 / 3666	1.01	[0.95-1.06]
	Multiple Aneurysms	Yes / No	0.010	1544 / 3955	1.05	[1.01-1.09]
Single aneurysm all cases (N=3,820)	ICA	ICA / other	0.0041	578 / 3242	0.92	[0.86-0.98]
	PCOM	PCOM / other	0.94	517 / 3303	1.00	[0.94-1.06]
	ACA	ACA / other	0.084	1377 / 2443	1.04	[1.00-1.08]
	MCA	MCA / other	0.78	918 / 2902	1.01	[0.96-1.06]
	PC	Posterior / anterior	0.75	430 / 3390	1.01	[0.94-1.08]
Single aneurysm ASAH only (N=2,751)	ICA	ICA / other	0.16	257 / 2494	0.94	[0.85-1.03]
	PCOM	PCOM / other	0.93	445 / 2306	1.00	[0.93-1.07]
	ACA	ACA / other	0.89	1131 / 1620	1.00	[0.95-1.05]
	MCA	MCA / other	0.53	597 / 2154	1.02	[0.96-1.08]
	PC	Posterior / anterior	0.72	321 / 2430	1.01	[0.94-1.09]
ASAH only (N=3,918)	Age at Rupture	Years per SD	1.82×10 ⁻⁸	N=3,893	-4.82×10 ⁻³	[-6.49×10 ⁻³ to -3.14×10 ⁻³]
	Age at Rupture (outcome)**	Years per SD	1.82×10 ⁻⁸	N=3,893	-1.70	[-2.30 to -1.11]
	Aneurysm Size at Rupture	Millimeter per SD	0.35	N=2859	2.61×10 ⁻³	[-2.87×10 ⁻³ to 8.10×10 ⁻³]

*OR/Beta: odds ratio for all phenotypes, or beta for age at rupture and aneurysm size and rupture. ISGC-IA: international stroke genetics consortium intracranial aneurysm working group. **Age at rupture was also used as outcome instead of dependent variable to obtain an interpretable effect size of the association between metaGRS and age at rupture. Cases/controls: number of persons with / without for the respective phenotype, or total sample size in the case of a quantitative phenotype. CI: confidence interval. ASAH: aneurysmal subarachnoid hemorrhage. ICA: internal carotid artery. PCOM: posterior communicating artery. ACA: anterior cerebral artery. MCA: middle cerebral artery. PC: posterior circulation. per SD: effect size per standard deviation of the independent variable.

In the multivariate model, the association of multiple IAs with metaGRS was not independent of smoking and hypertension (OR=1.03 [0.99–1.08], P=0.16). Upon including smoking and hypertension, the effect of location at the ICA slightly reduced and became nominally significant (OR=0.93 [0.87–0.99], P=0.021), while the association between age at

ASAH and metaGRS remained essentially the same ($\beta = -5.2 \times 10^{-3}$ per year $[-7.03 \times 10^{-3}$ to $-3.30 \times 10^{-3}]$, $P = 6.15 \times 10^{-8}$) (Supplementary Table 13-15). Since the mean metaGRS was higher in persons from Finland (Supplementary Figure 19) we performed the associations analyses on multiple IAs, location at the ICA, or age at ASAH excluding these persons but the effect sizes remained essentially the same (Supplementary Table 16).

Discussion

We created a metaGRS for IA based on GWAS summary statistics for IA and 17 IA-related traits. The metaGRS displayed added value in predicting ASAH incidence in combination with clinical risk factors, and added value compared to a GRS trained using only IA association data. Moreover, we demonstrated that prediction by the metaGRS for ASAH, which disease is seen more often in women than in men,^{4, 5} performs better in women than in men. Last we showed that the metaGRS was higher in patients who suffered ASAH at a younger age and lower in patients with an IA located at the ICA which associations were independent of hypertension and smoking.

In a previous study comparing genetic risk between 109 persons with UIA and 4,781 controls using a 10-SNP GRS no difference was found.¹⁴ This may be explained by the low number of patients and SNPs included in that GRS. Otherwise, it may be argued that the lack of association is caused by the fact that only UIAs were studied as in our study we were also unable to show predictive ability of the metaGRS for IA presence (combined group of UIA or ASAH). However, we think that in our study we were unable to predict IA presence because many UIAs are likely to be left undetected since UIAs are often incidental findings and therefore have a high chance of not being diagnosed in participants of observational population cohorts as used in our study.⁴¹ This probably resulted in low statistical power for prediction of UIA alone or in combination with ASAH. The argument that UIAs may be left undetected does not apply to the previous study on 109 persons with UIA and 4,781 controls as all participants were systematically screened with for UIAs using brain MRI.¹⁴

A metaGRS has been developed for other cardiovascular diseases being ischemic stroke and coronary artery disease.^{9, 10} In the present study, we found a hazard ratio per standard deviation of metaGRS for prediction of ASAH which was lower than assessed for coronary artery disease (HR=1.71, 95% CI=0.168-1.73), but higher than assessed for ischemic stroke (HR=1.26, 95% CI=1.22-1.31).^{9, 10}

A previous study indicated that persons with IA at the MCA had higher genetic risk than persons with IA at other locations, while no associations were found for aneurysmal size at ASAH, patient age at ASAH, or family history of UIA/ASAH.^{13, 14, 42} We did not replicate the increased genetic load for patients with an IA at the MCA. Since this effect was found in participants from Finland and the Netherlands, and we included additional countries, this might indicate population-dependent heterogeneity. Alternatively, due to the smaller sample size (N=1,613) the previous study may have been more sensitive to false positive findings, meaning there is in no true effect. Instead, we found a decreased genetic load in patients with an IA at the ICA, which location was not analyzed in the previous study.¹³ Interestingly, IAs at the ICA also have the lowest rupture risk compared to IA at other locations.⁴³ Overall, location-specific rupture risks and genetic risks followed similar patterns (highest risk for ACA, PCOM and PC, medium for MCA and lowest for ICA),⁴³ albeit not statistically significant. This could mean that location-specific rupture risks are in part a downstream result of genetic risk factors, but this remains to be confirmed in future studies.

The predictive performance of the metaGRS was in part captured by the inclusion of clinical risk factors smoking and SBP. This further supports the importance of genetic predisposition

for smoking and blood pressure in the risk for ASAH.² Whether the remaining added value of the metaGRS is driven by additional genetic causes independent of smoking and SBP, or whether the metaGRS better captures lifelong exposure to smoking and SBP than clinical measurements of these phenotypes, remains subject for future studies.

Prevalence of IA and incidence of ASAH is higher in women than men, in contrast to most other cardiovascular diseases.^{5, 41} Here, we found improved prediction of ASAH, when the metaGRS was trained and validated in women, and reduced when trained and validated in men. Predictive value of clinical risk factors was also better in women compared to in men. Sex differences have also been described for the number and location of IAs, for which characteristics we also showed differences in genetic load.⁴⁴ To understand the difference in genetic mechanisms of IA between men and women future investigations of genetic risk factors for IA and ASAH need to emphasize on sex differences and interactions between genetic variants and sex.

In summary, we developed a metaGRS which showed predictive ability for ASAH with modest added value over clinical risk factors, but no improved prediction of IA presence (UIA and ASAH combined). Genetic risk prediction was more effective in women, warranting further study on the sex-specific genetic causes of IA. Calibration of the metaGRS combined with clinical risk factors in independent high-risk and population cohorts are the necessary next steps before the metaGRS can be implemented for clinical use. The metaGRS was associated with age at ASAH and IA location, showing further evidence for a role of genetic risk in clinical heterogeneity of IA.

Acknowledgements

We thank the ICBP consortium, the MEGASTROKE consortium, and the ISGC for providing summary statistics. This research has been conducted using the UK Biobank Resource under application number 2532. MKB had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

The Trøndelag Health Study (HUNT) is a collaboration between HUNT Research Centre (Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology NTNU), Trøndelag County Council, Central Norway Regional Health Authority, and the Norwegian Institute of Public Health. The genotyping was financed by the National Institute of health (NIH), University of Michigan, The Norwegian Research council, and Central Norway Regional Health Authority and the Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology (NTNU). The genotype quality control and imputation has been conducted by the K.G. Jebsen center for genetic epidemiology, Department of public health and nursing, Faculty of medicine and health sciences, Norwegian University of Science and Technology (NTNU).

Data availability

The metaGRS SNP weights are available under here:

<https://doi.org/10.6084/m9.figshare.19672272>

(including UK Biobank for all samples, male only, and female only) and here:

<https://doi.org/10.6084/m9.figshare.19672269> (excluding UK Biobank for all samples, male

only, and female only). Data can be used and shared upon citation of this paper and adapted if the contributions are distributed under the same conditions.

References

1. Nieuwkamp DJ, Setz LE, Algra A, Linn FH, de Rooij NK, Rinkel GJ. Changes in case fatality of aneurysmal subarachnoid haemorrhage over time, according to age, sex, and region: a meta-analysis. *Lancet Neurol* 2009;8:635-642.
2. Bakker MK, van der Spek RAA, van Rheenen W, et al. Genome-wide association study of intracranial aneurysms identifies 17 risk loci and genetic overlap with clinical risk factors. *Nat Genet* 2020;52:1303-1313.
3. Bakker MK, Ruigrok YM. Genetics of Intracranial Aneurysms. *Stroke* 2021;52:3004-3012.
4. Vlak MH, Rinkel GJ, Greebe P, Algra A. Independent risk factors for intracranial aneurysms and their joint effect: a case-control study. *Stroke* 2013;44:984-987.
5. Etminan N, Chang HS, Hackenberg K, et al. Worldwide Incidence of Aneurysmal Subarachnoid Hemorrhage According to Region, Time Period, Blood Pressure, and Smoking Prevalence in the Population: A Systematic Review and Meta-analysis. *JAMA Neurol* 2019;76:588-597.
6. Algra AM, Lindgren A, Vergouwen MDI, et al. Procedural Clinical Complications, Case-Fatality Risks, and Risk Factors in Endovascular and Neurosurgical Treatment of Unruptured Intracranial Aneurysms: A Systematic Review and Meta-analysis. *JAMA Neurol* 2019;76:282-293.
7. Lewis CM, Vassos E. Polygenic risk scores: from research tools to clinical instruments. *Genome Med* 2020;12:44.
8. Lloyd-Jones LR, Zeng J, Sidorenko J, et al. Improved polygenic prediction by Bayesian multiple regression on summary statistics. *Nat Commun* 2019;10:5086.
9. Inouye M, Abraham G, Nelson CP, et al. Genomic Risk Prediction of Coronary Artery Disease in 480,000 Adults: Implications for Primary Prevention. *J Am Coll Cardiol* 2018;72:1883-1893.
10. Abraham G, Malik R, Yonova-Doing E, et al. Genomic risk score offers predictive performance comparable to clinical risk factors for ischaemic stroke. *Nat Commun* 2019;10:5819.
11. Nieuwkamp DJ, Algra A, Velthuis BK, Rinkel GJ. Clinical and radiological heterogeneity in aneurysmal sub-arachnoid haemorrhage according to risk-factor profile. *Int J Stroke* 2014;9:1052-1056.
12. Slot EMH, Rinkel GJE, Algra A, Ruigrok YM. Patient and aneurysm characteristics in familial intracranial aneurysms. A systematic review and meta-analysis. *PLoS One* 2019;14:e0213372.
13. van 't Hof FN, Kurki MI, Kleinloog R, et al. Genetic risk load according to the site of intracranial aneurysms. *Neurology* 2014;83:34-39.

14. Peymani A, Adams HH, Cremers LG, et al. Genetic Determinants of Unruptured Intracranial Aneurysms in the General Population. *Stroke* 2015;46:2961-2964.
15. Feigin VL, Rinkel GJ, Lawes CM, et al. Risk factors for subarachnoid hemorrhage: an updated systematic review of epidemiological studies. *Stroke* 2005;36:2773-2780.
16. Can A, Castro VM, Ozdemir YH, et al. Alcohol Consumption and Aneurysmal Subarachnoid Hemorrhage. *Transl Stroke Res* 2018;9:13-19.
17. Can A, Castro VM, Ozdemir YH, et al. Association of intracranial aneurysm rupture with smoking duration, intensity, and cessation. *Neurology* 2017;89:1408-1415.
18. Evangelou E, Warren HR, Mosen-Ansorena D, et al. Genetic analysis of over 1 million people identifies 535 new loci associated with blood pressure traits. *Nat Genet* 2018;50:1412-1425.
19. Liu M, Jiang Y, Wedow R, et al. Association studies of up to 1.2 million individuals yield new insights into the genetic etiology of tobacco and alcohol use. *Nat Genet* 2019;51:237-244.
20. Algra AM, Klijn CJ, Helmerhorst FM, Algra A, Rinkel GJ. Female risk factors for subarachnoid hemorrhage: a systematic review. *Neurology* 2012;79:1230-1236.
21. Perry JR, Day F, Elks CE, et al. Parent-of-origin-specific allelic associations among 106 genomic loci for age at menarche. *Nature* 2014;514:92-97.
22. Day FR, Ruth KS, Thompson DJ, et al. Large-scale genomic analyses link reproductive aging to hypothalamic signaling, breast cancer susceptibility and BRCA1-mediated DNA repair. *Nat Genet* 2015;47:1294-1303.
23. Barban N, Jansen R, de Vlaming R, et al. Genome-wide analysis identifies 12 loci influencing human reproductive behavior. *Nat Genet* 2016;48:1462-1472.
24. Kim JH, Jeon J, Kim J. Lower risk of subarachnoid haemorrhage in diabetes: a nationwide population-based cohort study. *Stroke Vasc Neurol* 2021.
25. Lindbohm JV, Kaprio J, Korja M. Cholesterol as a Risk Factor for Subarachnoid Hemorrhage: A Systematic Review. *PLoS One* 2016;11:e0152568.
26. Willer CJ, Schmidt EM, Sengupta S, et al. Discovery and refinement of loci associated with lipid levels. *Nat Genet* 2013;45:1274-1283.
27. Shungin D, Winkler TW, Croteau-Chonka DC, et al. New genetic loci link adipose and insulin biology to body fat distribution. *Nature* 2015;518:187-196.
28. Scott RA, Scott LJ, Magi R, et al. An Expanded Genome-Wide Association Study of Type 2 Diabetes in Europeans. *Diabetes* 2017;66:2888-2902.
29. Locke AE, Kahali B, Berndt SI, et al. Genetic studies of body mass index yield new insights for obesity biology. *Nature* 2015;518:197-206.
30. Kurth T, Kase CS, Schurks M, Tzourio C, Buring JE. Migraine and risk of haemorrhagic stroke in women: prospective cohort study. *BMJ* 2010;341:c3659.
31. Carter KN, Anderson N, Jamrozik K, Hankey G, Anderson CS, Australasian Co-operative Research on Subarachnoid Haemorrhage Study G. Migraine and risk of subarachnoid haemorrhage: a population-based case-control study. *J Clin Neurosci* 2005;12:534-537.
32. Gormley P, Anttila V, Winsvold BS, et al. Meta-analysis of 375,000 individuals identifies 38 susceptibility loci for migraine. *Nat Genet* 2016;48:856-866.

33. International League Against Epilepsy Consortium on Complex Epilepsies. Genome-wide mega-analysis identifies 16 loci and highlights diverse biological mechanisms in the common epilepsies. *Nat Commun* 2018;9:5269.
34. Broderick JP, Viscoli CM, Brott T, et al. Major risk factors for aneurysmal subarachnoid hemorrhage in the young are modifiable. *Stroke* 2003;34:1375-1381.
35. Rietveld CA, Medland SE, Derringer J, et al. GWAS of 126,559 individuals identifies genetic variants associated with educational attainment. *Science* 2013;340:1467-1471.
36. Malik R, Chauhan G, Traylor M, et al. Multiancestry genome-wide association study of 520,000 subjects identifies 32 loci associated with stroke and stroke subtypes. *Nat Genet* 2018;50:524-537.
37. Woo D, Falcone GJ, Devan WJ, et al. Meta-analysis of genome-wide association studies identifies 1q22 as a susceptibility locus for intracerebral hemorrhage. *Am J Hum Genet* 2014;94:511-521.
38. Jones GT, Tromp G, Kuivaniemi H, et al. Meta-Analysis of Genome-Wide Association Studies for Abdominal Aortic Aneurysm Identifies Four New Disease-Specific Risk Loci. *Circ Res* 2017;120:341-353.
39. Georges A, Yang M-L, Berrandou T-E, et al. Genetic association studies of fibromuscular dysplasia identify new risk loci and shared genetic basis with more common vascular diseases. *medRxiv* 2020.
40. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis. *Am J Hum Genet* 2011;88:76-82.
41. Vlak MH, Algra A, Brandenburg R, Rinkel GJ. Prevalence of unruptured intracranial aneurysms, with emphasis on sex, age, comorbidity, country, and time period: a systematic review and meta-analysis. *Lancet Neurol* 2011;10:626-636.
42. Kleinloog R, van 't Hof FN, Wolters FJ, et al. The association between genetic risk factors and the size of intracranial aneurysms at time of rupture. *Neurosurgery* 2013;73:705-708.
43. Greving JP, Wermer MJ, Brown RD, Jr., et al. Development of the PHASES score for prediction of risk of rupture of intracranial aneurysms: a pooled analysis of six prospective cohort studies. *Lancet Neurol* 2014;13:59-66.
44. Turan N, Heider RA, Zaharieva D, Ahmad FU, Barrow DL, Pradilla G. Sex Differences in the Formation of Intracranial Aneurysms and Incidence and Outcome of Subarachnoid Hemorrhage: Review of Experimental and Human Studies. *Transl Stroke Res* 2016;7:12-19.

Figure legends

Figure 1. Overview of constructing the metaGRS. A) Steps to create the metaGRS used for prediction in the HUNT study. Dark grey boxes with diagonal line indicate traits without genetic correlation with IA and were not used for the next step. B) Steps to create IA GWAS summary statistics to be included in an adjusted metaGRS version for phenotype-genotype

correlation analysis. Single IA GWAS strata were excluded from the IA GWAS in a leave-one-out manner prior to the ‘Best model selection’ step in panel A. IA: intracranial aneurysm, SBP: systolic blood pressure, CPD: cigarettes per day, DBP: diastolic blood pressure, BMI: body-mass index, YoE: years of education, WHR: waist-to-hip ratio, FMD (mf): multifocal fibromuscular dysplasia, FMD (any) any fibromuscular dysplasia, TG: triglycerides, HDL: high density lipoprotein, AAA: abdominal aortic aneurysm, ICH (deep): deep intracerebral hemorrhage, MIG (any): any migraine, IS (any): any ischemic stroke, Age Mrch: age at menarche, T2D: type II diabetes, Vert AD: vertebral artery dissection, LDL: low density lipoprotein, EPI (gen): generalized epilepsy, N births: number of births, EPI (focal): focal epilepsy, DPW: alcoholic drinks per week, TC: total cholesterol, Age Mnp: age at menopause, sCEU: stratum of mixed European ancestry, sFC: French Canada, sFIN: Finnish, sFRA: France, sNL1, sNL2: Netherlands, sPOL: Poland, sUK1, sUK2: United Kingdom, sUSA: United States of America, HUNT: Nordic HUNT study.

Figure 2. Prediction of ASAH using the metaGRS in the HUNT study. C-index according to different combinations of clinical risk factors and metaGRS are shown on the horizontal axis. Error bars denote 95% confidence intervals. HR: hazard ratio per standard deviation of the IA-only GRS or metaGRS in the model. Reference: model including only sex. Clinical: model including sex, SBP, and smoking. IA: intracranial aneurysm. GRS: genetic risk score.

Figure 3. Association of metaGRS with patient characteristics. Phenotypes with at least nominally significant effect on metaGRS are shown, except hypertension and smoking status, of which proxies were included in the metaGRS. The metaGRS was transformed to

mean 0, variance 1. A) metaGRS according to single or multiple IAs. Horizontal lines correspond to population mean (middle line), and mean \pm one population standard deviation. B) Effect of age at ASAH on metaGRS. Line denotes regression line, and shaded area is 95% confidence interval of the regression. C) Effect of IA at the ICA versus other locations, on the metaGRS. Format same as 3A. IA: intracranial aneurysm, ASAH: aneurysmal subarachnoid hemorrhage, ICA: internal carotid artery.

Article information

Conflicts of interest:

JHV reports to have sponsored research agreements with Biogen.

Sources of funding

We acknowledge the support from the Netherlands Cardiovascular Research Initiative: An initiative with support of the Dutch Heart Foundation, CVON2015-08 ERASE. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No. 852173). The project was funded in part by NIH grant R35HL161016 and University of Michigan Taubman Institute.

Role of the funders

The funders had no role in the design and conduct of the study, nor were they involved in preparing or approving the manuscript.

HUNT All-In Stroke:

Anne Hege Aamodt¹, Anne Heidi Skogholt², Ben M Brumpton², Cristen J Willer³, Else C Sandset^{4,5}, Espen S Kristoffersen^{6,7,8}, Hanne Ellekjær^{9,10}, Ingrid Heuch⁸, John-Anker Zwart^{8,11,2}, Jonas B Nielsen^{2,3,12}, Knut Hagen⁹, Kristian Hveem^{2,13,14}, Lars G Fritsche¹⁵, Laurent F Thomas^{2,16,17,18}, Linda M Pedersen⁸, Maiken E Gabrielsen², Oddgeir L Holmen¹³, Sigrid Børte^{11,2,19}, Wei Zhou^{20,21}

¹Department of Neurology, Oslo University Hospital, Oslo, Norway

²K. G. Jebsen Center for Genetic Epidemiology, Department of Public Health and Nursing, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

³Department of Internal Medicine, Division of Cardiovascular Medicine, University of Michigan, Ann Arbor, MI, 48109, USA

⁴Stroke Unit, Department of Neurology, Oslo University Hospital, Oslo, Norway

⁵Research and Development, The Norwegian Air Ambulance Foundation, Norway

⁶Department of General Practice, University of Oslo, Oslo, Norway

⁷Department of Neurology, Akershus University Hospital, Lørenskog, Norway

⁸Department of Research and Innovation, Division of Clinical Neuroscience, Oslo University Hospital, Oslo, Norway

⁹Department of Neuromedicine and Movement Science, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

¹⁰Stroke Unit, Department of Internal Medicine, St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway

¹¹Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway

¹²Department of Epidemiology Research, Statens Serum Institut, Copenhagen, Denmark

¹³HUNT Research Center, Department of Public Health and Nursing, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

¹⁴Department of Research, Innovation and Education, St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway

¹⁵Center for Statistical Genetics, Department of Biostatistics, University of Michigan, Ann Arbor, MI, 48109, USA

¹⁶Department of Clinical and Molecular Medicine, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

¹⁷BioCore - Bioinformatics Core Facility, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

¹⁸Clinic of Laboratory Medicine, St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway

¹⁹Research and Communication Unit for Musculoskeletal Health (FORMI), Department of Research and Innovation, Division of Clinical Neuroscience, Oslo University Hospital, Oslo, Norway

²⁰Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI, 48109, USA

²¹Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, USA

CADISP Group

Belgium: Departments of Neurology, Erasmus University Hospital, Brussels and Laboratory of Experimental Neurology, ULB, Brussels (Shérine Abboud, Massimo Pandolfo); Department of Neurology, Leuven University Hospital (Vincent Thijs).

France: Departments of Neurology, Lille University Hospital-Inserm U1171 (Didier Leys, Marie Bodenant), Sainte-Anne University Hospital, Paris (Fabien Louillet, Emmanuel Touzé, Jean-Louis Mas), Pitié-Salpêtrière University Hospital, Paris (Yves Samson, Sara Leder, Anne Léger, Sandrine Deltour, Sophie Crozier, Isabelle Méresse), Amiens University Hospital (Sandrine Canaple, Olivier Godefroy), Dijon University Hospital (Maurice Giroud, Yannick Béjot), Besançon University Hospital (Pierre Decavel, Elizabeth Medeiros, Paola Montiel, Thierry Moulin, Fabrice Vuillier); Inserm U744, Pasteur Institute, Lille (Jean Dallongeville).

Finland: Department of Neurology, Helsinki University Central Hospital, Helsinki (Antti J Metso, Tiina Metso, Turgut Tatlisumak); Germany: Departments of Neurology, Heidelberg University Hospital (Caspar Grond-Ginsbach, Christoph Lichy, Manja Kloss, Inge Werner, Marie-Luise Arnold), University Hospital of Ludwigshafen (Michael Dos Santos, Armin Grau); University Hospital of München (Martin Dichgans); Department of Rehabilitation: Schmieder-Klinik, Heidelberg (Constanze Thomas-Feles, Ralf Weber, Tobias Brandt).

Italy: Departments of Neurology: Brescia University Hospital (Alessandro Pezzini, Valeria De Giuli, Filomena Caria, Loris Poli, Alessandro Padovani), Milan University Hospital (Anna Bersano, Silvia Lanfranconi), University of Milano Bicocca, San Gerardo Hospital, Monza, Italy (Simone Beretta, Carlo Ferrarese), Milan Scientific Institute San Raffaele University Hospital (Giacomo Giacolone); Department of Rehabilitation, Santa Lucia Hospital, Rome (Stefano Paolucci).

Switzerland: Department of Neurology, Basel University Hospital (Philippe Lyrer, Stefan Engelter, Felix Fluri, Florian Hatz, Dominique Gisler, Leo Bonati, Henrik Gensicke, Margareth Amort).

UK: Clinical Neuroscience, St George's University of London (Hugh Markus). USA: Department of

Neurology, Salt Lake City, USA (Jennifer Majersik); Department of Neurology, University of Virginia, Charlottesville, USA (Bradford Worrall, Andrew Southerland); Department of Neurology, Baltimore, USA (John Cole, Steven Kittner)

International Consortium for Blood Pressure (ICBP):

Evangelos Evangelou^{1,2}, Helen R Warren^{3,4}, He Gao^{1,5}, Georgios Ntritsos², Niki Dimou², Tonu Esko^{16,17}, Reedik Mägi¹⁶, Lili Milani¹⁶, Peter Almgren¹⁸, Thibaud Boutin¹⁹, Stéphanie Debette^{20,21}, Jun Ding²², Franco Giulianini²³, Elizabeth G Holliday²⁴, Anne U Jackson²⁵, Ruifang Li-Gao²⁶, Wei-Yu Lin²⁷, Jian'an Luan²⁸, Massimo Mangino^{29,30}, Christopher Oldmeadow²⁴, Bram Peter Prins³¹, Yong Qian²², Muralidharan Sargurupremraj²¹, Nabi Shah^{32,33}, Praveen Surendran²⁷, Sébastien Thériault^{34,35}, Niek Verweij^{17,36,37}, Sara M Willems²⁸, Jing-Hua Zhao²⁸, Philippe Amouyel³⁸, John Connell³⁹, Renée de Mutsert²⁶, Alex SF Doney³², Martin Farrall^{40,41}, Cristina Menni²⁹, Andrew D Morris⁴², Raymond Noordam⁴³, Guillaume Paré³⁴, Neil R Poulter⁴⁴, Denis C Shields⁴⁵, Alice Stanton⁴⁶, Simon Thom⁴⁷, Gonçalo Abecasis⁴⁸, Najaf Amin⁴⁹, Dan E Arking⁵⁰, Kristin L Ayers^{51,52}, Caterina M Barbieri⁵³, Chiara Batini⁵⁴, Joshua C Bis⁵⁵, Tineka Blake⁵⁴, Murielle Bochud⁵⁶, Michael Boehnke²⁵, Eric Boerwinkle⁵⁷, Dorret I Boomsma⁵⁸, Erwin P Bottinger⁵⁹, Peter S Braund^{60,61}, Marco Brumat⁶², Archie Campbell^{62,64}, Harry Campbell⁶⁵, Aravinda Chakravarti⁵⁰, John C Chambers^{1,5,66-68}, Ganesh Chauhan⁶⁹, Marina Ciullo^{70,71}, Massimiliano Cocca⁷², Francis Collins⁷³, Heather J Cordell⁵¹, Gail Davies^{74,75}, Martin H de Borst⁷⁶, Eco J de Geus⁵⁸, Ian J Deary^{74,75}, Joris Deelen⁷⁷, Fabiola Del Greco M⁷⁸, Cumhur Yusuf Demirkale⁷⁹, Marcus Dörr^{80,81}, Bernard D K Keavney^{123,124}, Georg B Ehret^{50,82}, Roberto Elosua^{83,84}, Stefan Enroth⁸⁵, A Mesut Erzurumluoglu⁵⁴, Teresa Ferreira^{86,87}, Mattias Fränberg⁸⁸⁻⁹⁰, Oscar H Franco⁹¹, Ilaria Gandin⁶², Paolo Gasparini^{62,72}, Vilmantas Giedraitis⁹², Christian Gieger⁹³⁻⁹⁵, Giorgia Grotto^{62,72}, Anuj Goel^{40,41}, Alan J Gow^{74,96}, Vilmundur Gudnason^{97,98}, Xiuqing Guo⁹⁹, Ulf Gyllenstein⁸⁵, Anders Hamsten^{88,89}, Tamara B Harris¹⁰⁰, Sarah E Harris^{63,74}, Catharina A Hartman¹⁰¹, Aki S Havulinna^{102,103}, Andrew A Hicks⁷⁸, Edith Hofer^{104,105}, Albert Hofman^{91,106}, Jouke-Jan Hottenga⁵⁸, Jennifer E Huffman^{19,107,108}, Shih-Jen Hwang^{109,110}, Erik Ingelsson^{111,112}, Alan James^{111,112}, Rick Jansen¹¹³, Marjo-Riitta Jarvelin^{1,5,114-116}, Roby Joehanes^{107,117}, Åsa Johansson⁸⁵, Andrew D Johnson^{107,118}, Peter K Joshi⁶⁵, Pekka Jousilahti¹⁰², J Wouter Jukema¹¹⁹, Antti Jula¹⁰², Mika Kähönen^{120,121}, Sekar Kathiresan^{17,36,122}, Bernard D K Keavney^{123,124}, Kay-Teo Khaw¹²⁵, Paul Knekt¹⁰², Joanne Knight¹²⁶, Ivana Kolcic¹²⁷, Jaspal S Kooner^{5,67,68,128}, Seppo Koskinen¹⁰², Kati Kristiansson¹⁰², Zoltan Kutalik^{56,129}, Maris Laan¹³⁰, Marty Larson¹⁰⁷, Lenore J Launer¹⁰⁰, Benjamin Lehne¹, Terho Lehtimäki^{131,132}, David CM Liewald^{74,75}, Li Lin⁸², Lars Lind¹³³, Cecilia M Lindgren^{40,87,134}, YongMei Liu¹³⁵, Ruth JF Loos^{28,59,136}, Lorna M Lopez^{74,137,138}, Yingchang Lu⁵⁹, Leo-Pekka Lyytikäinen^{131,132}, Anubha Mahajan⁴⁰, Chrysovalanto Mamasoula¹³⁹, Jaume Marrugat⁸³, Jonathan Marten¹⁹, Yuri Milaneschi¹⁴⁰, Anna Morgan⁶², Andrew P Morris^{40,141}, Alanna C Morrison¹⁴², Peter J Munson⁷⁹, Mike A Nalls^{143,144}, Priyanka Nandakumar⁵⁰, Christopher P Nelson^{60,61}, Teemu Niiranen^{102,145}, Ilja M Nolte¹⁴⁶, Teresa Nutile⁷⁰, Albertine J Oldehinkel¹⁴⁷, Ben A Oostra⁴⁹, Paul F O'Reilly¹⁴⁸, Elin Org¹⁶, Sandosh Padmanabhan^{64,149}, Walter Palmas¹⁵⁰, Aarno Palotie^{103,151,152}, Alison Pattie⁷⁵, Brenda WJH Penninx¹⁴⁰, Markus Perola^{102,103,153}, Annette Peters^{94,95,154}, Ozren Polasek^{127,155}, Peter P Pramstaller^{78,156,157}, Quang Tri Nguyen⁷⁹, Olli T Raitakari^{158,159}, Rainer Rettig¹⁶¹, Kenneth Rice¹⁶², Paul M Ridker^{23,163}, Janina S Ried⁹⁴, Harriette Riese¹⁴⁷, Samuli Ripatti^{103,164}, Antonietta Robino⁷², Lynda M Rose²³, Jerome I Rotter⁹⁹, Igor Rudan¹⁶⁵, Daniela Ruggiero^{70,71}, Yasaman Saba¹⁶⁶, Cinzia F Sala⁵³, Veikko Salomaa¹⁰², Nilesh J Samani^{60,61}, Antti-Pekka Sarin¹⁰³, Reinhold Schmidt¹⁰⁴, Helena Schmidt¹⁶⁶, Nick Shrine⁵⁴, David Siscovick¹⁶⁷, Albert V Smith^{97,98}, Harold Snieder¹⁴⁶, Siim Söber¹³⁰, Rossella Sorice⁷⁰, John M Starr^{74,168}, David J Stott¹⁶⁹, David P Strachan¹⁷⁰, Rona J Strawbridge^{88,89}, Johan Sundström¹³³, Morris A Swertz¹⁷¹, Kent D Taylor⁹⁹, Alexander Teumer^{81,172}, Martin D Tobin⁵⁴, Maciej Tomaszewski^{123,124}, Daniela Toniolo⁵³, Michela Traglia⁵³, Stella Trompet^{119,173}, Jaakko Tuomilehto¹⁷⁴⁻¹⁷⁷, Christophe Tzourio²¹, André G Uitterlinden^{91,178}, Ahmad Vaez^{146,179}, Peter J van der Most¹⁴⁶, Cornelia M van Duijn⁴⁹, Germaine C Verwoert⁹¹, Veronique Vitart¹⁹, Uwe Völker^{81,180}, Peter Vollenweider¹⁸¹, Dragana Vuckovic^{62,182}, Hugh Watkins^{40,41}, Sarah H Wild¹⁸³, Gonneke Willemsen⁵⁸, James F Wilson^{19,65}, Alan F Wright¹⁹, Jie Yao⁹⁹, Tatjana Zemunik¹⁸⁴, Weihua Zhang¹⁶⁷, John R Attia²⁴, Adam S Butterworth^{27,185}, Daniel I Chasman^{23,163}, David Conen^{186,187}, Francesco Cucca^{188,189}, John Danesh^{27,185}, Caroline Hayward¹⁹, Joanna MM Howson²⁷, Markku Laakso¹⁹⁰, Edward G Lakatta¹⁹¹, Claudia Langenberg²⁸, Olle Melander¹⁸, Dennis O Mook-Kanamori^{26,192}, Colin NA Palmer³², Lorenz Risch¹⁹³⁻¹⁹⁵, Robert A Scott²⁸, Rodney J Scott²⁴, Peter Sever³²⁸, Tim D Spector²⁹, Pim van der Harst¹⁹⁶, Nicholas J Wareham²⁸, Eleftheria Zeggini³¹, Daniel Levy^{107,118}, Patricia B Munroe^{3,4}, Christopher Newton-Cheh^{134,197,198}, Morris J Brown^{3,4}, Andres Metspalu¹⁶, Bruce M. Psaty^{201,202}, Louise V Wain⁵⁴, Paul Elliott^{1,5,203-205}, Mark J Caulfield^{3,4}

1. Department of Epidemiology and Biostatistics, Imperial College London, London, UK.
2. Department of Hygiene and Epidemiology, University of Ioannina Medical School, Ioannina, Greece.
3. William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK.
4. National Institute for Health Research, Barts Cardiovascular Biomedical Research Center, Queen Mary University of London, London, UK.
5. MRC-PHE Centre for Environment and Health, Imperial College London, London, UK.

7. Division of Epidemiology, Department of Medicine, Institute for Medicine and Public Health, Vanderbilt Genetics Institute, Vanderbilt University Medical Center, Tennessee Valley Healthcare System (626)/Vanderbilt University, Nashville, TN, USA.
8. Vanderbilt Genetics Institute, Vanderbilt Epidemiology Center, Department of Obstetrics and Gynecology, Vanderbilt University Medical Center; Tennessee Valley Health Systems VA, Nashville, TN, USA.
9. Department of Epidemiology, Emory University Rollins School of Public Health, Atlanta, GA, USA.
10. Department of Biomedical Informatics, Emory University School of Medicine, Atlanta, GA, USA.
11. Massachusetts Veterans Epidemiology Research and Information Center (MAVERIC), VA Boston Healthcare System, Boston, USA.
12. Division of Aging, Department of Medicine, Brigham and Women's Hospital, Boston, MA, Department of Medicine, Harvard Medical School, Boston, MA, USA.
13. Atlanta VAMC and Emory Clinical Cardiovascular Research Institute, Atlanta, GA, USA.
14. VA Palo Alto Health Care System; Division of Cardiovascular Medicine, Stanford University School of Medicine, CA, USA.
15. Nephrology Section, Memphis VA Medical Center and University of Tennessee Health Science Center, Memphis, TN, USA.
16. Estonian Genome Center, University of Tartu, Tartu, Estonia.
17. Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge, MA, USA.
18. Department Clinical Sciences, Malmö, Lund University, Malmö, Sweden.
19. MRC Human Genetics Unit, MRC Institute of Genetics and Molecular Medicine, University of Edinburgh, Western General Hospital, Edinburgh, Scotland, UK
20. Department of Neurology, Bordeaux University Hospital, Bordeaux, France.
21. Univ. Bordeaux, Inserm, Bordeaux Population Health Research Center, CHU Bordeaux, Bordeaux, France.
22. Laboratory of Genetics and Genomics, NIA/NIH, Baltimore, MD, USA.
23. Division of Preventive Medicine, Brigham and Women's Hospital, Boston, MA, USA.
24. Hunter Medical Research Institute and Faculty of Health, University of Newcastle, New Lambton Heights, New South Wales, Australia.
25. Department of Biostatistics and Center for Statistical Genetics, University of Michigan, Ann Arbor, MI, USA.
26. Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, the Netherlands.
27. MRC/BHF Cardiovascular Epidemiology Unit, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK.
28. MRC Epidemiology Unit, University of Cambridge School of Clinical Medicine, Cambridge, UK.
29. Department of Twin Research and Genetic Epidemiology, Kings College London, London, UK.
30. NIHR Biomedical Research Centre at Guy's and St Thomas' Foundation Trust, London, UK.
31. Wellcome Trust Sanger Institute, Hinxton, UK.
32. Division of Molecular and Clinical Medicine, School of Medicine, University of Dundee, UK.
33. Department of Pharmacy, COMSATS Institute of Information Technology, Abbottabad, Pakistan.
34. Department of Pathology and Molecular Medicine, McMaster University, Hamilton, Canada.
35. Institut universitaire de cardiologie et de pneumologie de Québec-Université Laval, Québec City, Canada.
36. Cardiovascular Research Center and Center for Human Genetic Research, Massachusetts General Hospital, Boston, Massachusetts, MA, USA.
37. University of Groningen, University Medical Center Groningen, Department of Cardiology, Groningen, The Netherlands.
38. University of Lille, Inserm, Centre Hosp. Univ Lille, Institut Pasteur de Lille, UMR1167 - RID-AGE - Risk factors and molecular determinants of aging-related diseases, Epidemiology and Public Health Department, Lille, France.
39. University of Dundee, Ninewells Hospital & Medical School, Dundee, UK.
40. Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, UK.
41. Division of Cardiovascular Medicine, Radcliffe Department of Medicine, University of Oxford, Oxford, UK.
42. Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, UK.
43. Department of Internal Medicine, Section Gerontology and Geriatrics, Leiden University Medical Center, Leiden, The Netherlands.
44. Imperial Clinical Trials Unit, Stadium House, 68 Wood Lane, London, UK.
45. School of Medicine, University College Dublin, Ireland.
46. Molecular and Cellular Therapeutics, Royal College of Surgeons in Ireland, Dublin, Ireland.
47. International Centre for Circulatory Health, Imperial College London, London, UK.
48. Center for Statistical Genetics, Dept. of Biostatistics, SPH II, Washington Heights, Ann Arbor, MI, USA.
49. Genetic Epidemiology Unit, Department of Epidemiology, Erasmus MC, Rotterdam, the Netherlands.
50. Center for Complex Disease Genomics, McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore, MD, USA.

51. Institute of Genetic Medicine, Newcastle University, Newcastle upon Tyne, UK.
52. Sema4, a Mount Sinai venture, Stamford, CT, USA.
53. Division of Genetics and Cell Biology, San Raffaele Scientific Institute, Milano, Italy.
54. Department of Health Sciences, University of Leicester, Leicester, UK.
55. Cardiovascular Health Research Unit, Department of Medicine, University of Washington, Seattle, WA, USA.
56. Institute of Social and Preventive Medicine, University Hospital of Lausanne, Lausanne, Switzerland.
57. Human Genetics Center, School of Public Health, The University of Texas Health Science Center at Houston and Human Genome Sequencing Center, Baylor College of Medicine, One Baylor Plaza, Houston, TX, USA.
58. Department of Biological Psychology, Vrije Universiteit Amsterdam, EMGO+ institute, VU University medical center, Amsterdam, the Netherlands.
59. The Charles Bronfman Institute for Personalized Medicine, Icahn School of Medicine at Mount Sinai, NY, USA.
60. Department of Cardiovascular Sciences, University of Leicester, Leicester, UK.
61. NIHR Leicester Biomedical Research Centre, Glenfield Hospital, Groby Road, Leicester, UK.
62. Department of Medical, Surgical and Health Sciences, University of Trieste, Trieste, Italy.
63. Medical Genetics Section, Centre for Genomic and Experimental Medicine, Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, UK.
64. Generation Scotland, Centre for Genomic and Experimental Medicine, University of Edinburgh, Edinburgh, UK.
65. Centre for Global Health Research, Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Edinburgh, Scotland, UK
66. Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore, Singapore.
67. Department of Cardiology, Ealing Hospital, Middlesex, UK.
68. Imperial College Healthcare NHS Trust, London, UK.
69. Centre for Brain Research, Indian Institute of Science, Bangalore, India.
70. Institute of Genetics and Biophysics "A. Buzzati-Traverso", CNR, Napoli, Italy.
71. IRCCS Neuromed, Pozzilli, Isernia, Italy.
72. Institute for Maternal and Child Health IRCCS Burlo Garofolo, Trieste, Italy.
73. Medical Genomics and Metabolic Genetics Branch, National Human Genome Research Institute, NIH, Bethesda, MD, USA.
74. Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, 7 George Square, Edinburgh, UK.
75. Department of Psychology, University of Edinburgh, 7 George Square, Edinburgh, UK.
76. Department of Internal Medicine, Division of Nephrology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.
77. Department of Molecular Epidemiology, Leiden University Medical Center, Leiden, the Netherlands.
78. Institute for Biomedicine, Eurac Research, Bolzano, Italy - Affiliated Institute of the University of Lübeck, Lübeck, Germany.
79. Mathematical and Statistical Computing Laboratory, Office of Intramural Research, Center for Information Technology, National Institutes of Health, Bethesda, MD, USA.
80. Department of Internal Medicine B, University Medicine Greifswald, Greifswald, Germany.
81. DZHK (German Centre for Cardiovascular Research), partner site Greifswald, Greifswald, Germany.
82. Cardiology, Department of Medicine, Geneva University Hospital, Geneva, Switzerland.
83. CIBERCV & Cardiovascular Epidemiology and Genetics, IMIM, Dr Aiguader 88, Barcelona, Spain.
84. Faculty of Medicine, Universitat de Vic-Central de Catalunya, Vic, Spain.
85. Department of Immunology, Genetics and Pathology, Uppsala Universitet, Science for Life Laboratory, Uppsala, Sweden.
86. Wellcome Centre for Human Genetics, University of Oxford, Roosevelt Drive, Oxford, UK.
87. Big Data Institute, Li Ka Shing Center for Health for Health Information and Discovery, Oxford University, Old Road, Oxford, UK.
88. Cardiovascular Medicine Unit, Department of Medicine Solna, Karolinska Institutet, Stockholm, Sweden.
89. Centre for Molecular Medicine, L8:03, Karolinska Universitetsjukhuset, Solna, Sweden.
90. Department of Numerical Analysis and Computer Science, Stockholm University, Stockholm, Sweden.
91. Department of Epidemiology, Erasmus MC, Rotterdam, the Netherlands.
92. Department of Public Health and Caring Sciences, Geriatrics, Uppsala, Sweden.
93. Research Unit of Molecular Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany.
94. Institute of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany.
95. German Center for Diabetes Research (DZD e.V.), Neuherberg, Germany.
96. Department of Psychology, School of Social Sciences, Heriot-Watt University, Edinburgh, UK.

97. Faculty of Medicine, University of Iceland, Reykjavik, Iceland.
98. Icelandic Heart Association, Kopavogur, Iceland.
99. The Institute for Translational Genomics and Population Sciences, Department of Pediatrics, LABioMed at Harbor-UCLA Medical Center, Torrance, CA, USA.
100. Intramural Research Program, Laboratory of Epidemiology, Demography, and Biometry, National Institute on Aging, Bethesda, MD, USA.
101. Department of Psychiatry, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.
102. Department of Public Health Solutions, National Institute for Health and Welfare (THL), Helsinki, Finland.
103. Institute for Molecular Medicine Finland (FIMM), University of Helsinki, Helsinki, Finland.
104. Clinical Division of Neurogeriatrics, Department of Neurology, Medical University of Graz, Graz, Austria.
105. Institute for Medical Informatics, Statistics and Documentation, Medical University of Graz, Graz, Austria.
106. Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, USA.
107. National Heart, Lung and Blood Institute's Framingham Heart Study, Framingham, MA, USA.
108. The Population Science Branch, Division of Intramural Research, National Heart Lung and Blood Institute national Institute of Health, Bethesda, MD, USA.
109. Department of Medical Sciences, Molecular Epidemiology and Science for Life Laboratory, Uppsala University, Uppsala, Sweden.
110. Division of Cardiovascular Medicine, Department of Medicine, Stanford University School of Medicine, Stanford, CA USA.
111. Department of Pulmonary Physiology and Sleep, Sir Charles Gairdner Hospital, Hospital Avenue, Nedlands, Australia.
112. School of Medicine and Pharmacology, University of Western Australia.
113. Department of Psychiatry, VU University Medical Center, Amsterdam Neuroscience, Amsterdam, the Netherlands.
114. Biocenter Oulu, University of Oulu, Oulu, Finland.
115. Center For Life-course Health Research, University of Oulu, Oulu Finland.
116. Unit of Primary Care, Oulu University Hospital, Oulu, Oulu, Finland.
117. Hebrew SeniorLife, Harvard Medical School, Boston, MA, USA.
118. Population Sciences Branch, National Heart, Lung and Blood Institute, National Institutes of Health, Bethesda, MD, USA.
119. Department of Cardiology, Leiden University Medical Center, Leiden, the Netherlands.
120. Department of Clinical Physiology, Tampere University Hospital, Tampere, Finland.
121. Department of Clinical Physiology, Finnish Cardiovascular Research Center - Tampere, Faculty of Medicine and Life Sciences, University of Tampere, Tampere, Finland.
122. Broad Institute of the Massachusetts Institute of Technology and Harvard University, Cambridge, MA, USA.
123. Division of Cardiovascular Sciences, Faculty of Biology, Medicine and Health, The University of Manchester, Manchester, UK.
124. Division of Medicine, Manchester University NHS Foundation Trust, Manchester Academic Health Science Centre, Manchester, UK
125. Department of Public Health and Primary Care, Institute of Public Health, University of Cambridge, Cambridge, UK.
126. Data Science Institute and Lancaster Medical School, Lancaster, UK.
127. Department of Public Health, Faculty of Medicine, University of Split, Croatia.
128. National Heart and Lung Institute, Imperial College London, London, UK.
129. Swiss Institute of Bioinformatics, Lausanne, Switzerland.
130. Institute of Biomedicine and Translational Medicine, University of Tartu, Tartu, Estonia.
131. Department of Clinical Chemistry, Fimlab Laboratories, Tampere, Finland.
132. Department of Clinical Chemistry, Finnish Cardiovascular Research Center - Tampere, Faculty of Medicine and Life Sciences, University of Tampere, Tampere, Finland
133. Department of Medical Sciences, Cardiovascular Epidemiology, Uppsala University, Uppsala, Sweden.
134. Program in Medical and Population Genetics, Broad Institute, Cambridge, MA, USA.
135. Division of Public Health Sciences, Wake Forest School of Medicine, Winston-Salem, NC, USA.
136. Mindich Child health Development Institute, The Icahn School of Medicine at Mount Sinai, New York, NY, USA.
137. Department of Psychiatry, Royal College of Surgeons in Ireland, Education and Research Centre, Beaumont Hospital, Dublin, Ireland.
138. University College Dublin, UCD Conway Institute, Centre for Proteome Research, UCD, Belfield, Dublin, Ireland.
139. Institute of Health and Society, Newcastle University, Newcastle upon Tyne, UK.

140. Department of Psychiatry, Amsterdam Public Health and Amsterdam Neuroscience, VU University Medical Center/GGZ inGeest, Amsterdam, The Netherlands.
141. Department of Biostatistics, University of Liverpool, Block F, Waterhouse Building, Liverpool, UK.
142. Department of Epidemiology, Human Genetics and Environmental Sciences, School of Public Health, University of Texas Health Science Center at Houston, Houston, TX, USA.
143. Data Tecnica International, Glen Echo, MD, USA.
144. Laboratory of Neurogenetics, National Institute on Aging, Bethesda, USA.
145. Department of Medicine, Turku University Hospital and University of Turku, Finland.
146. Department of Epidemiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.
147. Interdisciplinary Center Psychopathology and Emotion regulation (ICPE), University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.
148. SGDP Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK.
149. British Heart Foundation Glasgow Cardiovascular Research Centre, Institute of Cardiovascular and Medical Sciences, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, UK.
150. Department of Medicine, Columbia University Medical Center, New York, NY, USA.
151. Analytic and Translational Genetics Unit, Department of Medicine, Department of Neurology and Department of Psychiatry Massachusetts General Hospital, Boston, MA, USA.
152. The Stanley Center for Psychiatric Research and Program in Medical and Population Genetics, The Broad Institute of MIT and Harvard, Cambridge, MA, USA.
153. University of Tartu, Tartu, Estonia.
154. German Center for Cardiovascular Disease Research (DZHK), partner site Munich, Neuherberg, Germany.
155. Psychiatric hospital "Sveti Ivan", Zagreb, Croatia.
156. Department of Neurology, General Central Hospital, Bolzano, Italy.
157. Department of Neurology, University of Lübeck, Lübeck, Germany.
158. Department of Clinical Physiology and Nuclear Medicine, Turku University Hospital, Turku, Finland.
159. Research Centre of Applied and Preventive Cardiovascular Medicine, University of Turku, Turku, Finland.
161. Institute of Physiology, University Medicine Greifswald, Karlsburg, Germany.
162. Department of Biostatistics University of Washington, Seattle, WA, USA.
163. Harvard Medical School, Boston MA.
164. Public health, Faculty of Medicine, University of Helsinki, Finland
165. Centre for Global Health Research, Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Scotland, UK.
166. Gottfried Schatz Research Center for Cell Signaling, Metabolism & Aging, Molecular Biology and Biochemistry, Medical University of Graz, Graz, Austria.
167. The New York Academy of Medicine, New York, NY, USA.
168. Alzheimer Scotland Dementia Research Centre, University of Edinburgh, Edinburgh, UK.
169. Institute of Cardiovascular and Medical Sciences, Faculty of Medicine, University of Glasgow, United Kingdom.
170. Population Health Research Institute, St George's, University of London, London, UK.
171. Department of Genetics, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.
172. Institute for Community Medicine, University Medicine Greifswald, Greifswald, Germany.
173. Department of Gerontology and Geriatrics, Leiden University Medical Center, Leiden, the Netherlands.
174. Dasman Diabetes Institute, Dasman, Kuwait.
175. Chronic Disease Prevention Unit, National Institute for Health and Welfare, Helsinki, Finland.
176. Department of Public Health, University of Helsinki, Helsinki, Finland.
177. Saudi Diabetes Research Group, King Abdulaziz University, Jeddah, Saudi Arabia.
178. Department of Internal Medicine, Erasmus MC, Rotterdam, the Netherlands.
179. Research Institute for Primordial Prevention of Non-communicable Disease, Isfahan University of Medical Sciences, Isfahan, Iran.
180. Interfaculty Institute for Genetics and Functional Genomics, University Medicine Greifswald, Greifswald, Germany.
181. Department of Internal Medicine, University Hospital, CHUV, Lausanne, Switzerland.
182. Experimental Genetics Division, Sidra Medical and Research Center, Doha, Qatar.
183. Centre for Population Health Sciences, Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Scotland, UK
184. Department of Biology, Faculty of Medicine, University of Split, Croatia.
185. The National Institute for Health Research Blood and Transplant Research Unit in Donor Health and Genomics, University of Cambridge, UK.

186. Division of Cardiology, University Hospital, Basel, Switzerland.
187. Division of Cardiology, Department of Medicine, McMaster University, Hamilton, Canada.
188. Institute of Genetic and Biomedical Research, National Research Council (CNR), Monserrato, Cagliari, Italy.
189. Department of Biomedical Sciences, University of Sassari, Sassari, Italy.
190. Institute of Clinical Medicine, Internal Medicine, University of Eastern Finland and Kuopio University Hospital, Kuopio, Finland.
191. Laboratory of Cardiovascular Science, NIA/NIH, Baltimore, MD, USA.
192. Department of Public Health and Primary Care, Leiden University Medical Center, Leiden, the Netherlands.
193. Labormedizinisches Zentrum Dr. Risch, Schaan, Liechtenstein.
194. Private University of the Principality of Liechtenstein, Triesen, Liechtenstein.
195. University Institute of Clinical Chemistry, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland.
196. Department of Cardiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.
197. Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA.
198. Cardiovascular Research Center, Massachusetts General Hospital, Boston, MA, USA.
201. Cardiovascular Health Research Unit, Departments of Medicine, Epidemiology and Health Services, University of Washington, Seattle, WA, USA.
202. Kaiser Permanente Washington Health Research Institute, Seattle, WA, USA.
203. National Institute for Health Research Imperial Biomedical Research Centre, Imperial College Healthcare NHS Trust and Imperial College London, London, UK.
204. UK Dementia Research Institute (UK DRI) at Imperial College London, London, UK
205. Health Data Research-UK London substantive site, London, U.K

International Headache Genetic Consortium (IHGC):

Padhraig Gormley^{1-4,81}, Verner Anttila^{2,3,5,81}, Bendik S Winsvold⁶⁻⁸, Priit Palta⁹, Tonu Esko^{2,10,11}, Tune H Pers^{2,11-13}, Kai-How Farh^{2,5,14}, Ester Cuenca-Leon^{1-3,15}, Mikko Muona^{9,16-18}, Nicholas A Furlotte¹⁹, Tobias Kurth^{20,21}, Andres Ingason²², George McMahon²³, Lannie Ligthart²⁴, Gisela M Terwindt²⁵, Mikko Kallela²⁶, Tobias M Freilinger^{27,28}, Caroline Ran²⁹, Scott G Gordon³⁰, Anine H Stam²⁵, Stacy Steinberg²², Guntram Borck³¹, Markku Koiranen³², Lydia Quaye³³, Hieab H H Adams^{34,35}, Terho Lehtimäki³⁶, Antti-Pekka Sarin⁹, Juho Wedenoja³⁷, David A Hinds¹⁹, Julie E Buring^{21,38}, Markus Schürks³⁹, Paul M Ridker^{21,38}, Maria Gudlaug Hrafnisdottir⁴⁰, Hreinn Stefansson²², Susan M Ring²³, Jouke-Jan Hottenga²⁴, Brenda W J H Penninx⁴¹, Markus Färkkilä²⁶, Ville Artto²⁶, Mari Kaunisto⁹, Salli Vepsäläinen²⁶, Rainer Malik²⁸, Andrew C Heath⁴², Pamela A F Madden⁴², Nicholas G Martin³⁰, Grant W Montgomery³⁰, Mitja I Kurki^{1-3,9,43}, Mart Kals¹⁰, Reedik Mägi¹⁰, Kalle Pärn¹⁰, Eija Hämmäläinen⁹, Hailiang Huang^{2,3,5}, Andrea E Byrnes^{2,3,5}, Lude Franke⁴⁴, Jie Huang⁴, Evie Stergiakouli²³, Phil H Lee¹⁻³, Cynthia Sandor⁴⁵, Caleb Webber⁴⁵, Zameel Cader^{46,47}, Bertram Muller-Myhsok^{48,76,93}, Stefan Schreiber⁴⁹, Thomas Meitinger^{50,51}, Johan G Eriksson^{52,53}, Veikko Salomaa⁵³, Kauko Heikkilä⁵⁴, Elizabeth Loehrer^{34,55}, Andre G Uitterlinden⁵⁶, Albert Hofman³⁴, Cornelia M van Duijn³⁴, Lynn Cherkas³³, Linda M Pedersen⁶, Audun Stubhaug^{57,58}, Christopher S Nielsen^{57,59}, Minna Männikkö³², Evelin Mihailov¹⁰, Lili Milani¹⁰, Hartmut Göbel⁶⁰, Ann-Louise Esserlind⁶¹, Anne Francke Christensen⁶¹, Thomas Folkmann Hansen⁶², Thomas Werge⁶³⁻⁶⁵, International Headache Genetics Consortium⁶⁶, Jaakko Kaprio^{9,37,67}, Arpo J Aromaa⁵³, Olli Raitakari^{68,69}, M Arfan Ikram^{34,35,70}, Tim Spector³³, Marjo-Riitta Järvelin^{32,71-73}, Andres Metspalu¹⁰, Christian Kubisch⁷⁴, David P Strachan⁷⁵, Michel D Ferrari²⁵, Andrea C Belin²⁹, Martin Dichgans^{28,76}, Maija Wessman^{9,16}, Arn M J M van den Maagdenberg^{25,77}, John-Anker Zwart⁶⁻⁸, Dorret I Boomsma²⁴, George Davey Smith²³, Kari Stefansson^{22,78}, Nicholas Eriksson¹⁹, Mark J Daly^{2,3,5}, Benjamin M Neale^{2,3,5,82}, Jes Olesen^{61,82}, Daniel I Chasman^{21,38,82}, Dale R Nyholt^{79,82} & Aarno Palotie^{1-5,9,80,82}

¹Psychiatric and Neurodevelopmental Genetics Unit, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts, USA. ²Medical and Population Genetics Program, Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA. ³Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA. ⁴Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton, UK. ⁵Analytic and Translational Genetics Unit, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts, USA. ⁶FORMI, Oslo University Hospital, Oslo, Norway. ⁷Department of Neurology, Oslo University Hospital, Oslo, Norway. ⁸Institute of Clinical Medicine, University of Oslo, Oslo, Norway. ⁹Institute for Molecular Medicine Finland (FIMM), University of Helsinki, Helsinki, Finland. ¹⁰Estonian Genome Center, University of Tartu, Tartu, Estonia. ¹¹Division of Endocrinology, Boston Children's Hospital, Boston, Massachusetts, USA. ¹²Department of Epidemiology Research, Statens Serum Institut, Copenhagen, Denmark. ¹³Novo Nordisk Foundation Center for Basic Metabolic Research, University of

Copenhagen, Copenhagen, Denmark. ¹⁴Illumina, San Diego, California, USA. ¹⁵Pediatric Neurology, Vall d'Hebron Research Institute, Barcelona, Spain. ¹⁶Folkhälsan Institute of Genetics, Helsinki, Finland. ¹⁷Neuroscience Center, University of Helsinki, Helsinki, Finland. ¹⁸Molecular Neurology Research Program, Research Programs Unit, University of Helsinki, Helsinki, Finland. ¹⁹23andMe, Inc., Mountain View, California, USA. ²⁰Institute of Public Health, Charité–Universitätsmedizin Berlin, Berlin, Germany. ²¹Division of Preventive Medicine, Brigham and Women's Hospital, Boston, Massachusetts, USA. ²²deCODE Genetics, Reykjavik, Iceland. ²³Medical Research Council (MRC) Integrative Epidemiology Unit, University of Bristol, Bristol, UK. ²⁴Department of Biological Psychology, Vrije Universiteit, Amsterdam, the Netherlands. ²⁵Department of Neurology, Leiden University Medical Centre, Leiden, the Netherlands. ²⁶Department of Neurology, Helsinki University Central Hospital, Helsinki, Finland. ²⁷Department of Neurology and Epileptology, Hertie-Institute for Clinical Brain Research, University of Tuebingen, Tuebingen, Germany. ²⁸Institute for Stroke and Dementia Research, Klinikum der Universität München, Ludwig-Maximilians-Universität München, Munich, Germany. ²⁹Department of Neuroscience, Karolinska Institutet, Stockholm, Sweden. ³⁰Department of Genetics and Computational Biology, QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia. ³¹Institute of Human Genetics, Ulm University, Ulm, Germany. ³²Center for Life Course Epidemiology and Systems Medicine, University of Oulu, Oulu, Finland. ³³Department of Twin Research and Genetic Epidemiology, King's College London, London, UK. ³⁴Department of Epidemiology, Erasmus University Medical Center, Rotterdam, the Netherlands. ³⁵Department of Radiology, Erasmus University Medical Center, Rotterdam, the Netherlands. ³⁶Department of Clinical Chemistry, Fimlab Laboratories, School of Medicine, University of Tampere, Tampere, Finland. ³⁷Department of Public Health, University of Helsinki, Helsinki, Finland. ³⁸Harvard Medical School, Boston, Massachusetts, USA. ³⁹Department of Neurology, University Duisburg–Essen, Essen, Germany. ⁴⁰Landspítali University Hospital, Reykjavik, Iceland. ⁴¹Department of Psychiatry, VU University Medical Centre, Amsterdam, the Netherlands. ⁴²Department of Psychiatry, Washington University School of Medicine, St. Louis, Missouri, USA. ⁴³Department of Neurosurgery, NeuroCenter, Kuopio University Hospital, Kuopio, Finland. ⁴⁴Department of Genetics, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands. ⁴⁵MRC Functional Genomics Unit, Department of Physiology, Anatomy & Genetics, Oxford University, Oxford, UK. ⁴⁶Nuffield Department of Clinical Neuroscience, University of Oxford, Oxford, UK. ⁴⁷Oxford Headache Centre, John Radcliffe Hospital, Oxford, UK. ⁴⁸Max Planck Institute of Psychiatry, Munich, Germany. ⁴⁹Institute of Clinical Molecular Biology, Christian Albrechts University, Kiel, Germany. ⁵⁰Institute of Human Genetics, Helmholtz Zentrum München, Neuherberg, Germany. ⁵¹Institute of Human Genetics, Technische Universität München, Munich, Germany. ⁵²Department of General Practice and Primary Health Care, University of Helsinki and Helsinki University Hospital, Helsinki, Finland. ⁵³National Institute for Health and Welfare, Helsinki, Finland. ⁵⁴Institute of Clinical Medicine, University of Helsinki, Helsinki, Finland. ⁵⁵Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA. ⁵⁶Department of Internal Medicine, Erasmus University Medical Center, Rotterdam, the Netherlands. ⁵⁷Department of Pain Management and Research, Oslo University Hospital, Oslo, Norway. ⁵⁸Medical Faculty, University of Oslo, Oslo, Norway. ⁵⁹Department of Ageing and Health, Norwegian Institute of Public Health, Oslo, Norway. ⁶⁰Kiel Pain and Headache Center, Kiel, Germany. ⁶¹Danish Headache Center, Department of Neurology, Rigshospitalet, Glostrup Hospital, University of Copenhagen, Copenhagen, Denmark. ⁶²Institute of Biological Psychiatry, Mental Health Center Sct. Hans, University of Copenhagen, Roskilde, Denmark. ⁶³Institute of Biological Psychiatry, MHC Sct. Hans, Mental Health Services Copenhagen, Copenhagen, Denmark. ⁶⁴Institute of Clinical Sciences, Faculty of Medicine and Health Sciences, University of Copenhagen, Copenhagen, Denmark. ⁶⁵iPSYCH—The Lundbeck Foundation Initiative for Integrative Psychiatric Research, Copenhagen, Denmark. ⁶⁶<http://www.headachegenetics.org/>. ⁶⁷Department of Health, National Institute for Health and Welfare, Helsinki, Finland. ⁶⁸Research Centre of Applied and Preventive Cardiovascular Medicine, University of Turku, Turku, Finland. ⁶⁹Department of Clinical Physiology and Nuclear Medicine, Turku University Hospital, Turku, Finland. ⁷⁰Department of Neurology, Erasmus University Medical Center, Rotterdam, the Netherlands. ⁷¹Department of Epidemiology and Biostatistics, MRC Health Protection Agency (HPE) Centre for Environment and Health, School of Public Health, Imperial College London, London, UK. ⁷²Biocenter Oulu, University of Oulu, Oulu, Finland. ⁷³Unit of Primary Care, Oulu University Hospital, Oulu, Finland. ⁷⁴Institute of Human Genetics, University Medical Center Hamburg-Eppendorf, Hamburg, Germany. ⁷⁵Population Health Research Institute, St George's, University of London, London, UK. ⁷⁶Munich Cluster for Systems Neurology (SyNergy), Munich, Germany. ⁷⁷Department of Human Genetics, Leiden University Medical Centre, Leiden, the Netherlands. ⁷⁸Faculty of Medicine, University of Iceland, Reykjavik, Iceland. ⁷⁹Statistical and Genomic Epidemiology Laboratory, Institute of Health and Biomedical Innovation, Queensland University of Technology, Kelvin

Grove, Queensland, Australia. ⁸⁰Department of Neurology, Massachusetts General Hospital, Boston, Massachusetts, USA.


⁸¹These authors contributed equally to this work. ⁸²These authors jointly supervised this work.

International Stroke Genetics Consortium (ISGC) Intracranial Aneurysm









Working Group:

Mark K. Bakker¹, Romain Bourcier^{2,3}, Robin G. Walters^{4,5}, Rainer Malik⁶, Martin Dichgans^{6,7,8}, Muralidharan Sargurupremraj^{9,10}, Turgut Tatlisumak¹¹, Stéphanie Debette^{9,10}, Gabriel J.E. Rinkel¹, Bradford B. Worrall¹², Joanna Pera¹³, Agnieszka Slowik¹³, Joseph P. Broderick¹⁴, David J. Werring¹⁵, Daniel Woo¹⁴, Philippe Bijlenga¹⁶, Yoichiro Kamatani¹⁷, Ynte M. Ruigrok¹

¹Department of Neurology and Neurosurgery, University Medical Center Utrecht Brain Center, Utrecht University, Utrecht, The Netherlands. ²Université de Nantes, CHU Nantes, INSERM, CNRS, l'institut du thorax, Nantes, France. ³CHU Nantes, Department of Neuroradiology, Nantes, France. ⁴Clinical Trial Service Unit and Epidemiological Studies Unit, Nuffield Department of Population Health, University of Oxford, Oxford, U.K. ⁵Medical Research Council Population Health Research Unit, University of Oxford, Oxford, U.K. ⁶Institute for Stroke and Dementia Research, University Hospital, Ludwig-Maximilians-University, Munich. ⁷Munich Cluster for Systems Neurology (SyNergy), Munich, Germany. ⁸Deutsches Zentrum für Neurodegenerative Erkrankungen (DZNE), Munich, Germany. ⁹INSERM U1219 Bordeaux Population Health Research Center, University of Bordeaux, Bordeaux, France. ¹⁰Department of Neurology, Institute for Neurodegenerative Disease, Bordeaux University Hospital, Bordeaux, France. ¹¹Department of Clinical Neuroscience at Institute of Neuroscience and Physiology, University of Gothenburg, Sweden. ¹²Departments of Neurology and Public Health Sciences, University of Virginia School of Medicine, Charlottesville, VA, USA. ¹³Department of Neurology, Faculty of Medicine, Jagiellonian University Medical College, ul. Botaniczna 3, 31-503, Krakow, Poland. ¹⁴University of Cincinnati College of Medicine, Cincinnati, OH, USA. ¹⁵Stroke Research Centre, University College London Queen Square Institute of Neurology, London, UK. ¹⁶Neurosurgery Division, Department of Clinical Neurosciences, Faculty of Medicine, Geneva University Hospitals, Geneva, Switzerland. ¹⁷Graduate School of Frontier Sciences, The University of Tokyo, Tokyo, Japan.

A

medRxiv preprint doi: <https://doi.org/10.1101/2023.04.29.23294404>; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. All rights reserved. No reuse allowed without permission.

**B**

C-index per model

