Immunology of asymptomatic parasitemia

1 Asymptomatic carriage of *Plasmodium falciparum* in children living in a hyperendemic area occurs

- 2 independently of IgG responses but is associated with a balanced inflammatory cytokine ratio.
- 3 Balotin Fogang^{1,2}, Matthieu Schoenhals¹, Franklin M. Maloba¹, Marie Florence Biabi^{1,3}, Estelle
- 4 Essangui¹, Christiane Donkeu^{1,2}, Glwadys Cheteug^{1,4}, Marie Kapen¹, Rodrigue Keumoe¹, Sylvie
- 5 Kemleu¹, Sandrine Nsango^{1,5}, Douglas H. Cornwall⁶, Carole Eboumbou^{1,5}, Ronald Perraut¹, Rosette
- 6 Megnekou², Tracey J. Lamb^{*6} and Lawrence S. Ayong^{*1}

7 Affiliations:

- 8 ¹ Molecular Parasitology Laboratory, Centre Pasteur du Cameroun, BP 1274 Yaounde, Cameroon
- 9 ² Department of Animal Biology and Physiology of the University of Yaoundé I, BP 812 Yaounde,
- 10 Cameroon
- ³ Department of Biochemistry, University of Douala, BP 24157 Douala, Cameroon
- ⁴ Department of Medical Laboratory Sciences, University of Buea, BP 63 Buea, Cameroon
- 13 ⁵Faculty of Medicine and Pharmaceutical Sciences, University of Douala, BP 2701 Douala,
- 14 Cameroon
- ⁶ Department of Pathology, University of Utah, 15 N Medical Drive, Salt Lake City 84112, USA
- 16 * Co-corresponding authors
- 17 Correspondence to: Tracey Lamb tracey.lamb@path.utah.edu Tel: +1 801 213 2097

18

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Immunology of asymptomatic parasitemia

19 Footnote Page

- 20 All authors declare no conflict of interest in this study. This work was supported by a capacity
- 21 building grant from the Institute Pasteur International Division to LSA. This information has
- 22 previously been presented at the American Association for Tropical Medicine and Hygiene Annual
- 23 Meeting in 2019. Correspondence to: Tracey Lamb tracey.lamb@path.utah.edu

Immunology of asymptomatic parasitemia

24 Abstract

Background: Asymptomatic carriage of infected red blood cells (iRBCs) can be prevalent in communities regardless of transmission patterns and can occur with infection of different *Plasmodium* species. Clinical immunity dampens the inflammatory responses leading to disease symptoms in malaria. The aim of this study was to define the immunological correlates of asymptomatic carriage of *P. falciparum* in a highly exposed population.

Methods: 142 asymptomatic *Plasmodium*-infected individuals greater than 2 years of age without fever (body temperature <37.5°C) were followed weekly for 10 weeks before being treated with artemisinin-based combination treatment (ACT). Plasma levels of 38 cytokines were measured at baseline by Luminex and the quantity and growth inhibitory activities of circulating parasite-reactive antibodies measured. The *Plasmodium* antigen tested included *P. falciparum* merozoite extract (ME) and schizont extract (SE), and the recombinant proteins Erythrocyte binding antigen 175 (EBA-175) and merozoite surface protein 1 (MSP-1₁₉).

37 **Results:** Median levels of IgG against *P. falciparum* EBA-175 and MSP-1₁₉ at baseline were significantly higher in those older than 20 years of age compared with the younger age group and 38 39 appeared to correlate with better parasite control. Amongst all participants there were no discernible 40 changes in IgG levels over time. Parasite density was higher in the younger age group and associated with IL-10, TNF- α and MCP-1 levels. A balanced IL-10:TNF- α ratio was associated with 41 42 asymptomatic malaria regardless of age, and balanced ratios of IL-10/TNF- α and IL-10/IFN- γ were 43 the only significant correlate of maintenance of asymptomatic malaria over the course of the study in 44 individuals 20 years of age and younger.

45 Conclusion: The above findings indicate that asymptomatic carriage of *P. falciparum* in children 46 living in a hyperendemic area occurs independently of IgG but is associated with a balanced 47 inflammatory cytokine ratio.

Immunology of asymptomatic parasitemia

Key words: Asymptomatic malaria, *Plasmodium falciparum*, IgG and cytokine levels, TNF-α, IL10

50 Introduction

51 Malaria is still a significant global health problem with an estimated 249 million cases and 608 000 deaths in 2022 [1]. Although mortality rates from malaria have decreased considerably since the turn 52 53 of the century [2] this has been paralleled by high increases in proportions of asymptomatic infections, known to contribute significantly to persistent malaria transmission in most endemic countries [3.4]. 54 55 Asymptomatic infections can persist for weeks to years [5] and untreated *P. falciparum* infections 56 have been shown persist on average for 6 months, with fluctuations in parasite density over the course 57 of an infection [6,7]. The maintenance of chronic asymptomatic malaria will depend on the host 58 immune response to the blood stage of *Plasmodium* [3]. Anti-parasite immunity directly suppress 59 parasite growth resulting in low infection densities (below the pyrogenic threshold [3,8]) whilst anti-60 disease immunity modulates inflammation-mediated clinical manifestations of the infection.

61 Older children and adults have more cumulative exposure to parasite variants circulating within the 62 community, developing adaptive responses which mediate both anti-parasite and anti-disease 63 immunity [9,10]. However, asymptomatic infections have been documented in younger children < 5 64 years who are too young to have developed robust adaptive immunity [11]. This suggests that 65 asymptomatic malaria may be differentially controlled by innate and adaptive immune responses 66 depending on age.

Anti-parasite immunity which is acquired with age or exposure [12,13], is mediated mainly by antibodies which control parasite replication and mediate clearance of *Plasmodium*-infected red blood cells (iRBCs) [14–16]. Suppression of inflammation associated with iRBCs depends on the balance between anti- and pro-inflammatory cytokine responses [17,18] and it has been hypothesized that asymptomatic infections are associated with a shift in host immune responses towards an antiinflammatory profile after multiple exposures to malaria [19]. This could occur via a decrease in

Immunology of asymptomatic parasitemia

73 levels of pro-inflammatory immune mediators such as IFN- γ , TNF- α , or an increase in anti-

74 inflammatory mediators such as IL-10 or TGF- β [18,20–22].

75 The relative contributions of anti-parasite and anti-disease immunity to the maintenance of persistent asymptomatic *Plasmodium* infections in younger individuals and older individuals remain unclear. 76 77 To determine the immunological associations of chronic asymptomatic malaria in younger (≤ 20 78 years) and older (>20 years) individuals we have undertaken a longitudinal study of highly exposed 79 individuals of all ages with asymptomatic malaria over a 3-month period where we monitored the stability of IgG and cytokine responses. We also examined these responses upon clearance of 80 81 parasites to ascertain which responses depended on the presence of circulating parasites. Our study 82 demonstrates a lack of correlation of iRBC density and IgG responses in individuals ≤ 20 years old, 83 with an association of parasite carriage with IL-10 and an association of asymptomatic status with a 84 balanced ratio of IL-10:TNF- α and IL-10: IFN- γ .

85

86 Material and methods

87 Ethics statement

The study protocol was approved by the Cameroon National Ethics Committee for Human Health
Research (Ethical clearance N°: 2018/09/1104/CE/CNERSH/SP) and the Ministry of Public Health.
Written informed consent was obtained from each participant or parent/guardian of minor participants
(< 19 years old) prior to enrolment into the study. Assent was obtained from all children older than
12 years of age.

93 Study area and population

The study population consisted of individuals of both genders aged at least two years of age and permanently living in one of the five selected villages of the Esse health district, Mefou-et-Afamba Division, Centre Region of Cameroon where malaria transmission is perennial, stable and hyperendemic with *P. falciparum* as the predominant *Plasmodium* species. The study area has been

Immunology of asymptomatic parasitemia

described elsewhere [23–25]. The inclusion criteria were: i) obtain the consent of each older
participant or parent or legal guardian for the children, ii) living permanently in the study area, iii)
not pregnant.

101 Study design and participants

102 This was a longitudinal household-based study that involved 353 individuals. Participants were 103 considered asymptomatic for malaria if positive for *P. falciparum* using multiplex nested PCR in the 104 absence of fever at least forty-eight hours prior to enrolment. Being febrile was assessed based on 105 axillary temperature \geq 37.5 °C or based on the history of fever as independently confirmed by another 106 household member. All symptomatic individuals (fever + rapid diagnostic test (RDT) +) were treated 107 with ACT-based drugs following recommended national guidelines. Participants with asymptomatic 108 carriage of *Plasmodium* parasites were monitored weekly until they either developed a fever, or until 109 the end of the follow up period at week 16 post-enrolment. Participant follow-up comprised of weekly interviews regarding the occurrence of fever during the past week, measurement of axillary 110 111 temperatures on each visit, and laboratory testing for malaria at weeks 1, 4, 11 and 16 (PCR and 112 microscopy). All symptomatic cases occurring during the follow-up period were treated with ACT as 113 recommended and excluded from further follow-up. At week 11 post-enrolment all participants with 114 persisting asymptomatic parasitemia were treated as recommended, and monitored for five additional 115 weeks to assess the effect of ACT-based antimalarial treatment on selected host immune responses. Participants with persistent asymptomatic parasitemia 10 weeks following enrolment were 116 117 considered as persistent asymptomatic malaria whereas those who developed malaria-associated symptoms before week 11 post-enrolment were considered as developing uncomplicated malaria. 118

119 Blood sample collection, malaria diagnosis and parasite genotyping

Approximately 3 ml of venous blood was collected in EDTA-coated tubes and used for laboratory testing for malaria and blood hemoglobin levels as well as to obtain plasma for immunological analysis. Plasma was separated from the red blood cell (RBC) pellet by centrifugation (3500 RPM,

Immunology of asymptomatic parasitemia

123 5 min) and stored at -80 °C until use. Blood hemoglobin levels were assessed using a portable Mission

124 Hb haemoglobinometer (ACON Laboratories, USA).

Parasite density (parasites/ μ l) was determined by microscopic examination of Giemsa-stained thick blood smears. Parasite density was determined based on the number of parasites per at least 200 WBC count on a thick film, assuming a total white blood cell count of 8000 cells/ μ L of whole blood. A blood slide was considered positive when a concordant result was obtained by at least two independent microscopists and when a discrepancy occurred, a third microscopist was used for confirmation. A slide was considered negative if no parasites were detected after a count of 500 white blood cells.

Genomic DNA was extracted from dried blood spots using the Chelex method [26], and used to screen for *Plasmodium* infections by multiplex nested PCR targeting the 18S small subunit rRNA (ssrRNA) genes as previously described [24]. Genotyping of *msp2* gene was performed by nested PCR on *P. falciparum* PCR positive samples as previously described [24,27].

136 **Case definitions for the study**

Participants were defined as asymptomatic if positive for malaria parasites by PCR in the absence of 137 fever at least 48 hours prior to enrolment. Febrile malaria was determined based on RDT, light 138 139 microscopy or PCR positivity for malaria parasites and axillary temperature $\geq 37.5^{\circ}$ C or a history of 140 fever in the past 48 hours. Those developing uncomplicated malaria were defined as asymptomatic 141 individuals who had developed fever associated with *Plasmodium* infection during the 10-week 142 infection follow-up period, whereas the persistent asymptomatic malaria classification represented 143 the asymptomatic individuals who remained asymptomatic until anti-malarial treatment at the end of 144 the infection follow-up period.

Immunology of asymptomatic parasitemia

145 Anti-Plasmodium antibody levels and avidity assessment

Plasma IgG antibody levels and avidity were determined by indirect ELISA method using *P*. *falciparum* Merozoite extract (ME), schizont extract (SE) or the recombinant proteins erythrocyte
binding antigen-175 (EBA-175) and Merozoite Surface Protein-1₁₉ (MSP1₁₉) as coating antigens.
ME and SE were isolated from laboratory maintained *P. falciparum* 3D7 parasites as described
previously [28], and used for protein extraction by freeze-thaw fractionation [23,29]. The
recombinant protein EBA-175 (F3 region) was obtained from BEI Resources (MRA-1162), whereas *P. falciparum* MSP1₁₉ was produced using a baculovirus-insect cell expression system [30].

153

154 96-well microplates (F96 CERT-Maxisorp) were coated with each antigen (2 mg/ml of ME and SE, 155 0.20 mg/ml of EBA-175 and 0.5 mg/ml of MSP-1₁₉) in 0.1 M bicarbonate buffer (for ME, SE and 156 EBA-175) or PBS (for MSP-1₁₉) and incubated overnight at 4°C. Plasma samples were diluted 157 (1/250) in PBS-T/1%BSA and used to measure anti-*Plasmodium* IgG as previously described [23,31]. 158 Plasma from non-exposed European blood donors (10 samples) were used as naive controls and a pool of IgG fractions from adults living in high malaria endemic areas was used as positive controls. 159 Results were expressed as optical-density ratios (OD sample/mean OD naive control). For avidity 160 161 testing, 100 µL of 8M urea in wash buffer (treated wells) or 100 µL of wash buffer (untreated wells) 162 were added for 15 minutes at room temperature to the respective wells following primary antibody 163 binding as previously described [32]. Avidity indices (AI) were calculated as proportion of bound 164 antibodies following urea treatment (AI = [mean OD of treated wells /mean OD of untreated wells] 165 \times 100). Samples with OD ratios > 2, corresponding to mean OD naive control + 3SD were considered 166 as seropositive. In addition, only seropositive samples were used for avidity testing.

167

168 Growth inhibition assay (GIA)

169 GIA was performed using plasma from endemic and non-endemic individuals as described by

170 Duncan et al. [33]. Briefly, plasma samples were heat inactivated at 56 ° C for 20 min and a 1:20

Immunology of asymptomatic parasitemia

171 ratio of 50% RBC was added to each sample and incubated at room temperature for 1 h to remove

the RBC-reactive antibody. After spinning for 2 min at 13,000 x g to pellet the RBCs, the

- 173 supernatant was transferred to a 96-well V-bottom plate. Synchronized late trophozoite/early
- schizont (30-40 h) stages of 3D7 *P. falciparum* at 0.5% parasitemia were incubated with plasma
- 175 (1:10 ratio) for 48 h at 37 °C. The GIA harvest (parasite growth) was carried out by SYBR green-I
- 176 DNA quantification assay [34]. Parasite growth inhibition (GI) activities of plasma antibodies were
- presented as: GI indices (%) = $100 \times [1 ((OD \text{ sample OD RBC}) / (OD \text{ parasitized RBC OD})]$
- 178 RBC))].

179 Determination of plasma cytokine levels

180 Plasma concentrations of 38 immune mediators including interleukin (IL)-1a, IL-1B, IL-1RA, IL-2 181 IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-12p40, IL-p70, IL-13, IL-15, IL-17A, interferon (IFN)- α 2, IFN- γ , tumor necrosis factor (TNF)- α , lymphotoxin- α (LT- α), IFN- γ induced protein (IP-182 183 10), monocyte chemoattractant protein (MCP)-1, MCP-3, macrophage inflammatory protein (MIP)-184 1α, MIP-1β, Eotaxin, Fractalkine, human growth-regulated oncogene (GRO), Macrophage-derived chemokine (MDC), granulocyte-macrophage colony-stimulating factor (GM-CSF), granulocyte 185 186 colony-stimulating factor (G-CSF), sCD40L, FMS-like tyrosine kinase 3 ligand (Flt-3L), epidermal 187 growth factor (EGF), fibroblast growth factor 2 (FGF-2), transforming growth factor alpha (TGF- α) and vascular endothelial growth factor (VEGF) were determined by Luminex-based method using 188 189 the Human Cytokine/Chemokine Magnetic Bead Panel MILLIPLEX MAP Kit (Cat. No. 190 HCYTMAG-60K-PX38, EMD Millipore Corporation) according to the manufacturer's instructions. 191 Briefly, 25 µL of premixed magnetic capture beads were incubated with equal volumes of plasma or 192 diluted standard in a 96-well microplate placed on a horizontal plate shaker for 2 hours. After washing 193 using a magnetic plate, plates were incubated with 25 µL/well of biotinylated detection antibodies 194 for 1 hour at room temperature followed by 25 µL of streptavidin-PE in each well for 30 minutes at 195 room temperature. Median fluorescence intensity was read on a Luminex MAGPIX Analyser (XMAP 196 Technology) equipped with xPONENT software version 4.2. The relative concentration of each

Immunology of asymptomatic parasitemia

197 cytokine (pg/ml) in each sample was determined based on the automatically generated standard curve
198 for each analyte. Analyte amounts less than the limit of detection for each cytokine were attributed
199 the value of detection limit.

200 Statistical analysis

Statistical analyses were performed using RStudio version 4.1.1 or version 4.2.2 [35] and Graphpad 201 202 Prism version 8.4.3 (Graphpad software, San Diego, California USA). Differences in antibody and 203 cytokine distribution between two or more groups were determined using Wilcoxon signed rank test 204 and Kruskal-Wallis test, respectively. Spearman rank correlation tests or linear regression were used to determine the correlation between quantitative variables. Differences in categorical 205 206 variables were determined by chi-square tests. P-values < 0.05 were considered statistically 207 significant. Factor analysis of mixed data (FAMD) with FactoMineR and factoextra packages was 208 used to assess the association between cytokines levels and age and circulating parasite loads. The complexHeatmap and ggpubr packages were used to draw the heatmaps and the line plots, 209 210 respectively. Given that the aim of this study was to determine the host immunological 211 determinants in the maintenance of the asymptomatic infections, among the 353 individuals 212 included, only participants with complete information on the study outcome during the longitudinal 213 follow-up period were included in the analysis (n= 211). Participants who missed at least one of the 214 follow-up points (the predominant reason being farming far away from the enrolment site) were not included in our study. Three samples (2 symptomatic and 1 asymptomatic) were excluded because 215 216 no cytokines were detected in the plasma. For analysis of cytokines at the initial time point of our 217 study with respect to correlation with persistent asymptomatic malaria, we employed a mixed linear 218 model using the LME4 package [36]. Cytokine level was used as the response with conversion 219 status as the fixed-effect of the model. We used sex and age as random effects in the model and ran 220 the model separately for children ≤ 20 years old, adults > 20 years old and all participants, regardless of age. To obtain p-value estimates from the t-values we used the lmerTest package [37]. P values 221 were corrected for multiple comparisons using the Sattherthwaite correction. For analysis, parasite 222

Immunology of asymptomatic parasitemia

densities were separated into three groups as follows, 1. submicroscopic parasitemia, 2. low levels

- of microscopic parasitemia (parasitemia between 37.2 parasites/µl and 413.1 parasites/µl) and 3.
- high levels of parasitemia (parasitemia> 413.1 parasites/µl, representing the median parasitemia in
- the study population).
- 227

228 **Results**

229 Asymptomatic malaria prevalence in the study area

We carried out immunological analysis of a subset of 211 individuals from a previously reported 230 231 study of 353 individuals screened for malaria using PCR, where 328 (92.9%) were positive for 232 Plasmodium falciparum infection, of which 266 (81.1%) were asymptomatic and 62 (18.9%) were symptomatic [24]. This subset had complete sampling from the follow-up study, and included 142 233 234 individuals with asymptomatic infection, 61 symptomatic individuals and 8 non-infected individuals. At baseline when these individuals were first sampled, parasite densities were highly variable 235 236 between the symptomatic and asymptomatic groups, and no significant difference was observed in 237 the median parasite densities of the two groups at the initial time point of the study (Table 1). Age, 238 sex, and hemoglobin levels were also not different across the study groups at this initial time point 239 (Table 1).

Immunology of asymptomatic parasitemia

v ariables	Non de infe (n	ection = 8)	Asym] (n=	= 142)	=u) buňc	61)	r va	ue
Age groups	≤ 20	> 20	≤ 20	> 20	≤ 20	> 20	≤ 20	
n	л	З	94	48	42	19		
Sex ratio (M:F)	0.67 (2:3)	50 (1:2)	0.82 (42:51)	41.2 (14:34)	1.1 (22:20)	58.3 (7:12)	0.7114	0
Median age, years	14 (4-20)	68 (50-68)	9 (2–19)	50 (22-86)	9 (2-20)	28 (21-64)	0.2193	0
(range)								
Median Hb, g/dl	11.8	112.5	12 (4.9-14.3)	13.1(10.9-16)	11.1 (6.8-	12.8 (9.4-	0.1866	0.
(range)	(10.7-	(11.9-14)			14)	16.8)		
	12.7)							
% of submicroscopic	ı		22.3 (21)	60.4 (29)	9.5 (4)	57.9 (11)	0.0746	0.8
% (n)								
Geometric mean	ı		575.7 (37.4-	113.5 (38.3-	764.1 (38.1-	183.3(37.2-	0.5139	0.0
parasitemia,			68190.5)	660.2)	240457.1)	5058.8)		
narasites/III (range)								

lividuals with parasite densities at the detection limit of light microscopy. The sex ratio between groups were compared using chi-square test. Hb: hemoglobin. M: male. female.	licoxon signed-rank test and Kruskal-Wallis test when comparing two or three groups, respectively. The median parasitemia presented in this table is only for the infected	th complete data from the follow-up study to 11 weeks and does not include baseline data from individuals who dropped out of the study. P-value was determined using	e data presented in this table is from 211 individuals from our study at baseline. The subset of 142 asymptomatic individuals included in this table were those individuals
--	--	--	---

Immunology of asymptomatic parasitemia

245

Anti-parasite antibodies were inversely correlated with parasitemia but were not associated with asymptomatic parasite carriage

248 Consistent with the hypothesis that host immunity increases with repeated exposure to malaria 249 parasites, baseline parasite densities above the detection limit of microscopy decreased significantly 250 with age (r = -0.49, P< 0.0001) (Figure 1A) but only in individuals aged 20 years and younger (r = -0.39, P< 0.0001) (Figure 1B). Of the 142 participants who completed the follow-up study as 251 252 designed, 78 (54.9%) remained asymptomatic during the 10-weeks follow-up period, whereas 63 253 (44.4%) developed malaria-associated fevers and 1 (0.7\%) resolved the infection in the absence of 254 any known antimalarial treatment. Based on the parasitemia distribution of those with microscopic 255 *Plasmodium* infections at baseline the study population was divided into two groups aged ≤ 20 years 256 and > 20 years old for further analyses. The proportion of submicroscopic infections was significantly 257 higher (P<0.0001) in the older age group (age >20 years) at the baseline time point than in the younger 258 age group (Figure 1C). At the baseline time point, median IgG levels against *P. falciparum* merozoite 259 (ME) and schizont (SE) crude extracts were significantly higher in the >20-year age group than in 260 the younger age group (P < 0.0001) (Figure 2A). This was mirrored by IgG reactive to *P. falciparum* 261 recombinant proteins EBA-175 (P=0.0250) and MSP-1₁₉ (P<0.0001) (Figure 2A). Additionally, In 262 the younger age group IgG levels against merozoite (r= 0.22, P= 0.0077) and schizont (r= 0.29, P=263 0.0004) crude extracts as well as against MSP-1₁₉ (r=0.15, P=0.0764) but not EBA-175 (r=0.01, P=264 0.8525) were positively correlated with age at the baseline time point (Figure 2B). A negative 265 correlation was observed between IgG levels to P. falciparum merozoite and schizont extracts at 266 baseline and parasite density measured at the same time point (Supplementary Figure 1). Indeed, 267 the multilevel linear regression model adjusted for village and household levels, using IgG levels at 268 the baseline and age as predictors, showed that only age was the independent predictor of *Plasmodium* 269 parasite densities in the study population (coef= -0.02, P= 0.0004 for age; coef= 0.01, P= 0.9170 for

Immunology of asymptomatic parasitemia

270 ME; coef=-0.19, P=0.1529 for SE; coef=-0.02, P=0.4746 for EBA-175; coef=0.02, P=0.5362 for

272 Median IgG levels were not significantly different between participants with sub-microscopic 273 parasitemia and those with microscopic parasites at baseline (Supplementary Figure 2A), and no 274 significant difference was observed in the *in vitro* growth inhibition potential of inactivated plasma 275 from submicroscopic parasite carriers and that of microscopic carriers (Supplementary Figure 2B). 276 Both IgG antibody avidity (Supplementary Figure 3A&B) and growth inhibition potentials 277 (Supplementary Figure 2B) were similar in the two groups (P> 0.05). Together, these findings 278 suggest that anti-parasite antibodies may control parasite density levels by mechanisms other than 279 direct inhibition of parasite growth in vivo.

²⁷¹ MSP-1₁₉).

Immunology of asymptomatic parasitemia



283 Figure 1. Older individuals (>20 years) had a better control of parasite densities. A) Scatterplot showing 284 a correlation between baseline parasite densities and age in all the infected individuals with parasitemia at the 285 detection limit of light microscopy (n=138). Spearman rank correlation is presented as the best fit line and the 286 coefficient (r), as well as the p-value (p), are shown. The red straight line was used to divide the study 287 population into two groups aged ≤ 20 years and > 20 years old for further analyses. **B**) Scatterplot of correlation 288 between baseline parasitemia detected by microscopy and age in different age groups (< 20 years and > 20) 289 years) at the different time points of our study. Spearman rank correlation is presented as the best fit line and the coefficient (r), as well as the p-value (p), are shown for each age group. C) The pie chart shows the 290 291 proportion of sub-microscopic infections (PCR+/microscopy-) versus microscopic (microscopy+) infections 292 in each age group. PCR: Polymerase Chain Reaction.

293

294

295

296





299 Figure 2. Individuals >20 years old had more anti-parasite IgG at baseline. A) Data presented in the 300 scatterplot represent baseline IgG antibody levels against different P. falciparum antigens (ME, SE, EBA-175 301 and MSP-1₁₉) within age groups. Median levels between age groups were compared by Wilcoxon signed-rank 302 test and the p-values are shown on the graph. **B**) The scatterplots on B show the correlation between IgG levels 303 against different *P. falciparum* antigens (ME, SE, EBA-175 and MSP-119) and age in different age groups (≤ 304 20 years and > 20 years). Spearman rank correlation is presented as the best fit line and the coefficient (r), as 305 well as the p-values (p), are shown for each age group. SE: Schizont extract. ME: Merozoite extract. EBA-306 175: Erythrocyte binding antigen-175. MSP-1₁₉: Merozoite surface protein-1₁₉. OD: optical density.

307

Immunology of asymptomatic parasitemia

308 Similarly, no significant difference was observed in malarial IgG antibody levels and avidity between 309 participants with symptomatic or asymptomatic parasitemia (Supplementary Figure 4A), and no 310 significant difference was seen in the blood parasitemia at the detection limit of microscopy 311 (Supplemental Figure 4B; Table 1). However, the *in vitro* growth inhibition potential of plasma from 312 asymptomatic malaria was greater than that of symptomatic individuals (P= 0.0245) despite no significant differences in the parasitemia amongst those with parasite levels that were detectable by 313 314 microscopy (Supplementary Figure 4B). This may be a reflection of higher percentage of 315 asymptomatic children with submicroscopic parasitemias (77.7%) compared to symptomatic children 316 (90.5%) (Supplementary Figure 4C). Taken together, these data suggest that humoral responses other than IgG responses to the tested parasite antigens contribute to the maintenance of 317 318 asymptomatic parasitemia in the study area.

319 Persistence of fluctuating asymptomatic parasitemia occurred in highly exposed individuals 320 without alteration in IgG levels

321 Of the 142 participants who completed the follow-up study as designed, 72 (50.7%) remained asymptomatic during the 10-weeks follow-up period, whereas 69 (48.6%) developed malaria-322 323 associated fevers and 1 (0.7%) resolved the infection in the absence of any known antimalarial treatment. Surprisingly, the propensity to convert was not associated with age (Figure 3A). Of the 72 324 325 participants who maintained chronic asymptomatic P. falciparum carriage, the proportion of 326 submicroscopic infections remained similar in both age groups with most participants becoming microscopy positive on week 10 regardless of age (Figure 3B). Parasite densities fluctuated widely 327 328 over the follow-up period from ultra-low parasitemia (sub-microscopic) to high parasitemia (Figure 329 **3B**), possibly due to variation in the infecting clones from new infections as this is a high transmission area (Figure 3C). However, no changes in IgG levels (Figure 3D) against all antigens tested were 330 observed during persistent asymptomatic infections indicating that persistent asymptomatic 331 332 parasitemia does not affect anti-*Plasmodium* IgG levels over this short 3-month time period. These

Immunology of asymptomatic parasitemia

- 333 data suggests that asymptomatic carriage of *P. falciparum* occurs independently of circulating
- 334 parasite numbers, even within highly-exposed individuals.



335

336 Figure 3. Asymptomatic malaria persisted in some individuals despite changes in the circulating 337 parasite densities and population over time, but did not affect the antibody levels. A) The pie charts 338 show the proportion of individuals with initial asymptomatic infection (cross-sectional) who resolved the 339 infection (clearers), developed malaria-associated fever (uncomplicated malaria) or remained asymptomatic 340 (persistent asymptomatic) over the 10-week follow-up period, according to age groups. B) Panel B displays 341 the proportion of sub-microscopic/microscopic parasitemia in persistent asymptomatically infected 342 individuals at three different time points (weeks 1, 4 and 11) according to age groups. Individuals who were 343 RDT positive but without detectable micoroscopic parasitemia was detected are shown in pink. Of those who

Immunology of asymptomatic parasitemia

had microscopic parasitemia, the line plots parasitemia across the 3 time points (C). To generate the line

- 345 plots, the value 10 was assigned to submicroscopic parasitemia. The P values on the line plots refer to the
- 346 comparison of the median parasitemia between two time points using the Wilcoxon test. **D**) Pie charts show
- 347 the proportion of msp2 alleles (representing the clones) determined by nested PCR in individuals with
- 348 persistent asymptomatic infection per week. Numbers represent the band sizes obtained. Blue slices
- 349 represent the alleles only present at week 1. Red slices represent the alleles which appear at week4. Green
- 350 slices represent the allele only present at week 11. Grey slices represent the alleles that were present at all
- 351 three sampling time points (bands of 375, 400, 475, 500, 520, 534, 540, 600, 620, 633. 680, 700, 800 bp). **E**)
- 352 Data presented in the line plots represent the profile of IgG anti-*Plasmodium* antibody levels in each chronic
- asymptomatically infected individuals at three different time points (week 1, 4 and 11) in the two age groups.
- 354 SE: Schizont extract. ME: Merozoite extract. EBA-175: Erythrocyte binding antigen-175. MSP-1₁₉:
- 355 Merozoite surface protein-1₁₉. OD: optical density.

Immunology of asymptomatic parasitemia

357 Parasite density was associated with inflammatory responses amongst younger individuals

358 with asymptomatic malaria

359	Given that <i>Plasmodium</i> parasites are associated with induction of inflammatory responses we
360	measured the cytokine profiles of the 72 participants who had persistent asymptomatic carriage of
361	P. falciparum across the course of our study. There was no particular cytokine signature that
362	associated with asymptomatic carriage of <i>P. falciparum</i> in our study participants over the 11-week
363	follow-up period (Figure 4A). Using a factor analysis of mixed data (FAMD) on cytokine data
364	along with age and circulating parasite density, we found that levels of IL-10, TNF- α , the ratio of
365	IL-10:IFN- γ and MCP-1, along with age and parasite density explained most of the variation in the
366	peripheral cytokine response in asymptomatic individuals (Figure 4B). Multivariable analysis using
367	age and parasitemia as predictors showed that age independently predicted the levels of TNF- α
368	(coef= -0.31, P< 0.0001) peaking in younger study participants (Figure 4C). This is likely a
369	reflection of circulating parasite densities which were higher in the younger group (Figure 1) and
370	were also independently predictive of TNF- α levels (coef= 0.09, P< 0.0001) (Figure 4D).
371	Circulating parasite density also independently predicted the levels of IL-10 (coef= 0.21, P<
372	0.0001) and MCP-1 (coef= 0.06 , P= 0.01) with a significant increase in those who had high levels
373	of parasitemia (parasitemia> 413.1 parasites/µl) relative to those who had low levels of microscopic
374	parasitemia (parasitemia between 37.2 parasites/ μ l and 413.1 parasites/ μ l) (Figure 4D). This
375	indicates that during asymptomatic carriage of <i>P. falciparum</i> circulating parasites simultaneously
376	induce anti-inflammatory cytokine production to counteract the inflammatory effects of TNF- α . To
377	determine which cytokines were dependent on parasite carriage we treated participants at week 11
378	of the study with ACT. This resulted in a significant decline in the IL-10 levels, but not TNF- α or
379	MCP-1 levels, in the younger age group (Figure 5). Thus, the levels of IL-10 measured is a
380	reflection of parasite carriage in the younger age group.

Immunology of asymptomatic parasitemia



381

Figure 4. Cytokine responses were blunted with increased age and correlated with parasite densities at baseline. A) Hierarchical clustering of the standardized least-square means of 38 differentially expressed cytokines/chemokines/growth factors across the three sampling time points in participants with persistent asymptomatic infection (n= 72). B) Principal component of cytokines, age and parasitemia from FAMD (left graph) analysis. The right graph of B shows the contribution of each variable on the principal component 2 (dimension 2). **C** and **D**) box plots representing interquartile ranges (25th, median and 75th percentile, whisker

Immunology of asymptomatic parasitemia

- 388 percentile (1st and 99th)) of the levels (Log10 pg/ml) of some main regulatory cytokines measured by Luminex
- 389 according to age and parasite level groups at the baseline, respectively. Median levels of cytokines between
- 390 two or more groups were compared by Wilcoxon rank-sum test and Kruskal-Wallis test, respectively. In Panel
- 391 A, pairwise comparisons on neighboring groups were performed only when multiple comparisons were
- 392 significant at p< 0.05. Parasite densities were groups in three groups as follows: submicroscopic parasitemia,
- 393 low levels of microscopic parasitemia (parasitemia ≤ 413.1 parasites/µl, representing the median parasitemia
- in the study population) and high levels of parasitemia (parasitemia> 413.1 parasites/µl). FAMD: Factor
- 395 analysis of mixed data. *p< 0.05; **p< 0.001, ***p< 0.0001, ***p< 0.0001.
- 396

Immunology of asymptomatic parasitemia

398

399



400 Figure 5. IL-10 and TNF- α levels were a reflection of parasite carriage, and the balance of inflammatory 401 cytokines at baseline was associated with the maintenance of asymptomatic infections in the younger 402 age, ≤ 20 years. A) Box plots representing interquartile ranges (25th, median and 75th percentile, whisker 403 percentile (1st and 99th)) of cytokine levels (Log_{10} pg/ml) during persistent asymptomatic infections (week 1, 404 4 and 11) and after anti-malarial treatment (week 16). B) Box plots representing interquartile ranges (25th, 405 median and 75th percentile, whisker percentile (1st and 99th)) of cytokine levels (Log₁₀ pg/ml) at the baseline 406 between in those who developed uncomplicated malaria (UM; grey) and those who went on to display 407 persistent asymptomatic malaria throughout the course of the study (PAM; blue). Analyses were performed 408 including values from individuals with submicroscopic and microscopic parasitemia. Median cytokines levels 409 between two or more groups were compared by Wilcoxon rank-sum test and Kruskal-Wallis test, respectively 410 and significant between group differences indicated by the P value on the plot. In Panel A, pairwise 411 comparisons on neighboring groups were performed only when multiple comparisons between the 4 groups 412 were significant and are shown by the grey brackets. p < 0.05 and p < 0.0001.

Immunology of asymptomatic parasitemia

413 Maintenance of asymptomatic carriage of *P. falciparum* is associated with a more balanced

414 inflammatory cytokine ratio

415 There was no particular cytokine signature of asymptomatic individuals relative to symptomatic 416 individuals in our study (Figure 6A). However, given that anti-inflammatory cytokines may be 417 important in down-regulating the pathogenic inflammatory response in malaria, we tested the 418 hypothesis that asymptomatic infection would be associated with higher levels of IL-10. To our 419 surprise, at the initial time point of our study there was a non-significant trend towards higher levels of IL-10 in the symptomatic compared to asymptomatic participants (Figure 6B). This may be a 420 reflection of higher inflammation levels in symptomatic individuals as other pro-inflammatory 421 markers (IL-6 and MCP-1) were significantly higher in individuals with symptomatic malaria 422 423 compared to those with asymptomatic *P. falciparum* infections in the younger age group (Figure 6B). 424 There was an IL-10-skewed ratio of IL-10:TNF- α and IL-10:IFN- γ in the symptomatic group compared to the asymptomatic group in both age groups (Figure 6C). This is in agreement with 425 426 previously published data in individuals with uncomplicated malaria and the IL-10 skewed ratio of IL-10 to TNF- α that can be measured in those with severe symptoms of malaria 427 [38,39]. 428 Multivariable logistic regression analysis showed that the IL-10:TNF-α ratio and IL-1RA levels were 429 independently associated with developing symptoms during *P. falciparum* infection in this study 430 (Supplementary Table S1).

431





Figure 6. IL-10 at baseline was associated with clinical malaria in the younger age group (\leq 20 years). A) Hierarchical clustering (heatmap) of the standardized least-square means of 38 differentially expressed cytokines/chemokines/growth factors by infection status at the baseline of this study. **B** and **C** box plots represent interquartile ranges (25th, median and 75th percentile, whisker percentile (1st and 99th)) of the baseline cytokine levels (Log₁₀ pg/ml) measured by Luminex according to infection status. Analyses were performed including both individuals submicroscopic and microscopic parasitemia. Median levels of cytokines between two or three groups were compared by Wilcoxon rank-sum test and Kruskal-Wallis test, respectively.

Immunology of asymptomatic parasitemia

In Panel B and C, pairwise comparisons were performed only when multiple comparisons were significant. P
values refer to the Kruskal-Wallis test comparison between the three groups. NI: Non-infected. AS:
Asymptomatic. SY: Symptomatic. *p< 0.05; **p< 0.001; ***p< 0.0001.

444

445 To further test the hypothesis that asymptomatic carriage of *P. falciparum* may be related to IL-10 production, we compared the cytokine ratios in the younger group who converted to symptomatic 446 447 malaria throughout our study with those who remained asymptomatic. At the initial time point of the 448 study there was a trend towards increased IL-10 levels in those who subsequently went on to develop uncomplicated malaria compared to those who experienced persistent asymptomatic malaria in both 449 450 age groups. In the younger age group IL-10/pro-inflammatory cytokine ratios were significantly 451 higher and IL-10-skewed (P= 0.0207 and P= 0.0111 for IL-10:TNF- α and IL-10:IFN- γ ratios, 452 respectively) in the those developing uncomplicated malaria compared to those with persistent 453 asymptomatic malaria in younger age group (Figure 5B). In this group those with persistent 454 asymptomatic malaria had a ratios of 1.001±0.1355 for IL-10:TNF-a and 1.016±0.1794 for IL-455 10:IFN- γ) displaying an even balance of IL-10: proinflammatory cytokine. On the other hand adults did not have IL-10 skewed ratios with TNF- α or IFN- γ suggesting that a different or additional 456 457 mechanism may facilitate the persistence of asymptomatic malaria in the older age group. 458 Collectively this data suggests that IL-10 is not sufficient to dampen inflammatory responses in 459 Plasmodium infection that lead to the fevers that define uncomplicated malaria.

460

461 **Discussion**

Significant gaps exist in the understanding of the contribution of anti-parasite and anti-disease immunity in the persistent asymptomatic parasite carriage in areas with high transmission of *P*. *falciparum*. Mathematical modelling of existing data indicates that asymptomatic *P*. *falciparum* infections could last for as long as 13-years [40] and field data documents the longevity of asymptomatic infections ranges from at least 2-9 months [7,41,42]. As we have previously

Immunology of asymptomatic parasitemia

467 published, our study area has high rate of persistent asymptomatic parasite carriage over the 3 month 468 time period of study [24]. However, a proportion of individuals in our study converted to symptomatic 469 malaria as defined by the development of fever. Thus, we have assessed immune correlates of 470 asymptomatic carriage of *P. falciparum*, as well as the predictors of conversion to symptomatic 471 malaria.

Consistent with the acquisition of immune response with age and exposure [13,43] in our study, older 472 473 asymptomatic individuals had better control of circulating iRBCs with a greater proportion of sub-474 microscopic infections than younger individuals. This was associated with an increase in the quantity 475 of circulating anti-parasite IgG which appeared to saturate at around 20 years old. Our data is in agreement with an age-structured mathematical model of malaria transmission which showed that 476 477 anti-parasite immunity which results in more rapid clearance of parasitemia has an approximate half-478 life of 20 years [44]. With a focus on the younger age group, there did not appear to be any correlations 479 with respect to parasitemia levels and IgG levels or the avidity of the IgG. The in vitro invasionblocking capability of the plasma in young individuals had some functionality with respect to invasion 480 481 blocking, but only at higher parasitemia levels. In this assay plasma factors other than IgG, such as 482 IgM [45,46], may contribute to this observation and control of *P. falciparum* parasitemia in children. 483 Collectively these data support a role for IgG in exerting control of iRBCs in *P. falciparum* infections, but it may not be the principal factor that dictates asymptomatic carriage of parasites in younger 484 485 individuals.

Surprisingly the proportion of those developing uncomplicated malaria over the course of our study was comparable between the younger and the older age groups, suggesting that the conversion from an asymptomatic to symptomatic infection is independent of age in highly exposed individuals. The reasons for this finding are not clear but may be related to heterogeneity of the circulating *P*. *falciparum* parasite population and emergence of highly inflammatory parasite clonotypes [47,48] that overcome the pyrogenic threshold of individuals. Consistent with the oscillation of parasite densities over time reported in a Vietnam cohort [7], parasitemia in persistent asymptomatic

Immunology of asymptomatic parasitemia

493 infections in our study fluctuated widely and independently of IgG levels, and different clonotypes

494 of *P. falciparum* were observed at different time points in this study.

495 Cytokine profiles are considerably more dynamic than IgG responses. In line with the hypothesis that clinical immunity to malaria is mediated by a tempering of inflammation, previous studies have 496 497 shown that the clinical outcome of malaria infection is largely dictated by the balance between pro-498 inflammatory and anti-inflammatory markers [17,18]. Although our study highlighted that carriage 499 of *P. falciparum* across the duration of this study was associated with a consistent inflammatory 500 response characterized by TNF- α , IL-10 and MCP-1, we found no obvious cytokine signature 501 associated with the asymptomatic carriage of P. falciparum when compared to symptomatic 502 individuals. Similar to findings in a study of visceral leishmaniasis [49], where increased IL-10 levels 503 were the primary predictor of clinical conversion from asymptomatic to symptomatic 504 visceral leishmaniasis, the conversion from asymptomatic to symptomatic malaria in our study was linked to increased baseline levels of IL-10 skewed ratios with TNF- α and IFN- γ in children at the 505 506 initial time point of our study.

507 Although IL-10 is known to exert anti-inflammatory effects that include inhibition of monocytes and 508 macrophages in the production of inflammatory cytokines such as IL-6, TNF- α , and IL-1 [50], in agreement with a recent study from the Gambia [51] in our study IL-10 was associated with the 509 510 presence of *Plasmodium* parasites. Given that higher levels of IL-6, IL-1RA and MCP-1 were also associated with symptomatic infections in the younger age group of our study, IL-10 levels are likely 511 512 a consequence of increased inflammation. Notably, IL-10 induces IL1RA, a potent anti-inflammatory cytokine that regulates the pro-inflammatory actions of IL-1 [52] that has previously been shown to 513 be associated with severity of malarial disease [53]. At the baseline time point in our study, we found 514 515 that balanced inflammatory cytokine ratios were correlated with developing symptoms from P. 516 falciparum infection in our younger age group. Specifically, those who developed uncomplicated 517 malaria had a significant IL-10 skewed ratio at the initial time point compared to those who had

Immunology of asymptomatic parasitemia

- 518 persistent asymptomatic malaria. An IL-10 skewed ratio may indicate that an individual will shortly
- 519 develop symptoms of *P. falciparum* infection.
- Most of the association between biomarkers and clinical malaria in this study were only found in the younger age group, but not in adults, indicating that the mechanisms of the protection against clinical malaria may be different according to age. As such, more studies into how asymptomatic carriage of *Plasmodium* occurs are warranted and should include non-antibody mediated mechanisms of parasite control and clinical immunity which may be relevant in children.

525

526 Acknowledgements

We thank the field staff and, most importantly, residents of the Esse Health district in Cameroon whoparticipated in this study.

529

530 Author contributions

- 531 BF devised the experiments, performed the experiments, analyzed data, and wrote the paper; FM,
- 532 MFA, EE, CD, GC, MK, RK and SK performed fieldwork and sample processing; MS and RP
- 533 provided cytokine kits and recombinant *P. falciparum* antigens, respectively and reviewed paper. RM
- supervised the experiments and reviewed paper; LA and TL devised the experiments, supervised the
- 535 experiments, analyzed data, and wrote the paper.

536 Financial Support:

537 This work was supported by the Institute Pasteur International Division to LSA.

538 **Competing interests**

539 Authors declare no competing interests.

Immunology of asymptomatic parasitemia

541 References

- 542 1. WHO. World Malaria Report. Genova; 2023.
- 543 2. Cibulskis RE, Alonso P, Aponte J, et al. Malaria: Global progress 2000 2015 and future
 544 challenges. Infect Dis Poverty. 2016; 5(1):61.
- 545 3. Galatas B, Bassat Q, Mayor A. Malaria Parasites in the Asymptomatic: Looking for the Hay in
 546 the Haystack. Trends Parasitol. 2016; 32(4):296–308.
- Drakeley C, Gonçalves B, Okell L, Slater H. Understanding the Importance of Asymptomatic and Low- Density Infections for Malaria Elimination. In: Manguin S, Dev V, editors. Malar Elimin - Leap Forw [Internet]. InTech; 2018 [cited 2018 Sep 4]. Available from: http://www.intechopen.com/books/towards-malaria-elimination-a-leapforward/understanding-the-importance-of-asymptomatic-and-low-density-infections-for-
- 551 forward/understanding-the-importance-of-asymptomatic-and-low-density-infections-for-552 malaria-elimination
- 553 5. Ashley EA, White NJ. The duration of Plasmodium falciparum infections. Malar J. 2014;
 554 13(1):500.
- 555 6. Slater HC, Ross A, Felger I, et al. The temporal dynamics and infectiousness of subpatent
 556 Plasmodium falciparum infections in relation to parasite density. Nat Commun [Internet]. 2019
 557 [cited 2020 Jun 4]; 10. Available from:
 558 https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6440965/
- Nguyen T-N, Seidlein L von, Nguyen T-V, et al. The persistence and oscillations of
 submicroscopic Plasmodium falciparum and Plasmodium vivax infections over time in
 Vietnam: an open cohort study. Lancet Infect Dis. 2018; 18(5):565–572.
- Ademolue TW, Awandare GA. Evaluating antidisease immunity to malaria and implications for
 vaccine design. Immunology. 2018; 153(4):423–434.
- Ademolue TW, Awandare GA. Evaluating antidisease immunity to malaria and implications for vaccine design. Immunology. 2018; 153(4):423–434.
- Kijogi C, Kimura D, Bao LQ, et al. Modulation of immune responses by Plasmodium falciparum
 infection in asymptomatic children living in the endemic region of Mbita, western Kenya.
 Parasitol Int. 2018; 67(3):284–293.
- 569 11. Ouédraogo M, Samadoulougou S, Rouamba T, et al. Spatial distribution and determinants of
 570 asymptomatic malaria risk among children under 5 years in 24 districts in Burkina Faso. Malar
 571 J. 2018; 17:460.
- 572 12. Griffin JT, Hollingsworth TD, Reyburn H, Drakeley CJ, Riley EM, Ghani AC. Gradual
 573 acquisition of immunity to severe malaria with increasing exposure. Proc R Soc B Biol Sci.
 574 2015; 282(1801):20142657–20142657.
- 575 13. White M, Watson J. Age, exposure and immunity. eLife [Internet]. 2021 [cited 2021 Mar 31];
 576 7. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6103766/
- 577 14. Boyle MJ, Reiling L, Feng G, et al. Human Antibodies Fix Complement to Inhibit Plasmodium
 578 falciparum Invasion of Erythrocytes and Are Associated with Protection against Malaria.
 579 Immunity. 2015; 42(3):580–590.

Immunology of asymptomatic parasitemia

- 580 15. Duncan CJA, Hill AVS, Ellis RD. Can growth inhibition assays (GIA) predict blood-stage
 581 malaria vaccine efficacy? Hum Vaccines Immunother. 2012; 8(6):706–714.
- 582 16. Garcia-Senosiain A, Kana IH, Singh S, et al. Neutrophils dominate in opsonic phagocytosis of
 583 P. falciparum blood-stage merozoites and protect against febrile malaria. Commun Biol. 2021;
 584 4(1):984.
- 585 17. Rovira-Vallbona E, Moncunill G, Bassat Q, et al. Low antibodies against Plasmodium
 586 falciparum and imbalanced pro-inflammatory cytokines are associated with severe malaria in
 587 Mozambican children: a case-control study. Malar J. 2012; 11:181.
- 588 18. Andrade BB, Santos CJN, Camargo LM, et al. Hepatitis B Infection Is Associated with
 589 Asymptomatic Malaria in the Brazilian Amazon. PLOS ONE. 2011; 6(5):e19841.
- 19. Kimenyi KM, Wamae K, Ochola-Oyier LI. Understanding P. falciparum Asymptomatic
 Infections: A Proposition for a Transcriptomic Approach. Front Immunol [Internet]. 2019 [cited
 2019 Dec 12]; 10. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6803459/
- 593 20. Guiyedi V, Bécavin C, Herbert F, et al. Asymptomatic Plasmodium falciparum infection in children is associated with increased auto-antibody production, high IL-10 plasma levels and antibodies to merozoite surface protein 3. Malar J [Internet]. 2015 [cited 2018 Feb 5]; 14(1).
 596 Available from: http://www.malariajournal.com/content/14/1/162
- 597 21. Gonçalves RM, Scopel KK, Bastos MS, Ferreira MU. Cytokine balance in human malaria: does
 598 Plasmodium vivax elicit more inflammatory responses than Plasmodium falciparum? PLoS
 599 One. 2012; 7(9):e44394.
- Wilson NO, Bythwood T, Solomon W, et al. Elevated levels of IL-10 and G-CSF associated
 with asymptomatic malaria in pregnant women. Infect Dis Obstet Gynecol. 2010; 2010.
- Essangui E, Moukoko CEE, Nguedia N, et al. Demographical, hematological and serological risk factors for Plasmodium falciparum gametocyte carriage in a high stable transmission zone in Cameroon. PLOS ONE. 2019; 14(4):e0216133.
- Fogang B, Biabi MF, Megnekou R, et al. High Prevalence of Asymptomatic Malarial Anemia and Association with Early Conversion from Asymptomatic to Symptomatic Infection in a Plasmodium falciparum Hyperendemic Setting in Cameroon. Am J Trop Med Hyg. 2021;
 :tpmd210316.
- 609 25. Biabi MFAB, Fogang B, Essangui E, et al. High Prevalence of Polyclonal Plasmodium
 610 falciparum Infections and Association with Poor IgG Antibody Responses in a Hyper-Endemic
 611 Area in Cameroon. Trop Med Infect Dis. 2023; 8(8):390.
- 612 26. Plowe CV, Djimde A, Bouare M, Doumbo O, Wellems TE. Pyrimethamine and proguanil
 613 resistance-conferring mutations in Plasmodium falciparum dihydrofolate reductase: polymerase
 614 chain reaction methods for surveillance in Africa. Am J Trop Med Hyg. 1995; 52(6):565–568.
- Ranford-Cartwright LC, Taylor J, Umasunthar T, et al. Molecular analysis of recrudescent
 parasites in a Plasmodium falciparum drug efficacy trial in Gabon. Trans R Soc Trop Med Hyg. **1997**; 91(6):719–724.
- 618 28. Quintana M del P, Ch'ng J-H, Zandian A, et al. SURGE complex of Plasmodium falciparum in
 619 the rhoptry-neck (SURFIN4.2-RON4-GLURP) contributes to merozoite invasion. PLOS ONE.
 620 2018; 13(8):e0201669.

Immunology of asymptomatic parasitemia

647

- 621 29. Heiber A, Spielmann T. Preparation of parasite protein extracts and western blot analysis. Bio622 Protoc. 2014; 4:e1136.
- 30. Bonnet S, Pêtres S, Holm I, et al. Soluble and glyco-lipid modified baculovirus Plasmodium
 falciparum C-terminal merozoite surface protein 1, two forms of a leading malaria vaccine
 candidate. Vaccine. 2006; 24(33–34):5997–6008.
- 626 31. Perraut R, Varela M-L, Loucoubar C, et al. Serological signatures of declining exposure 627 following intensification of integrated malaria control in two rural Senegalese communities. 628 [cited] **PLoS** ONE [Internet]. 2017 2019 Sep 8]; 12(6). Available from: 629 https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5469466/
- Berry AA, Gottlieb ER, Kouriba B, et al. Immunoglobulin G subclass and antibody avidity
 responses in Malian children immunized with Plasmodium falciparum apical membrane antigen
 1 vaccine candidate FMP2.1/AS02 A. Malar J. 2019; 18(1):13.
- 33. Duncan EH, Bergmann-Leitner ES. Miniaturized Growth Inhibition Assay to Assess the Antiblood Stage Activity of Antibodies. In: Vaughan A, editor. Malar Vaccines [Internet]. New
 York, NY: Springer New York; 2015 [cited 2019 Apr 30]. p. 153–165. Available from:
 http://link.springer.com/10.1007/978-1-4939-2815-6_13
- 637 34. Efange NM, Lobe MMM, Keumoe R, Ayong L, Efange SMN. Spirofused
 638 tetrahydroisoquinoline-oxindole hybrids as a novel class of fast acting antimalarial agents with
 639 multiple modes of action. Sci Rep. 2020; 10:17932.
- 640 35. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. R
 641 Foundation for Statistical Computing; 2021. Available from: https://www.R-project.org/
- 642 36. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using Ime4. J
 643 Stat Softw. 2015;67(1):1-48.
- 645 37. Kuznetsova A, Brockhoff P, Christensen R. ImerTest: Tests in linear mixed effects models.
 646 https://cranr-projectorg/package=lmerTest. 2016.
- May J, Lell B, Luty AJ, Meyer CG, Kremsner PG. Plasma interleukin-10:Tumor necrosis factor
 (TNF)-alpha ratio is associated with TNF promoter variants and predicts malarial complications.
 J Infect Dis. 2000; 182(5):1570–1573.
- 39. Othoro C, Lal AA, Nahlen B, Koech D, Orago AS, Udhayakumar V. A low interleukin-10 tumor necrosis factor-alpha ratio is associated with malaria anemia in children residing in a holoendemic malaria region in western Kenya. J Infect Dis. 1999;179(1):279-82. doi: 10.1086/314548. PubMed PMID: 9841855.
- 40. Ashley EA, White NJ. The duration of Plasmodium falciparum infections. Malar J. 2014;
 13(1):500.
- 41. Felger I, Maire M, Bretscher MT, et al. The Dynamics of Natural Plasmodium falciparum
 Infections. Gosling RD, editor. PLoS ONE. 2012; 7(9):e45542.

42. Landier J, Kajeechiwa L, Thwin MM, et al. Safety and effectiveness of mass drug administration
to accelerate elimination of artemisinin-resistant falciparum malaria: A pilot trial in four villages
of Eastern Myanmar. Wellcome Open Res. 2017; 2:81.

Immunology of asymptomatic parasitemia

- 43. Rodriguez-Barraquer I, Arinaitwe E, Jagannathan P, et al. Quantification of anti-parasite and
 anti-disease immunity to malaria as a function of age and exposure. eLife. 2018; 7:e35832.
- 44. Filipe JAN, Riley EM, Drakeley CJ, Sutherland CJ, Ghani AC. Determination of the Processes
 Driving the Acquisition of Immunity to Malaria Using a Mathematical Transmission Model.
 PLOS Comput Biol. Public Library of Science; 2007; 3(12):e255.
- 45. Hopp CS, Sekar P, Diouf A, et al. Plasmodium falciparum-specific IgM B cells dominate in children, expand with malaria, and produce functional IgM. J Exp Med. 2021;
 218(4):e20200901.
- 46. Thouvenel CD, Fontana MF, Netland J, et al. Multimeric antibodies from antigen-specific
 human IgM+ memory B cells restrict Plasmodium parasites. J Exp Med. 2021;
 218(4):e20200942.
- 673 47. Corrigan RA, Rowe JA. Strain variation in early innate cytokine induction by Plasmodium
 674 falciparum. Parasite Immunol. 2010; 32(7):512–527.
- 48. Allan RJ, Rowe A, Kwiatkowski D. Plasmodium falciparum varies in its ability to induce tumor
 676 necrosis factor. Infect Immun. 1993; 61(11):4772–4776.
- 49. Topno RK, Kumar R, Pandey K, et al. Early diagnostic marker for the conversion of
 asymptomatic to symptomatic visceral leishmaniasis. J Med Allied Sci. 2019; 9(2):96–99.
- 50. Waal Malefyt R de, Abrams J, Bennett B, Figdor CG, Vries JE de. Interleukin 10(IL-10) inhibits
 cytokine synthesis by human monocytes: an autoregulatory role of IL-10 produced by
 monocytes. J Exp Med. 1991; 174(5):1209–1220.
- 51. Mooney JP, DonVito SM, Jahateh M, et al. "Bouncing Back" From Subclinical Malaria:
 Inflammation and Erythrocytosis After Resolution of P. falciparum Infection in Gambian
 Children. Front Immunol. 2022; 13:780525.
- 685 52. Arend WP, Malyak M, Guthridge CJ, Gabay C. Interleukin-1 receptor antagonist: role in
 686 biology. Annu Rev Immunol. 1998; 16:27–55.
- 53. John CC, Park GS, Sam-Agudu N, Opoka RO, Boivin MJ. Elevated serum levels of IL-1ra in children with P. falciparum malaria are associated with increased severity of disease. Cytokine.
 2008; 41(3):204–208.
- 690
- 691