

Dietary Intake by Diabetic Status

Differences in Dietary Intake Exist Among U.S. Adults by Diabetic Status Using NHANES 2009-2016

1 **Luotao Lin¹, Fengqing Zhu², Edward J. Delp², Heather A. Eicher-Miller^{1*}**

2
3 ¹Department of Nutrition Science, Purdue University, West Lafayette, IN, United States;
4 ²School of Electrical and Computer Engineering, Purdue University, West Lafayette, IN, United
5 States.

6
7 ***Correspondence:**
8 Heather Eicher-Miller
9 heicherm@purdue.edu

10
11 Number of words: 3557
12 Number of tables: 8

13
14 **Keywords: insulin takers, food frequency, energy contribution, food category, diet choice,**
15 **diabetes.**

16
17 **Abstract**
18 Objectives:
19 Insulin takers' diets have rarely been described although insulin dosing is highly dependent on
20 dietary intake. The objective of this paper was to determine the most frequently consumed food
21 or beverage items, food subcategories, and food categories, and those that contributed most to
22 total energy intake among U.S. insulin takers, those with type 2 diabetes (T2D) not taking insulin
23 and those without diabetes; the groups were also compared at the broad food category level.

24 Methods:
25 The Laboratory tests and questionnaires of the NHANES 2009–2016 classified 774 insulin
26 takers, 2,758 T2D not taking insulin, and 17,796 participants without diabetes. USDA assigned
27 8-digit food codes identifying each food item and its membership to a category and subcategory
28 of foods were used to categorize foods based on the WWEIA categories. Raw and weighted
29 frequency and energy contributions of each food item, food subcategory, and food category were
30 calculated and ranked. Comparisons among groups by broad food category used the Rao–Scott
31 modified chi-square test.

32 Results:
33 Diet soft drinks ranked as the 4th and 7th most consumed food subcategory in insulin takers and
34 T2D not taking insulin, respectively. Soft drinks ranked as the 8th and 6th most consumed food
35 subcategory in T2D not taking insulin and those without diabetes, and contributed 5th and 2nd
36 most to energy, respectively. Protein foods represented 4 of the top 10 highest energy
37 contributing food subcategories among insulin takers, 3 of the top 10 food subcategories among
38 those with T2D not taking insulin, and only 1 subcategory among those without diabetes. Insulin
39 takers had higher consumption frequency of grains, and lower consumption frequency of sweets
40 and alcohol, and a larger share of energy comprising protein, vegetables, and grains, and a
41 smaller share of energy comprising beverages and alcohol compared to participants without
42 diabetes.

43 Conclusions:

Dietary Intake by Diabetic Status

44 Differences in dietary intake exist among U.S. adults by diabetic status. Insulin takers are likely
45 to consume more protein foods and less regular soft drinks compared to other 2 groups. Lists of
46 the most frequently reported foods and foods contributing most to energy may be helpful for
47 nutrition education, prescribing diets, and digital-based dietary assessment for insulin takers.

48

49 **1 Introduction**

50 Diabetes is a chronic disease highly influenced by diet where individuals either do not produce
51 enough insulin to regulate blood glucose level or they are not able to use the insulin they produce
52 to effectively move blood glucose into body cells (1). Over time, diabetes can cause additional
53 serious health problems including heart disease, vision loss, and kidney disease. About 34.2
54 million U.S. individuals (10.5% of the U.S. population) diabetes including type 1 (T1D), type 2
55 (T2D), and gestational diabetes in 2020 (2).

56 Both those with T1D and T2D may use injectable or oral insulin to control their blood glucose
57 levels. Proper insulin dosing to control blood glucose levels and reduce risk of severe
58 consequences is dependent on dietary intake, particularly the frequency and energy from
59 macronutrients: carbohydrate, protein, and fat (3,4). Thus, quantification of the foods comprising
60 dietary intake may be used to inform insulin dosing algorithms to enhance blood glucose level
61 control (5,6) and knowledge of the foods comprising overall dietary intake may also aid the
62 creation of meal plans, inform substitutions, and other interventions. Furthermore, it is likely that
63 diabetes status may differentiate food and beverage choice and intake among those using insulin,
64 T2D not using insulin, and those without diabetes as one strategy for glucose control is through
65 the avoidance of certain foods and monitored intake of carbohydrates, for example sugar-
66 sweetened soft drinks (7). However, very little information on the food intake of those taking
67 insulin or those with T2D not taking insulin is available (8–10).

68 Therefore, the objectives of this study were to (1) determine the most frequently consumed food
69 items, food subcategories, and food categories and the food items, food subcategories, and food
70 categories that contributed most to energy intake, and (2) compare the differences of the most
71 frequently consumed broad food categories and the broad food categories that contributed most
72 to energy intake among insulin takers, T2D participants not taking insulin, and participants
73 without diabetes among U.S. adults aged 18 years old or older using data from the National
74 Health And Nutrition Examination Survey (NHANES) 2009–2016.

75 **2 Materials and methods**

76 **2.1 Participants and Dataset**

77 NHANES is a cross-sectional survey that is conducted by the National Center for Health
78 Statistics (NCHS) of the U.S. Centers for Disease Control and Prevention to assess the health
79 and nutritional status of the non-institutionalized civilian population in the U.S. (11).

80 Participants' sociodemographic information was obtained via questionnaires during the in-person
81 household interview, which included age, sex, race/ethnicity, and poverty to income ratio (PIR).
82 The NCHS Research Ethics Review Board approved the survey and all participants consented to
83 participate (12).

84 **2.2 Analytic Sample**

85 NHANES 2009-2016 was used to include the most recent available dietary data and sufficient
86 sample size. Adults in the United States aged 18 or older who had a reliable first 24-hour dietary
87 recall, reliable laboratory test results, and valid answers to diabetic-related questions made up the
88 analytic sample.

Dietary Intake by Diabetic Status

89 NHANES diabetic-related questions and laboratory test results were used to divide the sample
90 into three groups: insulin takers, participants with diabetes not taking insulin, and participants
91 without diabetes. The number of insulin takers was determined by the question “Are you/ is the
92 sample person now taking insulin?” Those participants with diabetes who were not taking
93 insulin were identified by subtracting those using insulin from the total number of participants
94 with diabetes. Diabetes was classified by self-reporting being told they had diabetes by a doctor
95 or taking medications that lowered glucose, or by fasting plasma glucose concentration ≥ 126
96 mg/dL or hemoglobin A1c $\geq 6.5\%$ (13). Since T1D was classified when a participant reported
97 being diagnosed with diabetes before the age of 30 years old and reporting continuous insulin
98 use since diagnosis (14), insulin takers encompass all those with T1D, so participants with
99 diabetes not taking insulin is equal to participants with T2D not taking insulin. Those
100 participants without diabetes were identified by subtracting the number of participants with
101 diabetes from the sample. As a result, the study sample included 774 insulin takers, 2,758 T2D
102 who did not use insulin, and 17,796 participants who did not have diabetes.

103 **2.3 Sociodemographic characteristics**

104 Sociodemographic characteristics included survey year (2009-2010, 2011-2012, 2013-2014, and
105 2015-2016) and self-reported sex (male or female), race/ethnicity (grouped as Hispanic, non-
106 Hispanic white, non-Hispanic black, and other including multi-race), age (grouped as 18-34, 35-
107 49, and 50-80 years), and PIR which was the reported household income divided by the federal
108 poverty guideline for household income, and grouped as 0-0.99 (below poverty threshold), 1-
109 1.99, 2-2.99, 3-3.99, and 4- 5 (15).

110 **2.4 Dietary Assessment**

111 The USDA Automated Multiple-Pass Method (16) was used to collect a dietary recall recorded
112 during the physical health examination where participants reported all of the foods consumed
113 during the previous 24-hour period, including information on the time of intake, amount and type
114 of each food, and detailed food descriptions (17). Each reported food item was then linked to the
115 USDA’s Food and Nutrient Database for Dietary Studies (FNDDS) (18–21) to assign an 8-digit
116 food codes. The USDA food codes were also used to sort the reported foods into the What We
117 Eat in America food categories and subcategories (22).

118 **2.5 Statistical Analysis**

119 The Rao-Scott modified chi-square test was used to compare sociodemographic variables created
120 from the survey data among the three groups.

121 The weighted frequencies of reported foods were computed by determining the weighted sum of
122 the number of food items, food subcategories, and food categories:

$$123 \sum_{i=1}^n (R_i W_i)$$

124 where n = the sample size, i = each participant, R_i = # of reports of individual food code for the i^{th}
125 participant, W_i = dietary day 1 sample weight for the i^{th} participant (23,24). The number of
126 reports of foods for each participant was multiplied by the dietary day 1 sample weight, and then
127 summed to yield the total weighted frequency of the particular food or beverage for each
128 respective group, which represented how often that food or beverage was consumed by the
129 respective group in the U.S. in one day. The weighted frequency of the food or beverage was
130 also divided by the total weighted frequency of all foods or beverages reported in a single day to
131 get the weighted percent of the frequency of that food or beverage being consumed out of all
132 reported foods and beverages consumed.

Dietary Intake by Diabetic Status

133 The weighted energy contribution of reported foods items, food subcategories, and food
134 categories were computed by determining the weighted sum as:

$$135 \sum_{i=1}^n (E_i W_i)$$

136 where n= the sample size, i= each participant E_i =total energy for an individual food code for the
137 i^{th} participant, W_i = dietary day 1 sample weight for the i^{th} participant (23,24). The energy
138 contributed by a particular food or beverage was first summed for each participant and then
139 multiplied by the dietary day 1 sample weight of the participant, then summed for all participants
140 in the sample for each group, which represented how much energy that food or beverage
141 contributed to the total energy for the respective group in the U.S. in one day. The weighted
142 energy of the particular food or beverage was also divided by the total weighted energy from all
143 reported foods and beverages to get the weighted percent of energy from that food or beverage
144 out of the energy estimated from all reported foods and beverages.

145 The Rao-Scott modified chi-square goodness-of-fit tests were used to compare the frequency and
146 energy contribution comparisons of food terms among the three groups. Significant differences
147 were indicated as when $p < 0.05/42$ or $p < 0.0012$ using a Bonferroni type adjustment for
148 multiple comparisons for food intake among 14 broad food categories \times 3 groups. Survey
149 weights, or the reciprocal of sample inclusion probability, were used in all computations to allow
150 for inference to the non-institutionalized U.S. population. Further adjustments were performed to
151 account for the clustering and stratification inherent to the survey design. All analyses were
152 completed in SAS 9.4 (SAS Institute Inc).

153 **3 Results**

154 **3.1 Sociodemographic characteristics**

155 The sociodemographic characteristics of participants taking insulin, with T2D not taking insulin,
156 and participants without diabetes are shown in **Table 1**. Sex, age, race/ethnicity, and PIR were
157 significantly different among the three groups.

158 Most insulin takers and those with T2D who were not using insulin were ages 50 to 80 years.

159 Participants without diabetes were more evenly distributed in three age groups (18-34, 35-49, 50-
160 80). About half of the group using insulin and with T2D but not taking insulin had PIR <2,
161 furthermore, 28.1% of insulin takers had PIR below the poverty threshold (PIR<1). The group
162 without diabetes had higher incomes with more than 25% of participants having a PIR greater
163 than 4.

164 **3.2 The most frequently consumed food items, food subcategories, and food categories**

165 The top 10 most frequently reported food items, food subcategories, and food categories for
166 insulin takers, those with T2D not using insulin, and participants without diabetes are shown in
167 **Tables 2-4**. Foods are presented in descending weighted frequency order by their FNDDS short
168 food descriptions. In frequency lists, 7 food items (Table 2), 6 food subcategories (Table 3), and
169 8 food categories (Table 4) are common across the three groups but their rankings were varied.
170 Notably, diet soft drink was the 4th most frequently consumed food subcategory among insulin
171 takers, the 7th among those with T2D not taking insulin, and did not appear in the top 10 food
172 subcategory frequency list among those without diabetes (Table3). Regular soft drink did not
173 appear in the top 10 food subcategory frequency list for insulin takers, but was the 8th most
174 frequently consumed food subcategory among those with T2D not taking insulin, and the 6th
175 most frequently consumed food subcategory among participants without diabetes (Table3).

176 **3.3 The highest energy contributing food items, food subcategories, and food categories**

Dietary Intake by Diabetic Status

177 The top 10 highest energy contributing food items, food subcategories, and food categories for
178 insulin takers, T2D not using insulin, and participants without diabetes are shown in **Tables 5-7**.
179 Foods are presented in descending weighted order according to their energy contribution by their
180 FNDDS short food descriptions. In energy contribution lists, 5 food items (Table 5), 6 food
181 subcategories (Table 6), and 5 food categories (Table 7) were similar among the three groups but
182 their rankings were different. Regular soft drinks did not rank in the top 10 most energy
183 contributing food subcategories list among insulin takers, but was ranked 5th among those with
184 T2D who did not use insulin, and 2nd among participants without diabetes (Table 6).
185 Furthermore, insulin takers consumed less energy from food subcategories with added sugars
186 compared with the other two groups (cakes and pies were 6th and cookies and brownies 7th
187 among insulin takers; cakes and pies were 3rd and cookies and brownies 6th among those with
188 T2D not taking insulin; cookies and brownies were 8th and cakes and pies 9th among participants
189 without diabetes). More protein food subcategories ranked high within the top 10 list (whole
190 pieces of chicken ranked 3rd place, eggs and omelets ranked 4th place, meat mixed dishes ranked
191 5th place, cold cuts and cured meats ranked 10th place) among insulin takers compared to the
192 other two groups (whole pieces of chicken ranked 4th place, eggs and omelets 9th place, meat
193 mixed dishes 10th place in those with T2D not using insulin, and whole pieces of chicken ranked
194 5th place in participants without diabetes) (Table 6).

195 **3.4 Broad Food Categories Intake by Frequency and Energy Contribution**

196 The results of the comparison of the frequency and energy contribution of the broad WWEIA
197 food or beverage categories among the three groups are shown in **Table 8**. Broad categories of
198 grains and alcohol were statistically significantly different in both intake frequency and energy
199 contribution across the three groups. From insulin takers to T2D not using insulin to participants
200 without diabetes, grains exhibited a decreasing trend in consumption frequency and energy
201 contribution, while alcohol showed an increasing trend. Snacks / sweets were statistically
202 significantly different in consumption frequency among the three groups, which showed an
203 increasing trend from insulin takers to T2D not using insulin to participants without diabetes.
204 Protein foods, vegetables, beverages, water, and other were statistically significantly different in
205 energy contribution across the three groups. From insulin takers to participants without diabetes,
206 protein foods and vegetables exhibited a decreasing trend in energy contribution, while
207 beverages, water, and other showed an increasing trend.

208 **4 Discussion**

209 Previously, few studies have investigated the diet of insulin takers (8–10) nor included
210 consideration of intake contributions by both energy and frequency. To our knowledge, this is
211 the first study to investigate and compare the diets of insulin takers, those who have T2D but do
212 not use insulin, and participants without diabetes. Participants with varying diabetes and insulin
213 using status showed significant differences in both the frequency of which certain foods or
214 beverages are consumed and the energy contributions attributed to certain foods or beverages,
215 implying that insulin using status may be linked to dietary behaviors and the importance of
216 certain foods or beverages in the overall diet. In some broad WWEIA food categories, the
217 statistically significant differences observed in the frequency were also observed in energy
218 contribution. Both consumption frequency and energy contribution of grains (wheat bread) and
219 alcoholic beverages (beer) were statistically significantly different among the three groups.
220 The results also included statistically significant findings where either frequency or energy
221 contribution differed among the groups but not both. Beverages, water, and the “other” (nutrition
222 powder) broad food categories did not exhibit significantly different frequency of intake across

Dietary Intake by Diabetic Status

223 the three groups, but the share of energy contributed by these broad food categories was lower
224 among insulin takers compared to participants without diabetes. The share of energy contributed
225 by protein (eggs and omelets, cold cuts and cured meats) and vegetables (tomatoes and lettuce)
226 was also greater among insulin takers compared with participants without diabetes. Snacks /
227 Sweets had lower consumption frequency among insulin takers compared with participants
228 without diabetes, but the energy intake was not significantly different across all three groups.
229 In this study, insulin takers exhibited a pattern of “higher diet quality” regarding the foods they
230 consumed frequently and that contributed most to energy compared to those with T2D not using
231 insulin and participants without diabetes. Since older adults accounted for the majority of the
232 insulin takers, this finding is consistent with previous studies showing that older adults have
233 higher-quality diets (10,23,25,26). In addition, that insulin takers have a higher diet quality is
234 also supported by a previous study (10), suggesting that those with diabetes who are using
235 insulin engage in dietary behaviors promoting a more healthful mix of dietary intake compared
236 with those who do not have diabetes and those with T2D. The group using insulin reported a
237 higher consumption frequency of grains, energy contribution from grains, protein foods and
238 vegetables; and lower consumption frequency of alcohol and sweets/snacks and energy
239 contributed by beverages and alcohol compared to participants without diabetes. One explanation
240 is that grains, protein foods, and vegetables have been promoted as foods to consume to help
241 manage blood glucose control for those taking insulin. Carbohydrate intake is highly related to
242 postprandial glycemia, whole grains are a source of carbohydrates that have a favorable
243 influence on postprandial blood glucose levels (27). Protein foods may also aid dietary
244 compliance and weight loss maintenance, which is beneficial for diabetes (28,29). Vegetables
245 consumption has been associated with the improvement of diabetes due to benefits offered by the
246 bioactive compounds comprising them including dietary fiber, resistant starch, antioxidant
247 vitamins, and minerals (30–32), which may explain why insulin takers select a higher percentage
248 of their calories from vegetables compared with those who do not have diabetes. Furthermore,
249 alcohol, sweets/snacks, and beverages (heavily influenced by sweetened beverages) are foods
250 that those taking insulin may try to avoid due to their negative impact on glycemic control. These
251 foods that may add solid fats or added sugars and intake among all Americans should be reduced
252 according to the Dietary Guidelines for American, 2020-2025 (33,34). Previous studies found
253 that those with diabetes drink less alcohol compared to those without diabetes (35,36), which
254 supports the findings here that those who used insulin and those with T2D not taking insulin,
255 consumed less alcohol than participants without diabetes. Even though the long-term effect of
256 alcohol consumption on glycemic management remains to be investigated (37), heavy alcohol
257 consumption is associated with an increased incidence of diabetes (37) and alcohol consumption
258 may be a negative indicator of diabetes self-care behavior (38). The findings related to lower
259 frequency of sweets/snacks and beverages among those using insulin compared to those without
260 diabetes is supported by a previous study where participants with diabetes consumed lower
261 amounts of sweets and beverage (juice) compared to those without diabetes (36). Sweets/snacks
262 and beverages especially sweetened beverages that heavily add to energy intake, are typically
263 high in added sugar. Added sugar intake is not only a significant contributor to weight gain that
264 can lead to increased risk of diabetes, but also increased dietary glycemic load and fructose
265 metabolism that leads to inflammation and insulin resistance (39). However, it should be noted
266 that the literature reviewed here includes studies focused on those with diabetes, not specifically
267 those using insulin, due to the extremely limited evidence of dietary intake among those using

Dietary Intake by Diabetic Status

268 insulin (31,32,36). Therefore, the findings here should be further explored among those taking
269 insulin.

270 Limitations of this analysis may be due to the aggregation of foods to subcategories and
271 categories that may obscure specific food items or types of food that could be responsible for
272 much of the differences among groups. The addition or removal of specific foods or beverages
273 from a food category has the potential to change statistically significant comparisons so the
274 results and interpretation should be viewed as one perspective on the dietary intake among these
275 three groups keeping in mind that analysis of alternative food categorization, splitting to
276 subcategories or itemization may yield differential results. However, the application of the
277 WWEIA food categories lends standardization and consistency with other reporting by USDA
278 and other researchers (40–42). In addition, the ranking lists (table 2-7) were generated from the
279 foods and beverages reported over all times of the day, so separating out the most frequently
280 consumed foods and foods that highly contributed to energy intake for different times or meals
281 of the day may be helpful to inform interventions or dosing algorithms as the timing of dietary
282 intake and combinations of foods consumed at the same time influences glucose metabolism
283 (43–47). Furthermore, the NHANES dietary data this study used is only a one-day dietary recall
284 of the participants, which may not be accurate due to participants' memory and may not
285 represent the usual dietary intake (48). Moreover, misreported dietary intake may have been
286 presented in this study sample resulting in errors in energy, frequency, food item, subcategory,
287 and category representation and comparisons (49). Those with diabetes and insulin takers may
288 exhibit different patterns of misreporting compared with other population groups (36). Previous
289 studies have shown that dietary intake misreporting and especially underreporting is more
290 prevalent in overweight and obese individuals compared to normal weight individuals (50–52),
291 and prevalence of underreporting in diabetic patients may be even higher (53), considering that
292 obesity is a risk factor of diabetes that 81% of diabetic patients are obese (54). Another previous
293 study pointed out that energy misreports are prone to misreport foods that highly contributed to
294 glycemic index and glycemic load such as underreporting foods including sugars, cookies, and
295 milk products and overreporting foods including fruit and vegetables (52). For future study, the
296 differences in misreporting by diabetic status should be evaluated.

297 There are several implications and applications for the results of this study. Diabetes is a chronic
298 disease that is highly related to diet (55). The lists of foods generated in this study may be used
299 to fill the gap in knowledge of the diets of U.S. adult insulin takers or diabetic patients (8,9),
300 specifically, menus and dietary interventions developed to improve diabetes management may be
301 informed by the lists. Although frequently consumed foods and those contributing most to
302 energy may not satisfy the preferences of every person, they are generally accepted as indicated
303 by their selection in the population. Those using insulin may be more likely to adhere to a diet
304 they accept, thus forming healthier eating habits. Second, the findings of this study may inform
305 future studies to gain a greater knowledge of the diets of those using insulin. For example, the
306 list of frequently consumed foods and those most contributing to energy may inform the
307 prioritization of foods and beverages to include in food frequency questionnaires. Third, the
308 findings of this study may aid the development of nutrition assessment tools as researchers
309 should concentrate efforts on developing the most accurate identification methods for the foods
310 and beverages that are consumed the most frequently and contribute the most to total energy
311 intake in the group they are tailored for (23).

312 **5 Conclusions**

Dietary Intake by Diabetic Status

313 Differences in dietary intake exist among U.S. adults by diabetic status using NHANES 2009-
314 2016. Insulin takers consumed grains more frequently, while consuming snack/sweet and alcohol
315 less frequently compared to participants without diabetes. Protein, grains, and vegetables
316 contributed more to insulin takers' daily total energy intake compared to participants without
317 diabetes, whereas beverage and alcohol contributed less. Frequently consumed foods and those
318 contributing to energy may inform diabetes control and management, tailored food recipe,
319 consumer education, questionnaire design, and dietary assessment.

320

321 **6 Conflict of interest**

322 The authors declare that the research was conducted in the absence of any commercial or
323 financial relationships that could be construed as a potential conflict of interest.

324

325 **7 Author Contributions**

326 HAE-M and LL designed research. LL analyzed data and drafted the manuscript. HAE-M, FZ,
327 and ED critically reviewed and edited the manuscript. HAE-M had primary responsibility for
328 final content. All authors read and approved the final manuscript.

329

330 **8 Funding**

331 This study was supported by Eli Lilly and Company.

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

Dietary Intake by Diabetic Status

359 Reference:

- 360 1. Centers for Disease Control and Prevention. What is Diabetes? (2020).
361 <https://www.cdc.gov/diabetes/basics/diabetes.html> [Accessed July 14, 2021].
- 362 2. Centers for Disease Control and Prevention. National Diabetes Statistics Report (2020).
363 <https://www.cdc.gov/diabetes/pdfs/data/statistics/national-diabetes-statistics-report.pdf>
364 [Accessed July 14, 2021].
- 365 3. Centers for Disease Control and Prevention. The Insulin Resistance – Diabetes Connection
366 (2020). <https://www.cdc.gov/diabetes/basics/insulin-resistance.html> [Accessed July 14,
367 2021].
- 368 4. Centers for Disease Control and Prevention. Types of Insulin (2021).
369 <https://www.cdc.gov/diabetes/basics/type-1-types-of-insulin.html> [Accessed July 14, 2021].
- 370 5. Joslin EP, Kahn CR. Joslin's Diabetes Mellitus. Lippincott Williams & Wilkins (2005).
371 1244 p.
- 372 6. Gillespie SJ, Kulkarni KD, Daly AE. Using Carbohydrate Counting in Diabetes Clinical
373 Practice. *J Am Diet Assoc* (1998) 98(8):897 – 905.
- 374 7. Schulze MB, Manson JE, Ludwig DS, Colditz GA, Stampfer MJ, Willett WC, et al. Sugar-
375 Sweetened Beverages, Weight Gain, and Incidence of Type 2 Diabetes in Young and
376 Middle-Aged Women. *JAMA* (2004) 292(8):927 – 34.
- 377 8. McClure ST, Schlechter H, Oh S, White K, Wu B, Pilla SJ, et al. Dietary intake of adults
378 with and without diabetes: results from NHANES 2013 – 2016. *BMJ Open Diabetes Res*
379 *Care* (2020) 8(1):e001681.
- 380 9. Maffeis C, Tomasselli F, Tommasi M, Bresadola I, Trandev T, Fornari E, et al. Nutrition
381 habits of children and adolescents with type 1 diabetes changed in a 10 years span. *Pediatr*
382 *Diabetes*. (2020) 21(6):960 – 8.
- 383 10. Muñoz-Pareja M, León-Muñoz LM, Guallar-Castillón P, Graciani A, López-García E,
384 Banegas JR, et al. The Diet of Diabetic Patients in Spain in 2008 – 2010: Accordance with
385 the Main Dietary Recommendations—A Cross-Sectional Study. *PLOS ONE* (2012)
386 7(6):e39454.
- 387 11. Centers for Disease Control and Prevention. NHANES - National Health and Nutrition
388 Examination Survey Homepage (2021). <https://www.cdc.gov/nchs/nhanes/index.htm>
389 [Accessed July 14, 2021].
- 390 12. Centers for Disease Control and Prevention. NHANES - NCHS Research Ethics Review
391 Board Approval (2019). <https://www.cdc.gov/nchs/nhanes/irba98.htm> [Accessed July 14,
392 2021].

Dietary Intake by Diabetic Status

- 393 13. American Diabetes Association. Diagnosis and Classification of Diabetes Mellitus.
394 Diabetes Care (2014) 37(Supplement_1):S81 – 90.
- 395 14. Nelson KM, Reiber G, Boyko EJ. Diet and Exercise Among Adults With Type 2 Diabetes:
396 Findings from the Third National Health and Nutrition Examination Survey (NHANES III).
397 Diabetes Care (2002) 25(10):1722 – 8.
- 398 15. The United States Census Bureau. How the Census Bureau Measures Poverty (2021).
399 <https://www.census.gov/topics/income-poverty/poverty/guidance/poverty-measures.html>
400 [Accessed October 15, 2021].
- 401 16. U.S. Department of Agriculture, Agricultural Research Service. AMPM - USDA
402 Automated Multiple-Pass Method (2021). [https://www.ars.usda.gov/northeast-](https://www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-group/docs/ampm-usda-automated-multiple-pass-method/#what)
403 [area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-](https://www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-group/docs/ampm-usda-automated-multiple-pass-method/#what)
404 [group/docs/ampm-usda-automated-multiple-pass-method/#what](https://www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-group/docs/ampm-usda-automated-multiple-pass-method/#what) [Accessed October 15,
405 2021].
- 406 17. Centers for Disease Control and Prevention. NHANES Dietary Data (2021).
407 <https://wwwn.cdc.gov/nchs/nhanes/Search/DataPage.aspx?Component=Dietary> [Accessed
408 July 14, 2021].
- 409 18. U.S. Department of Agriculture, Agricultural Research Service. USDA Food and Nutrient
410 Database for Dietary Studies, 5.0. Food Surveys Research Group Home Page (2012).
411 <http://www.ars.usda.gov/ba/bhnrc/fsrg> [Accessed July 14, 2021].
- 412 19. U.S. Department of Agriculture, Agricultural Research Service. USDA Food and Nutrient
413 Database for Dietary Studies 2011-2012. Food Surveys Research Group Home Page
414 (2014). <http://www.ars.usda.gov/ba/bhnrc/fsrg> [Accessed July 14, 2021].
- 415 20. U.S. Department of Agriculture, Agricultural Research Service. USDA Food and Nutrient
416 Database for Dietary Studies 2013-2014. Food Surveys Research Group Home Page
417 (2016). <http://www.ars.usda.gov/nea/bhnrc/fsrg> [Accessed July 14, 2021].
- 418 21. U.S. Department of Agriculture, Agricultural Research Service. USDA Food and Nutrient
419 Database for Dietary Studies 2015-2016. Food Surveys Research Group Home Page
420 (2018). [https://www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-](https://www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-group/docs/fndds-download-databases/)
421 [nutrition-research-center/food-surveys-research-group/docs/fndds-download-databases/](https://www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-group/docs/fndds-download-databases/)
422 [Accessed July 14, 2021].
- 423 22. U.S. Department of Agriculture, Agricultural Research Service. What We Eat in America
424 Food Categories 2015-2016 (2021).
425 [https://www.ars.usda.gov/ARUserFiles/80400530/pdf/1516/Food_categories_2015-](https://www.ars.usda.gov/ARUserFiles/80400530/pdf/1516/Food_categories_2015-2016.pdf)
426 [2016.pdf](https://www.ars.usda.gov/ARUserFiles/80400530/pdf/1516/Food_categories_2015-2016.pdf) [Accessed July 14, 2021].
- 427 23. Eicher-Miller HA, Boushey CJ. How Often and How Much? Differences in Dietary Intake
428 by Frequency and Energy Contribution Vary among U.S. Adults in NHANES 2007 – 2012.
429 Nutrients. (2017) 9(1):86.

Dietary Intake by Diabetic Status

- 430 24. Eicher-Miller HA, Boushey CJ, Bailey RL, Yang YJ. Frequently Consumed Foods and
431 Energy Contributions among Food Secure and Insecure U.S. Children and Adolescents.
432 *Nutrients*. (2020) 12(2):304.
- 433 25. Hiza HAB, Casavale KO, Guenther PM, Davis CA. Diet Quality of Americans Differs by
434 Age, Sex, Race/Ethnicity, Income, and Education Level. *J Acad Nutr Diet*. (2013)
435 113(2):297 – 306.
- 436 26. Forshee RA, Storey ML. Demographics, not beverage consumption, is associated with diet
437 quality. *Int J Food Sci Nutr*. (2006) 57(7 – 8):494 – 511.
- 438 27. Fung TT, Hu FB, Pereira MA, Liu S, Stampfer MJ, Colditz GA, et al. Whole-grain intake
439 and the risk of type 2 diabetes: a prospective study in men. *Am J Clin Nutr*. (2002)
440 76(3):535 – 40.
- 441 28. Clifton P. High-protein and low-glycaemic diets improve dietary compliance and
442 maintenance of weight loss in overweight adults who have lost weight on a low-calorie diet.
443 *BMJ Evid-Based Med*. (2011) 16(4):112 – 3.
- 444 29. Westerterp-Plantenga MS, Nieuwenhuizen A, Tomé D, Soenen S, Westerterp KR. Dietary
445 Protein, Weight Loss, and Weight Maintenance. *Annu Rev Nutr*. (2009) 29(1):21 – 41.
- 446 30. Dias JS. Nutritional Quality and Health Benefits of Vegetables: A Review. *Food Nutr Sci*.
447 (2012) 3(10):1354 – 74.
- 448 31. Carlos J, Dias J, Imai S. Vegetables Consumption and its Benefits on Diabetes. *J Nutr Ther*.
449 (2017) 6:1 – 10.
- 450 32. Platel K, Srinivasan K. Plant foods in the management of Diabetes mellitus: Vegetables as
451 potential hypoglycaemic agents. *Food Nahr*. (1997) 41(2):68 – 74.
- 452 33. Nicklas TA, O'Neil CE. Development of the SoFAS (Solid Fats and Added Sugars)
453 Concept: The 2010 Dietary Guidelines for Americans. *Adv Nutr*. (2015) 6(3):368S-375S.
- 454 34. U.S. Department of Agriculture and U.S. Department of Health and Human Services.
455 *Dietary Guidelines for Americans, 2020-2025. 9th Edition* (2020).
456 <https://DietaryGuidelines.gov> [Accessed July 14, 2021].
- 457 35. Mackenzie T, Brooks B, O'Connor G. Beverage Intake, Diabetes, and Glucose Control of
458 Adults in America. *Ann Epidemiol*. (2006) 16(9):688 – 91.
- 459 36. Nöthlings U, Boeing H, Maskarinec G, Sluik D, Teucher B, Kaaks R, et al. Food intake of
460 individuals with and without diabetes across different countries and ethnic groups. *Eur J*
461 *Clin Nutr*. (2011) 65(5):635 – 41.
- 462 37. Howard AA, Arnsten JH, Gourevitch MN. Effect of Alcohol Consumption on Diabetes
463 Mellitus. *Ann Intern Med*. (2004) 140(3):211 – 9.

Dietary Intake by Diabetic Status

- 464 38. Ahmed AT, Karter AJ, Liu J. Alcohol consumption is inversely associated with adherence
465 to diabetes self-care behaviours. *Diabet Med.* (2006) 23(7):795 – 802.
- 466 39. Malik VS, Popkin BM, Bray GA, Després J-P, Hu FB. Sugar-Sweetened Beverages,
467 Obesity, Type 2 Diabetes Mellitus, and Cardiovascular Disease Risk. *Circulation.* (2010)
468 121(11):1356 – 64.
- 469 40. Hess JM, Cifelli CJ, Agarwal S, Fulgoni VL. Comparing the cost of essential nutrients from
470 different food sources in the American diet using NHANES 2011 – 2014. *Nutr J.* (2019)
471 18(1):68.
- 472 41. Murphy MM, Scrafford CG, Barraj LM, Bi X, Higgins KA, Jaykus L-A, et al. Potassium
473 chloride-based replacers: modeling effects on sodium and potassium intakes of the US
474 population with cross-sectional data from NHANES 2015 – 2016 and 2009 – 2010. *Am J*
475 *Clin Nutr.* (2021) 114(1):220 – 30.
- 476 42. Slining MM, Popkin BM. Trends in intakes and sources of solid fats and added sugars
477 among U.S. children and adolescents: 1994 – 2010. *Pediatr Obes.* (2013) 8(4):307 – 24.
- 478 43. Lowden A, Moreno C, Holmbäck U, Lennernäs M, Tucker P. Eating and shift work —
479 effects on habits, metabolism, and performance. *Scand J Work Environ Health.* (2010)
480 36(2):150 – 62.
- 481 44. Aqeel M, Forster A, Richards EA, Hennessy E, McGowan B, Bhadra A, et al. The Effect of
482 Timing of Exercise and Eating on Postprandial Response in Adults: A Systematic Review.
483 *Nutrients.* (2020) 12(1):221.
- 484 45. Johnston JD. Physiological responses to food intake throughout the day. *Nutr Res Rev.*
485 (2014) 27(1):107 – 18.
- 486 46. Jacobs DR Jr, Steffen LM. Nutrients, foods, and dietary patterns as exposures in research: a
487 framework for food synergy. *Am J Clin Nutr.* (2003) 78(3):508S-513S.
- 488 47. Winham DM, Hutchins AM, Thompson SV. Glycemic Response to Black Beans and
489 Chickpeas as Part of a Rice Meal: A Randomized Cross-Over Trial. *Nutrients.* (2017)
490 9(10):1095.
- 491 48. Klesges RC, Eck LH, Ray JW. Who underreports dietary intake in a dietary recall?
492 Evidence from the Second National Health and Nutrition Examination Survey. *J Consult*
493 *Clin Psychol.* (1995) 63(3):438 – 44.
- 494 49. Freedman LS, Commins JM, Moler JE, Arab L, Baer DJ, Kipnis V, et al. Pooled Results
495 From 5 Validation Studies of Dietary Self-Report Instruments Using Recovery Biomarkers
496 for Energy and Protein Intake. *Am J Epidemiol.* (2014) 180(2):172 – 88.

Dietary Intake by Diabetic Status

- 497 50. Lissner L, Troiano RP, Midthune D, Heitmann BL, Kipnis V, Subar AF, et al. OPEN about
498 obesity: recovery biomarkers, dietary reporting errors and BMI. *Int J Obes.* (2007)
499 31(6):956 – 61.
- 500 51. Ocké MC, Larrañaga N, Grioni S, van den Berg SW, Ferrari P, Salvini S, et al. Energy
501 intake and sources of energy intake in the European Prospective Investigation into Cancer
502 and Nutrition. *Eur J Clin Nutr.* (2009) 63(4):S3 – 15.
- 503 52. Livingstone MBE, Black AE. Markers of the Validity of Reported Energy Intake. *J Nutr.*
504 (2003) 133(3):895S-920S.
- 505 53. Salle A, Ryan M, Ritz P. Underreporting of food intake in obese diabetic and nondiabetic
506 patients. *Diabetes Care.* (2006) 29(12):2726-7.
- 507 54. Gregg EW, Cheng YJ, Narayan KMV, Thompson TJ, Williamson DF. The relative
508 contributions of different levels of overweight and obesity to the increased prevalence of
509 diabetes in the United States: 1976 – 2004. *Prev Med.* (2007) 45(5):348 – 52.
- 510 55. Nuttall FQ. Diet and the Diabetic Patient. *Diabetes Care.* (1983) 6(2):11.
511

Dietary Intake by Diabetic Status

512 **Table 1.** Sociodemographic characteristics of U.S. adults 18 y or older taking insulin, with Type 2 diabetes but not taking insulin, and
 513 without diabetes as drawn from the NHANES, 2009-2016¹.

	Insulin takers (n=774)	T2D w/o insulin (n=2,758)	Participants w/o diabetes (n=17,796)	P-value ²
Sex				0.0003*
Male	403 (52.1%)	1416 (51.3%)	8578 (48.2%)	
Female	371 (47.9%)	1342 (48.7%)	9218 (51.8%)	
Age in Years				<0.0001*
18-34	44 (5.6%)	121 (4.4%)	6247 (35.1%)	
35-49	91 (11.8%)	470 (17.0%)	4633 (26.0%)	
50-80	639 (82.6%)	2167 (78.6%)	6916 (38.9%)	
Survey year				0.4
2009-2010	193 (24.9%)	766 (27.8%)	5005 (28.1%)	
2011-2012	181 (23.4%)	647 (23.5%)	4168 (23.4%)	
2013-2014	181 (23.4%)	660 (23.9%)	4387 (24.7%)	
2015-2016	219 (28.3%)	685 (24.8%)	4236 (23.8%)	
Race/Ethnicity				<0.0001*
Hispanic	205 (27.7%)	859 (31.1%)	4448 (25.0%)	
White	276 (35.7%)	905 (32.8%)	7517 (42.2%)	
Black	229 (29.6%)	692 (25.1%)	3640 (20.5%)	
Other	54 (7.0%)	302 (11.0%)	2191 (12.3%)	
Poverty Income Ratio				<0.0001*
0.00-0.99	201 (28.1%)	623 (25.0%)	3776 (23.2%)	
1.00-1.99	220 (30.7%)	748 (30.0%)	4219 (25.9%)	
2.00-2.99	106 (14.8%)	384 (15.4%)	2353 (14.4%)	
3.00-3.99	66 (9.2%)	277 (11.1%)	1793 (11.0%)	
4.00-5.00	123 (17.2%)	463 (18.5%)	4156 (25.5%)	

514 ¹ Total numbers do not always add to sample size due to missing values. Percentages do not always add to 100 due to rounding.

515 ² Rao–Scott modified chi-square p-value is shown. Analyses were adjusted for survey design including stratification and clustering.
 516 Sample weights were appropriately constructed and applied to this analysis as directed by NCHS. Weights were rescaled so that the
 517 sum to the weights matched the survey population at the midpoint of the 8 years, 2009–2016. Statistical significance for differences
 518 among three groups when p < 0.05 is indicated by “*”.

519

Dietary Intake by Diabetic Status

520 **Table 2.** Top 10 most frequently consumed foods items or beverages¹, unweighted and weighted frequency of reported foods items or
 521 beverages, percent of reported foods or beverages, and standard error of percent of reported foods or beverages among all reported
 522 foods or beverages for U.S. adults 18 y or older taking insulin, with Type 2 diabetes but not taking insulin, and without diabetes drawn
 523 from 2009–2016 NHANES data.

Rank	Insulin takers (n=774)				T2D w/o taking insulin (n=2,758)				Participants w/o diabetes (n=17,796)			
	What We Eat In America item ¹	Weighted Frequency ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (SE)	What We Eat In America item ¹	Weighted Frequency ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (SE)	What We Eat In America item ¹	Weighted Frequency ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (SE)
Total		98,959,094	12,095			376,469,886	43,421			3,245,557,852	277,244	
1	tap water	5,294,037	579	5.3 (0.4)	Tap water	20,867,328	2,231	5.5 (0.2)	Tap water	202,005,485	15,400	6.2 (0.2)
2	Unsweetened Bottled Water	4,341,351	600	4.4 (0.3)	Unsweetened bottled water	15,815,995	2,157	4.2 (0.3)	Unsweetened bottled water	114,654,421	11,820	3.5 (0.1)
3	Coffee	2,867,410	336	2.9 (0.2)	Coffee	10,982,634	1,212	2.9 (0.1)	Coffee	92,688,711	6,906	2.9 (0.1)
4	Tomatoes	1,583,968	179	1.6 (0.2)	Lettuce	5,520,635	597	1.5 (0.1)	White sugar	48,766,824	5,271	1.5 (0.1)
5	2% reduced fat milk	1,483,662	187	1.5 (0.2)	Tomatoes	5,499,873	598	1.5 (0.1)	Lettuce	45,653,422	3,684	1.4 (0.0)
6	Soft drink (cola type, sugar free)	1,367,532	155	1.4 (0.2)	2% reduced fat milk	5,377,691	613	1.4 (0.1)	Soft drink (cola-type)	43,801,567	4,172	1.3 (0.1)
7	Lettuce	1,277,944	156	1.3 (0.1)	Soft drink (cola-type, sugar-free)	4,446,739	344	1.2 (0.1)	Tomatoes	43,656,646	3,468	1.3 (0.0)
8	Soft drink (cola-type, decaffeinated)	867,827	86	0.9 (0.2)	White sugar	4,285,836	633	1.1 (0.1)	2% Fat reduced milk	37,409,699	3,298	1.2 (0.0)
9	Sugar substitute (saccharin-based)	850,553	124	0.9 (0.1)	Banana	3,928,098	530	1.0 (0.1)	Banana	31,398,643	2,737	1.0 (0.0)

Dietary Intake by Diabetic Status

10	Banana	818,469	125	0.8 (0.1)	Soft drink (cola-type)	3,726,509	460	1.0 (0.1)	Whole milk	23,685,000	2,338	0.7 (0.0)
----	--------	---------	-----	--------------	---------------------------	-----------	-----	--------------	---------------	------------	-------	--------------

524

525 ¹ 6,772 food or beverage items were reported by participants from NHANES 2009-2016;

526 ² The sum of the estimated weighted frequencies for all type foods or beverages reported in one day among adult participants ≥ 18
527 years of NHANES 2009–2016;

528 ³ Survey weights and adjustments for the complex survey design were applied to represent the non-institutionalized U.S. population; ⁴
529 The frequency that a food or beverages was reported without sample weights;

530 ⁵ Derived from the weighted frequency of the individual food or beverage divided by the total weighted frequency of all foods and
531 beverages (n) reported by whole sample in a single day, where n = 98,959,094 for insulin takers, n = 376,469,886 for T2D w/o taking
532 insulin, and n = 3,245,557,852 for participants w/o diabetes. Estimated weighted percent has been abbreviated by “Wtd %”; SE stands
533 for standard error.

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

Dietary Intake by Diabetic Status

555 **Table 3.** Top 10 most frequently consumed What We Eat in America foods or beverage subcategories¹, unweighted and weighted
 556 frequency of reported foods or beverage subcategories, percent of the reported foods or beverage subcategory out of the total, and
 557 standard error of the percent of reported foods or beverage represented by the subcategory among all reported foods or beverage
 558 subcategories for U.S. adults 18 y or older taking insulin, with Type 2 diabetes but not taking insulin, and without diabetes drawn from
 559 2009–2016 NHANES data.
 560

Rank	Insulin takers (n=774)				T2D w/o taking insulin (n=2,758)				Participants w/o diabetes (n=17,796)			
	What We Eat In America subcategory ¹	Weighted Frequency ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (S E)	What We Eat In America subcategory ¹	Weighted Frequency ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (S E)	What We Eat In America subcategory ¹	Weighted Frequency ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (S E)
Total		98,959,094	12,095			376,469,886	43,421			3,245,557,852	277,244	
1	Tap water	5,356,421	585	5.4 (0.4)	Tap water	21,157,722	2,272	5.6 (0.2)	Tap water	205,305,226	15,673	6.3 (0.2)
2	Coffee	4,409,221	560	4.5 (0.3)	Coffee	17,061,047	2,097	4.5 (0.1)	Coffee	136,685,004	11,227	4.2 (0.1)
3	Bottled water	4,341,351	600	4.4 (0.3)	Bottled water	15,815,995	2,157	4.2 (0.3)	Bottled water	114,654,421	11,820	3.5 (0.1)
4	Diet soft drinks	3,718,166	391	3.8 (0.4)	Yeast breads	13,091,450	1,529	3.5 (0.1)	Cheese	95,913,887	7,342	3.0 (0.1)
5	Yeast bread	3,652,103	449	3.7 (0.2)	Cheese	9,577,396	969	2.5 (0.1)	Yeast breads	95,137,901	8,275	2.9 (0.1)
6	Tea	2,584,194	293	2.6 (0.3)	Tea	9,519,279	1,070	2.5 (0.2)	Soft drinks	92,330,988	8,918	2.8 (0.1)
7	Cheese	2,261,411	265	2.3 (0.2)	Diet soft drinks	8,358,603	685	2.2 (0.1)	Tea	73,796,069	6,327	2.3 (0.1)
8	Sugar substitutes	2,220,376	326	2.2 (0.3)	Soft drinks	7,651,814	957	2.0 (0.1)	Other vegetables and combinations	61,835,369	5,004	1.9 (0.0)
9	Eggs and omelets	2,000,524	255	2.0 (0.2)	Other vegetables	7,539,753	823	2.0 (0.1)	Sugars and honey	60,595,579	6,302	1.9 (0.1)

Dietary Intake by Diabetic Status

					and combinati ons							
10	Cold cuts and cured meats	1,973,622	222	2.0 (0.2)	Cream and cream substitutes	7,361,081	858	2.0 (0.1)	Lettuce and lettuce salads	60,358,697	4,650	1.9 (0.0)

561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587

¹ The What We Eat in America Food Categories were applied to categorize all foods and beverages to 151 subcategories;
² The sum of the estimated weighted frequencies for all type foods or beverages reported in one day among adult participants ≥ 18 years of NHANES 2009–2016;
³ Survey weights and adjustments for the complex survey design were applied to represent the non-institutionalized U.S. population;
⁴ The frequency that a food or beverages was reported without sample weights;
⁵ Derived from the weighted frequency of the food or beverage subcategories divided by the total weighted frequency of all foods and beverages subcategories (n) reported by whole sample in a single day, where n = 98,959,094 for insulin takers, n = 376,469,886 for T2D w/o taking insulin, and n = 3,245,557,852 for participants w/o diabetes. Estimated weighted percent has been abbreviated by “Wtd %”; SE stands for standard error.

Dietary Intake by Diabetic Status

588 **Table 4.** Top 10 most frequently consumed What We Eat In America food or beverage categories¹, unweighted and weighted
 589 frequency of reported food or beverage categories, percent of the reported foods or beverage category out of total, and standard error
 590 of the percent of reported foods or beverage represented by the category among all reported foods or beverage categories for U.S.
 591 adults 18 y or older taking insulin, with Type 2 diabetes but not taking insulin, and without diabetes drawn from 2009–2016 NHANES
 592 data.
 593

Rank	Insulin takers (n=774)				T2D w/o taking insulin (n=2,758)				Participants w/o diabetes (n=17,796)			
	What We Eat In America Category ¹	Weighted Frequency ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (S E)	What We Eat In America Category ¹	Weighted Frequency ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (S E)	What We Eat In America Category ¹	Weighted Frequency ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (S E)
Total		98,578,506	12,039			374,707,908	43,155			3,232,070,662	275,796	
1	Plain water	9,697,772	1,185	9.8 (0.4)	Plain water	36,973,717	4,429	9.9 (0.3)	Plain water	319,959,647	27,493	9.9 (0.2)
2	Vegetables w/o white potato	9,059,302	982	9.2 (0.6)	Vegetables w/o white potato	33,386,898	3,683	8.9 (0.3)	Vegetables w/o white potato	274,695,816	22,064	8.5 (0.1)
3	Coffee & tea	6,993,415	853	7.1 (0.3)	Coffee & tea	26,580,326	3,167	7.1 (0.2)	Coffee & tea	210,481,073	17,554	6.5 (0.1)
4	Fats & oil	6,975,164	799	7.1 (0.4)	Fats & oils	24,735,148	2,599	6.6 (0.2)	Fats & oils	196,833,742	15,227	6.1 (0.1)
5	Breads, rolls, tortillas	5,814,047	704	5.9 (0.3)	Breads, rolls, tortillas	21,868,818	2,632	5.8 (0.1)	Breads, rolls, tortillas	161,270,439	14,243	5.0 (0.1)
6	Fruits	4,539,100	604	4.6 (0.3)	Fruits	17,789,900	2,199	4.7 (0.2)	Fruits	150,617,266	12,803	4.7 (0.1)
7	Diet beverages	4,191,552	449	4.3 (0.4)	Condiments and sauces	15,875,817	1,788	4.2 (0.2)	Condiments and sauces	148,905,014	12,777	4.6 (0.1)
8	Condiments and sauces	4,117,112	470	4.2 (0.3)	Sugars	14,422,141	1,877	3.8 (0.2)	Sweetened beverages	144,772,013	14,108	4.5 (0.1)

Dietary Intake by Diabetic Status

9	Sugar	3,757,238	562	3.8 (0.3)	Milk	13,242,566	1,459	3.5 (0.2)	Sugars	110,810,638	10,222	3.4 (0.1)
10	Cured meats/poultry	3,366,292	397	3.4 (0.3)	Sweet bakery product	12,011,790	1,339	3.2 (0.1)	Milk	103,668,977	8,770	3.2 (0.1)

594 ¹ The What We Eat in America Food Categories were applied to categorize all foods and beverages to 46 categories;
 595 ² The sum of the estimated weighted frequencies for all type foods or beverages reported in one day among adult participants ≥ 18
 596 years of NHANES 2009–2016;
 597 ³ Survey weights and adjustments for the complex survey design were applied to represent the non-institutionalized U.S. population; ⁴
 598 The frequency that a food or beverages was reported without sample weights;
 599 ⁵ Derived from the weighted frequency of the food or beverage categories divided by the total weighted frequency of all foods and
 600 beverages categories (n) reported by whole sample in a single day, where n = 98,578,506 for insulin takers, n = 374,707,908 for T2D
 601 w/o taking insulin, and n = 3,232,070,662 for participants w/o diabetes. Estimated weighted percent has been abbreviated by
 602 “Wtd %”; SE stands for standard error.
 603
 604
 605
 606
 607
 608
 609
 610
 611
 612
 613
 614
 615
 616
 617
 618
 619
 620

Dietary Intake by Diabetic Status

621 **Table 5.** Top 10 foods items or beverages contributing the most to total unweighted and weighted energy, percent of the total
 622 represented by the reported food or beverage item¹, and standard error of the percent of the total represented by the reported food or
 623 beverage item among all reported foods or beverage items for U.S. adults 18 y or older taking insulin, with Type 2 diabetes but not
 624 taking insulin, and without diabetes drawn from 2009–2016 NHANES data.
 625

Rank	Insulin takers (n=774)				T2D w/o taking insulin (n=2,758)				Participants w/o diabetes (n=17,796)			
	What We Eat In America item ¹	Weighted Energy Contribution ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (SE)	What We Eat In America item ¹	Weighted Energy Contribution ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (SE)	What We Eat In America item ¹	Weighted Energy Contribution ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (SE)
Total		11,173,326,140	10,655			45,994,398,876	38,344			438,552,576,789	246,981	
1	2% reduced fat milk	175,158,078	187	1.5 (0.3)	Soft drink, cola-type	620,925,847	460	1.4 (0.1)	Soft drink, cola-type	7,727,219,654	4,172	1.8 (0.1)
2	Wheat or cracked wheat bread	107,505,626	85	1.0 (0.6)	2% reduced fat milk	542,124,845	613	1.2 (0.1)	Beer (include ale)	7,624,446,189	1,643	1.7 (0.1)
3	Soft white roll	100,546,094	54	0.9 (0.2)	Whole milk	495,105,225	345	1.1 (0.2)	French fries	4,536,464,186	1,608	1.0 (0.0)
4	Regular ice cream (not chocolate)	97,758,809	28	0.9 (0.3)	Beer (include ale)	446,611,096	133	1.0 (0.2)	2% reduced fat milk	4,370,770,940	3,298	1.0 (0.1)
5	White bread	92,313,363	100	0.8 (0.1)	French fries	390,088,888	171	0.8 (0.1)	Beer, lite	4,119,958,979	1,005	0.9 (0.1)
6	Whole wheat bread	88,551,645	59	0.8 (0.2)	Banana	370,150,703	530	0.8 (0.1)	Ice cream (not	3,839,785,911	1,133	0.9 (0.1)

Dietary Intake by Diabetic Status

									chocolate)			
7	Soft drink, cola-type	85,779,001	77	0.8 (0.2)	Ice cream (Not chocolate)	358,175,595	157	0.8 (0.1)	Banana	3,167,442,419	2,737	0.7 (0.0)
8	White potato	79,297,997	56	0.7 (0.1)	White bread	339,050,117	328	0.7 (0.1)	Whole milk	3,063,855,299	2,338	0.7 (0.0)
9	Banana	77,782,509	125	0.7 (0.1)	Soft white roll	293,239,277	242	0.6 (0.1)	White bread	3,028,693,277	1,902	0.7 (0.0)
10	Whole milk	77,426,678	85	0.7 (0.1)	Wheat or cracked wheat bread	267,359,710	275	0.6 (0.1)	Soft drink, fruit-flavored, caffeine free	2,649,165,951	2,007	0.6 (0.0)

¹ 6,772 food or beverage items were reported by participants from NHANES 2009-2016;

² The sum of the estimated weighted energy contribution for all type foods or beverages reported in one day among adult participants ≥ 18 years of NHANES 2009–2016;

³ Survey weights and adjustments for the complex survey design were applied to represent the non-institutionalized U.S. population;

⁴ The frequency that a food or beverages was reported without sample weights;

⁵ Derived from the weighted energy contribution of the individual food or beverage divided by the total weighted energy contribution of all foods and beverages (n) reported by whole sample in a single day, where n = 11,173,326,140 for insulin takers, n = 45,994,398,876 for T2D w/o taking insulin, and n = 438,552,576,789 for participants w/o diabetes. Estimated weighted percent has been abbreviated by “Wtd %”; SE stands for standard error.

626
627
628
629
630

Dietary Intake by Diabetic Status

631 **Table 6.** Top 10 highest energy contributing What We Eat in America food and beverage subcategories¹ contributing the most to
 632 unweighted and weighted energy, percent of the total represented by the reported food or beverage subcategory, and standard error of
 633 the percent of the total represented by the reported food or beverage subcategory among all reported food or beverage subcategories
 634 for U.S. adults 18 y or older taking insulin, with Type 2 diabetes but not taking insulin, and without diabetes drawn from 2009–2016
 635 NHANES data.
 636

Rank	Insulin takers (n=774)				T2D w/o taking insulin (n=2,758)				Participants w/o diabetes (n=17,796)			
	What we Eat in America subcategory ¹	Weighted Energy Contribution ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (SE)	What we Eat in America subcategory ¹	Weighted Energy Contribution ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (SE)	What we Eat in America subcategory ¹	Weighted Energy Contribution ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (SE)
Total		11,173,326,140	10,655			45,994,398,876	38,344			438,552,576,789	246,981	
1	Yeast breads	542,059,395	449	4.9 (0.3)	Yeast breads	2,015,776,561	1,529	4.4 (0.2)	Pizza	17,155,217,138	2,325	3.9 (0.2)
2	Pizza	352,519,439	74	3.2 (0.6)	Pizza	1,630,091,499	249	3.5 (0.5)	Soft drinks	16,878,162,486	8,918	3.8 (0.1)
3	Chicken, whole pieces	344,627,978	182	3.1 (0.5)	Cakes and pies	1,406,511,677	386	3.1 (0.3)	Yeast breads	15,025,292,371	8,275	3.4 (0.1)
4	Eggs and omelets	322,535,347	255	2.9 (0.3)	Chicken, whole pieces	1,276,775,627	731	2.8 (0.2)	Beer	12,606,044,121	2,790	2.9 (0.1)
5	Meat mixed dishes	279,742,218	105	2.5 (0.4)	Soft drinks	1,266,355,047	957	2.8 (0.2)	Chicken, whole pieces	11,830,893,925	4,719	2.7 (0.1)
6	Cakes and pies	258,928,280	91	2.3 (0.4)	Cookies and brownies	1,132,693,357	630	2.5 (0.2)	Nuts and seeds	11,385,283,950	3,925	2.6 (0.1)
7	Cookies and brownies	245,466,831	185	2.2 (0.2)	Nuts and seeds	1,129,967,899	563	2.5 (0.2)	Burritos and tacos	10,936,204,042	1,480	2.5 (0.1)

Dietary Intake by Diabetic Status

8	Cheese	241,035,357	265	2.2 (0.3)	Burritos and tacos	1,116,149,548	203	2.4 (0.4)	Cookies and brownies	10,449,344,792	4,386	2.4 (0.1)
9	Burritos and tacos	238,780,285	49	2.1 (0.5)	Eggs and omelets	1,062,625,029	770	2.3 (0.1)	Cakes and pies	10,399,896,397	2,373	2.4 (0.1)
10	Cold cuts and cured meats	218,354,393	222	2.0 (0.4)	Meat mixed dishes	980,422,000	371	2.1 (0.2)	Cheese	9,908,844,842	7,342	2.3 (0.1)

637

638

¹ The What We Eat in America Food Categories were applied to categorize all foods and beverages to 151 subcategories;

639

² The sum of the estimated weighted energy contribution for all type foods or beverages reported in one day among adult participants ≥ 18 years of NHANES 2009–2016;

640

³ Survey weights and adjustments for the complex survey design were applied to represent the non-institutionalized U.S. population;

641

⁴ The frequency that a food or beverages was reported without sample weights;

642

⁵ Derived from the weighted energy contribution of the food or beverage subcategories divided by the total weighted energy contribution of all foods and beverages subcategories (n) reported by whole sample in a single day, where n = 11,173,326,140 for insulin takers, n = 45,994,398,876 for T2D w/o taking insulin, and n = 438,552,576,789 for participants w/o diabetes. Estimated weighted percent has been abbreviated by “Wtd %”; SE stands for standard error.

643

644

645

646

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

Dietary Intake by Diabetic Status

662 **Table 7.** Top 10 highest energy contributing What We Eat in America food and beverage categories¹ contributing the most to
 663 unweighted and weighted energy, percent of the total represented by the reported food or beverage category, and standard error of the
 664 percent of the total represented by the reported food or beverage category among all reported food or beverage categories for U.S.
 665 adults 18 y or older taking insulin, with Type 2 diabetes but not taking insulin, and without diabetes drawn from 2009–2016 NHANES
 666 data.
 667

Rank	Insulin takers (n=774)				T2D w/o taking insulin (n=2,758)				Participants w/o diabetes (n=17,796)			
	What We Eat In America category ¹	Weighted Energy Contribution ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (S E)	What We Eat In America category ¹	Weighted Energy Contribution ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (S E)	What We Eat In America category ¹	Weighted Energy Contribution ^{2,3}	Frequency ⁴	Wtd % ^{3,5} (S E)
Total		11,067,583,100	10,599			45,507,634,682	38,078			434,779,439,791	245,533	
1	Breads, rolls, tortillas	923,994,769	704	8.3 (0.4)	Breads, rolls, tortillas	3,653,085,399	2,632	8.0 (0.2)	Sweet bakery product	27,904,006,131	9,010	6.4 (0.1)
2	Sweet bakery product	629,785,636	341	5.7 (0.5)	Sweet bakery product	3,257,752,964	1,339	7.2 (0.4)	Breads, rolls, tortillas	27,497,056,987	14,243	6.3 (0.1)
3	Mixed dishes-sandwiches	550,338,991	155	5.0 (0.6)	Sweetened beverages	2,041,686,370	1,518	4.5 (0.2)	Sweetened beverages	26,867,482,295	14,106	6.2 (0.1)
4	Mixed dishes-meat, poultry, seafood	487,328,523	194	4.4 (0.5)	Mixed dishes-sandwiches	1,918,027,605	477	4.2 (0.3)	Alcoholic beverages	21,928,960,736	5,463	5.0 (0.2)
5	Fats & oils	449,849,893	799	4.1 (0.3)	Mixed dishes-meat, poultry, seafood	1,846,139,541	703	4.1 (0.3)	Mixed dishes-sandwiches	18,966,260,477	3,297	4.4 (0.1)

Dietary Intake by Diabetic Status

6	Poultry	441,972,521	233	4.0 (0.5)	White potatoes	1,762,408,5 54	842	3.9 (0.2)	Mixed dishes- pizza	17,155,217,1 38	2,325	3.9 (0.2)
7	Cured meats/poul try	427,895,678	397	3.9 (0.4)	Plant based protein foods	1,711,721,0 04	1,041	3.8 (0.2)	Mixed dishes- Mexican	16,503,304,1 78	2,545	3.8 (0.2)
8	White potatoes	377,782,656	216	3.4 (0.4)	Poultry	1,654,562,1 56	915	3.6 (0.2)	Poultry	16,063,883,3 80	6,066	3.7 (0.1)
9	Mixed dishes- Mexican	362,937,339	96	3.3 (0.6)	Mixed dishes- pizza	1,630,091,4 99	249	3.6 (0.5)	Plant based protein foods	15,960,314,7 84	6,477	3.7 (0.1)
10	Sweetened beverages	361,977,279	336	3.3 (0.3)	Mixed dishes- Mexican	1,602,969,8 44	361	3.5 (0.4)	Mixed dishes- grain based	15,334,421,3 26	2,991	3.5 (0.1)

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

¹ The What We Eat in America Food Categories were applied to categorize all foods and beverages to 46 categories;

² The sum of the estimated weighted energy contribution for all type foods or beverages categories reported in one day among adult participants ≥ 18 years of NHANES 2009–2016;

³ Survey weights and adjustments for the complex survey design were applied to represent the non-institutionalized U.S. population;

⁴ The frequency that a food or beverages was reported without sample weights;

⁵ Derived from the weighted energy contribution of the food or beverage category divided by the total weighted energy contribution of all foods and beverage categories (n) reported by whole sample in a single day, where n = 11,067,583,100 for insulin takers, n = 45,507,634,682 for T2D w/o taking insulin, and n = 434,779,439,791 for participants w/o diabetes. Estimated weighted percent has been abbreviated by “Wtd %”; SE stands for standard error.

Dietary Intake by Diabetic Status

684 **Table 8.** Broad What We Eat In America food and beverage category¹ intake comparisons by frequency and energy among U.S.
 685 adults 18 y or older taking insulin, with Type 2 diabetes but not taking insulin, and without diabetes drawn from NHANES 2009–
 686 2016²
 687

Broad WWEIA food category ¹	Insulin takers (n=774)		T2DM w/o insulin (n=2,758)		Participants without diabetes (n=17,796)		X ² P-Value	
	Wtd % ³ of reported foods (SE)	Wtd % ³ of reported energy (SE)	Wtd % ³ of reported foods (SE)	Wtd % ³ of reported energy (SE)	Wtd % ³ of reported foods (SE)	Wtd % ³ of reported energy (SE)	Frequency ⁴	Energy ⁴
Milk/Dairy ⁵	6.9 (0.4)	6.9 (0.5)	7.1 (0.2)	6.6 (0.3)	7.4 (0.1)	6.3 (0.1)	0.1	0.3
Protein ⁶	11.4 (0.5)	18.8 (1.0)	11.2 (0.2)	17.2 (0.4)	10.6 (0.1)	15.7 (0.2)	0.01	<0.0001*
Mixed Dish ⁷	6.9 (0.4)	22.2 (1.2)	6.9 (0.2)	21.7 (0.6)	7.5 (0.1)	22.7 (0.3)	0.004	0.3
Grain ⁸	9.7 (0.3)	15.2 (0.6)	9.6 (0.2)	14.3 (0.4)	8.8 (0.1)	12.1 (0.2)	<0.0001*	<0.0001*
Snack/Sweet ⁹	8.5 (0.4)	13.8 (0.8)	9.5 (0.2)	15.4 (0.3)	10.2 (0.1)	15.1 (0.2)	<0.0001*	0.2
Fruit ¹⁰	4.6 (0.3)	2.8 (0.2)	4.7 (0.2)	2.7 (0.1)	4.6 (0.1)	2.5 (0.1)	0.9	0.05
Vegetable ¹¹	11.0 (0.6)	6.1 (0.5)	11.0 (0.3)	6.3 (0.2)	10.4 (0.1)	5.4 (0.1)	0.1	0.0009*

Dietary Intake by Diabetic Status

Beverage ¹²	14.8 (0.3)	6.1 (0.4)	13.9 (0.3)	7.3 (0.3)	13.8 (0.1)	9.5 (0.2)	0.09	<0.0001*
Alcohol ¹³	0.9 (0.2)	2.4 (0.5)	1.2 (0.1)	2.9 (0.3)	2.1 (0.1)	5.0 (0.2)	<0.0001*	<0.0001*
Water ¹⁴	10.2 (0.4)	0.0 (0.0)	10.1 (0.4)	0.0 (0.0)	10.2 (0.2)	0.1 (0.0)	0.9	0.0011*
Fat/Oil ¹⁵	7.1 (0.4)	4.0 (0.3)	6.6 (0.2)	3.5 (0.1)	6.1 (0.1)	3.2 (0.1)	0.002	0.003
Cond ¹⁶ /Sauce ¹⁷	4.2 (0.3)	0.9 (0.1)	4.2 (0.2)	0.9 (0.1)	4.6 (0.1)	1.0 (0.0)	0.09	0.6
Sugars ¹⁸	3.8 (0.3)	0.8 (0.1)	3.8 (0.2)	1.1 (0.1)	3.4 (0.1)	1.2 (0.0)	0.05	0.08
Other	0.2 (0.1)	0.1 (0.0)	0.2 (0.0)	0.1 (0.0)	0.4 (0.0)	0.3 (0.0)	0.006	<0.0001*

688

689 ¹ The What We Eat in America broad food categories were applied to categorize all foods and beverages reported in a single day to 14
690 broad food groups;

691 ² Survey weights and adjustments for the complex survey design were applied to represent the non-institutionalized U.S. population.
692 Total numbers and percentages do not always add up to sample size due to missing values and rounding;

693 ³ Wtd % stands for the estimated weighted percent of all reports of foods or beverages or energy from reported foods or beverages
694 reported in a single day that are included in a food group;

695 ⁴ p-values were calculated using the Rao – Scott modified chi-square statistic; “*” indicates $p < 0.05/42$ or $p < 0.0012$ using a
696 Bonferroni type adjustment for multiple comparisons for food group intake among 14 broad food categories \times 3 groups;

697 ⁵ Milk, flavored milk, dairy drinks and substitutes, cheese and yogurt;

698 ⁶ Meats, poultry, seafood, eggs, cured meats/poultry, and plant-based protein foods;

699 ⁷ Mixed dishes containing meat, poultry seafood; grain-based; Asian; Mexican; pizza; sandwiches, and soups;

Dietary Intake by Diabetic Status

- 700 ⁸ Cooked grains, breads, rolls, tortillas, quick breads and bread products, ready-to-eat cereals, and cooked cereals;
- 701 ⁹ Savory snacks, crackers, snack/meal bars, sweet bakery products, candy and other desserts;
- 702 ¹⁰ Fresh fruits, dried fruits, and fruit salads;
- 703 ¹¹ Vegetables and white potatoes;
- 704 ¹² 100% juice, diet beverages, sweetened beverages, coffee and tea;
- 705 ¹³ Beer, wine, liquor and cocktails;
- 706 ¹⁴ Plain water and flavored or enhanced water;
- 707 ¹⁵ Butter and animal fats, margarine, cream cheeses, cream, mayonnaise, salad dressings and vegetable oils;
- 708 ¹⁶ Condiment;
- 709 ¹⁷ Tomato-based, soy-based, mustard, olives, pickled vegetables, pasta sauces, dips, gravies, and other sauces;
- 710 ¹⁸ Sugars, honey, jams, syrups, and toppings