

New England Journal of Medicine – Original Research

Alzheimer’s disease in retired elite collision sports athletes: cohort study

G. David Batty, PhD DSc (E. david.batty@ucl.ac.uk | ORCID: 0000-0003-1822-5753)
Department of Epidemiology and Public Health, University College London, London, UK

Urho M. Kujala, MD PhD (urho.m.kujala@fimnet.fi | 0000-0002-9262-1992)
Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland

Seppo J. Sarna, PhD (seppo.sarna@helsinki.fi | 0000-0003-3458-1627)
Department of Public Health, University of Helsinki, Helsinki, Finland

Carlos A. Valencia-Hernández, MD PhD (c.valencia-hernandez@imperial.ac.uk | 0000-0002-7596-4846)
National Heart & Lung Institute, Imperial College London, London, UK

Jaakko Kaprio, MD PhD (jaakko.kaprio@helsinki.fi | 0000-0002-3716-2455)
Institute for Molecular Medicine FIMM, University of Helsinki, Helsinki, Finland

Corresponding author: David Batty, Department of Epidemiology and Public Health, University College London, 1-19 Torrington Place, London, UK, WC1E 6BT. E. david.batty@ucl.ac.uk

Manuscript characteristics: 2678 words; 2 tables and 3 figures; 1 supplemental file

Contributions: GDB generated the idea for the paper, formulated the analytical plan, and drafted the manuscript. UMK and SJS designed data collection and edited the manuscript. CA V-H prepared the figures and edited the manuscript. JK designed data collection, formulated the analytical plan, conducted the data analyses, and edited the manuscript.

Conflict of interest: None to declare.

Acknowledgement: The preparation of this manuscript received no direct funding. GDB is supported by the UK Medical Research Council (MR/P023444/1) and the US National Institute on Aging (1R56AG052519-01; 1R01AG052519-01A1), and JK by the Academy of Finland Centre of Excellence in Complex Disease Genetics (336823).

Abstract

Background: Neuropathology consistent with chronic traumatic encephalopathy has been reported at autopsy in retired collision sports athletes, but whether these individuals also experience greater rates of dementia over their lifetime is uncertain.

Methods: Cohort study of 2005 male retired athletes who had competed internationally for Finland between 1920 and 1965, and 1386 male controls. At follow-up, data were captured via questionnaire administration and linkage to mortality, hospitalisation, and medication registries.

Results: A mean of 31 years of event surveillance (range 1-46) of a sample of 3391 men gave rise to 488 cases of neurodegenerative disease. In multiple-adjusted analyses, relative to the general population, boxing was associated with a quadrupling (4.10; 2.55, 6.61), and soccer (2.07; 1.23, 3.46) and wrestling (2.11; 1.28, 3.48) a doubling of the risk of Alzheimer's disease. Boxers with extended careers had twice the risk of Alzheimer's disease (2.07; 0.95, 4.51) relative to those who competed for shorter periods. For all collision sports, disease risk was most pronounced for those who competed after the second World War. Computation of competing risks and a series of sensitivity analyses did not appreciably alter these results.

Conclusions: In the present study, men with an elite-level background in soccer, wrestling, and boxing subsequently experienced poorer brain health than general population and non-collision sports controls.

Introduction

That former elite athletes have a longer life expectancy and a lower risk of cardiovascular disease relative to the general population is frequently used as evidence of the long-term health benefits of physical activity.¹⁻⁵ More recently, however, there has been suggestions of an increased burden of neurodegenerative disorders in retired athletes, in particular in those with a background in collision sports.^{1,4,6,7} This is consistent with autopsy evidence of neuropathology, including chronic traumatic encephalopathy, among some former ice hockey⁸ and American football⁹ players.

Although the occurrence of brain injury in collisions sports participants¹⁰ and the notion that this may lead to dementia in boxers was first mooted a century ago,¹¹ there has been few epidemiological investigations of brain health in former athletes. American football is perhaps the most scrutinised sport in this context^{1,12,13} but, strikingly, there is an absence of similar studies comparing rates of diagnosis in boxers or wrestlers relative to unexposed individuals. Moreover, for the most popular global collision sport, soccer, with one exception,⁴ existing reports universally focus on the risk of amyotrophic lateral sclerosis,^{5,14-16} a rare neurodegenerative disorder relative to the more burdensome dementias. Most studies in this field are also too underpowered to detect a relationship should one exist, and the use of a modest range of covariate data complicates the attribution of poor brain health to the sport in question rather than other risk factors.

We were able to address this gap in knowledge by utilising data from a well-characterised cohort study of men drawn from an array of elite sporting backgrounds entering their 11th decade.^{17,18} Given that boxing and wrestling are known to have a higher incidence of head trauma relative to other activities such as soccer,^{19,20} in this cross-sport comparison with the general population and non-collision sports groups, we anticipated that the occurrence of neurodegenerative disease, including dementia, would be highest in the combat sports.

Methods

This is an exposure-based cohort study originally established to examine the relationship between elite sports performance and longevity.^{21,22} Established in 1978, the sample was generated from 2613 Finnish male former elite athletes (640 Olympians) who competed internationally between 1920 and 1965 (supplemental figure 1). Identified using a combination of sports yearbooks, association records, and periodicals, these athletes represent a diverse array of sporting backgrounds. Based on the occurrence of concussion-causing head injury,^{19,20,23} we denoted sports as collision (soccer, boxing, Greco-Roman/classic wrestling, ice hockey, and basketball) or non-collision (track and field, cross-country skiing, and weight-lifting), with the former group disaggregated as the numbers of neurodegenerative disorders at follow-up allowed.

For the purposes of the present analyses, participants in the non-collision sports represented the unexposed athletic control group. A general population-based comparison group comprising apparently healthy men was also generated using records from a population-wide medical examination routinely administered to all Finnish males at around 20 years of age as part of national service (N=1712). Data collection was approved by the ethics committee of the Hospital Districts of Helsinki and Uusimaa, and all participants provided informed consent. The present manuscript conforms to the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement of guidelines for reporting observational studies.²⁴

Assessment of covariates

In 1985, surviving study members and population controls (N=2851; 66% of the original cohort) were mailed a self-completion questionnaire (N=1917; response 67%) (supplemental figure 1). In addition to enquiries regarding career duration, a series of covariate data were collected including smoking habit, alcohol intake (amount and pattern), marital status, sports participation in early adulthood, and height and weight from which body mass index was computed using the usual formulae ($\text{weight}[\text{kg}]/[\text{height}(\text{m})]^2$).²⁵

Men also responded to questions about their occupation and, in combination with data extracted from the Finnish Central Population Registry, longest held job was used as our indicator of socioeconomic status (executives [highest status], clerical staff, skilled workers, unskilled workers; or farmers).²⁶ Additionally, the presence of diabetes, hypertension, and coronary heart disease was derived from linkage of study members to the National Social Insurance Institution register which has records of individuals prescribed approved drug therapies for these and other conditions for the purposes of reimbursement.

Ascertainment of neurodegenerative disorders

Study members were linked to three health registries to ascertain the occurrence of a neurodegenerative disorders during follow-up: death, including cause; hospitalisations, including cause; and the Finnish Drug Prescription Register for medications specifically used to treat Alzheimer's disease (the Anatomical Therapeutic Chemical code observed were: N06DA02, donepezil; N06DA03, rivastigmine; N06DA04, galantamine; and N06DX01, memantine). We then generated five neurodegenerative disease outcomes for use in the present analyses (for the International Classification of Disease codes (versions 8-10), see supplemental table 1): all neurodegenerative disease combined, all dementias, Alzheimer's disease, non-Alzheimer's dementia, and Parkinson's disease. Deaths were based on any mention of a neurodegenerative disease on the death certificate. As demonstrated for other risk factors,^{27,28} and in the present analyses for collision sport participation, associations with neurodegenerative disorders were little-changed whether endpoint data were based on underlying cause only versus any mention on the death certificate (supplemental table 2), and whether we used deaths only (supplemental table 3) versus a combination of death, hospital admission, and drug treatment for Alzheimer's disease (main analyses).

Statistical analyses

After exclusion of study members owing to record-linkage failure and death prior to the beginning of follow-up (supplemental figure 1), the analytical sample comprised 3,391 men (2005 former athletes, 1386 population controls). Born 1884-1949 (mean 1925), event surveillance was from 1st January 1970 until the occurrence of a neurodegenerative disease event or the end of surveillance (31st December, 2015) –

whichever came first. Having ascertained that the proportional hazards assumption had not been violated, we used Cox regression to compute hazard ratios with accompanying 95% confidence intervals to summarise the relationship of sport participation with the risk of neurodegenerative disorder.²⁹ Age was used as the time covariate in statistical models that focused on former boxing, wrestling, and soccer athletes – there were insufficient degenerative disease endpoints in other sports to facilitate analyses. All analyses were computed using Stata 15 (StataCorp, College Station, TX).

Results

In general, relative to the general population, elite sports participants had somewhat more favourable baseline risk factor profiles (supplemental tables 4 and 5). These differentials were most consistently evident for socio-economic disadvantage and cigarette smoking. In contrast, heavy alcohol consumption was more common in the athletes. As anticipated, several of these baseline characteristics were related to an elevated rate of dementia (supplemental table 6), although statistical significance at conventional levels was not always apparent. Thus, there was around a 50% increase in the risk of future dementia if study members came from poor social circumstances (age-adjusted hazard ratio; 95% confidence interval: 1.70; 1.25, 2.31), had a physical morbidity such as diabetes (1.50; 1.15, 1.97) or hypertension (1.41; 1.15, 1.73), smoked cigarettes (1.41; 1.01, 1.98), or lost consciousness at least annually owing to heavy alcohol intake (1.69; 1.23, 2.33). There was, however, an absence of association for existing heart disease, being unmarried, and being overweight or obese.

Compared to the general population, individuals in the athlete groups typically had a lower risk of mortality from all-cause and cardiovascular disease (table 1). The only exception was for former boxers and we investigated this further by generating a mortality outcome that did not include neurodegenerative disease. In these analyses, boxing was also related to a lower risk of death, although statistical significance at conventional levels was not apparent for this combat sport as it was not for other athletic groups.

In the main analyses, a mean of 30.6 years of surveillance (range 1-46), corresponding to 103,768 person-years in the analytical sample of 3391 men, gave rise to 488 neurodegenerative disease events (265 ascribed to Alzheimer's disease, 333 to non-Alzheimer's disease, and 92 to Parkinson disease). Figures 1-3 (and supplemental tables 7-11) depict the occurrence of these outcomes in relation to collision sports participation. In age-adjusted analyses, boxers experienced three times the rate of neurodegenerative disorder combined relative to the general population (3.08; 2.31, 4.10) (supplemental table 7). While there was also a raised risk amongst soccer players (1.58; 1.15, 2.19) and a more modest relationship for wrestling (1.18; 0.84, 1.67), neurodegenerative disease was not convincingly related to participation in other collision sports. Adjusting for an array of potential explanatory factors which included socioeconomic status, comorbidity, and alcohol had modest impact on the point estimates for former boxers (3.11; 2.17, 4.45), soccer players (1.40; 0.93, 2.10), and wrestlers (1.28; 0.84, 1.95) (figure 1 and supplemental table 7).

The association of participation in boxing and soccer with neurodegenerative disease was largely generated by the link with dementias (supplemental table 8) and, of these, effects were strongest for Alzheimer's disease (figure 2 and supplemental table 9) relative to other subtypes (supplemental table 10). That is, in minimally-adjusted analyses, boxing was related to a more than quadrupling in the risk of later Alzheimer's disease (4.45; 3.07, 6.45), whereas the disease rates in wrestlers (1.90; 1.24, 2.90) and soccer players (2.43; 1.62, 3.64) was around double that of general population controls. When we took into account explanatory factors additional to age, the pattern of association was essentially unchanged (figure 2). That the association of collision sports with neurodegenerative disease were lower than that apparent for the dementias was due to the absence of association between any of the individual collision sports and Parkinson's disease (supplemental table 11).

We further explored the effects for specific collision sports and Alzheimer's disease. In analyses of career duration, there was a suggestion that boxers who fought competitively for 15 years or more had twice the rate of this disorder (2.07; 0.95, 4.51) relative to those with shorter exposures; no such differential was apparent for wrestling or soccer, however (supplemental table 12). We also investigated era-specific

changes to rules or equipment designed to protect the players might have an impact on the sports– Alzheimer’s disease relation. We in fact found a markedly stronger relation with Alzheimer’s disease for athletes who participated in each of the collision sports post-world war II relative to earlier (table 2). For instance, soccer players active 1948-1965 experienced around three times the risk of Alzheimer’s disease (2.96; 1.82, 4.82), whereas for those who competed earlier the corresponding risk was lower (1.43; 0.66, 3.12).

We also conducted some sensitivity analyses. First, owing to missing data, the more complex statistical models comprise fewer study members. We therefore recomputed the main effects in a non-missing dataset and results were essentially unchanged (supplemental table 13). Second, given the amateur era in which study members competed internationally (1920-1965), it is plausible that, if sufficiently skilled, they would have participated at elite level in more than one of the sports linked to neurodegenerative disorder in our analyses. Mutual control for each of the three sports associated with neurodegenerative disease did not, however, impact upon the existing pattern of association (supplemental table 14). Third, given the maturity of this cohort, we subjected the outcome of Alzheimer’s disease to a competing-risks analysis whereby the existence of the most common cause of death and hospitalisation herein (cardiovascular disease) precludes the occurrence of the primary event of interest (Alzheimer’s disease). There was no suggestion that this was the case (supplemental table 15). Lastly, it is plausible that the general population controls contained some elite sports people, although this misclassification would have had the impact of diminishing the risk of neurodegenerative disease in the elite sports groups. A total of 66 of the 765 men in the control group who returned a questionnaire in 1985 indicated they participated as young adults in one of three collision sports associated with neurodegenerative disease in the present study, but when they were excluded from the control group (supplemental table 16) there was again no impact on the results.

Discussion

Our main finding was that, relative to control groups comprising either non-collision sports participants or the general population, former elite-level collision athletes had a subsequently raised risk of

neurodegenerative disorders. This was particularly evident for Alzheimer's disease where boxers experienced a post-retirement quadrupling of risk and soccer players and wrestlers a doubling. These post-retirement elevated disease rates were found despite the contrasting head impact profiles of these activities.^{19,20} That is, concussions are more common in boxing and wrestling^{19 20} and result largely from person-on-person contact, whereas in soccer, the insult is more likely to be sub-concussive and attributable to contact with equipment – the ball – when attempting to redirect it using the head.

That boxers with extended careers had around twice the risk of Alzheimer's disease relative to those who competed for shorter periods may signal the additional harms of accumulated exposure, although the same was not apparent for former soccer players and wrestlers. We also found that, for all these sports, the risk of Alzheimer's disease was particularly pronounced in those men taking part post-war. Improved diagnosis of Alzheimer's disease will have led to more cases being identified in younger study members, although this would need to be differential with respect to the exposure for it to influence our associations. If the greater risk of collision sports post war is not artefactual, it could be ascribed to a more intensive regime of training and competition in the modern era, with a concomitant increase in the occurrence of head injury. Another possibility is risk compensation whereby an athlete engages in more damaging behaviour in response to a perception of improved protection from, for instance, the introduction of head gear in boxing.^{30,31}

That we were able to recapitulate several known risk factor–dementia relationships in this cohort for low socioeconomic status, comorbidities, and smoking³² gives us some confidence in these novel results for collision sports. Our findings are also supported by results from cohorts of people exposed to head injury of greater severity where an increased future burden of dementia has been reported in people with a history of hospitalised for traumatic brain injury typically due to vehicular accidents or falls.³³⁻³⁶ In the sporting realm, most empirical evidence comes from two groups: studies of former American football players and the occurrence of neurodegenerative disease, including dementia, with mixed results;^{1,12,13} and

amyotrophic lateral sclerosis in retired soccer players where disease risk seems to be consistently elevated.^{5,14,15}

In mechanistically-orientated research that may be brought to bear on our observational data, autopsy studies reveal chronic traumatic encephalopathy-type neuropathological change in former soccer players.³⁷ There is also evidence that heading produces an acute if temporary lowering of cognitive function,³⁸ while potentially more permanent structural changes to the brain are evident upon imaging.³⁹

Our study is not without its limitations. Our sample exclusively comprises men, and although we can think of no reason for the present results not applying to women given that other risk factors predict dementia equally well in sex-specific analyses,^{40,41} this clearly requires empirical testing. If the expected relationship does exist, the population impact may in fact be greater given the apparent higher prevalence of head injury, including concussion, in female athletes relative to males in the same sports.⁴² Whether a background in collision sports such as rugby union or league, lacrosse, basketball, and mixed martial arts, is also related to dementia risk is currently unknown – where we had some such participants in the present study, there was an insufficiently high number of disease events to facilitate analyses. There are reasons to anticipate that pre-adult exposure to head injury may be compensated by a greater degree of neurological plasticity;⁴³ equally it could be argued that the impact on brain health could be more damaging owing programming effects. We did not, however, have data on head injury in childhood or adolescence with which to test these propositions.

Participants in boxing represent a miniscule proportion of the population and, while dementia is a pernicious condition, the population impact of greater safety legislation in that sport will therefore be low despite the high dementia risk found herein. By contrast, wrestling is relative common at high school and collegiate level in selected countries, most notably the USA⁴⁴ and, with around a quarter of a billion participants, soccer is the most popular global sport⁴⁵ – the same order as the estimated number of people with a diagnosis of diabetes worldwide,⁴⁶ a recognised risk factor for dementia.³²

In conclusion, in the present study, a history of elite level engagement in soccer, wrestling, and boxing was related to later risk of Alzheimer's disease independently of many recognised risk factors. The global burden of soccer participation for brain health may be considerable.

References

1. Lehman EJ, Hein MJ, Baron SL, Gersic CM. Neurodegenerative causes of death among retired National Football League players. *Neurology* 2012;79(19):1970-4. (In eng). DOI: 10.1212/WNL.0b013e31826daf50.
2. Baron SL, Hein MJ, Lehman E, Gersic CM. Body mass index, playing position, race, and the cardiovascular mortality of retired professional football players. *Am J Cardiol* 2012;109(6):889-96. (In eng). DOI: 10.1016/j.amjcard.2011.10.050.
3. Lincoln AE, Vogel RA, Allen TW, et al. Risk and Causes of Death among Former National Football League Players (1986-2012). *Med Sci Sports Exerc* 2018;50(3):486-493. DOI: 10.1249/MSS.0000000000001466.
4. Mackay DF, Russell ER, Stewart K, MacLean JA, Pell JP, Stewart W. Neurodegenerative Disease Mortality among Former Professional Soccer Players. *N Engl J Med* 2019;381(19):1801-1808. (In eng). DOI: 10.1056/NEJMoa1908483.
5. Taioli E. All causes of mortality in male professional soccer players. *Eur J Public Health* 2007;17(6):600-4. (In eng). DOI: 10.1093/eurpub/ckm035.
6. Morales JS, Valenzuela PL, Saco-Ledo G, et al. Mortality Risk from Neurodegenerative Disease in Sports Associated with Repetitive Head Impacts: Preliminary Findings from a Systematic Review and Meta-Analysis. *Sports Med* 2021 (In eng). DOI: 10.1007/s40279-021-01580-0.
7. Blecher R, Elliott MA, Yilmaz E, et al. Contact Sports as a Risk Factor for Amyotrophic Lateral Sclerosis: A Systematic Review. *Global Spine J* 2019;9(1):104-118. (In eng). DOI: 10.1177/2192568218813916.
8. Schwab N, Wennberg R, Grenier K, Tartaglia C, Tator C, Hazrati LN. Association of Position Played and Career Duration and Chronic Traumatic Encephalopathy at Autopsy in Elite Football and Hockey Players. *Neurology* 2021;96(14):e1835-e1843. (In eng). DOI: 10.1212/wnl.0000000000011668.
9. Mez J, Daneshvar DH, Kiernan PT, et al. Clinicopathological Evaluation of Chronic Traumatic Encephalopathy in Players of American Football. *Jama* 2017;318(4):360-370. (In eng). DOI: 10.1001/jama.2017.8334.
10. Nichols EH, Smith HB. The physical aspect of American football. *The Boston Medical and Surgical Journal* 1906;154(1):1-8.
11. Martland HS. Punch drunk. *Journal of the American Medical Association* 1928;91(15):1103-1107.
12. Savica R, Parisi JE, Wold LE, Josephs KA, Ahlskog JE. High school football and risk of neurodegeneration: a community-based study. *Mayo Clin Proc* 2012;87(4):335-40. (In eng). DOI: 10.1016/j.mayocp.2011.12.016.
13. Janssen PH, Mandrekar J, Mielke MM, et al. High School Football and Late-Life Risk of Neurodegenerative Syndromes, 1956-1970. *Mayo Clin Proc* 2017;92(1):66-71. (In eng). DOI: 10.1016/j.mayocp.2016.09.004.
14. Belli S, Vanacore N. Proportionate mortality of Italian soccer players: is amyotrophic lateral sclerosis an occupational disease? *Eur J Epidemiol* 2005;20(3):237-42. (In eng). DOI: 10.1007/s10654-004-6879-7.
15. Pupillo E, Bianchi E, Vanacore N, et al. Increased risk and early onset of ALS in professional players from Italian Soccer Teams. *Amyotroph Lateral Scler Frontotemporal Degener* 2020;21(5-6):403-409. (In eng). DOI: 10.1080/21678421.2020.1752250.
16. Chio A, Calvo A, Dossena M, Ghiglione P, Mutani R, Mora G. ALS in Italian professional soccer players: the risk is still present and could be soccer-specific. *Amyotroph Lateral Scler* 2009;10(4):205-9. (In eng). DOI: 10.1080/17482960902721634.
17. Kujala UM, Sarna S, Kaprio J, Koskenvuo M. Hospital care in later life among former world-class Finnish athletes. *Jama* 1996;276(3):216-20. (In eng).
18. Kujala UM, Tikkanen HO, Sarna S, Pukkala E, Kaprio J, Koskenvuo M. Disease-specific mortality among elite athletes. *Jama* 2001;285(1):44-5. (In eng). DOI: 10.1001/jama.285.1.44.
19. Putukian M, D'Alonzo BA, Campbell-McGovern CS, Wiebe DJ. The Ivy League-Big Ten Epidemiology of Concussion Study: A Report on Methods and First Findings. *Am J Sports Med* 2019;47(5):1236-1247. DOI: 10.1177/0363546519830100.

20. Prien A, Grafe A, Rossler R, Junge A, Verhagen E. Epidemiology of Head Injuries Focusing on Concussions in Team Contact Sports: A Systematic Review. *Sports Med* 2018;48(4):953-969. DOI: 10.1007/s40279-017-0854-4.
21. Sarna S, Sahi T, Koskenvuo M, Kaprio J. Increased life expectancy of world class male athletes. *Med Sci Sports Exerc* 1993;25(2):237-44. (In eng).
22. Kujala UM, Kaprio J, Taimela S, Sarna S. Prevalence of diabetes, hypertension, and ischemic heart disease in former elite athletes. *Metabolism* 1994;43(10):1255-60. (In eng). DOI: 10.1016/0026-0495(94)90219-4.
23. Dick R, Agel J, Marshall SW. National Collegiate Athletic Association Injury Surveillance System commentaries: introduction and methods. *J Athl Train* 2007;42(2):173-82. (<https://www.ncbi.nlm.nih.gov/pubmed/21714302>).
24. von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet* 2007;370(9596):1453-7. DOI: 10.1016/S0140-6736(07)61602-X.
25. Status WECOP. The use and interpretation of anthropometry. Geneva: World Health Organisation, 1995.
26. Kujala UM, Sarna S, Kaprio J, Koskenvuo M. Asthma and other pulmonary diseases in former elite athletes. *Thorax* 1996;51(3):288-92. DOI: 10.1136/thx.51.3.288.
27. Batty GD, Gale CR, Kivimaki M, Bell S. Assessment of Relative Utility of Underlying vs Contributory Causes of Death. *JAMA Open Network* 2019.
28. Crews DE, Stamler J, Dyer A. Conditions other than underlying cause of death listed on death certificates provide additional useful information for epidemiologic research. *Epidemiology* 1991;2(4):271-5. (In eng). DOI: 10.1097/00001648-199107000-00006.
29. Cox DR. Regression models and life-tables. *J R Stat Soc [Ser B]* 1972;34:187-220.
30. Richens J, Imrie J, Copas A. Condoms and seat belts: the parallels and the lessons. *Lancet* 2000;355(9201):400-3. DOI: 10.1016/S0140-6736(99)09109-6.
31. Thompson DC, Thompson RS, Rivara FP. Risk compensation theory should be subject to systematic reviews of the scientific evidence. *Inj Prev* 2001;7(2):86-8. DOI: 10.1136/ip.7.2.86.
32. Livingston G, Huntley J, Sommerlad A, et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet* 2020;396(10248):413-446. (In eng). DOI: 10.1016/s0140-6736(20)30367-6.
33. Nordström A, Nordström P. Traumatic brain injury and the risk of dementia diagnosis: A nationwide cohort study. *PLoS Med* 2018;15(1):e1002496. (In eng). DOI: 10.1371/journal.pmed.1002496.
34. Fann JR, Ribe AR, Pedersen HS, et al. Long-term risk of dementia among people with traumatic brain injury in Denmark: a population-based observational cohort study. *Lancet Psychiatry* 2018;5(5):424-431. (In eng). DOI: 10.1016/s2215-0366(18)30065-8.
35. Raj R, Kaprio J, Korja M, Mikkonen ED, Jousilahti P, Siironen J. Risk of hospitalization with neurodegenerative disease after moderate-to-severe traumatic brain injury in the working-age population: A retrospective cohort study using the Finnish national health registries. *PLoS Med* 2017;14(7):e1002316. DOI: 10.1371/journal.pmed.1002316.
36. Raj R, Kaprio J, Jousilahti P, Korja M, Siironen J. Risk of Dementia After Hospitalization Due to Traumatic Brain Injury: A Longitudinal, Population-Based Study. *Neurology* 2022. DOI: 10.1212/WNL.000000000000200290.
37. Lee EB, Kinch K, Johnson VE, Trojanowski JQ, Smith DH, Stewart W. Chronic traumatic encephalopathy is a common co-morbidity, but less frequent primary dementia in former soccer and rugby players. *Acta Neuropathol* 2019;138(3):389-399. (In eng). DOI: 10.1007/s00401-019-02030-y.
38. Di Virgilio TG, Hunter A, Wilson L, et al. Evidence for Acute Electrophysiological and Cognitive Changes Following Routine Soccer Heading. *EBioMedicine* 2016;13:66-71. (In eng). DOI: 10.1016/j.ebiom.2016.10.029.
39. Lipton ML, Kim N, Zimmerman ME, et al. Soccer heading is associated with white matter microstructural and cognitive abnormalities. *Radiology* 2013;268(3):850-7. DOI: 10.1148/radiol.13130545.

40. Gong J, Harris K, Peters SAE, Woodward M. Sex differences in the association between major cardiovascular risk factors in midlife and dementia: a cohort study using data from the UK Biobank. *BMC Med* 2021;19(1):110. DOI: 10.1186/s12916-021-01980-z.
41. Chatterjee S, Peters SA, Woodward M, et al. Type 2 Diabetes as a Risk Factor for Dementia in Women Compared With Men: A Pooled Analysis of 2.3 Million People Comprising More Than 100,000 Cases of Dementia. *Diabetes Care* 2016;39(2):300-7. DOI: 10.2337/dc15-1588.
42. McGroarty NK, Brown SM, Mulcahey MK. Sport-Related Concussion in Female Athletes: A Systematic Review. *Orthop J Sports Med* 2020;8(7):2325967120932306. (In eng). DOI: 10.1177/2325967120932306.
43. Viner RM, Ross D, Hardy R, et al. Life course epidemiology: recognising the importance of adolescence. *J Epidemiol Community Health* 2015;69(8):719-20. DOI: 10.1136/jech-2014-205300.
44. Association NCA. Sports Sponsorship and Participation Rates Database (<https://www.ncaa.org/sports/2018/10/10/ncaa-sports-sponsorship-and-participation-rates-database.aspx> - retrieved March 2022)2021.
45. Hulteen RM, Smith JJ, Morgan PJ, et al. Global participation in sport and leisure-time physical activities: A systematic review and meta-analysis. *Preventive medicine* 2017;95:14-25.
46. Saeedi P, Petersohn I, Salpea P, et al. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9(th) edition. *Diabetes Res Clin Pract* 2019;157:107843. DOI: 10.1016/j.diabres.2019.107843.

Table 1. Association of participation in collision sports with mortality

	Number of total deaths / Number at risk	All-causes	Non-neurodegenerative disorders	Cardiovascular disease
		Hazard ratios (95% CI)	Hazard ratios (95% CI)	Hazard ratios (95% CI)
Boxing	188/230	1.14 (0.97, 1.33)	0.93 (0.78, 1.11)	1.11 (0.88, 1.39)
Wrestling	188/247	0.87 (0.74, 1.02)	0.86 (0.73, 1.02)	0.98 (0.79, 1.22)
Soccer	175/248	0.87 (0.74, 1.03)	0.78 (0.65, 0.93)	0.86 (0.68, 1.09)
Other collision sports	117/236	0.58 (0.48, 0.70)	0.57 (0.46, 0.70)	0.56 (0.42, 0.75)
Non-collision sports	756/1044	0.75 (0.66, 0.82)	0.74 (0.66, 0.82)	0.77 (0.67, 0.89)
General population (controls)	1005/1386	1.0 (ref)	1.0 (ref)	1.0 (ref)


Hazard ratios are age-adjusted. CI, confidence interval

Table 2. Association of participation in collision sports with Alzheimer’s disease according to era of competition

	Hazard ratios (95% CI)	
	1909-1947	1948-1965
Number of events / Number at risk	82/1417	183/1974
Boxing	2.72 (1.42, 5.20)	6.53 (4.12, 10.36)
Wrestling	1.05 (0.46, 2.38)	2.15 (1.29, 3.58)
Soccer	1.43 (0.66, 3.12)	2.96 (1.82, 4.82)
Other collision sports	0.33 (0.04, 2.40)	0.79 (0.40, 1.53)
Non-collision sports	0.74 (0.43, 1.28)	0.81 (0.52, 1.25)
General population (controls)	1.0 (ref)	1.0 (ref)


Hazard ratios are age-adjusted. CI, confidence interval

Figure 1. Association of participation in collision sports with all neurodegenerative disorders




Hazard ratios (HR) are adjusted for age, socioeconomic status, co-morbidities (diabetes, hypertension, and coronary heart disease), and lifestyle factors (alcohol intake, passing out due to alcohol, body mass index, and smoking status). Analytical sample is based on study members with questionnaire data and follow-up starting 1986.

Figure 2. Association of participation in collision sports with Alzheimer's disease



Hazard ratios (HR) are adjusted for age, socioeconomic status, co-morbidities (diabetes, hypertension, and coronary heart disease), and lifestyle factors (alcohol intake, passing out due to alcohol, body mass index, and smoking status). Analytical sample is based on study members with questionnaire data and follow-up starting 1986.

Figure 3. Association of participation in collision sports with non-Alzheimer’s dementia



Hazard ratios (HR) are adjusted for age, socioeconomic status, co-morbidities (diabetes, hypertension, and coronary heart disease), and lifestyle factors (alcohol intake, passing out due to alcohol, body mass index, and smoking status). Analytical sample is based on study members with questionnaire data and follow-up starting 1986.