

Estimating the replicability of highly cited clinical research (2004-2018)

Gabriel Gonçalves da Costa (ORCID: 0000-0002-1141-7827), Kleber Neves (ORCID: 0000-0001-9519-4909), Olavo B. Amaral (ORCID: 0000-0002-4299-8978)

Affiliations:

Institute of Medical Biochemistry Leopoldo de Meis, Federal University of Rio de Janeiro

Funding:

G.G.C. receives funding from CNPq. O.B.A. receives funding from FAPERJ (E-26/200.824/2021), CNPq (308624/2018-1) and the Serrapilheira Institute.

Correspondence: Gabriel Gonçalves da Costa (gabriel.costa@bioqmed.ufrj.br)

Author Contributions:

Conceptualization: All authors

Methodology: All authors

Project Administration: All authors

Investigation (data collection): GGC and KN

Writing – Original Draft: GGC

Writing – Review and Editing: OBA and KN

Supervision: OBA

Conflicts of Interest:

None declared.

Abstract

Introduction: Previous studies about the replicability of clinical research based on the published literature have suggested that highly cited articles are often contradicted or found to have inflated effects. Nevertheless, there are no recent updates of such efforts, and this situation may have changed over time.

Methods: We searched the Web of Science database for articles studying medical interventions with more than 2000 citations, published between 2004 and 2018 in high-impact medical journals. We then searched for replications of these studies in PubMed using the PICO framework. We evaluated whether replications were successful by the presence of a statistically significant effect in the same direction and by overlap of their effect sizes confidence intervals (CIs) with those of the original studies. We also analyzed evidence of effect size inflation and potential predictors or replicability.

Results: We found a total of 89 eligible studies, of which 24 had valid replications. Of these, 21 (88%) had effect sizes with overlapping CIs. Of 15 highly cited studies with a statistically significant difference in the primary outcome, 13 (87%) had a significant effect in the replication as well. When both criteria were considered together, the replicability rate in our sample was 83%. When comparing effect sizes, there was little evidence of systematic inflation in these highly cited studies. Due to the small number of contradicted results, our analysis was underpowered to detect predictors of replicability.

Conclusion: Although most studies did not have eligible replications, the replicability rate of highly cited clinical studies in our sample was higher than previous estimates, with little evidence of systematic effect size inflation.

Introduction

The replicability of published research has been recently questioned in different scientific fields, with replication rates shown to be variable and often low (1–8). Whether this represents a “reproducibility crisis” is open to debate (9), as defining what constitutes a successful replication is not trivial (10). Moreover, systematic replication efforts have mostly focused on restricted samples of the literature, and data on the subject is still lacking in most areas of science.

The replicability of highly cited clinical research was studied by Ioannidis in 2005, based on available published replications of a sample of articles between 1990 and 2003 (11). The study focused on the reproducibility of study conclusions, typically assessed by statistical significance, as well as on effect size comparisons between studies. It concluded that 44% of highly cited studies in the sample had been successfully replicated, 16% had been contradicted, 16% had found effects that were larger than those of subsequent studies, and 24% remained unchallenged.

A similar effort for highly cited psychiatry research between 2000 and 2002 found lower estimates, with 19% of studies replicated, 19% contradicted, 13% initially inflated and 48% unchallenged (12). Another study on critical care research found that 18% of interventions published in high-profile journals between 1946 and 2016 had their results replicated by a subsequent study, whereas 22% were contradicted, 2% had replications in progress and 58% remained unchallenged (13).

Clinical research has changed in some aspects over the last two decades. *A priori* registration of study protocols has become more common and mandatory for clinical trials in many countries (23–25). Although publication bias has not been eliminated (26), the likelihood of null results has increased (27). Reporting guidelines have become more widely used and underwent reevaluations and updates (28,29). The push for full reporting of results and availability of individual patient data has also gained ground (23,24). Thus, the replicability panorama in high-impact clinical research may have changed during this period (30–32).

The goal of this study is to estimate the replicability of highly cited clinical studies published between 2004 and 2018. Our primary outcome is the rate of successful replication in these studies, as measured both by statistical significance in the same direction and by overlap of CIs for the main effect measure in both studies. We also explore effect size inflation and potential predictors of replicability in this sample of articles.

Methods

An overview of the project, datasets and analysis code can be found at <https://osf.io/a8zug/>. The protocol for the study was preregistered at <https://osf.io/nh965/>, with a step-by-step methodology available at <https://osf.io/2qncz/> and updates and amendments described at

<https://osf.io/26d98/>. All statistical analyses were performed in R version 4.1.2 (34). Data and analysis code are available at <https://osf.io/m6djb/>.

Search for highly cited studies

We searched the Web of Science database for articles with more than 2000 citations published between January 1st, 2004 and December 31st, 2018 in medical journals with an impact factor above 14 in the 2020 Journal Citation Reports (see list at <https://osf.io/2qncz/>). General journals were searched on February 7th, 2020, and specialty journals on March 4th, 2020. The cutoffs for citation and impact factors were twice as large as those used by Ioannidis (11), accounting for the growth in the total number of articles in PubMed during the period (<https://osf.io/t9xu7/> for details).

Within this sample, one author (K.N.) screened titles and abstracts for articles that addressed the efficacy of therapeutic or preventive interventions with primary data (i.e. excluding reviews, meta-analyses or articles that combined two or more previous studies). Two evaluators (G.G.C. and K.N.) then selected the primary outcome in each study, or the main conclusion in the abstract if the study had no primary outcome. In the case of co-primary outcomes or equally emphasized conclusions (35–37), we chose the outcome that was deemed more clinically relevant (e.g. mortality over progression or neurologic improvement over reperfusion). In the case of trials with more than two arms (38), we selected the most effective drug as the intervention and randomly chose an active comparator among the other arms. Studies with no control group (e.g. phase 1 studies) were also considered eligible when the abstract clearly stated that an intervention was clinically effective. When both benefits and harms or caveats were presented, focus was given on the net conclusion of whether the experimental intervention merited consideration for use in clinical practice. Disagreements in outcome selection were solved by consensus with the help of a third investigator (O.B.A.).

For each article, we extracted the study design, sample size, journal name and category (general or specialty). We also extracted the selected outcome measure – i.e. odds ratio (OR), relative risk (RR), hazard ratio (HR) or objective response rate (ORR) – with its effect size and respective CI. For controlled studies, results were classified as positive or negative according to the authors' stated statistical significance threshold. Non-inferiority trials were classified as positive only if the intervention was found to be superior (e.g. not merely non-inferior) to the comparator.

For each result, we extracted the population, intervention, comparator and outcome (PICO) (39), both in specific (e.g. “myocardial infarction, ischemic stroke, unstable angina, or cardiovascular surgery”) and general forms (e.g. “cardiovascular events”) (23). Two evaluators (K.N. and G.G.C.) described PICO components independently and resolved disagreements by consensus. Results of the independent extraction of the PICO components and consensus decisions can be found at <https://osf.io/sfdxv/>.

Search for replications

After agreement was reached on PICO components, two evaluators (G.G.C. and K.N.) performed independent searches for replications of highly cited studies in PubMed. Search terms were defined independently by each evaluator and included the name of the drug or intervention, the general form of the outcome, and the population (i.e. clinical condition) as described in the article's title, along with corresponding MeSH terms. The comparator was included in the search strategy only if it was an active intervention (i.e. not a placebo or sham). A spreadsheet containing the terms used in the searches, along with the dates when they were performed, can be found at <https://osf.io/zv65u/>.

A study was considered a replication of the highly cited study when it shared the same PICO general components, namely (a) the drug or intervention, without considering dose or regimen (except for studies performing dose or regimen comparisons), (b) the general form of the outcome (as described in 23), (c) the population/clinical condition as described in the highly cited article's title and (d) the comparator. When geographical information was included as a descriptor of the population (e.g. "European patients"), we did not include this information as part of the population component.

Replications were included for analysis when they were either (a) a study type with higher strength of evidence (40) (i.e. randomized controlled trials (RCTs) over cohort studies over uncontrolled studies) or (b) a similar study type with a sample size equal to or larger than the original study. Meta-analyses were considered as eligible replications if the highly cited study accounted for less than half of its sample size. For network meta-analyses, only the sample size of the direct comparison between the intervention and comparator was considered for this purpose. If more than one replication was found, the one with the largest sample size for the specific comparison was considered. For each included replication, we recorded the same information extracted from highly cited studies.

Replications selected by both evaluators in this first step were automatically included. When different replications were selected by each evaluator, both were made available for the two evaluators to choose the best option independently. Disagreements in this step were solved by consensus with the participation of a third author (O.B.A). Agreement in the initial selection was 36%, but rose to 91% when selected replications were made available to both evaluators. A detailed spreadsheet containing the studies selected by both evaluators in each step can be found at <https://osf.io/qz6u9/>.

After identifying the best available replication, both evaluators independently selected the outcome and effect size from the replication that corresponded most closely to the one in the original study. Disagreements were solved by consensus among all three authors. For network meta-analyses, direct comparisons were favored over indirect ones when both were available. A spreadsheet detailing the outcomes selected by both evaluators and consensus decisions in this stage can be found at <https://osf.io/2c6jx/>.

Evaluating Replication Success

Pairs of highly cited studies and their replications were analyzed to evaluate whether results were successfully replicated on the basis of two criteria: (a) statistical significance (an effect in the replication with $p < 0.05$ in the same direction as that observed in the highly cited study) and (b) confidence interval overlap (an overlap of the CIs for the outcome of interest in both studies). When the highly cited study presented a non-significant effect or did not include a statistical comparison (e.g. phase 1 trials), only the second criterion was used. The primary outcome was the rate of successful replication in our sample by both criteria (or by confidence interval overlap alone when statistical significance was not applicable). As additional criteria, we analyzed whether the replication point estimate was contained in the CI of the highly cited study and vice versa. We also performed a sensitivity analysis applying the statistical significance criterion to initially non-significant studies as well. In one case where the replication was a Bayesian meta-analysis, we compared the original study's confidence interval with the replication's credible interval, as the meta-analysis used a minimally informative prior (41).

When outcome measures differed between highly cited studies and replications (e.g., RR vs. OR) and the replication was a primary study, the replication measure was converted to the one in the highly cited study using the data available in the article. When the replication was a meta-analysis that included the highly cited study, we chose the risk measure that was used for data synthesis, using the original study's effect size as included in the meta-analysis. If the highly cited study was not included in the meta-analysis (as in the case when the meta-analysis included a follow-up), we manually converted the outcome measure of the highly cited study to the one in the meta-analysis using the original data. When the original study was a phase 1 study measuring ORR, we manually calculated this measure in the replication, whether it was an RCT or meta-analysis, with CIs based on the Clopper-Pearson exact method. In the meta-analysis by Hamid et al. (42), ORRs for both RCTs that were eligible replications of the highly cited study (a phase 1 trial) were calculated manually based on the combined data. These calculations can be found at <https://osf.io/mfwv2/>.

Based on the primary outcome, studies were classified as replicated if both criteria were met, or contradicted if at least one of them was not met. For studies not reporting a statistical comparison (e.g. phase 1 trials) or those in which the original study reported a non-significant result, only the confidence interval overlap criterion was used to define replication success. The replicability rate was defined as the percentage of studies that were replicated successfully by each criterion among all studies in which that criterion was applicable.

Effect size inflation

The effect sizes of highly cited studies and replications were used to estimate effect size inflation ratios. For unfavorable outcomes (e.g., death, tumor progression), in which effectiveness increases as the outcome measure decreases, the inflation ratio was defined as the point estimate of the replication divided by that of the original study. For

favorable outcomes (e.g., neurologic improvement), in which effectiveness increases along with the outcome measure, it was defined as the point estimate of the original study divided by that of the replication. Order of publication was considered in this calculation: thus, when the replication was a meta-analysis in which the pooled sample size of studies preceding the highly cited study was larger than that of those that followed it, we inverted the ratios, considering the highly cited study as the replication and vice-versa for this purpose. CIs for mean effect size inflation were calculated by the Wilson score interval.

As these two necessary adjustments had not been pre-specified in the protocol, we performed sensitivity analyses using different ways to deal with positive/negative outcomes and study order within meta-analyses. For the former, coining of the effects was performed to convert all favorable outcomes to unfavorable outcomes: overall response rates were subtracted from one ($1 - \text{ORR}$), odds ratios were inverted ($1/\text{OR}$) and relative risks for the complementary outcome were calculated based on the original data (see <https://osf.io/5tmus/> and <https://osf.io/7wtr8/>). For the latter, we analyzed data considering the highly cited study as the original one, independent of study order in the meta-analysis.

For analysis of effect size inflation, we used log-transformed inflation ratios for each study pair (including those with initially negative results) to calculate the mean and CI of these ratios both for the whole sample and for phase 1 studies and RCTs separately. For the whole sample, we performed a one-sample t-test against a theoretical mean of 0, which would indicate absence of systematic inflation. Although we performed these calculations using log-transformed values to correct for the inherent asymmetry of ratios, we transformed means and CIs back to a linear scale for the sake of clarity when describing results.

Predictors of Replicability

Finally, we analyzed if studies with contradicted results – i.e., those failing in one or both replication criteria – differed from successfully replicated ones in the following aspects: (a) study design (RCTs vs. other designs); (b) nature of intervention, (pharmacological vs. non-pharmacological); (c) sample size; (d) p-value of the original study; and (e) citations per year. To compare these aspects between replicated and contradicted studies, we used Fisher's exact test for categorical variables (a and b) and Mann-Whitney's U test for continuous variables (c through e). We had also planned to use the effect size of the highly cited study as a predictor, but due to the heterogeneity in outcome measures, which included both proportions (i.e. overall response rates) and measures of association (i.e. odds ratios, relative risks and hazard ratios) converting them to a single effect size measure turned out to be unfeasible.

Results

Results from our systematic search of the literature are shown as a flowchart in **Figure 1**. A total of 89 highly cited studies met our inclusion criteria. Of these, 24 had an eligible replication according to our criteria.




Figure 1. PRISMA flowchart. 89 eligible highly cited studies were found, of which 26 had an eligible replication. A complete list of the studies can be found at <https://osf.io/ub38r/>. A more detailed list of reasons for inclusion/exclusion is available at <https://osf.io/ma9bn/>.

As shown on **Table 1**, included studies received a median of 2842 citations, and were mostly RCTs of pharmacological interventions in cancer or heart disease, with some phase 1 cancer trials as well.

Description		Total (n = 89)	%	Replication found (n = 24)	%
Highly cited study design	RCT	81	91%	17	71%
	Phase 1 trial	7	8%	7	29%
	Cohort study	1	1%	0	0%
Type of intervention	Pharmacological	72	81%	16	67%
	Non- pharmacological	17	19%	8	33%
Journal	<i>New England Journal of Medicine</i>	80	90%	22	92%
	<i>Lancet</i>	4	4%	0	0%
	<i>Lancet Oncology</i>	4	4%	2	8%
	<i>JAMA</i>	1	1%	0	0%
Condition	Cancer	53	60%	15	63%
	Cardiovascular	27	30%	8	33%
	Other	9	10%	1	4%
Citations	< 3,000	45	51%	13	54%
	3,000 - 4,000	26	29%	8	33%
	4,000 - 6,000	12	13%	1	4%
	> 6,000	6	7%	2	8%

Table 1. Features of highly cited studies. Columns show the numbers of highly cited studies and of those for which an eligible replication was found in each category. Percentages refer to the total number of highly cited studies (n=89) or studies with replications (n=24), respectively.

Most replications were meta-analyses, followed by network meta-analyses, RCTs and 1 phase 2 trial (**Table 2**), with RCTs more commonly representing replications of phase 1 trials. All phase 1 trials had available replications in the literature, while the only cohort study in our sample had no eligible replication.

Design of highly cited study	Replication study design				
	Replication found (%)	Phase 2 Trial	RCT	Meta-analysis	Network meta-analysis
Cohort study	0/1 (0%)	0	0	0	0
Phase 1 trial	7/7 (100%)	1	3	2	1
RCT	17/81 (21%)	0	1	11	5
Total	24/89 (27%)	1	4	13	6

Table 2. Features of replications. Percentages refer to the total number of highly cited studies in each category. For studies with a replication, the table describes the type of study; most studies, however, did not have a valid replication.

A list of claims from highly cited studies is shown in **Table 3**. With few exceptions, most of them made clear claims of efficacy in their abstracts. Efficacy in phase 1 studies was measured by ORR (n=7), while differences in outcomes in RCTs were measured by HR (n=8), RR (n=6) or OR (n=3). All phase 1 trials made clear claims of efficacy based on tumor regression. Among RCTs, 2 were negative, with a p-value above the standard 0.05 cutoff.

Highly Cited Study	Replication	PICO General Components	Original conclusion	Effect Measure	Highly Cited ES [95% CI]	Replication ES [95%CI]
Brahmer et al., 2012 (43)	Zhang et al., 2016 (44)	Advanced cancer, Nivolumab, NA, Response	Nivolumab induced tumor regression and prolonged stabilization of disease in advanced cancers	ORR	0.13 [0.06-0.17]	0.27 [0.21-0.33]
Topalian. et al. 2012 (45)	Tie et al., 2017 (46)	Cancer, Nivolumab, NA, Tumour Response	Nivolumab produced objective responses in cancer regression.	ORR	0.21 [0.16-0.26]	0.26 [0.21-0.31]
TAXUS-IV (Stone et al., 2004) (47)	Bangalore et al., 2013 (48)	Coronary artery disease, Paclitaxel stent, Metal stent, Revascularization	A paclitaxel-eluting stent reduced the rate of clinical and angiographic restenosis at nine months	RR	0.39 [0.26-0.59]	0.66 [0.59-0.74]
SYNTAX (Serruys et al., 2009) (49)	Ali et al., 2018 (50)	Severe coronary artery disease, Percutaneous coronary intervention (PCI), Coronary-artery bypass grafting (CABG), Cardiovascular and cerebrovascular events	CABG resulted in lower rates of major adverse cardiac or cerebrovascular events than PCI after 1 year	OR	1.44 [1.11-1.89]	1.42 [1.27-1.59]
HERA (Piccart-Gebhart et al., 2005) (51)	Genuino et al., 2019 (52)	HER2-positive breast cancer after adjuvant chemotherapy, Trastuzumab, Observation, Progression	Trastuzumab after adjuvant chemotherapy improved disease-free survival in HER2-positive breast cancer	HR	0.54 [0.43-0.67]	0.65 [0.55-0.75]
CATIE (Lieberman et al., 2005) (38)	Soares-Weiser et al., 2013 (53)	Schizophrenia, Olanzapine, Quetiapine, Treatment discontinuation	Time to discontinuation of treatment for any cause was longer for olanzapine than for quetiapine	HR	0.63 [0.52-0.76]	0.68 [0.56-0.83]
SHARP (Llovet et al., 2008) (54)	Niu et al., 2016 (55)	Advanced hepatocellular carcinoma, Sorafenib, Placebo, Survival	Survival was longer with sorafenib than with placebo	HR	0.69 [0.55-0.87]	0.69 [0.60-0.79]
ERSPC (Schröder et al., 2009) (56)	Ilic et al., 2018 (57)	Middle- to old-age men, PSA screening, Control (no screening), Prostate cancer death	Periodic PSA-based screening reduced the rate of death from prostate cancer after a median follow-up of 9 years	RR	0.80 [0.65-0.98]	0.96 [0.85-1.08]

Highly Cited Study	Replication	PICO General Components	Original conclusion	Effect Measure	Highly Cited ES [95% CI]	Replication ES [95%CI]
PROFILE 1007 (Shaw et al., 2013) (58)	Elliott et al., 2020 (59)	Advanced ALK-positive lung cancer, Crizotinib, Chemotherapy, Progression	Crizotinib was superior to standard chemotherapy in preventing cancer progression	HR	0.49 [0.37-0.64]	0.46 [0.39-0.54]
ACCORD (Action to Control Cardiovascular Risk in Diabetes Study Group et al., 2008) (60)	Fang et al., 2016 (61)	Type 2 diabetes, Intensive glucose control, Standard therapy, Cardiovascular events or cardiovascular death	As compared with standard therapy, intensive therapy did not reduce major cardiovascular events after a mean follow-up of 3.5 years	RR	0.95 [0.82-1.09]	0.92 [0.85-1.00]
Cheng et al., 2009 (62)	Niu et al., 2016 (55)	Advanced hepatocellular carcinoma, Sorafenib, Placebo, Survival	Sorafenib increased survival in advanced hepatocellular carcinoma when compared with placebo	HR	0.68 [0.50-0.93]	0.69 [0.54-0.87]
EXTEND-IA (Campbell et al., 2015) (37)	Goyal et al., 2016 (63)	Ischemic stroke, Endovascular thrombectomy + Alteplase, Alteplase, Disability	Early thrombectomy with the Solitaire FR stent retriever improved disability in ischemic stroke as compared with alteplase alone	OR	6 [2-18]	4.04 [2.75-5.93]
PARTNER A (Smith et al., 2011) (64)	Pagnesi et al., 2017 (65)	Aortic stenosis, Transcatheter aortic-valve implantation (TAVI), Surgical replacement, Mortality	Transcatheter and surgical procedures for aortic valve replacement were associated with similar rates of survival at 1 year in high-risk patients	RR	0.98 [0.75-1.26]	0.86 [0.70-1.04]
MR CLEAN (Berkhemer et al., 2015) (66)	Rodrigues et al., 2016 (67)	Acute ischemic stroke, Intraarterial treatment + Usual care, Usual care, Disability	Intraarterial treatment administered within 6 hours after stroke onset was effective in reducing disability assessed at 90 days post-intervention.	RR	1.73 [1.27-2.35]	1.37 [1.14-1.64]
ECASS III (Hacke et al., 2008) (68)	Wardlaw et al., 2012 (69)	Acute ischemic stroke, Alteplase, Placebo, Disability	Intravenous alteplase improved disability in patients with acute ischemic stroke assessed 90 days after the intervention	OR	1.34 [1.02-1.76]	1.29 [1.16-1.43]

Highly Cited Study	Replication	PICO General Components	Original conclusion	Effect Measure	Highly Cited ES [95% CI]	Replication ES [95%CI]
EURTAC (Rosell et al., 2012) (70)	Zhao et al., 2019 (71)	Advanced EGFR mutation-positive non-small-cell lung cancer, Erlotinib, Standard chemotherapy, Progression	Erlotinib increased progression-free survival when compared to standard chemotherapy in Asian and European patients with advanced EGFR mutation-positive non-small-cell-lung cancer	HR	0.37 [0.25-0.54]	0.33 [0.28-0.40]
ESCAPE (Goyal et al., 2015) (72)	Rodrigues et al., 2016 (67)	Ischemic stroke, Standard care + Endovascular treatment, Standard care, Disability	Rapid endovascular treatment (thrombectomy) improved functional outcomes in up to 12 hours after symptom onset	RR	1.86 [1.39-2.47]	1.37 [1.14-1.64]
NEJSG (Maemondo et al., 2010) (73)	Greenhalgh et al., 2016 (74)	Non-small-cell lung cancer with mutated EGFR, Gefitinib, Carboplatin-paclitaxel, Progression	First-line gefitinib improved progression-free survival as compared with standard chemotherapy in patients with non-small-cell lung cancer with EGFR mutations.	HR	0.32 [0.22-0.41]	0.39 [0.32-0.48]
Hamid et al., 2013 (42)	Pyo et al., 2017 (75)	Melanoma, Lambrolizumab, NA, Tumor response	In advanced melanoma, treatment with lambrolizumab resulted in a high rate of sustained tumor regression	ORR	0.38 [0.25-0.44]	0.29 [0.26-0.32]
KEYNOTE-024 (Reck et al., 2016) (76)	KEYNOTE-042 (Mok et al., 2019) (77)	PD-L1-positive non-small-cell lung cancer, Pembrolizumab, Chemotherapy, Survival	Pembrolizumab led to longer overall survival than platinum-based chemotherapy for PD-L1-positive non-small-cell lung cancer	HR	0.50 [0.37-0.68]	1.07 [0.94-1.21]
KEYNOTE-001 (Garon et al., 2015) (78)	KEYNOTE-010 (Herbst et al., 2016) (79)	Non-small-cell lung cancer, Pembrolizumab, NA, Tumor Response	Pembrolizumab showed antitumor activity and led to objective responses in patients with advanced non-small-cell lung cancer	ORR	0.19 [0.16-0.23]	0.25 [0.22-0.29]
Wolchock et al., 2013 (80)	Checkmate 067 (Larkin et al., 2015) (36)	Advanced Melanoma, Nivolumab + Ipilimumab, NA, Response	Concurrent therapy with nivolumab and ipilimumab led to tumor regression in a substantial proportion of patients	ORR	0.40 [0.27-0.55]	0.44 [0.38-0.49]
Flaherty et al., 2010 (81)	BRIM-3 (Chapman et al., 2011) (35)	Metastatic melanoma with activated BRAF, PLX4032 (Vemurafenib), NA, Response	Treatment of metastatic melanoma carrying the V600E BRAF mutation with PLX4032 (Vemurafenib) resulted in tumor regression	ORR	0.81 [0.63-0.93]	0.48 [0.42-0.55]

Highly Cited Study	Replication	PICO General Components	Original conclusion	Effect Measure	Highly Cited ES [95% CI]	Replication ES [95%CI]
Fong et al., 2009 (82)	Kaufman et al., 2015 (83)	BRCA1- or BRCA2-mutated resistant cancers, Olaparib (AZD2281), NA, Tumor Response	Olaparib had antitumor activity and led to objective responses in cancers associated with BRCA1 or BRCA2 mutations	ORR	0.47 [0.24-0.71]	0.26 [0.21-0.32]

Table 3. Summary of the 24 highly cited studies with replications. Table shows the references for both studies, the general PICO components, the original study's conclusion, the outcome measure and the effect sizes (with CIs) found in the original study and replication.

When a meta-analysis was selected as a replication, the highly cited study could come after some (or most) of the studies in the meta-analysis, and thus consist of a replication of previous literature itself. This is analyzed in **Figure 2**, which shows the relative sample sizes of the highly cited study and of the sum of studies that precede or follow it. On average, highly cited studies corresponded to 22% of the total sample size of the meta-analyses, but this number ranged from 6% to 47%. Most meta-analyses had a larger number of patients after the highly cited study than before it, with four exceptions, including one in which all other studies in the meta-analysis preceded the highly cited one.




Figure 2. Relative contribution of highly cited studies to the sample size of replication meta-analyses. For studies with meta-analyses as replications, blue bars show the fraction of the total sample size that corresponds to the highly cited study, while gray bars represent the rest of the sample size. Fractions are arranged in temporal order from left to right, so that gray bars to the left of the blue ones represents studies that came before the highly cited one. The meta-analysis by Hamid et al., 2013 (42) is not shown because the highly cited study is not included in it, due to the lack of a control group.

Replication rates using different criteria are shown in **Table 4**. Among the 15 highly cited studies with statistically significant results, only 2 (13%) had a non-significant result in the replication, whereas out of the 2 highly cited studies with negative results, 1 had a significant result in the replication. We did not consider the latter as a replication failure in our main analysis, as lack of significance in null hypothesis tests should not be taken as evidence of equivalence, especially considering that sample size was higher in the replications. However, we did perform a sensitivity analysis using the statistical significance criterion for negative studies as well.

Criterion	Total	Replicated	% Replicated [95% CI]
Statistical significance	15	13	87% [62, 96]
Confidence interval overlap	24	21	88% [69 - 96]
Confidence interval overlap and statistical significance	24	20	83% [64 - 93]
Statistical significance (including negative studies)	17	14	82% [59 - 94]
Replication estimate within highly cited study's CI	24	18	75% [55 - 88]
Highly cited study estimate within replication CI	24	11	46% [28 - 65]

Table 4. Rate of successful replication as measured by different criteria. Results are shown as the percentage of studies with eligible replications in which the original result was replicated by each of 5 different criteria. For the statistical significance criterion, we excluded 7 phase 1 trials that did not report p-values and 2 studies with non-significant results. For these studies, the aggregate criterion (confidence interval overlap + statistical significance) only considers confidence interval overlap. The 2 negative studies are included in the “statistical significance (including negative studies)” criterion as a sensitivity analysis.

Phase 1 studies with no control group were also not considered for the statistical significance criterion. That said, among the 7 phase 1 trials, 6 had replications with statistically significant results when comparing the intervention to a control group in an RCT; the remaining one was replicated by an uncontrolled phase 2 trial (83).

Concerning the effect size criterion, 21 out of 24 studies (88%) had overlapping confidence intervals, with only 2 phase 1 trials and 1 RCT not meeting this criterion. In total, 18 out of 24 replications (75%) had point estimates that were contained in the confidence intervals of the highly cited studies; conversely, only 11 (46%) of the original point estimates were included in the replication's confidence intervals. That said, as confidence intervals get narrower with increasing sample size, the latter criterion is excessively strict and should not be considered as a measure of replicability.

Effect size inflation was not particularly marked in this sample of highly cited clinical research (**Figure 3** and **Table 5**). Inflation increased slightly in phase 1 trials when effect sizes were coined (i.e. when the percentage of non-responders was used as the outcome) and in RCTs when publication order was not taken into account (e.g. when meta-analyses were always considered as replications), but remained low on average and failed to reach statistical significance in the main analysis.




Figure 3. Effect size inflation. Effect size inflation was calculated for each study pair considering the effect sizes of highly cited studies and their respective replications. For unfavorable outcomes (e.g., death, tumor response), inflation was calculated as the ratio between the replication and highly cited point estimates. For favorable outcomes (e.g., neurologic improvement), it was calculated as the ratio between the highly cited and the replication point estimates. When the highly cited study came after most of the studies in the meta-analysis, we considered the highly cited study as the replication and inverted the ratio. Left panel shows the results in a linear scale (in which distribution is expected to be skewed upwards even in the absence of inflation because of the nature of ratios), while right panel shows them in a log-scale (in which values should be distributed symmetrically around 1 in the absence of inflation). Lines indicate the corresponding means of the plotted values. Different colors indicate Phase 1 studies (red) and RCTs (orange).

Analysis	All	p-value	Phase 1 trials	RCTs
Main analysis	1.06 [0.92 - 1.23]	0.38	1.01 [0.64 - 1.57]	1.09 [0.95 - 1.25]
Publication order	1.11 [0.97 - 1.28]	0.12	1.01 [0.64 - 1.57]	1.16 [1.03 - 1.32]
Effect coining	1.10 [0.97 - 1.26]	0.13	1.17 [0.80 - 1.71]	1.08 [0.94 - 1.24]
Publication order + coining	1.15 [1.02 - 1.30]	0.03	1.17 [0.80 - 1.71]	1.14 [1.01 - 1.29]

Table 5. Analyses of effect size inflation. Analyses to detect evidence of effect size inflation using different ways to handle favorable/unfavorable outcomes and meta-analyses with most of the sample size preceding the highly cited study. In “Publication order”, we did not invert the ratio if a replication meta-analysis had most of the sample size before the highly cited study. In “effect coining”, we used unfavorable outcomes for all effect measures, thus avoiding the need to reverse ratios for favorable ones. The bottom row combines both approaches. P-values refer to a one-sample Student’s t test against a theoretical mean of 0 for the log-transformed inflation ratios for each study pair. Values for means and 95% CIs were calculated for these log-transformed ratios and exponentiated back to the original scale.

Potential predictors of replicability are shown in **Table 6**. Although this analysis was planned in the protocol, it is underpowered due to the low number of contradicted studies in our sample. Both study design and the p-value of the highly cited study show trends of association in the expected direction, with higher replicability rates for RCTs and studies with lower p-values, but low power prevents us from drawing any definite conclusions.

Predictor	Criteria	Median (IQR) Replicated	Median (IQR) Contradicted	p-value
Citations/year of the highly cited study	Significance ($p < 0.05$)	292 (250 - 454)	441 (338 - 543)	0.69
	CI overlap	329 (250 - 454)	494 (368 - 570)	0.35
	Significance ($p < 0.05$) & CI overlap	337 (270 - 459)	368 (241 - 532)	0.79
p-value of the highly cited study	Significance ($p < 0.05$)	3×10^{-5} (6×10^{-7} - 2×10^{-3})	0.02 (8×10^{-3} - 0.02)	0.48
	CI overlap	1×10^{-3} (2×10^{-6} - 0.02)	1×10^{-5} (1×10^{-5} - 1×10^{-5})	0.82
	Significance ($p < 0.05$) & CI overlap	5×10^{-4} (2×10^{-6} - 0.01)	0.02 (8×10^{-3} - 0.02)	0.72
Sample size of the highly cited study	Significance ($p < 0.05$)	500 (230 - 821)	91152 (45729 - 136576)	0.57
	CI overlap	602 (296 - 1314)	256 (232 - 280)	0.23
	Significance ($p < 0.05$) & CI overlap	551 (280 - 944)	305 (256 - 91152)	0.96
Highly cited study design	CI overlap	Phase 1 trial: 5/7 RCT: 16/17	Phase 1 trial: 2/7 RCT: 1/17	0.19
	Significance ($p < 0.05$) & CI overlap	Phase 1 trial: 5/7 RCT: 15/17	Phase 1 trial: 2/7 RCT: 2/17	0.55
Type of intervention	Significance ($p < 0.05$)	Pharmacological: 7/8 Other: 5/6	Pharmacological: 1/8 Other: 1/6	1.00
	CI overlap	Pharmacological: 13/16 Other: 8/8	Pharmacological: 3/16 Other: 0/8	0.53
	Significance ($p < 0.05$) & CI overlap	Pharmacological: 13/16 Other: 7/8	Pharmacological: 3/16 Other: 1/8	1.00

Table 6. Predictors of replication success. Table shows comparisons of potential predictors of replicability between replicated and contradicted studies using different definitions of replication success. P-values are shown for a Mann-Whitney test (for continuous variables) or Fisher's exact test (for categorical variables) comparing replicated and contradicted studies. For the significance criterion, only 15 studies (i.e. initially positive RCTs) were considered. Study design is not included as a predictor of statistical significance, as phase 1 trials did not use significance testing. Sample size for each group varies according to the specific criteria: statistical significance: 13 replicated, 2 contradicted; confidence interval overlap: 21 replicated, 3 contradicted; both criteria: 20 replicated, 4 contradicted.

Discussion

Replicability of the literature

The replicability of highly cited clinical studies in our sample was high, with 83% of studies replicated when considering our primary outcome of overlapping confidence intervals along with statistical significance in the same direction for initially significant studies. Using alternative definitions of a successful replication led to different estimates, but these remained reasonably high, with 71% of replication point estimates included in the original confidence interval, for instance. Moreover, evidence of systematic effect size inflation was very limited, both for phase 1 trials and RCTs.

These replication success rates are higher than those found in previous studies of highly cited clinical literature, where rates of 59% in general medicine (11) and 37% in psychiatry (12) were described when both statistical significance and effect size were considered (although the criteria for comparing effect sizes differed in each study). For statistical significance alone (a more homogeneous criterion), the successful replication rate for studies with significant results was 87% in our study, as compared to 79% and 63% in the previous two studies, respectively.

Although the differences in these estimates could be due to changes in the replicability of the published literature, methodological discrepancies between studies should be considered. Ioannidis' study (11), used a broader definition of replication; thus, many article pairs in his sample would not have been considered to have matching PICO components by our criteria. Moreover, among the contradicted or inflated studies in his sample, 4 were cohort studies whose replications were RCTs or meta-analyses of RCTs. The author himself acknowledges that it is not always possible to validate exposures as interventions, and other studies comparing observational research with RCTs have used the term "concordance" instead of replicability (84). If one considers only interventional studies (i.e. RCTs and case series) in Ioannidis' sample – a scenario more compatible with our study –, the replication rate is 67% when considering both significance and effect sizes, or 87% – exactly the same as ours – when considering statistical significance alone.

Defining replication boundaries

Considering a study as a replication of another inevitably requires establishing the boundary conditions of a claim (10). We opted to define replications as studies that had matching PICO general components (85). This led most highly cited studies to be classified as unchallenged, with replications being found for only 27% of our sample (as opposed to 76% in Ioannidis (11), 52% in Tajika et al., (12) and 42% in Niven et al. (13), in which criteria were less stringent). Many studies that could have been considered replications by looser criteria were thus excluded.

For instance, one study (86) in our sample only included male subjects, while subsequent studies included both men and women. Another one (87) studied an intervention in untreated melanoma, while its closest replication (88) investigated the same intervention in advanced melanoma. Other studies had different comparators: the ANCHOR trial (89) was not replicated because its two closest studies compared the intervention with verteporfin (90,91) instead of a sham procedure. Some studies had very specific controls, such as one that compared ipilimumab with gp100, a peptide vaccine (87), or another that compared vemurafenib and dacarbazine (35). Other studies had specific composite outcomes, such as composites of hospitalization and death from any cause (92) or of major macrovascular and microvascular events (93), while subsequent studies only analyzed them separately.

Even though we were more conservative than previous studies in defining replication boundaries, our study pairs were still not perfect replicas of each other. In some cases, definitions for clinical conditions were very broad, such as “cancer” in Brahmer et al. 2012 (43) and Topalian et al. 2012 (45), meaning that replication samples could potentially be quite distinct from the original one. These discrepancies thus remain as possible explanations for observed contradictions. Heterogeneity between study populations and interventions presents challenges to studying reproducibility within clinical research, and methodological differences between the original study and replications seem to be the norm in previous studies as well (13).

Contradicted studies

Regarding our primary outcome, two phase 1 studies and one RCT were classified as contradicted. Both phase 1 trials had confidence intervals that did not overlap with those of the replication – in one case, the effect was larger in the original study, while in the other it was larger in the replication. As phase 1 trials typically have small sample sizes and are likely to be more prone to publication/citation bias (i.e. a negative phase 1 trial is unlikely to become highly cited), their replicability is expected to be lower than that of RCTs. Nevertheless, the majority of phase 1 trials in our sample were successfully replicated, although replication criteria were less stringent for these studies as (a) they were not subject to the statistical significance criteria and (b) confidence intervals for their effect sizes were broader. Still, it’s worth noting that all RCTs replicating phase 1 studies in our sample showed a statistically significant benefit of the intervention when compared to a control group.

Concerning RCTs, the two replication failures for initially positive studies were observed for ERSPC (56) (a large prostate cancer screening trial) and KEYNOTE-024 (76) (a trial of the checkpoint inhibitor pembrolizumab in lung cancer). ERSPC (56) was contradicted because it showed a statistically significant effect ($p=0.03$) while the meta-analysis did not reach significance, although effect sizes do overlap. Some methodological issues have been proposed to account for this discrepancy, such as the large degree of control group contamination in the PLCO trial (94) and the lower screening intensity in the CAP trial (95), two negative studies that account for most of the weight in the replication meta-analysis.

KEYNOTE-024 (76), meanwhile, was considered contradicted by our replication criteria, which specifically addressed the primary outcome in the highly cited study – in this case, progression-free survival. Nevertheless, both the original study and the replication (KEYNOTE-042) (77) show an overall survival benefit, and the lack of effect on progression in the replication seems to be due to differences in the early stage of the trial, even though the intervention group fared better on the long run. Thus, the result was replicated when the more clinically relevant outcome of survival is considered, and the lack of replication for our chosen outcome appears to be a statistical accident.

Of note, we did not use the statistical significance criterion to evaluate studies with non-significant results (as lack of statistical significance should not be taken as evidence of equivalence between treatments). Accordingly, of the two non-significant studies in our sample, one of them – the ACCORD trial (60) – had a replication with a significant result – in this case, a large meta-analysis that reached marginal significance ($p=0.044$). Nevertheless, other meta-analyses arrived at different conclusions depending on the outcome of interest (61,93,96–98). Effect sizes were similar between studies, suggesting that lack of agreement in this criterion was a consequence of lower statistical power in the highly cited study. The second non-significant study, PARTNER A (64), had a non-significant result in the replication as well.

Effect size inflation

Many studies analyzing replications have found evidence that published effects are systematically inflated, a fact that is expected when statistical significance thresholds are used as a criterion for publication (99). Nevertheless, strategies to measure effect size inflation vary widely across studies. Ioannidis found initially stronger effects in 7 out of 27 replicated studies, using the criteria of a decrease in risk reduction of at least 50%, or a benefit of shorter duration or limited generalizability in the replication (11). Tajika et al. reported standardized mean differences of initial studies to be 2.3 times larger than those of replications (12), while Niven et al. reported a mean absolute risk difference of 16% between original studies and replications. Similar evidence of effect size inflation has also been found both in systematic replication initiatives (5,8) and meta-analyses of effect sizes over time (100,101).

Contrary to these studies, we found little evidence of systematic effect size inflation in the highly cited clinical literature between 2004 and 2018. This suggests that publication/citation bias might be more limited in the highly cited clinical literature than in other fields of research. That said, our approach to measure effect size inflation is different from those in previous attempts (11,12,101), and our ability to detect it may have been reduced by the use of meta-analyses as replications, as the replication sample typically includes the highly cited study, as well as studies that came before it. We tried to correct for this by taking study order into account in these cases, but some degree of circularity inevitably remains.

Another limitation is that using ratios for measuring effect size inflation leads to variation in estimates depending on the outcome used. When nonresponse rates or response odds were used instead of ORRs for phase 1 studies, for example, estimates of

inflation increased to 17% or 18%, respectively – although confidence intervals were still wide and compatible with absence of systematic inflation. One can also make the case that relative differences in effect sizes may be less relevant than absolute ones for clinical practice. Nevertheless, the fact that different outcome measures were used across studies makes absolute differences non-commensurable between them and prevents us from analyzing the sample as a whole in this manner.

Although evidence for systematic inflation was limited, this does not mean that initially stronger effects were not found in some studies, as in the case of KEYNOTE-024 (76), in which a risk reduction in progression disappeared in the replication, and TAXUS-IV (47), in which risk reduction decreased from 61% to 34%. Nevertheless, the fact that increases in effect size were found in other studies, such as Brahmer et al., 2012 (43), in which the ORR doubled from 13% to 27%, suggests that some or most of these discrepancies can be explained by statistical fluctuation, and that systematic bias in favor of positive studies is smaller in this literature than in other fields.

Replication criteria

Different replication criteria complement each other by capturing distinct aspects of replicability (6,8). Statistical significance alone does not distinguish between magnitude and precision, and thus says little about how two effects compare directly (102). Comparing effect sizes, meanwhile, avoids emphasis on statistical thresholds (8); on the other hand, conclusions based on these thresholds may be different even when effect sizes are similar, due to different levels of precision. For this reason, our primary outcome was a combination of statistical significance and confidence interval overlap of effect sizes.

Highly cited studies were expected to predominantly present statistically significant results for their primary outcomes, as this literature is enriched in studies with high power and high prior probabilities (103). Most replications also yielded significant results, something that would be expected if the primary findings represent true effects. In fact, considering that the replication rate for this criterion was 87% for initially positive studies, even replication failures could represent vibration around statistical thresholds (as might have been the case for the KEYNOTE-024 replication (76), for example), as studies in clinical medicine are often powered around that level.

Replicability based on confidence interval overlap was similarly high (88%), although this is a rather loose threshold for effect size similarity: absence of CI overlap for two identical effects is expected to occur by chance alone in around 0.6% of cases (105), and this high bar for type 1 error comes at the cost of lower power to detect differences in effect sizes. A more stringent criteria of having the replication effect size included in the original confidence interval leads to a lower but still reasonably high (75%) replication rate, despite not considering the potential variability in replication estimates. Inclusion of the highly cited study's point estimate in the replication confidence interval was less frequent (46%), but this is an overly strict criterion, especially when replications are large. Calculating prediction intervals for the original effect given the replication sample size would likely represent the fairest way to assess replication of effect sizes

(106). That said, in the absence of raw data for the original studies, calculating these intervals would not be feasible for all chosen outcomes.

General limitations

Defining what constitutes a replication is not trivial: even though we followed a predefined protocol to define PICO components, their abstraction inevitably involves a degree of subjectivity. Moreover, as was the case in previous studies of replications in the published literature (11,12,101), there was no way to develop a systematic search strategy that was applicable for every replication. Perhaps because of these factors, our independent searches for replications had low agreement, and a second step was needed to reach consensus. Still, it is possible that our searches could have missed some valid replication candidates.

In at least one case – the ACCORD trial on intensive glucose control (60) – there were candidate replications that reached different results (60,61,96–98). Although we used an objective criterion to define the replication selected for analysis (i.e. sample size), a replication with a different result might have been chosen if we used other criteria. Of note, we did not evaluate risk of bias or methodological quality in replications, opening up the possibility that the largest replication available might not be necessarily the most reliable one.

An important caveat in our analysis is the fact that meta-analyses were considered as replications, even though most of them included the highly cited study. This leads to a degree of circularity in the analysis that may have biased our reproducibility estimate upwards. Overlap was limited for most studies, but may have had a more pronounced effect in cases such as PARTNER-A (64), PROFILE 1007 (58) and NEJSG (73), where the highly cited studies had larger weights in the meta-analyses. Nevertheless, performing all meta-analyses from scratch excluding the highly cited studies would be unfeasible in many cases, due to insufficient data and/or methodological information to replicate the analyses. On the other hand, excluding meta-analyses as valid replications would have left us with few eligible replication studies.

Even after including meta-analyses, our stringent criteria to consider studies as having matching PICO components left us with a small sample, and the high replicability rate led to an even lower number of contradicted studies. Thus, our analysis was markedly underpowered to detect predictors of replicability. Even though none of our predictors reached statistical significance, it seems likely that factors such as lower p values or higher sample sizes would be associated with a higher replicability rate, had a larger sample been available.

Lastly, we relied on replications that were published in the literature. As the existence and publication of these replications are subject both to the interest of researchers to perform them and to that of editors and reviewers to publish them, the approach in our study is not directly comparable to the systematic replication attempts that have been performed in other areas of the literature (1–8). This is particularly important given that the majority of highly cited studies in our sample had no available replications according to our criteria – thus, selectiveness in performing or publishing replications

may have biased our replicability rates. It is also possible that successfully replicated studies receive more citations in the long run, biasing the reproducibility rate of highly cited studies upwards.

Conclusions

Despite the high rate of unchallenged studies, we found the replicability rate of the highly cited clinical literature between 2004 and 2018 to be higher than previous estimates, with little evidence of effect size inflation. Naturally, these numbers are valid for a narrow, very influential subsample of articles, and cannot be generalized to medical research at large. Nevertheless, they run counter to the assertion that there is a widespread reproducibility crisis in science, and suggest that this may not be the case for every scientific field.

The higher replication rate found in our study when compared to estimates based on earlier samples of the clinical literature could also be taken as a sign of improvement over time; nevertheless, this conclusion is tentative at best, as differences in methodology (such as the definition of effect size inflation) and samples (such as the frequency of different study designs) do not warrant direct comparisons between studies.

Further research is warranted to examine whether the high replicability of highly cited clinical research is related to particular research practices that are not as widely used in other areas of biomedical science, such as randomization, blinding or prospective protocol registration (14,23,27,104,107,108). If such links can be reliably established, they can be used to inform attempts to improve replicability in different research fields.

References

1. Neves K, Carneiro CF, Wasilewska-Sampaio AP, Abreu M, Valério-Gomes B, Tan PB, et al. Two years into the Brazilian Reproducibility Initiative: reflections on conducting a large-scale replication of Brazilian biomedical science. *Mem Inst Oswaldo Cruz*. 2020;115:e200328.
2. Klein RA, Vianello M, Hasselman F, Adams BG, Adams RB, Alper S, et al. Many Labs 2: Investigating Variation in Replicability Across Samples and Settings. *Advances in Methods and Practices in Psychological Science*. 2018 Dec 1;1(4):443–90.
3. Ebersole CR, Atherton OE, Belanger AL, Skulborstad HM, Allen JM, Banks JB, et al. Many Labs 3: Evaluating participant pool quality across the academic semester via replication. *Journal of Experimental Social Psychology*. 2016 Nov 1;67:68–82.
4. Errington TM, Mathur M, Soderberg CK, Denis A, Perfito N, Iorns E, et al. Investigating the replicability of preclinical cancer biology. *Elife*. 2021 Dec 7;10:e71601.

5. Camerer CF, Dreber A, Holzmeister F, Ho TH, Huber J, Johannesson M, et al. Evaluating the replicability of social science experiments in Nature and Science between 2010 and 2015. *Nat Hum Behav.* 2018 Sep;2(9):637–44.
6. Cova F, Strickland B, Abatista A, Allard A, Andow J, Attie M, et al. Estimating the Reproducibility of Experimental Philosophy. *RevPhilPsych.* 2021 Mar;12(1):9–44.
7. Errington TM, Iorns E, Gunn W, Tan FE, Lomax J, Nosek BA. An open investigation of the reproducibility of cancer biology research. Rodgers P, editor. *eLife.* 2014 Dec 10;3:e04333.
8. Open Science Collaboration. Estimating the reproducibility of psychological science. *Science.* 2015 Aug 28;349(6251):aac4716–aac4716.
9. Fanelli D. Opinion: Is science really facing a reproducibility crisis, and do we need it to? *Proc Natl Acad Sci USA.* 2018 Mar 13;115(11):2628–31.
10. Nosek BA, Errington TM. What is replication? *PLoS Biol.* 2020 Mar 27;18(3):e3000691.
11. Ioannidis JPA. Contradicted and Initially Stronger Effects in Highly Cited Clinical Research. *JAMA.* 2005 Jul 13;294(2):218.
12. Tajika A, Ogawa Y, Takeshima N, Hayasaka Y, Furukawa TA. Replication and contradiction of highly cited research papers in psychiatry: 10-year follow-up. *Br J Psychiatry.* 2015 Oct;207(4):357–62.
13. Niven DJ, McCormick TJ, Straus SE, Hemmelgarn BR, Jeffs L, Barnes TRM, et al. Reproducibility of clinical research in critical care: a scoping review. *BMC Medicine.* 2018 Feb 21;16(1):26.
14. Chan AW, Altman DG. Epidemiology and reporting of randomised trials published in PubMed journals. *Lancet.* 2005 Apr 26;365(9465):1159–62.
15. Jadad AR, Moore RA, Carroll D, Jenkinson C, Reynolds DJ, Gavaghan DJ, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials.* 1996 Feb;17(1):1–12.
16. Schulz KF, Chalmers I, Hayes RJ, Altman DG. Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *JAMA.* 1995 Feb 1;273(5):408–12.
17. Schulz KF, Grimes DA, Altman DG, Hayes RJ. Blinding and exclusions after allocation in randomised controlled trials: survey of published parallel group trials in obstetrics and gynaecology. *BMJ.* 1996 Mar 23;312(7033):742–4.
18. Hotopf M, Lewis G, Normand C. Putting trials on trial--the costs and consequences of small trials in depression: a systematic review of methodology. *J Epidemiol Community Health.* 1997 Aug;51(4):354–8.

19. Dickinson K, Bunn F, Wentz R, Edwards P, Roberts I. Size and quality of randomised controlled trials in head injury: review of published studies. *BMJ*. 2000 May 13;320(7245):1308–11.
20. Thornley B, Adams C. Content and quality of 2000 controlled trials in schizophrenia over 50 years. *BMJ*. 1998 Oct 31;317(7167):1181–4.
21. Moher D, Pham B, Jones A, Cook DJ, Jadad AR, Moher M, et al. Does quality of reports of randomised trials affect estimates of intervention efficacy reported in meta-analyses? *Lancet*. 1998 Aug 22;352(9128):609–13.
22. Begg C, Cho M, Eastwood S, Horton R, Moher D, Olkin I, et al. Improving the quality of reporting of randomized controlled trials. The CONSORT statement. *JAMA*. 1996 Aug 28;276(8):637–9.
23. Zarin DA, Tse T, Williams RJ, Rajakannan T. Update on Trial Registration 11 Years after the ICMJE Policy Was Established. *N Engl J Med*. 2017 Jan 26;376(4):383–91.
24. Taichman DB, Sahni P, Pinborg A, Peiperl L, Laine C, James A, et al. Data Sharing Statements for Clinical Trials — A Requirement of the International Committee of Medical Journal Editors. *N Engl J Med*. 2017 Jun 8;376(23):2277–9.
25. Ioannidis JPA. The Mass Production of Redundant, Misleading, and Conflicted Systematic Reviews and Meta-analyses: Mass Production of Systematic Reviews and Meta-analyses. *The Milbank Quarterly*. 2016 Sep;94(3):485–514.
26. Goldacre B, Drysdale H, Dale A, Milosevic I, Slade E, Hartley P, et al. COMPare: a prospective cohort study correcting and monitoring 58 misreported trials in real time. *Trials*. 2019 Dec;20(1):118.
27. Kaplan RM, Irvin VL. Likelihood of Null Effects of Large NHLBI Clinical Trials Has Increased over Time. *PLOS ONE*. 2015 Aug 5;10(8):e0132382.
28. Altman DG. The Revised CONSORT Statement for Reporting Randomized Trials: Explanation and Elaboration. *Ann Intern Med*. 2001 Apr 17;134(8):663.
29. Schulz KF, Altman DG, Moher D. CONSORT 2010 Statement: updated guidelines for reporting parallel group randomised trials. 2010;9.
30. Kim C, Prasad V. Cancer Drugs Approved on the Basis of a Surrogate End Point and Subsequent Overall Survival: An Analysis of 5 Years of US Food and Drug Administration Approvals. *JAMA Intern Med*. 2015 Dec 1;175(12):1992.
31. Ebrahim S, Sohani ZN, Montoya L, Agarwal A, Thorlund K, Mills EJ, et al. Reanalyses of Randomized Clinical Trial Data. *JAMA*. 2014 Sep 10;312(10):1024.
32. Jager LR, Leek JT. An estimate of the science-wise false discovery rate and application to the top medical literature. *Biostatistics*. 2014 Jan;15(1):1–12.

33. Leek JT, Jager LR. Is Most Published Research Really False? *Annu Rev Stat Appl*. 2017 Mar 7;4(1):109–22.
34. R Core Team. R: A language and environment for statistical computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; Available from: <https://www.R-project.org/>
35. Chapman PB, Hauschild A, Robert C, Haanen JB, Ascierto P, Larkin J, et al. Improved Survival with Vemurafenib in Melanoma with BRAF V600E Mutation. *N Engl J Med*. 2011 Jun 30;364(26):2507–16.
36. Larkin J, Chiarion-Sileni V, Gonzalez R, Grob JJ, Cowey CL, Lao CD, et al. Combined Nivolumab and Ipilimumab or Monotherapy in Untreated Melanoma. *N Engl J Med*. 2015 Jul 2;373(1):23–34.
37. Campbell BCV, Mitchell PJ, Kleinig TJ, Dewey HM, Churilov L, Yassi N, et al. Endovascular Therapy for Ischemic Stroke with Perfusion-Imaging Selection. *N Engl J Med*. 2015 Mar 12;372(11):1009–18.
38. Lieberman JA, Stroup TS, McEvoy JP, Swartz MS, Rosenheck RA, Perkins DO, et al. Effectiveness of Antipsychotic Drugs in Patients with Chronic Schizophrenia. *N Engl J Med*. 2005 Sep 22;353(12):1209–23.
39. Cochrane Handbook for Systematic Reviews of Interventions | Cochrane Training [Internet]. [cited 2021 Aug 2]. Available from: <https://training.cochrane.org/handbook/current>
40. Murad MH, Asi N, Alsawas M, Alahdab F. New evidence pyramid. *Evid Based Med*. 2016 Aug;21(4):125–7.
41. Higgins JPT, Thompson SG, Spiegelhalter DJ. A re-evaluation of random-effects meta-analysis. *Journal of the Royal Statistical Society Series A, (Statistics in Society)*. 2009 Jan;172(1):137.
42. Hamid O, Robert C, Daud A, Hodi FS, Hwu WJ, Kefford R, et al. Safety and Tumor Responses with Lambrolizumab (Anti-PD-1) in Melanoma. *N Engl J Med*. 2013 Jul 11;369(2):134–44.
43. Brahmer JR, Tykodi SS, Chow LQM, Hwu WJ, Topalian SL, Hwu P, et al. Safety and Activity of Anti-PD-L1 Antibody in Patients with Advanced Cancer. *N Engl J Med*. 2012 Jun 28;366(26):2455–65.
44. Zhang T, Xie J, Arai S, Wang L, Shi X, Shi N, et al. The efficacy and safety of anti-PD-1/PD-L1 antibodies for treatment of advanced or refractory cancers: a meta-analysis. *Oncotarget*. 2016 Nov 8;7(45):73068–79.
45. Topalian SL, Hodi FS, Brahmer JR, Gettinger SN, Smith DC, McDermott DF, et al. Safety, Activity, and Immune Correlates of Anti-PD-1 Antibody in Cancer. *N Engl J Med*. 2012 Jun 28;366(26):2443–54.

46. Tie Y, Ma X, Zhu C, Mao Y, Shen K, Wei X, et al. Safety and efficacy of nivolumab in the treatment of cancers: A meta-analysis of 27 prospective clinical trials: Safety and efficacy of nivolumab. *Int J Cancer*. 2017 Feb 15;140(4):948–58.
47. Stone GW, Ellis SG, Cox DA, Hermiller J, O’Shaughnessy C, Mann JT, et al. A Polymer-Based, Paclitaxel-Eluting Stent in Patients with Coronary Artery Disease. *N Engl J Med*. 2004 Jan 15;350(3):221–31.
48. Bangalore S, Toklu B, Amoroso N, Fusaro M, Kumar S, Hannan EL, et al. Bare metal stents, durable polymer drug eluting stents, and biodegradable polymer drug eluting stents for coronary artery disease: mixed treatment comparison meta-analysis. *BMJ*. 2013 Nov 8;347(nov08 1):f6625–f6625.
49. Serruys PW, Morice MC, Kappetein AP, Colombo A, Holmes DR, Mack MJ, et al. Percutaneous Coronary Intervention versus Coronary-Artery Bypass Grafting for Severe Coronary Artery Disease. *N Engl J Med*. 2009 Mar 5;360(10):961–72.
50. Ali WE, Vaidya SR, Ejeh SU, Okoroafor KU. Meta-analysis study comparing percutaneous coronary intervention/drug eluting stent versus coronary artery bypass surgery of unprotected left main coronary artery disease: Clinical outcomes during short-term versus long-term (> 1 year) follow-up. *Medicine*. 2018 Feb;97(7):e9909.
51. Piccart-Gebhart MJ, Procter M, Leyland-Jones B, Goldhirsch A, Untch M, Smith I, et al. Trastuzumab after Adjuvant Chemotherapy in HER2-Positive Breast Cancer. *N Engl J Med*. 2005 Oct 20;353(16):1659–72.
52. Genuino AJ, Chaikledkaew U, The DO, Reungwetwattana T, Thakkinstian A. Adjuvant trastuzumab regimen for HER2-positive early-stage breast cancer: a systematic review and meta-analysis. *Expert Rev Clin Pharmacol*. 2019 Aug;12(8):815–24.
53. Soares-Weiser K, Bécharad-Evans L, Howard Lawson A, Davis J, Ascher-Svanum H. Time to all-cause treatment discontinuation of olanzapine compared to other antipsychotics in the treatment of schizophrenia: A systematic review and meta-analysis. *European Neuropsychopharmacology*. 2013 Feb 1;23(2):118–25.
54. Llovet JM, Ricci S, Mazzaferro V, Hilgard P, Gane E, Blanc JF, et al. Sorafenib in Advanced Hepatocellular Carcinoma [Internet]. <http://dx.doi.org/10.1056/NEJMoa0708857>. Massachusetts Medical Society; 2009 [cited 2021 Oct 6]. Available from: <https://www.nejm.org/doi/full/10.1056/nejmoa0708857>
55. Niu M, Hong D, Ma TC, Chen XW, Han JH, Sun J, et al. Short-term and long-term efficacy of 7 targeted therapies for the treatment of advanced hepatocellular carcinoma: a network meta-analysis: Efficacy of 7 targeted therapies for AHCC. *Medicine (Baltimore)*. 2016 Dec;95(49):e5591.

56. Schröder FH, Hugosson J, Roobol MJ, Tammela TLJ, Ciatto S, Nelen V, et al. Screening and Prostate-Cancer Mortality in a Randomized European Study. *N Engl J Med*. 2009 Mar 26;360(13):1320–8.
57. Ilic D, Djulbegovic M, Jung JH, Hwang EC, Zhou Q, Cleves A, et al. Prostate cancer screening with prostate-specific antigen (PSA) test: a systematic review and meta-analysis. *BMJ* [Internet]. 2018 Sep 5 [cited 2021 Oct 6];362. Available from: <https://www.bmj.com/content/362/bmj.k3519>
58. Shaw AT, Kim DW, Nakagawa K, Seto T, Crinó L, Ahn MJ, et al. Crizotinib versus Chemotherapy in Advanced ALK-Positive Lung Cancer [Internet]. <http://dx.doi.org/10.1056/NEJMoa1214886>. Massachusetts Medical Society; 2013 [cited 2021 Oct 6]. Available from: <https://www.nejm.org/doi/10.1056/NEJMoa1214886>
59. Elliott J, Bai Z, Hsieh SC, Kelly SE, Chen L, Skidmore B, et al. ALK inhibitors for non-small cell lung cancer: A systematic review and network meta-analysis. *PLOS ONE*. 2020 Feb 19;15(2):e0229179.
60. Effects of Intensive Glucose Lowering in Type 2 Diabetes. *N Engl J Med*. 2008 Jun 12;358(24):2545–59.
61. Fang HJ, Zhou YH, Tian YJ, Du HY, Sun YX, Zhong LY. Effects of intensive glucose lowering in treatment of type 2 diabetes mellitus on cardiovascular outcomes: A meta-analysis of data from 58,160 patients in 13 randomized controlled trials. *International Journal of Cardiology*. 2016 Sep;218:50–8.
62. Al C, Yk K, Z C, Cj T, S Q, Js K, et al. Efficacy and safety of sorafenib in patients in the Asia-Pacific region with advanced hepatocellular carcinoma: a phase III randomised, double-blind, placebo-controlled trial. *The Lancet Oncology* [Internet]. 2009 Jan [cited 2021 Mar 11];10(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/19095497/>
63. Goyal M, Menon BK, van Zwam WH, Dippel DWJ, Mitchell PJ, Demchuk AM, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet*. 2016 Apr 23;387(10029):1723–31.
64. Smith CR, Leon MB, Mack MJ, Miller DC, Moses JW, Svensson LG, et al. Transcatheter versus Surgical Aortic-Valve Replacement in High-Risk Patients. *N Engl J Med*. 2011 Jun 9;364(23):2187–98.
65. Pagnesi M, Chiarito M, Stefanini GG, Testa L, Reimers B, Colombo A, et al. Is Transcatheter Aortic Valve Replacement Superior to Surgical Aortic Valve Replacement?: A Meta-Analysis of Randomized Controlled Trials. *JACC Cardiovasc Interv*. 2017 Sep 25;10(18):1899–901.
66. Berkhemer OA, Fransen PSS, Beumer D, van den Berg LA, Lingsma HF, Yoo AJ, et al. A Randomized Trial of Intraarterial Treatment for Acute Ischemic Stroke

[Internet]. <https://doi.org/10.1056/NEJMoa1411587>. Massachusetts Medical Society; 2014 [cited 2021 Oct 6]. Available from: <https://www.nejm.org/doi/full/10.1056/nejmoa1411587>

67. Rodrigues FB, Neves JB, Caldeira D, Ferro JM, Ferreira JJ, Costa J. Endovascular treatment versus medical care alone for ischaemic stroke: systematic review and meta-analysis. *BMJ* [Internet]. 2016 Apr 18 [cited 2021 Mar 11];353. Available from: <https://www.bmj.com/content/353/bmj.i1754>

68. Hacke W, Kaste M, Bluhmki E, Brozman M, Dávalos A, Guidetti D, et al. Thrombolysis with Alteplase 3 to 4.5 Hours after Acute Ischemic Stroke [Internet]. <http://dx.doi.org/10.1056/NEJMoa0804656>. Massachusetts Medical Society; 2009 [cited 2021 Mar 11]. Available from: <https://www.nejm.org/doi/full/10.1056/nejmoa0804656>

69. Wardlaw JM, Murray V, Berge E, del Zoppo G, Sandercock P, Lindley RL, et al. Recombinant tissue plasminogen activator for acute ischaemic stroke: an updated systematic review and meta-analysis. *Lancet*. 2012 Jun 23;379(9834):2364–72.

70. Rosell R, Carcereny E, Gervais R, Vergnenegre A, Massuti B, Felip E, et al. Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EURTAC): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol*. 2012 Mar;13(3):239–46.

71. Zhao Y, Liu J, Cai X, Pan Z, Liu J, Yin W, et al. Efficacy and safety of first line treatments for patients with advanced epidermal growth factor receptor mutated, non-small cell lung cancer: systematic review and network meta-analysis. *BMJ* [Internet]. 2019 Oct 7 [cited 2021 Oct 6];367. Available from: <https://www.bmj.com/content/367/bmj.l5460>

72. Goyal M, Demchuk AM, Menon BK, Eesa M, Rempel JL, Thornton J, et al. Randomized Assessment of Rapid Endovascular Treatment of Ischemic Stroke [Internet]. <http://dx.doi.org/10.1056/NEJMoa1414905>. Massachusetts Medical Society; 2015 [cited 2021 Mar 11]. Available from: <https://www.nejm.org/doi/full/10.1056/nejmoa1414905>

73. Maemondo M, Inoue A, Kobayashi K, Sugawara S, Oizumi S, Isobe H, et al. Gefitinib or Chemotherapy for Non-Small-Cell Lung Cancer with Mutated EGFR [Internet]. <http://dx.doi.org/10.1056/NEJMoa0909530>. Massachusetts Medical Society; 2010 [cited 2021 Mar 11]. Available from: <https://www.nejm.org/doi/full/10.1056/nejmoa0909530>

74. Greenhalgh J, Dwan K, Boland A, Bates V, Vecchio F, Dundar Y, et al. First-line treatment of advanced epidermal growth factor receptor (EGFR) mutation positive non-squamous non-small cell lung cancer. *Cochrane Database of Systematic Reviews* [Internet]. 2016 [cited 2021 Oct 6];(5). Available from:

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD010383.pub2/full>

75. Pyo JS, Kang G. Immunotherapy in advanced melanoma: a network meta-analysis. *Immunotherapy*. 2017 May;9(6):471–9.
76. Reck M, Rodríguez-Abreu D, Robinson AG, Hui R, Csósz T, Fülöp A, et al. Pembrolizumab versus Chemotherapy for PD-L1-Positive Non-Small-Cell Lung Cancer [Internet]. <https://doi.org/10.1056/NEJMoa1606774>. Massachusetts Medical Society; 2016 [cited 2021 Oct 6]. Available from: <https://www.nejm.org/doi/full/10.1056/nejmoa1606774>
77. Mok TSK, Wu YL, Kudaba I, Kowalski DM, Cho BC, Turna HZ, et al. Pembrolizumab versus chemotherapy for previously untreated, PD-L1-expressing, locally advanced or metastatic non-small-cell lung cancer (KEYNOTE-042): a randomised, open-label, controlled, phase 3 trial. *Lancet*. 2019 May 4;393(10183):1819–30.
78. Garon EB, Rizvi NA, Hui R, Leighl N, Balmanoukian AS, Eder JP, et al. Pembrolizumab for the Treatment of Non-Small-Cell Lung Cancer [Internet]. <http://dx.doi.org/10.1056/NEJMoa1501824>. Massachusetts Medical Society; 2015 [cited 2021 Oct 6]. Available from: <https://www.nejm.org/doi/full/10.1056/NEJMoa1501824>
79. Herbst RS, Baas P, Kim DW, Felip E, Pérez-Gracia JL, Han JY, et al. Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomised controlled trial. *The Lancet*. 2016 Apr 9;387(10027):1540–50.
80. Wolchok JD, Kluger H, Callahan MK, Postow MA, Rizvi NA, Lesokhin AM, et al. Nivolumab plus Ipilimumab in Advanced Melanoma [Internet]. <https://doi.org/10.1056/NEJMoa1302369>. Massachusetts Medical Society; 2013 [cited 2021 Oct 6]. Available from: <https://www.nejm.org/doi/10.1056/NEJMoa1302369>
81. Flaherty KT, Puzanov I, Kim KB, Ribas A, McArthur GA, Sosman JA, et al. Inhibition of Mutated, Activated BRAF in Metastatic Melanoma. *N Engl J Med*. 2010 Aug 26;363(9):809–19.
82. Fong PC, Boss DS, Yap TA, Tutt A, Wu P, Mergui-Roelvink M, et al. Inhibition of Poly(ADP-Ribose) Polymerase in Tumors from BRCA Mutation Carriers [Internet]. <http://dx.doi.org/10.1056/NEJMoa0900212>. Massachusetts Medical Society; 2009 [cited 2021 Oct 6]. Available from: <https://www.nejm.org/doi/full/10.1056/nejmoa0900212>
83. Kaufman B, Shapira-Frommer R, Schmutzler RK, Audeh MW, Friedlander M, Balmaña J, et al. Olaparib Monotherapy in Patients With Advanced Cancer and a Germline *BRCA1/2* Mutation. *JCO*. 2015 Jan 20;33(3):244–50.

84. Moorthy D, Chung M, Lee J. Volume 6. Concordance Between the Findings of Epidemiological Studies and Randomized Trials in Nutrition: An Empirical Evaluation and Citation Analysis. :63.
85. Schwingshackl L, Balduzzi S, Beyerbach J, Bröckelmann N, Werner SS, Zähringer J, et al. Evaluating agreement between bodies of evidence from randomised controlled trials and cohort studies in nutrition research: meta-epidemiological study. *BMJ*. 2021 Sep 15;n1864.
86. Marin JM, Carrizo SJ, Vicente E, Agusti AG. Long-term cardiovascular outcomes in men with obstructive sleep apnoea-hypopnoea with or without treatment with continuous positive airway pressure: an observational study. *The Lancet*. 2005 Mar;365(9464):1046–53.
87. Hodi FS, O'Day SJ, McDermott DF, Weber RW, Sosman JA, Haanen JB, et al. Improved Survival with Ipilimumab in Patients with Metastatic Melanoma. *N Engl J Med*. 2010 Aug 19;363(8):711–23.
88. Larkin J, Chiarion-Sileni V, Gonzalez R, Grob JJ, Rutkowski P, Lao CD, et al. Five-Year Survival with Combined Nivolumab and Ipilimumab in Advanced Melanoma. *N Engl J Med*. 2019 Oct 17;381(16):1535–46.
89. Brown DM, Kaiser PK, Michels M, Soubrane G, Heier JS, Kim RY, et al. Ranibizumab versus Verteporfin for Neovascular Age-Related Macular Degeneration. *N Engl J Med*. 2006 Oct 5;355(14):1432–44.
90. Regillo CD, Brown DM, Abraham P, Yue H, Ianchulev T, Schneider S, et al. Randomized, Double-Masked, Sham-Controlled Trial of Ranibizumab for Neovascular Age-related Macular Degeneration: PIER Study Year 1. *American Journal of Ophthalmology*. 2008 Feb;145(2):239-248.e5.
91. Rosenfeld PJ, Brown DM, Heier JS, Boyer DS, Kaiser PK, Chung CY, et al. Ranibizumab for Neovascular Age-Related Macular Degeneration. *N Engl J Med*. 2006 Oct 5;355(14):1419–31.
92. Cleland JGF, Daubert JC, Erdmann E, Freemantle N, Gras D, Kappenberger L, et al. The Effect of Cardiac Resynchronization on Morbidity and Mortality in Heart Failure. *N Engl J Med*. 2005 Apr 14;352(15):1539–49.
93. Group TAC. Intensive Blood Glucose Control and Vascular Outcomes in Patients with Type 2 Diabetes [Internet]. <http://dx.doi.org/10.1056/NEJMoa0802987>. Massachusetts Medical Society; 2009 [cited 2021 Mar 11]. Available from: <https://www.nejm.org/doi/full/10.1056/nejmoa0802987>
94. Tsodikov A, Gulati R, Heijnsdijk EAM, Pinsky PF, Moss SM, Qiu S, et al. Reconciling the Effects of Screening on Prostate Cancer Mortality in the ERSPC and PLCO Trials. *Ann Intern Med*. 2017 Oct 3;167(7):449–55.

95. Martin RM, Donovan JL, Turner EL, Metcalfe C, Young GJ, Walsh EI, et al. Effect of a Low-Intensity PSA-Based Screening Intervention on Prostate Cancer Mortality: The CAP Randomized Clinical Trial. *JAMA*. 2018 Mar 6;319(9):883–95.
96. Hemmingsen B, Lund SS, Gluud C, Vaag A, Almdal T, Hemmingsen C, et al. Intensive glycaemic control for patients with type 2 diabetes: systematic review with meta-analysis and trial sequential analysis of randomised clinical trials. *BMJ*. 2011 Jan 9;343(nov24 1):d6898–d6898.
97. Marso SP, Kennedy KF, House JA, McGuire DK. The effect of intensive glucose control on all-cause and cardiovascular mortality, myocardial infarction and stroke in persons with type 2 diabetes mellitus: a systematic review and meta-analysis. *Diabetes and Vascular Disease Research*. 2010 Apr;7(2):119–30.
98. Sardar P, Udell JA, Chatterjee S, Bansilal S, Mukherjee D, Farkouh ME. Effect of Intensive Versus Standard Blood Glucose Control in Patients With Type 2 Diabetes Mellitus in Different Regions of the World: Systematic Review and Meta-analysis of Randomized Controlled Trials. *JAMA [Internet]*. 2015 May 14 [cited 2021 Sep 8];4(5). Available from: <https://www.ahajournals.org/doi/10.1161/JAHA.114.001577>
99. Ioannidis JPA. Why most discovered true associations are inflated. *Epidemiology*. 2008 Sep;19(5):640–8.
100. Fanelli D, Costas R, Ioannidis JPA. Meta-assessment of bias in science. *Proc Natl Acad Sci USA*. 2017 Apr 4;114(14):3714–9.
101. Cristea I, Ioannidis john, Georgescu R. Are effect sizes reported in highly cited emotion research overestimated relative to larger studies and meta-analyses addressing the same questions? [Internet]. *PsyArXiv*; 2021 Feb [cited 2021 Aug 2]. Available from: <https://osf.io/6t7pw>
102. Amrhein V, Korner-Nievergelt F, Roth T. The earth is flat ($p > 0.05$): significance thresholds and the crisis of unreplicable research. *PeerJ*. 2017 Jul 7;5:e3544.
103. Ioannidis JPA. Why Most Published Research Findings Are False. *PLoS Med*. 2005 Aug 30;2(8):e124.
104. Protzko J, Krosnick J, Nelson LD, Nosek BA, Axt J, Berent M, et al. High Replicability of Newly-Discovered Social-behavioral Findings is Achievable [Internet]. *PsyArXiv*; 2020 Sep [cited 2021 Aug 2]. Available from: <https://osf.io/n2a9x>
105. Austin PC, Hux JE. A brief note on overlapping confidence intervals. *J Vasc Surg*. 2002 Jul;36(1):194–5.
106. Spence JR, Stanley DJ. Prediction Interval: What to Expect When You're Expecting ... A Replication. *PLOS ONE*. 2016 Sep 19;11(9):e0162874.

107. Serghiou S, Contopoulos-Ioannidis DG, Boyack KW, Riedel N, Wallach JD, Ioannidis JPA. Assessment of transparency indicators across the biomedical literature: How open is open? PLOS Biology. 2021 Mar 1;19(3):e3001107.
108. Macleod MR, McLean AL, Kyriakopoulou A, Serghiou S, Wilde A de, Sherratt N, et al. Risk of Bias in Reports of In Vivo Research: A Focus for Improvement. PLOS Biology. 2015 Oct 13;13(10):e1002273.