

Socioeconomic status, lifestyle, and hypertension prevalence of the older adults in China

Jun Li¹, Meng Wei², Xiao Han¹, Hongman Wang^{1*}

¹School of Health Humanities, Peking University, Haidian District, Beijing, China

²National Institute of Social Development, Chinese Academy of Social Sciences

Co-first author:

Meng Wei

* Correspondence:

Hongman Wang

Peking University Health Science Center, 100191, 38 Xueyuan Rd, Haidian District, Beijing, China

cde@pku.edu.cn

Abstract

Aims: To examine the relationship between socioeconomic status (SES) and hypertension prevalence among the older adults in China. **Method:** We used the panel data of the three rounds (2006, 2009, and 2011) of the China Nutrition and Health Survey to examine the relationship adjusted for lifestyle factors by multi-level logistic regression. **Results:** Income and urbanicity levels are associated with hypertension in men. Men with low and middle income and living in low and middle urbanicity areas are more likely to have hypertension. Income and education are associated with hypertension in women. Women with low or middle income are more likely to have hypertension; women with lower education levels are less likely to have hypertension; women who never attended school are nearly as prone to get the disease as those who had high school or above education. **Conclusion:** Income, and community urbanicity level are reversely correlated with hypertension prevalence among older Chinese men and women; hypertension prevalence rates of older Chinese women vary by education levels without a gradient.

Keywords: older adults; socioeconomic status; lifestyle; hypertension; gender

With 11.4% of the Chinese population aged over 60 years old¹ and an increased old-age dependency ratio caused by lower birth rates, the social care and health care of the older adults in the society had attracted great attention in China. In most middle-income countries, where the situation is very similar, hypertension ranks high among the non-communicable diseases that are replacing communicable diseases as the main causes of mortality and hypertension has a high rate of prevalence and low rates of awareness, treatment, and control in the older population.²⁻⁴

Previous studies have explored the relationship between SES and health in the general population and the older adult population in some regions in China.⁵ Health disparity,

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

by SES among this population has been reported.⁶⁻¹² However, the relationship between hypertension and SES in this population in nationally representative samples remains unclear. Previous research also contains potential inaccuracy in the construction of some lifestyle factors, for example, using work-related PA as a proxy of overall PA.¹³

We used nationally representative longitudinal panel survey data to examine the relationship, controlling for lifestyle factors pinpointed by the World Health Organization for non-communicable diseases control—alcohol consumption, tobacco smoking, physical activity (PA, in MET-hours), total energy intake (TEI), and proportion of fat in total energy intake (as proportion in total calories, FIT) and demographic variables. The research question is what are the effects of the SES factors on hypertension in older men and older women in China?

We used the data of the 2006, 2009, and 2011 rounds of the China Nutrition and Health Surveys (CNHS, project website: <http://www.cpc.unc.edu/projects/china>), which had been conducted for 10 rounds between 1989 and 2015. The project investigated more than 7,200 urban and rural households and more than 30,000 individuals in 15 provinces chosen by multi-stage randomized sampling. We analyzed the 9,495 subjects aged 60 and over in the 9 provinces (Liaoning, Heilongjiang, Jiangsu, Shandong, Henan, Hubei, Hunan, Guangxi, Guizhou) covered by the CHNS in the three rounds. There were 2,876 subjects, 3,193 subjects and 3,426 subjects eligible in the three rounds, respectively. The total repeated measure rate of subjects (survey for at least two rounds) is 52.6% (82.34% in the 2006 and 2009 rounds, and 41.27% in the 2009 and 2011 rounds).

We represent subject SES by their annual per capita household income (PCHI), education level, and community urbanization index (CUI). Lifestyle factors controlled were smoking status, drinking status, TEI, FIT, and PA.

Education: 4 groups (1=Never attended school; 2=Primary school; 3=Junior middle school; 4=High school or above).

PCHI groups: Three groups divided by the tertile cut-off points, with income adjusted against the 2011 Consumer Price Index.

CUI groups: Three groups by tertile cut-off points of CUI, a composite urbanicity scale of 12 aspects, each scoring between 0 and 10.¹⁴

Smoking status: yes or no, reported by subjects when the answer to the question of "whether there is a history of smoking."

Drinking status: yes or no, reported by subjects when the answer to the question of "whether you drank last year."

TEI: Three groups by tertile cut-off points. The average total energy intake per day was calculated against the food ingredient table. The diet data was obtained by 24 hours recall method.

FIT groups: Three groups by tertile cut-off points. The proportion of fat in total energy intake= average calorie from fat per day/average total energy intake per day in the past three days (kcal). One gram of fat is calculated as 9 kcal of calories according to the method proposed by Xue.¹⁵

PA groups: Three groups by tertile cut-off points. We used the “Compendium of Physical Activities 2011”¹⁶ to calculate weekly PA in metabolic equivalent values of the self-reported PAs (sum of “MET/hour of each activity” * “hours lasted for the activities”).

Hypertension diagnosis: Subjects’ right arm blood pressure was measured three times in a sitting position after a 5-minute rest. The mean value of three measurements of the diastolic (and systolic) pressure was used as the diastolic (and systolic) pressure. Systolic blood pressure ≥ 140 mm Hg or diastolic blood pressure ≥ 90 mm Hg and/or taking antihypertensive drugs reported by the subjects is considered as having hypertension, according to the Chinese experts’ consensus for hypertension diagnosis and treatment in the older adults.³

The demographic control variables were age, ethnicity, and family size. For the selected characteristics of the subjects, see **Table 1**.

The data in this study is nested and the between-group correlation coefficient values (men=0.74, women=0.75) and the design effect values (1108.78, 1250.5) indicated that a multi-level model is fit for use.¹⁷⁻¹⁹

We examined the relationship between SES and hypertension, after adjusting for the demographic and lifestyle factors, in multi-level logistic regression models (generalized linear models). One model for men and one model for women, respectively.

SPSS 23.0 was used for statistical analysis.

In the three rounds of the survey, 4,496 men (on average, 1,499 surveyed per round) and 5,000 women were surveyed (on average, 1,667 per round).

There is little gender difference in PCHI and CUI as subjects in the same households, regardless of gender, have the same PCHI, and subjects who reside in the same community have the same CUI value.

There is a vast difference between genders, and little difference between survey years in terms of education level, age, and family size. More women survived men and lived alone.

Women's proportion of energy from fat is higher than that of men. More men had a history of smoking and reported drinking last year. Men's daily cigarette consumption and drinking frequency were higher than women. Men's smoking rate declined in the three rounds of the survey but was consistently higher than women (Table 1).

From 2006 to 2009, the prevalence of hypertension, the past diagnosis rate, and the rate of treatment in men and women increased (Table 1).

The accurate prediction value of the model increased with the addition of individual, family, community, and lifestyle variables until we obtained a satisfactory model for men and women respectively (Table 2). The model fit was judged based on the -2 log pseudo-fitness data. The smaller the value, the higher the model fit. The same applies for both modeling for men and women.

After adjusting for the demographic and lifestyle factors, men with low income and middle income (compared with high income, OR= 1.90(1.36~2.67), $p < 0.001$ and 1.39(1.02~1.90), $p < 0.05$) and with low and middle CUI (compared with high CUI, OR= 1.70(1.20~2.42), $p < 0.001$, and 1.48(1.07~2.04), $p < 0.05$) are more likely to have the disease (Table 2).

For women, we found the association between income (PCHI) and hypertension (OR= 2.07*** (1.28~3.11), $p < 0.001$ for low income, and 1.96*** (1.33~2.90), $p < 0.001$ for middle income) and the association between education level. Women with a lower education (primary schools) level are less likely to have hypertension (OR= 0.49(0.26~0.89), $p < 0.05$). But it is worth noting that women who never attended school, very probably illiterate, are almost equal probability to have hypertension as those who had high school or above education (OR=0.94(0.50~1.79), $p < 0.05$) (Table 2).

We found that SES factors—income, urbanicity, and education, are related to hypertension, and that gender differences exist in the relation between SES factors and hypertension in the older adults in China.

Our study found that higher per capita household income functions as a protective factor of hypertension in the older adults. Similarly, Kaplan et al.²⁰ had reported a significant inverse linear relationship between household income and the hypertension prevalence rate in the United States, but no evidence of such a relationship in Canada, and argued that social disparities and barriers to health care access and primary prevention in the United States may have played a role in it. We agree with Kaplan et al.'s explanation.²⁰ As China had adopted a market-oriented healthcare system since

its reform and opening-up at the end of the 1970s, income has become a much more important factor in obtaining living and healthcare resources. Many resources were obtained via a market mechanism, instead of rationing. Market-oriented policies stimulate efficiency in its economy, but at the cost of equity, as demonstrated here in health/disease disparity. To realize health equity, probably the strikingly severe income inequity as demonstrated in this study should be tackled first.

Our study confirms that people living in communities with higher urbanicity tend to have lower hypertension prevalence. Similar findings had been reported by previous studies. Buys et al.²¹ reported that the older adults with poorer community conditions have a higher prevalence of hypertension. Isaac and Laveist²² had reported that older people with convenience stores and recreational facilities have lower rates of hypertension, even if they live in communities of lower and medium SES. One possible mechanism is that with the decline of the mobility of older people, restrictions through neighborhood conditions became more prominent, rendering the older adults less capable of avoiding adverse socioeconomic conditions.²¹ Another possible mechanism might be that better income and living conditions facilitate the older adults to access quality preventive and curative health care and reduce anxiety and stress.²³

Therefore, we argue that geriatric health management, especially the management of chronic diseases, depends on an array of conditions, including transportation, community health care services, built community environment. Future research could test such a hypothesis.

We found that women with a lower education (primary schools) level are less likely to have hypertension; women who never attended school are almost as likely to have hypertension as women who had high school or above education.

The findings of previous research on education level and hypertension in the older adults in China are inconsistent, possibly influenced by the age groups, locality, and ethnic groups studied. Lei et al.²⁴ reported no gradient between the level of education and the prevalence of hypertension with a nationally representative sample data of the middle-aged and older adult population. Zuo et al.²⁵ reported no gradient between education level and hypertension among the older adults of 12 counties in Hubei. However, in Xinjiang, Zhang et al.²⁶ found a reversed correlation between education and hypertension among the older adults. Xinjiang is a northwestern region of China, where the proportion of Han Chinese is lower than that in other regions of China.

As for the possible mechanism underlying our finding, we speculate that older people with lower education have lower health literacy, which is essential in the management of their chronic diseases,²⁷ and those with higher education tend to have lower physical activity level (which we had found in our study, would be published soon in another article), which could be a predictive factor of hypertension in the older adults

in China. You et al.²⁸ found that middle-aged and older adults in China who usually participated in the moderate-to-vigorous activity for more than 10 minutes were less likely to have hypertension and Miao et al.²⁹ found similar results for the older adults in Beijing. More research is needed to verify the relationship between education and hypertension in the older adults.

The present study contributes to the literature by providing gender-specific results between SES and hypertension. It has been reported that the older adults with low SES in such countries as China^{7,8,30,31}, Germany³², UK³³, South Korea³⁴, and the United States^{20,35}, have worse health. However, previous studies ignore the gender difference in such a connection between SES and health. This study finds that the relationship between SES and hypertension in older men and women differ, which indicates that when considering the relation between social stratification and health of the older adults, gender should not be ignored.

We did not control the genetic variants, trace elements, diet structure, sleep, limitation of daily activities, cognitive abilities (which may affect their medication compliance and lifestyle), healthcare (such as medical security availability and access to medical services), SES of early life, pre-retirement occupation, and comorbidity in the study. The composite PA index that we use may obscure the effect difference of PA of high, medium, and low intensity. Future research could also consider dose, starting age and frequency of smoking and drinking and exposure to secondhand smoke, compare the accuracy of accelerometer-measured and self-recalled PA, and, if possible, use Mendelian randomization to examine the causality between SES and hypertension and the effect of lifestyle in the relationship.

Socioeconomic factors significantly correlate with hypertension in older men and women in China and that there are gender differences.

Abbreviations:

CNHS China Nutrition and Health Surveys

CUI Community Urbanization Index

FIT proportion of fat in total energy intake

HHS household size

MET-hours metabolic equivalent hours

OR odds ratio

PA physical activity

PCHI Per capita household income

SD standard deviation

SE standard error

SES socioeconomic status

TEI total energy intake

Declaration:

1. Ethics approval and consent to participate

This study used the data from the China Health and Nutrition Survey, which has obtained the ethics approval and subjects' consent to participate. We consulted the ethics committee office at Peking University and was informed that no ethical review was required since we used a public database for extracting our data and no personal information was revealed in the article. Therefore, we ask for waiving of the ethics approval and consent to participate.

2. Consent for publication

Not applicable

3. Competing interests

None.

4. Funding

This study receives funds from "Study on the construction of elderly care system under integrating health and social care arrangement" of the Society for Aging and Health of the China Health Economics Association (SAHCHEA-2021-01).

5. Authors' contributions

Jun Li: conceptualization, methodology, formal analysis, resources, data curation, and writing-original draft.

Meng Wei: conceptualization, methodology, formal analysis, resources; data curation; writing-original draft.

Xiao Han: writing-review & editing, submission.

Hongman Wang: conceptualization, methodology, resources, writing-review & editing, supervision, project administration, and funding acquisition.

All authors have read and approved the manuscript.

6. Acknowledgements

Acknowledgments: This research uses data from the China Health and Nutrition Survey (CHNS). We thank the National Institute of Nutrition and Food Safety, China Center for Disease Control and Prevention, Carolina Population Center, the University of North Carolina at Chapel Hill, the NIH (R01-HD30880, DK056350, and R01-HD38700) and the Fogarty International Center, NIH

for financial support for the CHNS data collection and analysis of files from 1989 to 2006 and both parties plus the China-Japan Friendship Hospital, Ministry of Health for support for CHNS 2009 and future surveys. We thank Professor Yun Zhou at Department of Sociology, Peking University, and Dr. Yan Ning at China Health Education Center for their suggestions for the study.

7. Availability of data and materials

The data used by this study is accessible at <https://www.cpc.unc.edu/projects/china/data>.

References

1. Data from: China Yearbook of Statistics 2017. *Population structure and dependency ratio*. 2018. *China*.
2. Lloyd-Sherlock P, Beard J, Minicuci N, Ebrahim S, Chatterji S. Hypertension among older adults in low- and middle-income countries: prevalence, awareness and control. *Int J Epidemiol*. 2014;43(1):116-128. doi:10.1093/ije/dyt215
3. Feng Y, Sun N, Li X, Chen L, Lin Z, Huang P. The characteristics of hypertension of the older adults and expert recommendations on clinical diagnosis and treatment process. *Chin J Hypertens*. 2014;22(7):620-628. doi:10.16439/j.cnki.1673-7245.2014.07.007
4. Fan T, Cao Q, Jiang L, Xu Z. Explaining the influencing factors on chronic diseases of the older adults using the ecological health model. *Chin Gen Pract*. 2012;15(1):33-36+40. doi:CNKI:SUN:QKYX.0.2012-01-015
5. Tu S. *Socioeconomic inequalities in prevalence and control of hypertension of the rural elderly in Shandong Province, China*. Doctor. Shandong University; 2009. <https://kns.cnki.net/KCMS/detail/detail.aspx?dbname=CDFD0911&filename=2009248858.nh>
6. Jiao K. Inequality of healthy life expectancy in the Chinese elderly. *Sociol Stud*. 2018;33(1):116-141+244-245. doi:10.19934/j.cnki.shxyj.2018.01.006
7. He Z. Socio-economic status, social support network, and the physical and mental status of rural elderly. *Soc Sci China*. 2002;23(3):135-148+207. doi:CNKI:SUN:ZSHK.0.2002-03-011
8. Han T. *Analysis of the influence of socioeconomic status on the health of the middle-aged and the older adults in China*. Master. Shandong University; 2017. <https://kns.cnki.net/KCMS/detail/detail.aspx?dbname=CMFD201702&filename=1017070268.nh>
9. Ai B, Wang S, Hoshi T. The mechanism of relationships between socioeconomic status and health among the older adults: take the example of the nine-year follow-up study in Shenyang city. *Popul Econ*. 2014;203(2):48-56. doi:CNKI:SUN:RKJJ.0.2014-02-007
10. Xia C. *The impact of socioeconomic status on the prevalence of chronic diseases in middle-aged and older adults*. Doctor. Peking University; 2016.

<https://thesis.lib.pku.edu.cn/docinfo.action?id1=554fcdcc27243b1819737e3880abb695&id2=kQb1vv%252BfK%252FE%253D>

11. Xu L, Yu L. The influence of social and economic status on elderly health: an empirical analysis based on CGSS 2013 data. *Stat Inf Forum*. 2016;31(3):52-59+67. doi:CNKI:SUN:TJLT.0.2016-03-008
12. Xue X, Ge K. The effect of SES on the health of the older adults in China: evidence from the Chinese longitudinal healthy longevity survey. *Popul Dev*. 2017;23(2):61-69. doi:CNKI:SUN:SCRK.0.2017-02-007
13. Ma Y. *The Impact of Socioeconomic Status and Dietary-related factors on Blood Pressure among Adults in 9 Provinces of China (1991-2009)*. Doctor. Chinese Center for Disease Control and Prevention; 2011. <https://kns.cnki.net/KCMS/detail/detail.aspx?dbname=CDFD1214&filename=1011210931.nh>
14. Jones-Smith JC, Popkin BM. Understanding community context and adult health changes in China: Development of an urbanicity scale. *Soc Sci Med*. 2010;71(8):1436-1446. doi:10.1016/j.socscimed.2010.07.027
15. Xue J. *Food Nutrition and Health*. China University of Science and Technology Press; 2004.
16. Ainsworth BE, Haskell WL, Whitt MC, et al. Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc*. 2000;32(9 Suppl):S498-504. doi:10.1097/00005768-200009001-00009
17. Ouyang Y, Wang Y, Wang H, Wang Z, Zhang B. A multi-level longitudinal study on the influence of dietary cholesterol intake on blood pressure and hypertension in Chinese adults. *Chin J Hypertens*. 2017;25(3):239-246. doi:10.16439/j.cnki.1673-7245.2017.03.010
18. Peugh JL. A practical guide to multilevel modeling. *J Sch Psychol*. 2010;48(1):85-112. doi:10.1016/j.jsp.2009.09.002
19. Sommet N, Morselli D. Keep Calm and Learn Multilevel Logistic Modeling: A Simplified Three-Step Procedure Using Stata, R, Mplus, and SPSS. *Int Rev Soc Psychol*. 2017;30(1):203-218. doi:10.5334/irsp.90
20. Kaplan MS, Huguet N, Feeny DH, McFarland BH. Self-reported hypertension prevalence and income among older adults in Canada and the United States. *Soc Sci Med*. 2010;70(6):844-849. doi:10.1016/j.socscimed.2009.11.019
21. Buys DR, Howard VJ, McClure LA, et al. Association Between Neighborhood Disadvantage and Hypertension Prevalence, Awareness, Treatment, and Control in Older Adults: Results From the University of Alabama at Birmingham Study of Aging. *Am J Public Health*. 2015;105(6):1181-1188. doi:10.2105/ajph.2014.302048
22. Isaac L, Laveist T. Neighborhood Assets and Hypertension in Disadvantaged Communities. presented at: APHA Annual Meeting and Exposition; 2010;
23. McMaughan DJ, Oloruntoba O, Smith ML. Socioeconomic Status and Access to Healthcare: Interrelated Drivers for Healthy Aging. 10.3389/fpubh.2020.00231. *Front Public Health*. 2020;8:231. doi:10.3389/fpubh.2020.00231

24. Lei X, Yin N, Zhao Y. Socioeconomic status and chronic diseases: The case of hypertension in China. *China Econ Rev.* 2012;23(1):105-121. doi:10.1016/j.chieco.2011.08.004
25. Zuo D, Chen J, Zhang Q, et al. The prevalence study on hypertension and risk factors among the middle- and the old-aged people in some areas of Hubei Province. *J Math Med.* 2008;21(6):698-701. doi:CNKI:SUN:SLYY.0.2008-06-026
26. Zhang X, Ma Y, Yang Y, et al. Prevalence, awareness, treatment, and control status of hypertension in the older adults population in Xinjiang. *Chin J Hypertens.* 2014;22(1):31-35. doi:10.16439/j.cnki.1673-7245.2014.01.010
27. Rajda C, George NM. The Effect of Education and Literacy Levels on Health Outcomes of the Elderly. *J Nurse Pract.* 2009;5(2):115-119. doi:10.1016/j.nurpra.2008.08.020
28. You Y, Teng W, Wang J, et al. Hypertension and physical activity in middle-aged and older adults in China. *Sci Rep.* 2018;8(1):16098. doi:10.1038/s41598-018-34617-y
29. Miao H, Weng X, Wu X, et al. Impact of physical activity on the prevalence of hypertension among the older adults in Beijing communities. *Adv Behav Health.* 2019;2(1):41-48. doi:10.25082/AHB.2019.01.003
30. Zimmer Z, Kwong J. Socioeconomic Status and Health among Older Adults in Rural and Urban China. *J Aging Health.* 2004;16(1):44-70. doi:10.1177/0898264303260440
31. Li Y, Sun X, Liu K. Review on the Relationship between Socioeconomic Status and Health Status of the Elderly at Home and Abroad. *Chin J Soc Med.* 2015;32(4):278-281.
32. Trachte F, Geyer S, Sperlich S. Impact of physical activity on self-rated health in older people: do the effects vary by socioeconomic status? *J Public Health.* 2016;38(4):754-759. doi:10.1093/pubmed/fdv198
33. Lang IA, Hubbard RE, Andrew MK, Llewellyn DJ, Melzer D, Rockwood K. Neighborhood Deprivation, Individual Socioeconomic Status, and Frailty in Older Adults. *J Am Geriatr Soc.* 2009;57(10):1776-1780. doi:10.1111/j.1532-5415.2009.02480.x
34. Lee S-G, Jeon S-Y. The relations of socioeconomic status to health status, health behaviors in the elderly. *J Prev Med Public Health.* 2005;38(2):154-162.
35. Huguét N, Kaplan MS, Feeny D. Socioeconomic status and health-related quality of life among elderly people: Results from the Joint Canada/United States Survey of Health. *Soc Sci Med.* 2008;66(4):803-810. doi:10.1016/j.socscimed.2007.11.011

Table 1 Hypertension status and lifestyle of the subjects

	Men						Women					
	2006		2009		2011		2006		2009		2011	
	n	prpt (%)	n	prpt (%)	n	prpt (%)	n	prpt (%)	n	prpt (%)	n	prpt (%)
hypertensive	369	33.21	571	50.49	610	45.97	419	32.38	662	52	735	48.77
diagnosed	991	89.76	972	75.35	1002	73.19	1104	87.2	1016	70.02	1070	68.37
medicated	76	36.36	261	83.92	301	82.69	117	43.17	378	87.3	423	85.98
ever smoke	760	64.35	761	58.99	807	58.95	158	11.57	108	7.44	129	8.24
ever drank	520	45.98	598	46.36	635	46.38	111	8.61	101	6.96	121	7.73
FIT \geq 30%	567	50.44	668	51.98	760	56.05	669	52.51	816	56.86	885	57.92
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
TEI	2135.44	706.68	2143.7	660.88	2024.92	802.32	1811.43	601.2	1816.21	638.12	1728.47	709.97
FIT	29.62	11.97	31.18	11.03	31.99	11.42	30.68	12.1	32.12	11.41	33.19	12.21
PAs	67.22	116.46	82.05	132.13	92.07	132.92	96.87	109.57	86.04	102.32	90.88	108.62

Notes: n=number, prpt=proportion, SD=standard deviation, diagnosed=had been previously diagnosed, FIT=fat in total energy intake, TEI=total energy intake, PAs= physical activities

Table 2 Multi-level regression models

	Men	Women
Fixed effect [OR (95% CI)]		
Intercept	0.65 (0.34~1.21)	1.20(0.44~3.20)
Age (Ref = \geq 80 years old)		
60-65 years old	2.71*** (1.64~4.47)	5.68***(3.20~9.82)
65-70 years old	1.58 (0.96~2.61)	4.26***(2.50~7.25)
70-75 years old	1.36 (0.82~2.25)	2.26***(1.35~3.79)
75-80 years old	1.58 (0.95~2.12)	0.80(0.49~1.30)
Education (Ref=high school)		
Never attended school	1.17 (0.75~1.82)	0.94*(0.50~1.79)
Primary school	0.97 (0.68~1.38)	0.49*(0.26~0.89)
Junior middle school	0.90 (0.62~1.31)	0.50(0.26~0.97)
PCHI (Ref= high)		
Low	1.90*** (1.36~2.67)	2.07***(1.28~3.11)
Middle	1.39* (1.02~1.90)	1.96***(1.33~2.90)
Family size (Ref= \geq 2 persons)		
1 person	0.79 (0.48~1.30)	0.78(0.50~1.20)
Urbanization level (Ref= high)		
Low	1.70*** (1.20~2.42)	1.41(0.93~2.16)
Middle	1.48* (1.07~2.04)	1.00(0.68~1.44)
Smoking (Ref= Yes)	0.77* (0.60~0.99)	0.71(0.43~1.19)
Drinking (Ref= Yes)	1.21 (0.96~1.53)	0.68(0.42~1.13)
PA (Ref= high)		
Low	0.76 (0.56~1.04)	0.88(0.60~1.29)
Middle	0.89 (0.68~1.17)	0.65**(0.47~0.90)
TEI (Ref = high)		
Low	1.11 (0.86~1.43)	0.74(0.53~1.05)
Middle	0.98 (0.76~1.25)	0.65*(0.47~0.91)
FIT (Ref= high)		
Low	0.80 (0.61~1.04)	1.68***(1.25~2.25)
Middle	0.85 (0.67~1.08)	1.17(0.89~1.54)
Randomized effect [variance (SE)]		
Intercept	redundant	2.07*(0.16)
Age	2.08***(0.30)	4.97(0.57)
Educational level	NS, O	NS, O
PCHI	1.56***(0.35)	2.07*(0.16)
Family size	NS, O	NS, O
Urbanization	1.54***(0.32)	4.97(0.57)
Drinking		3.09***(0.43)
PAs		1.73***(0.37)
Smoking	1.08***(0.30)	3.649***(0.33)
Residual effect [variance (SE)]		
2006	0.35***(0.06)	0.21***(0.04)

2009	0.51***(0.05)	0.23***(0.03)
2011	0.40***(0.05)	0.27***(0.03)
Summary		
Accurate prediction rate	96.00%	98.70%
-2 log pseudo-fitting	16289	20384

Note: Ref: control group; *= $p < 0.05$, **= $p < 0.01$, ***= $p < 0.001$; NS, O=not significant, omitted; SE=standard error, PCHI=per capita household income; PA=physical activity; TEI=total energy intake; FIT=proportion of fat in total energy.