1	Causal associations of education, lifestyle behaviors, and cardiometabolic traits
2	with epigenetic age acceleration: a Mendelian randomization study
3	
4	Lijie Kong, ^{1,2} * Chaojie Ye, ^{1,2} * Yiying Wang, ^{1,2} Jie Zheng, ³ Shuangyuan Wang, ^{1,2}
5	Hong Lin, ^{1,2} Zhiyun Zhao, ^{1,2} Mian Li, ^{1,2} Yu Xu, ^{1,2} Jieli Lu, ^{1,2} Yuhong Chen, ^{1,2} Min
6	Xu, ^{1,2} Weiqing Wang, ^{1,2} Guang Ning, ^{1,2} Yufang Bi, ^{1,2} Tiange Wang ^{1,2}
7	
8	*Lijie Kong and Chaojie Ye contributed equally as first authors.
9	
10	Author Affiliations:
11	¹ Department of Endocrine and Metabolic Diseases, Shanghai Institute of Endocrine
12	and Metabolic Diseases, Ruijin Hospital, Shanghai Jiao Tong University School of
13	Medicine, Shanghai, China;
14	² Shanghai National Clinical Research Center for Metabolic Diseases, Key Laboratory
15	for Endocrine and Metabolic Diseases of the National Health Commission of the PR
16	China, Shanghai Key Laboratory for Endocrine Tumor, State Key Laboratory of
17	Medical Genomics, Ruijin Hospital, Shanghai Jiao Tong University School of
18	Medicine, Shanghai, China;
19	³ MRC Integrative Epidemiology Unit (IEU), Bristol Medical School, University of
20	Bristol, Oakfield House, Oakfield Grove, Bristol, BS8 2BN, United Kingdom.
21	
22	Correspondence:
23	Tiange Wang, MD, PhD. Shanghai National Clinical Research Center for Endocrine
24	and Metabolic Diseases, Shanghai Institute of Endocrine and Metabolic Diseases,
25 NO	Ruijin Hospital, Shanghai Jiao Tong University School of Medicine, 197 Rujin 2nd TE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Road, Shanghai, China. E-mail: wtg@live.cn. Phone number: 86-021-64370045.

27 Key Points

28 •	Genetically p	predicted higher	educational a	attainment is	the strongest	protective
		4 1			4.7	

- 29 factor for both GrimAgeAccel and PhenoAgeAccel, independent of causal
- 30 lifestyle and cardiometabolic risk factors.
- Smoking is the leading causal lifestyle risk factor for GrimAgeAccel or
- 32 PhenoAgeAccel, followed by alcohol intake and daytime napping. Adiposity
- traits are the leading causal cardiometabolic risk factors for GrimAgeAccel and
- 34 PhenoAgeAccel, followed by type 2 diabetes, triglycerides, and C-reactive
- 35 protein.
- This study provides novel quantitative evidence on modifiable causal risk factors
- 37 for accelerated epigenetic ageing, indicating underlying contributors to the ageing

38 process and promising intervention targets to promote healthy longevity.

39 Abstract

40	Background: GrimAge acceleration (GrimAgeAccel) and PhenoAge acceleration
41	(PhenoAgeAccel) are DNA methylation-based markers of accelerated biological
42	ageing, standing out in predicting mortality and age-related cardiometabolic
43	morbidities. Causal risk factors for GrimAgeAccel and PhenoAgeAccel are unclear.
44	Objective: To evaluate causal associations of 18 common modifiable socioeconomic,
45	lifestyle, and cardiometabolic factors with GrimAgeAccel and PhenoAgeAccel.
46	Methods: We performed two-sample univariable and multivariable Mendelian
47	randomization (MR), using summary-level data for GrimAgeAccel and
48	PhenoAgeAccel derived from a genome-wide association study of 34,710 European
49	participants. We used the inverse-variance weighted method as the main analysis,
50	supplemented by three sensitivity analyses.
51	Results: Eleven and eight factors were causally associated with GrimAgeAccel and
52	PhenoAgeAccel, respectively. Smoking initiation was the strongest risk factor (β
53	[SE]: 1.299 [0.107] years) for GrimAgeAccel, followed by higher alcohol intake,
54	higher waist circumference, daytime napping, higher body fat percentage, higher
55	BMI, higher C-reactive protein, higher triglycerides, childhood obesity, and type 2
56	diabetes; whereas education was the strongest protective factor (β [SE] per 1-SD
57	increase in years of schooling: -1.143 [0.121] years). Higher waist circumference (β
58	[SE]: 0.850 [0.269] years) and education (β [SE]: -0.718 [0.151] years) were the
59	leading causal risk and protective factors for PhenoAgeAccel, respectively. Sensitivity
60	analyses strengthened the robustness of these causal associations, and the
61	multivariable MR analyses demonstrated independent direct effects of the strongest
62	risk and protective factors on GrimAgeAccel and PhenoAgeAccel, respectively.
63	Conclusion: Our findings provide novel quantitative evidence on modifiable causal

- risk factors for epigenetic ageing, and hint at underlying contributors and intervention
- 65 targets to the ageing process.
- 66
- 67 **Keywords:** Epigenetic ageing; Education; Lifestyle behaviors; Cardiometabolic
- 68 traits; Mendelian randomization.

69 Introduction

Ageing involves the gradual accumulation of a decline in multiple biological 70 functions over time, leading to increased risks of developing age-related diseases and 71 mortality [1,2]. Although chronological ageing is uniform and unchangeable, the rate 72 of biological ageing is variable and modifiable depending on individual genetics, 73 environmental exposures, and health-related behaviors [3]. Of several potential types 74 75 of biological age predictors (e.g., epigenetic clock, leukocyte telomere length, and transcriptomic predictors), the epigenetic clock that composed of DNA methylation at 76 77 multiple cytosine-phosphate-guanine (CpG) sites is currently the best one, as it correlates well with age and predicts mortality across populations [4]. The epigenetic 78 age acceleration is the difference between chronological age and epigenetic age, and 79 represents accelerated biological ageing [5]. The second-generation epigenetic age 80 acceleration indicators, namely GrimAge acceleration (GrimAgeAccel) and 81 PhenoAge acceleration (PhenoAgeAccel), have been evolved to incorporate ageing-82 related traits, and stand out in terms of predicting mortality and age-related 83 cardiometabolic morbidities [6-9]. 84 Limited observational evidence has suggested that certain socioeconomic, 85 lifestyle behaviors, and cardiometabolic traits may be related to GrimAgeAccel and 86 PhenoAgeAccel [6,7,10,11]. Whether and to what extent modifiable risk factors 87 influence GrimAgeAccel and PhenoAgeAccel, if causally established, could shed 88 light on potential contributors to the ageing process and elucidate promising targets 89 for preventing age-related diseases and improving healthy longevity [12,13]. Thus far, 90 such causal evidence is scarce. 91 To fill this knowledge gap, we applied Mendelian randomization (MR) to 92

93 evaluate the causal associations of 18 common modifiable socioeconomic, lifestyle

94	and cardiometabolic factors with GrimAgeAccel and PhenoAgeAccel. The MR
95	method uses genetic variants as instrumental variables (IVs) to infer causality among
96	correlated traits. Since genetic variants are randomly allocated at conception, the MR
97	study is less susceptible to confounding and reverse causality than conventional
98	observational studies [14]. We conducted both univariable and multivariable MR
99	analyses to discern if modifiable factors have independent direct causal effects on
100	GrimAgeAccel and PhenoAgeAccel.
101	
102	Methods
103	Study design
104	This MR study design is presented in Figure 1. We performed two-sample
105	univariable and multivariable MR strictly following the STROBE-MR guidelines
106	(Supplementary Table 1) [15]. To obtain unbiased estimates of the causal effects,
107	the MR analysis should adhere to three fundamental assumptions [16]: first, the IVs
108	are truly associated with the exposures; second, the IVs are independent of
109	confounders of the relationship between exposures and outcomes (GrimAgeAccel and
110	PhenoAgeAccel); third, the IVs influence the outcomes only through the exposures,
111	but not any direct or indirect pathways. All data used in this MR study are publicly
112	available. Ethical approval and informed consent had been obtained in all original
113	studies.
114	
115	Selection rationale and data sources of genetic instruments
116	We selected 18 common modifiable factors, including educational attainment,
117	lifestyle behaviors (smoking initiation, alcohol intake, coffee consumption, daytime

napping, sleep duration, and moderate-to vigorous physical activity [MVPA]), and

119	cardiometabolic traits (body mass index [BMI], waist circumference, body fat
120	percentage [BF%], childhood obesity, type 2 diabetes, low-density lipoprotein [LDL]
121	cholesterol, high-density lipoprotein [HDL] cholesterol, triglycerides, systolic blood
122	pressure [SBP], diastolic blood pressure [DBP], and C-reactive protein [CRP]).
123	Definitions of the 18 modifiable factors are shown in Supplementary Table 2 .
124	We extracted genetic variants for each of the modifiable factors from the largest
125	available genome-wide association studies (GWASs) of European ancestry [17-29],
126	ensuring minimum sample overlap with the genetic variants for GrimAgeAccel and
127	PhenoAgeAccel (Supplementary Table 3). We included single nucleotide
128	polymorphisms (SNPs) robustly associated with the 18 modifiable factors at the
129	genome-wide significance (P $< 5 \times 10-8$). To select independent genetic variants, a
130	stringent condition (linkage disequilibrium threshold of $r^2 < 0.01$) was set to minimize
131	the influence of linkage disequilibrium which may bias the results of randomized
132	allele allocation. Where SNPs for the exposures were not available in the GWAS
133	summary statistics of GrimAgeAccel or PhenoAgeAccel, we used proxies of SNPs
134	with $r^2 > 0.8$ as substitutes, by using the LDproxy search on the online platform
135	LDlink (https://ldlink.nci.nih.gov/) [30].
136	
137	Data source for epigenetic age acceleration

138 The genetic associations with GrimAgeAccel and PhenoAgeAccel were extracted

139 from a recent GWAS meta-analysis (summary statistics available at

140 https://datashare.is.ed.ac.uk/handle/10283/3645), which included 34,710 European

141 participants from 28 cohorts [31]. GrimAgeAccel and PhenoAgeAccel are two-

142 generation epigenetic age acceleration indicators, expressing the biological ageing

143 rate in years, of which GrimAgeAccel is more strongly associated with mortality than

PhenoAgeAccel [11]. Detailed definitions of GrimAgeAccel and PhenoAgeAccel and
data preparation in GWAS are shown in Supplementary Table 2.

146

147 Statistical analyses

In the main analysis, we used the inverse-variance weighted (IVW) method to 148 determine MR causal estimates (B coefficients with standard errors [SEs]) for 149 150 associations of each modifiable factor with GrimAgeAccel and PhenoAgeAccel. The IVW combined the Wald ratio estimates of every single SNP in the set of IVs into one 151 152 causal estimate using the random-effects meta-analysis approach [32]. To evaluate the robustness of the IVW estimates under different assumptions and to detect possible 153 pleiotropy, we performed three sensitivity analyses, including the MR weighted 154 median, the MR Egger, and the MR pleiotropy residual sum and outlier (MR-155 PRESSO) methods. The weighted median method selected the median MR estimate 156 157 as the causal estimate, and provided a consistent causal estimate if over 50% of the weight in the analysis was derived from valid IVs [33]. The MR Egger method which 158 allowed the intercept to be freely estimated as an indicator of pleiotropy was used to 159 identify and adjust for the potential directional pleiotropic bias, but has limited 160 precision [34]. The MR-PRESSO method was applied to detect and correct for any 161 outlier SNP reflecting likely horizontal pleiotropic biases for MR causal estimates 162 [35]. We evaluated the heterogeneity for the IVW estimates using the Cochran's Q 163 test [36], and identified the horizontal pleiotropy based on the p-value for the intercept 164 in the MR-Egger model [34]. A false discovery rate (FDR) method was used to 165 correct results for multiple testing, and FDR q-values were provided. In this study, 166 strong causal evidence was defined as an association supported by the main analysis 167 (FDR q-value <0.05) and at least one sensitivity analysis. Suggestive causal evidence 168

169	was defined as a suggest	association with P	< 0.05 and FDR o	γ value ≥ 0.05 in the main
-----	--------------------------	--------------------	------------------	--

- analysis. Null causal evidence was defined as no statistically significant association
- 171 revealed from the main analysis (P \ge 0.05).
- 172 We further conducted multivariable MR (MVMR) analyses to assess whether the
- 173 causal effects of the strongest protective factor and the strongest risk factor on
- 174 GrimAgeAccel and PhenoAgeAccel were independent of other exposure factors [37].
- 175 Taking into account the effect size and significance of causal associations, we
- selected exposure factors with β coefficients >0.5 and P <0.05 in the main analysis as
- 177 covariates in the adjustment models.
- 178 The two-sample MR analyses were conducted with the R packages

179 'TwoSampleMR' and 'MRPRESSO' in R software (version 4.0.3). FDR q-values were

180 estimated using the R package '*fdrtool*'.

- 181
- 182 **Results**

183 MR results for associations between 18 modifiable factors and GrimAgeAccel

- 184 Ten out of the 18 risk factors showed strong associations with increased
- 185 GrimAgeAccel after FDR adjustment for multiple comparisons (Table 1). Smoking
- initiation was the strongest risk factor (β [SE]: 1.299 [0.107] years) for
- 187 GrimAgeAccel, followed by higher alcohol intake (β [SE] per 1-SD increase: 0.899
- 188 [0.361] years), higher waist circumference (β [SE] per 1-SD increase: 0.815 [0.184]
- 189 years), daytime napping (β [SE]: 0.805 [0.355] years), higher BF% (β [SE] per 1-SD
- 190 increase: 0.748 [0.120] years), higher BMI (β [SE] per 1-SD increase: 0.592 [0.079]
- 191 years), higher CRP (β [SE] per 1-SD increase: 0.345 [0.073] years), higher
- triglycerides (β [SE] per 1-SD increase: 0.249 [0.091] years), childhood obesity (β
- 193 [SE]: 0.200 [0.075] years), and type 2 diabetes (β [SE]: 0.095 [0.041] years; Figure

194 **2a**). By contrast, educational attainment in years of schooling was the only one

195 protective factor associated with decreased GrimAgeAccel (β [SE] per 1-SD increase:

196 -1.143 [0.121] years). There was little evidence to support a causal association of

197 other modifiable factors with GrimAgeAccel.

198 Associations of the above 11 modifiable factors with GrimAgeAccel were robust

across sensitivity analyses with consistent effect directions and P < 0.05 in at least one

200 sensitivity analysis. The MR-Egger intercept tests indicated potential pleiotropy for

201 CRP (P_{intercept} <0.05; Supplementary Table 4). Cochran's Q-test showed possible

202 heterogeneity for educational attainment, alcohol intake, BMI, BF%, type 2 diabetes,

LDL cholesterol, HDL cholesterol, and CRP (P_h<0.05; **Supplementary Table 4**).

204 However, with the exclusion of outlying SNPs, the MR-PRESSO analysis showed

consistent results with the IVW analysis (**Table 1**).

206

207 MR results for associations between 18 modifiable factors and PhenoAgeAccel

208 Eight modifiable factors showed strong associations with PhenoAgeAccel after FDR

adjustment for multiple comparisons (**Table 2**). Higher waist circumference was the

strongest risk factor (β [SE] per 1-SD increase: 0.850 [0.269] years) for

211 PhenoAgeAccel, followed by higher BF% (β [SE] per 1-SD increase: 0.711 [0.152]

years), higher BMI (β [SE] per 1-SD increase: 0.586 [0.102] years), smoking

initiation (β [SE]: 0.519 [0.142] years), higher CRP (β [SE] per 1-SD increase: 0.349

[0.095] years), childhood obesity (β [SE]: 0.229 [0.095] years), and type 2 diabetes (β

[SE]: 0.125 [0.051] years). Whereas the genetically predicted higher educational

attainment in years of schooling was associated with decreased PhenoAgeAccel (β

[SE] per 1-SD increase: -0.718 [0.151] years; Figure 2b). Suggestive causality was

218 identified for the association between genetically predicted higher DBP and increased

PhenoAgeAccel (β [SE] per 1-mmHg increase: 0.049 [0.024] years). No significant 219 association was observed with PhenoAgeAccel for the other modifiable factors. 220 Associations for the above eight modifiable factors were robust in sensitivity 221 analyses with consistent effect directions and P <0.05 in at least one sensitivity 222 analysis. There was no evidence of pleiotropy for these risk factors except for CRP 223 (Pintercept <0.05; Supplementary Table 5). Potential heterogeneity was observed for 224 225 educational attainment, smoking initiation, BMI, waist circumference, BF%, type 2 diabetes, and CRP (Ph<0.05). One to four outliers were detected in the MR-PRESSO 226 227 analyses; however, the associations remained consistent after removal of these outliers (Table 2). For SBP, with excluding the outlying SNP rs62523863, the MR-228 PRESSO analysis revealed a potentially positive association between genetically 229 predicted higher SBP and PhenoAgeAccel (ß [SE] per 1-mmHg increase: 0.038 230 [0.016] years). 231

232

233 MVMR results for associations of the strongest protective and risk factors with

234 GrimAgeAccel and PhenoAgeAccel

In MVMR analyses, the association between educational attainment (the strongest
 protective factor) and GrimAgeAccel remained significant with adjustment for

smoking initiation, alcohol intake, waist circumference, daytime napping, BF%, or

BMI (Figure 3a). Likewise, the significant association between smoking initiation

239 (the strongest risk factor) and GrimAgeAccel persisted after adjustment for education,

alcohol intake, waist circumference, daytime napping, BF%, or BMI. All covariates in

- 241 MVMR analyses were selected based on their considerable effects on GrimAgeAccel
- in terms of size (β coefficients >0.5) and significance (P <0.05 in the main analysis).

The associations of education (the strongest protective factor) and waist
circumference (the strongest risk factor) with PhenoAgeAccel remained significant
with adjustment for waist circumference, BF%, BMI, or smoking initiation, which
had considerable causal effects on PhenoAgeAccel (β coefficients >0.5; Figure 3b).

This MR study for the first time delineated potential causal relationships of 18

common modifiable factors with GrimAgeAccel and PhenoAgeAccel, the robust

251 second-generation epigenetic age acceleration indicators. We identified strong

evidence for 11 and eight factors associated with GrimAgeAccel and PhenoAgeAccel,

253 respectively. Smoking initiation exhibited the greatest effect on increased

254 GrimAgeAccel (1.299 years), followed by higher alcohol intake, higher waist

circumference, daytime napping, higher BF%, higher BMI, higher CRP, higher

triglycerides, childhood obesity, and type 2 diabetes; whereas educational attainment

showed the greatest effect on decreased GrimAgeAccel (-1.143 years per 1-SD

258 increase in years of schooling). Higher waist circumference and educational

attainment were the leading causal risk and protective factors associated with

260 PhenoAgeAccel, respectively; BF%, BMI, smoking initiation, CRP, childhood

261 obesity, and type 2 diabetes were also associated with increased PhenoAgeAccel.

262 Multiple sensitivity analyses further strengthened the robustness of these causal

263 relationships, and the MVMR analyses demonstrated the independent direct effects of

the strongest risk and protective factors on GrimAgeAccel and PhenoAgeAccel,

respectively. Suggestive causality was identified for the association between higher

266 DBP and increased PhenoAgeAccel.

In this study, educational attainment was the major protective factor for both 267 GrimAgeAccel and PhenoAgeAccel, and this causal effect was largely independent of 268 other causal factors, such as smoking and adiposity-related traits. Educational 269 attainment is a strong proxy for socioeconomic status and a more upstream 270 determinant of health, with broad implications for a person's life-long lifestyle 271 behaviors and health-promoting resources [38]. A recent study using UK Biobank 272 273 data has documented that each 1-year increase in genetically determined educational attainment was associated with equivalently 4.2 years of age-related increases in 274 275 telomere length [39]. Telomere length and epigenetic age acceleration metrics (e.g., GrimAgeAccel and PhenoAgeAccel) point towards distinct mechanisms of the ageing 276 process that are marked by telomeres and the DNA methylation-based epigenetic 277 clocks, both of which are independently associated with chronological age and 278 mortality risk [40]. Our findings, together with those of the UK Biobank, highlight the 279 important impact of educational attainment on biological ageing rates from two 280 different aspects of the ageing process. Therefore, public health strategies aimed at 281 reducing educational inequalities and improving educational attainment may slow the 282 biological ageing rate and help reduce age-related health burdens. 283 This MR study identified several common lifestyle behaviors, including smoking 284

initiation, alcohol intake, and daytime napping, causally associated with increased GrimAgeAccel or PhenoAgeAccel (smoking initiation only), which were consistent with previous evidence from observational studies [6,7]. Smoking methylation proxy is a component of the GrimAge clock [7], thus it is not surprising that smoking initiation exhibited a large effect on increased GrimAgeAccel in this study, and the MVMR analysis further confirmed that the effect of smoking on GrimAgeAccel was independent of other causal lifestyle behaviors and adiposity-related traits. Similarly,

292	genetically determined smoking has been associated with shorter telomere length in
293	the UK Biobank [41], and evidence from the Danish Health Interview Survey
294	suggested that the life expectancy of a heavy smoker was a little more than seven
295	years shorter than that of a never smoker [42]. Our study also provided strong MR
296	evidence that genetically predicted higher alcohol intake was associated with
297	increased GrimAgeAccel. A study composed of the Hannum cohort and Family and
298	Community Health Studies cohort found that the relationship between alcohol use and
299	the first-generation epigenetic clocks seemed to be nonlinear [42]. However, given the
300	dose-dependent relationship of alcohol intake with all-cause mortality and cancers
301	[43], our findings suggest that reducing alcohol intake is necessary to decrease
302	GrimAgeAccel and retard the overall health loss. Previous observational studies also
303	reported positive or inverse correlations of other lifestyle behaviors such as MVPA,
304	coffee, and sleep duration with GrimAgeAccel, PhenoAgeAccel, or mortality
305	[11,44,45]. Nevertheless, in this study, there was little evidence supporting these
306	associations were causal. The discrepancy between our findings and previous
307	observations may result from the potential confounding or reverse causation in
308	conventional observational studies. Moreover, the non-linear association patterns, as
309	in the case of sleep duration and mortality, might also partially explain the
310	inconsistent results [46]. Therefore, our null findings should be cautiously interpreted.
311	Interestingly, of all cardiometabolic traits (i.e., adiposity traits, type 2 diabetes,
312	triglycerides, and CRP) which causally increased GrimAgeAccel or PhenoAgeAccel,
313	adiposity traits were the most dominant traits. Emerging observational studies have
314	pronounced the positive associations of BMI with GrimAgeAccel and
315	PhenoAgeAccel [11], and a meta-analysis of 87 observational studies showed each 5-
316	kg/m ² higher BMI corresponded to about 1 year of age-related decrease in telomere

length [46]. Our study further provided strong evidence for causal associations of 317 various adiposity phenotypes, including waist circumference, BF%, BMI, and 318 childhood obesity, with increased GrimAgeAccel and PhenoAgeAccel. In this study, 319 type 2 diabetes, triglycerides, and CRP showed strong but modest effects (ß 320 coefficients ≤ 0.5) on GrimAgeAccel or PhenoAgeAccel, which was consistent with 321 322 the findings of the UK Biobank study on telomere length [39]. 323 This MR study yields new insights into modifiable causal factors, including education, lifestyle behaviors, and cardiometabolic traits, associated with accelerating 324 325 or decelerating epigenetic ageing. This study brings us one step closer to understanding the potential contributors to the ageing process and provides promising 326 intervention targets for healthy ageing. Given the enormous burden induced by age-327 related morbidity and mortality, strategies to reduce educational inequalities, promote 328 healthy lifestyles primarily through reducing smoking, alcohol intake, and daytime 329 napping, and improve cardiometabolic traits, specifically adiposity, type 2 diabetes, 330 triglycerides, and CRP, to slow biological ageing rate are imminent. 331 In this study, we included independent and genome-wide significant SNPs as 332 instruments for each of the modifiable factors to ensure the first MR assumption was 333 fulfilled. Moreover, we applied strict criteria strengthened by the FDR-corrected 334 significance and the cross-validations by main and sensitivity analyses to draw robust 335 causal conclusions. However, several limitations merited consideration. First, we 336 found potential pleiotropy from the MR-Egger intercept test for CRP. However, we 337 conducted MR-PRESSO analysis, and the association remained consistent after 338 removal of outlying SNPs. Second, we could not rule out the possibility that the 339 associations of certain modifiable risk factors such as alcohol intake and sleep 340 duration with GrimAgeAccel or PhenoAgeAccel may be non-linear. Future studies 341

542 with marviauar revertatia are warranted to commit the intear of non-inte	al of non-inteal
--	------------------

- relationships. Third, to ensure the consistency of genetic background, this MR study
- 344 was performed only in European-ancestry participants, thus the generalization of our
- results to other ethnic groups should be cautious.
- 346

347 Conclusions

- 348 This MR study provided novel quantitative evidence on modifiable causal
- 349 socioeconomic, lifestyle, and cardiometabolic factors for accelerated epigenetic
- ageing. Our findings shed light on the underlying contributors to the biological ageing
- 351 process and point toward promising intervention targets to slow the biological ageing
- 352 rate and promote healthy longevity.

353 List of abbreviations

BF%	Body fat percentage
BMI	Body mass index
CI	Confidence interval
CpG	Cytosine-phosphate-guanine
CRP	C-reactive protein
DBP	Diastolic blood pressure
FDR	False discovery rate
GrimAgeAccel	GrimAge acceleration
GWAS	Genome-wide association study
HDL	High-density lipoprotein
IV	Instrumental variable
IVW	Inverse variance weighted
LDL	Low-density lipoprotein
MVPA	Moderate-to vigorous physical activity
MR	Mendelian randomization
MVMR	Multivariable Mendelian randomization
PhenoAgeAccel	PhenoAge acceleration
PRESSO	Pleiotropy residual sum and outlier
SBP	Systolic blood pressure
SD	Standard deviation
SE	Standard errors
SNP	Single nucleotide polymorphism
WM	Weighted median

355 Conflict-of-interest statement

- 356 The authors declare that they have no competing interests.
- 357

358 Ethics committee approval

- 359 This study is based on publicly available summarized data. Ethical approval and
- informed consent had been obtained in all original studies.
- 361

362 Availability of data and materials

- 363 The genetic association data of the 18 modifiable factors are available in
- 364 Supplementary Table 3. The GWAS summary statistics for GrimAgeAccel and
- 365 PhenoAgeAccel are available at https://datashare.is.ed.ac.uk/handle/10283/3645. The
- analytical script of the MR analyses conducted in this study is available via the
- 367 GitHub repository of the "TwoSampleMR" R package
- 368 (https://github.com/MRCIEU/TwoSampleMR/).
- 369

370 Funding

- 371 This work was supported by the grants from the National Natural Science Foundation
- of China (82022011, 81970706, 81730023, 82088102, 81970728, 81941017), the
- 373 Chinese Academy of Medical Sciences (2018PT32017, 2019PT330006), the
- 374 "Shanghai Municipal Education Commission–Gaofeng Clinical Medicine Grant
- 375 Support" from Shanghai Jiao Tong University School of Medicine (20171901 Round
- 2), the Shanghai Shenkang Hospital Development Center (SHDC12019101,
- 377 SHDC2020CR1001A, SHDC2020CR3064B), the Shanghai Jiao Tong University
- 378 School of Medicine (DLY201801), and the Ruijin Hospital (2018CR002).

379

380 Authors' contributions

381	LK and TW	contributed to	the concer	otion and	design	of the stu	ıdv. LK	and C	ĽY
001									

- 382 contributed to statistical analysis. LK contributed to drafting of the manuscript. TW
- 383 guaranteed this work and take responsibility for the integrity of the data. All authors
- 384 contributed to acquisition or interpretation of data, critical revision of the manuscript
- 385 for important intellectual content, and final approval of the version to be published.
- 386

387 Acknowledgements

- 388 We gratefully acknowledge the investigators and participants of all genome-wide
- association studies from which we used data.

390 **References**

- 1. Niccoli T, Partridge L. Ageing as a risk factor for disease. *Curr Biol.*
- 392 2012;22(17):R741-R52.
- 2. Schumacher B, Pothof J, Vijg J, Hoeijmakers JHJ. The central role of DNA damage
- in the ageing process. *Nature*. 2021;592(7856):695-703.
- 395 3. Warner HR. Current status of efforts to measure and modulate the biological rate of
- aging. J Gerontol A Biol Sci Med Sci. 2004;59(7):692-6.
- 4. Jylhävä J, Pedersen NL, Hägg S. Biological Age Predictors. *EBioMedicine*.
- 398 2017;21:29-36.
- 399 5. Horvath S, Raj K. DNA methylation-based biomarkers and the epigenetic clock
- 400 theory of ageing. *Nat Rev Genet.* 2018;19(6):371-84.
- 401 6. Levine ME, Lu AT, Quach A, Chen BH, Assimes TL, Bandinelli S, et al. An
- 402 epigenetic biomarker of aging for lifespan and healthspan. *Aging (Albany NY)*.
- 403 2018;10(4):573-91.
- 404 7. Lu AT, Quach A, Wilson JG, Reiner AP, Aviv A, Raj K, et al. DNA methylation
- 405 GrimAge strongly predicts lifespan and healthspan. *Aging (Albany NY)*.
- 406 2019;11(2):303-27.
- 407 8. Wang C, Ni W, Yao Y, et al. DNA methylation-based biomarkers of age
- 408 acceleration and all-cause death, myocardial infarction, stroke, and cancer in two
- 409 cohorts: The NAS, and KORA F4. *EBioMedicine*. 2021;63:103151.
- 410 9. Li A, Koch Z, Ideker T. Epigenetic aging: Biological age prediction and informing
- a mechanistic theory of aging [published online ahead of print, 2022 Jun 20]. J Intern

- 412 *Med*.
- 413 10. Quach A, Levine ME, Tanaka T, Lu AT, Chen BH, Ferrucci L, et al. Epigenetic
- 414 clock analysis of diet, exercise, education, and lifestyle factors. *Aging (Albany NY)*.
- 415 2017;9(2):419-46.
- 416 11. Oblak L, van der Zaag J, Higgins-Chen AT, Levine ME, Boks MP. A systematic
- 417 review of biological, social and environmental factors associated with epigenetic
- 418 clock acceleration. *Ageing Res Rev.* 2021;69:101348.
- 419 12. Sierra F, Caspi A, Fortinsky RH, Haynes L, Lithgow GJ, Moffitt TE, et al. Moving
- 420 geroscience from the bench to clinical care and health policy. *J Am Geriatr Soc.*
- 421 2021;69(9):2455-63.
- 422 13. Campisi J, Kapahi P, Lithgow GJ, Melov S, Newman JC, Verdin E. From
- 423 discoveries in ageing research to therapeutics for healthy ageing. *Nature*.
- 424 2019;571(7764):183-92.
- 425 14. Davies NM, Holmes MV, Davey Smith G. Reading Mendelian randomisation
- studies: a guide, glossary, and checklist for clinicians. *BMJ*. 2018;362:k601.
- 427 15. Skrivankova VW, Richmond RC, Woolf BAR, Davies NM, Swanson SA,
- 428 VanderWeele TJ, et al. Strengthening the reporting of observational studies in
- 429 epidemiology using mendelian randomisation (STROBE-MR): explanation and
- 430 elaboration. *BMJ*. 2021;375:n2233.
- 16. Zheng J, Baird D, Borges M-C, Bowden J, Hemani G, Haycock P, et al. Recent
- 432 Developments in Mendelian Randomization Studies. *Curr Epidemiol Rep.*
- 433 2017;4(4):330-45.

- 434 17. Lee JJ, Wedow R, Okbay A, et al. Gene discovery and polygenic prediction from a
- 435 genome-wide association study of educational attainment in 1.1 million individuals.
- 436 *Nat Genet*. 2018;50(8):1112-1121.
- 437 18. Liu M, Jiang Y, Wedow R, et al. Association studies of up to 1.2 million
- 438 individuals yield new insights into the genetic etiology of tobacco and alcohol use.
- 439 *Nat Genet*. 2019;51(2):237-244.
- 440 19. Zhong VW, Kuang A, Danning RD, et al. A genome-wide association study of
- 441 bitter and sweet beverage consumption [published correction appears in Hum Mol
- 442 Genet. 2019 May 02;:]. *Hum Mol Genet*. 2019;28(14):2449-2457.
- 443 20. Dashti HS, Daghlas I, Lane JM, et al. Genetic determinants of daytime napping
- and effects on cardiometabolic health. *Nat Commun.* 2021;12(1):900.
- 445 21. Dashti HS, Jones SE, Wood AR, et al. Genome-wide association study identifies
- genetic loci for self-reported habitual sleep duration supported by accelerometer-
- 447 derived estimates. *Nat Commun.* 2019;10(1):1100.
- 448 22. Klimentidis YC, Raichlen DA, Bea J, et al. Genome-wide association study of
- habitual physical activity in over 377,000 UK Biobank participants identifies multiple
- 450 variants including CADM2 and APOE. Int J Obes (Lond). 2018;42(6):1161-1176.
- 451 23. Yengo L, Sidorenko J, Kemper KE, et al. Meta-analysis of genome-wide
- 452 association studies for height and body mass index in \sim 700000 individuals of
- 453 European ancestry. *Hum Mol Genet*. 2018;27(20):3641-3649.
- 454 24. Shungin D, Winkler TW, Croteau-Chonka DC, et al. New genetic loci link adipose
- and insulin biology to body fat distribution. *Nature*. 2015;518(7538):187-196.

- 456 25. Bradfield JP, Taal HR, Timpson NJ, et al. A genome-wide association meta-
- 457 analysis identifies new childhood obesity loci. *Nat Genet.* 2012;44(5):526-531.
- 458 26. Mahajan A, Taliun D, Thurner M, et al. Fine-mapping type 2 diabetes loci to
- 459 single-variant resolution using high-density imputation and islet-specific epigenome
- 460 maps. *Nat Genet*. 2018;50(11):1505-1513.
- 461 27. Klarin D, Damrauer SM, Cho K, et al. Genetics of blood lipids among ~300,000
- 462 multi-ethnic participants of the Million Veteran Program. *Nat Genet*.
- 463 2018;50(11):1514-1523.
- 464 28. Evangelou E, Warren HR, Mosen-Ansorena D, et al. Genetic analysis of over 1
- 465 million people identifies 535 new loci associated with blood pressure traits [published
- 466 correction appears in Nat Genet. 2018 Dec;50(12):1755]. Nat Genet.
- 467 2018;50(10):1412-1425.
- 468 29. Han X, Ong JS, An J, Hewitt AW, Gharahkhani P, MacGregor S. Using Mendelian
- randomization to evaluate the causal relationship between serum C-reactive protein
- 470 levels and age-related macular degeneration. *Eur J Epidemiol*. 2020;35(2):139-146.
- 471 30. Machiela MJ, Chanock SJ. LDlink: a web-based application for exploring
- 472 population-specific haplotype structure and linking correlated alleles of possible
- 473 functional variants. *Bioinformatics*. 2015;31(21):3555-7.
- 474 31. McCartney DL, Min JL, Richmond RC, Lu AT, Sobczyk MK, Davies G, et al.
- 475 Genome-wide association studies identify 137 genetic loci for DNA methylation
- biomarkers of aging. *Genome Biol.* 2021;22(1):194.
- 477 32. Lawlor DA, Harbord RM, Sterne JA, Timpson N, Davey Smith G. Mendelian

- 478 randomization: using genes as instruments for making causal inferences in
- 479 epidemiology. *Stat Med.* 2008;27(8):1133-1163.
- 480 33. Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent Estimation in
- 481 Mendelian Randomization with Some Invalid Instruments Using a Weighted Median
- 482 Estimator. *Genet Epidemiol*. 2016;40(4):304-314.
- 483 34. Burgess S, Bowden J, Fall T, Ingelsson E, Thompson SG. Sensitivity Analyses for
- 484 Robust Causal Inference from Mendelian Randomization Analyses with Multiple
- 485 Genetic Variants. *Epidemiology (Cambridge, Mass)*. 2017;28(1):30-42.
- 486 35. Verbanck M, Chen C-Y, Neale B, Do R. Detection of widespread horizontal
- 487 pleiotropy in causal relationships inferred from Mendelian randomization between
- 488 complex traits and diseases. *Nat Genet.* 2018;50(5):693-8.
- 489 36. Bowden J, Del Greco M F, Minelli C, Davey Smith G, Sheehan N, Thompson J. A
- 490 framework for the investigation of pleiotropy in two-sample summary data Mendelian
- 491 randomization. *Stat Med.* 2017;36(11):1783-802.
- 492 37. Burgess S, Thompson SG. Multivariable Mendelian randomization: the use of
- 493 pleiotropic genetic variants to estimate causal effects. *Am J Epidemiol*.
- 494 2015;181(4):251-260.
- 495 38. Figgins E, Pieruccini-Faria F, Speechley M, Montero-Odasso M. Potentially
- 496 modifiable risk factors for slow gait in community-dwelling older adults: A systematic
- 497 review. Ageing Res Rev. 2021;66:101253.
- 498 39. Bountziouka V, Musicha C, Allara E, et al. Modifiable traits, healthy behaviours,
- and leukocyte telomere length: a population-based study in UK Biobank. Lancet

- 500 *Healthy Longev.* 2022;3(5):e321-e331.
- 40. Marioni RE, Harris SE, Shah S, et al. The epigenetic clock and telomere length
- are independently associated with chronological age and mortality [published
- 503 correction appears in Int J Epidemiol. 2018 Feb 1;47(1):356]. *Int J Epidemiol*.
- 504 2018;45(2):424-432.
- 41. Brønnum-Hansen H, Juel K. Abstention from smoking extends life and
- 506 compresses morbidity: a population based study of health expectancy among smokers
- and never smokers in Denmark. *Tob Control*. 2001;10(3):273-278.
- 42. Beach SR, Dogan MV, Lei MK, et al. Methylomic Aging as a Window onto the
- 509 Influence of Lifestyle: Tobacco and Alcohol Use Alter the Rate of Biological Aging. J
- 510 *Am Geriatr Soc.* 2015;63(12):2519-2525.
- 43. GBD 2016 Alcohol and Drug Use Collaborators. The global burden of disease
- attributable to alcohol and drug use in 195 countries and territories, 1990-2016: a
- 513 systematic analysis for the Global Burden of Disease Study 2016 [published
- 514 correction appears in Lancet Psychiatry. 2019 Jan;6(1):e2]. *Lancet Psychiatry*.
- 515 2018;5(12):987-1012.
- 44. Loftfield E, Cornelis MC, Caporaso N, Yu K, Sinha R, Freedman N. Association
- of Coffee Drinking With Mortality by Genetic Variation in Caffeine Metabolism:
- 518 Findings From the UK Biobank. JAMA Intern Med. 2018;178(8):1086-1097.
- 45. Wang C, Bangdiwala SI, Rangarajan S, et al. Association of estimated sleep
- duration and naps with mortality and cardiovascular events: a study of 116 632 people
- 521 from 21 countries. *Eur Heart J.* 2019;40(20):1620-1629.

- 522 46. Gielen M, Hageman GJ, Antoniou EE, et al. Body mass index is negatively
- 523 associated with telomere length: a collaborative cross-sectional meta-analysis of 87
- 524 observational studies. *Am J Clin Nutr.* 2018;108(3):453-475.

		Main analysis			Sensitivity analyses										
		IVW			Wei	Weighted median			MR-Egger			MR-PRESSO			
	No. of										No. of				
Modifiable factor	SNP	β (SE)	P value	q value	β (SE)	P value	q value	β (SE)	P value	q value	outlier	β (SE)	P value	q value	
Socioeconomic factor															
Educational attainment	751	-1.143 (0.121)	3.31E-21	2.98E-20	-1.129 (0.172)	5.47E-11	4.92E-10	-0.829 (0.452)	0.067	0.277	4	-1.116 (0.116)	6.54E-21	5.89E-20	
Lifestyle behavior															
Smoking initiation	299	1.299 (0.107)	4.90E-34	8.82E-33	1.257 (0.157)	1.42E-15	2.56E-14	1.321 (0.446)	0.003	0.027	0	1.276 (0.103)	7.63E-29	1.37E-27	
Alcohol intake	80	0.899 (0.361)	0.013	0.026	0.380 (0.567)	0.502	0.602	0.481 (0.738)	0.516	0.771	0	0.967 (0.357)	0.008	0.018	
Coffee consumption	12	0.003 (0.005)	0.589	0.624	-0.003 (0.006)	0.615	0.692	-0.008 (0.010)	0.464	0.771	0	0.003 (0.005)	0.600	0.635	
Sleep															
Daytime napping	115	0.805 (0.355)	0.023	0.038	0.524 (0.514)	0.308	0.449	0.411 (1.235)	0.740	0.833	0	0.813 (0.346)	0.020	0.033	
Sleep duration	77	-0.301 (0.266)	0.257	0.356	-0.403 (0.400)	0.313	0.449	-0.077 (1.017)	0.940	0.940	0	-0.275 (0.261)	0.294	0.378	
MVPA	6	-0.186 (0.903)	0.837	0.837	-0.036 (1.198)	0.976	0.976	-9.279 (5.557)	0.170	0.510	0	-0.186 (0.763)	0.817	0.817	
Cardiometabolic trait															
Adiposity															
BMI	941	0.592 (0.079)	9.16E-14	5.50E-13	0.490 (0.124)	7.92E-05	3.56E-04	0.790 (0.231)	0.001	0.018	2	0.595 (0.078)	6.30E-14	3.78E-13	
Waist circumference	44	0.815 (0.184)	9.26E-06	2.78E-05	0.776 (0.272)	0.004	0.014	0.913 (0.504)	0.077	0.277	0	0.815 (0.184)	6.31E-05	1.89E-04	
BF%	641	0.748 (0.120)	4.12E-10	1.85E-09	0.740 (0.182)	4.87E-05	2.92E-04	0.843 (0.400)	0.036	0.216	3	0.796 (0.115)	1.19E-11	5.36E-11	
Childhood obesity	5	0.200 (0.075)	0.007	0.016	0.142 (0.098)	0.146	0.292	0.789 (0.554)	0.249	0.560	0	0.204 (0.057)	0.016	0.029	
Type 2 diabetes	232	0.095 (0.041)	0.020	0.036	0.084 (0.071)	0.236	0.425	0.043 (0.092)	0.644	0.773	0	0.100 (0.040)	0.013	0.026	
Lipids															
LDL cholesterol	145	-0.094 (0.096)	0.327	0.392	-0.064 (0.163)	0.695	0.736	0.016 (0.142)	0.908	0.940	1	-0.069 (0.094)	0.463	0.542	
HDL cholesterol	223	-0.181 (0.095)	0.056	0.084	-0.267 (0.138)	0.053	0.136	-0.078 (0.162)	0.632	0.773	2	-0.175 (0.090)	0.054	0.081	
Triglycerides	173	0.249 (0.091)	0.006	0.015	0.239 (0.144)	0.096	0.216	0.152 (0.152)	0.319	0.638	0	0.254 (0.091)	0.006	0.015	
Blood pressure															
SBP	222	0.013 (0.012)	0.279	0.359	0.015 (0.018)	0.421	0.541	0.029 (0.049)	0.557	0.771	0	0.013 (0.012)	0.282	0.378	
DBP	264	0.011 (0.019)	0.565	0.624	0.027 (0.027)	0.324	0.449	0.083 (0.066)	0.213	0.548	0	0.013 (0.018)	0.482	0.542	
CRP	299	0.345 (0.073)	2.31E-06	8.32E-06	0.245 (0.107)	0.022	0.066	0.060 (0.100)	0.553	0.771	1	0.348 (0.071)	1.64E-06	5.90E-06	

Table 1. MR results of the associations between 18 modifiable factors and GrimAgeAccel

GrimAgeAccel represents epigenetic-age acceleration obtained using the GrimAge clock. The q value represents the false discovery rate (FDR)-adjusted *P* value. β represents the associations with GrimAgeAccel of respectively: 1-SD increase in years of educational attainment; Ever smoked regularly compared to never smoked; 1-SD increase in log-transformed alcoholic drinks per week; 1%-change in coffee consumption; 1-unit increase in napping category (never, sometimes, usually); 1-hour/day increase in sleep duration; 1-SD increase in MET-minutes/week of MVPA; 1-SD increase in BMI; 1-SD increase in waist circumference; 1-SD increase in BF%; 1-unit increase in log-transformed odds of type 2 diabetes; 1-SD increase in LDL cholesterol; 1-SD increase in HDL cholesterol; 1-SD increase in triglycerides; 1-mmHg increase in DBP; 1-SD increase in serum CRP levels.

Abbreviations: BF%=body fat percentage; BMI=body mass index; CRP=C-reactive protein; DBP=diastolic blood pressure; HDL=high-density lipoprotein; IVW=Inverse variance weighted; LDL=low-density lipoprotein; MVPA=moderate-to vigorous physical activity; MR=Mendelian randomization; MR-PRESSO=Mendelian randomization pleiotropy residual sum and outlier; No=number; SBP=systolic blood pressure; SD=standard deviation; SNP=single nucleotide polymorphism.

		Main analysis			Sensitivity analyses										
		IVW			Wei	Weighted median			MR-Egger			MR-PRESSO			
	No. of										No. of				
Modifiable factor	SNP	β (SE)	P value	q value	β (SE)	P value	q value	β (SE)	P value	q value	outlier	β (SE)	P value	q value	
Socioeconomic factor															
Educational attainment	751	-0.718 (0.151)	1.83E-06	1.65E-05	-0.633 (0.225)	0.005	0.030	-0.361 (0.565)	0.523	0.930	3	-0.689 (0.145)	2.52E-06	2.27E-05	
Lifestyle behavior															
Smoking initiation	299	0.519 (0.142)	2.43E-04	0.001	0.404 (0.205)	0.049	0.221	1.222 (0.589)	0.039	0.240	1	0.545 (0.135)	7.20E-05	3.24E-04	
Alcohol intake	80	0.669 (0.441)	0.130	0.213	1.134 (0.656)	0.084	0.252	1.693 (0.923)	0.070	0.252	0	0.737 (0.432)	0.092	0.151	
Coffee consumption	12	0.004 (0.008)	0.600	0.735	0.002 (0.007)	0.773	0.991	-0.003 (0.015)	0.861	0.930	0	0.004 (0.008)	0.610	0.705	
Sleep															
Daytime napping	115	0.115 (0.500)	0.818	0.818	0.014 (0.665)	0.984	0.991	0.152 (1.742)	0.931	0.931	0	0.083 (0.488)	0.866	0.866	
Sleep duration	77	-0.472 (0.357)	0.187	0.281	-0.526 (0.462)	0.255	0.417	2.771 (1.324)	0.040	0.240	0	-0.319 (0.357)	0.375	0.519	
MVPA	6	0.368 (1.138)	0.747	0.791	0.262 (1.347)	0.846	0.991	-2.565 (6.947)	0.731	0.930	0	0.368 (0.649)	0.596	0.705	
Cardiometabolic trait															
Adiposity															
BMI	941	0.586 (0.102)	1.08E-08	1.94E-07	0.680 (0.170)	6.33E-05	0.001	0.510 (0.298)	0.087	0.261	4	0.605 (0.099)	1.53E-09	2.75E-08	
Waist circumference	44	0.850 (0.269)	0.002	0.006	0.632 (0.339)	0.062	0.223	0.739 (0.740)	0.324	0.704	1	0.771 (0.249)	0.003	0.009	
BF%	641	0.711 (0.152)	2.88E-06	1.73E-05	0.734 (0.227)	0.001	0.009	1.666 (0.509)	0.001	0.018	4	0.664 (0.145)	5.75E-06	3.45E-05	
Childhood obesity	5	0.229 (0.095)	0.016	0.036	0.154 (0.123)	0.211	0.380	0.140 (0.724)	0.859	0.930	0	0.223 (0.071)	0.025	0.050	
Type 2 diabetes	232	0.125 (0.051)	0.014	0.036	0.102 (0.080)	0.205	0.380	0.056 (0.116)	0.628	0.930	0	0.120 (0.049)	0.016	0.036	
Lipids															
LDL cholesterol	145	0.055 (0.123)	0.653	0.735	-0.002 (0.174)	0.991	0.991	0.169 (0.181)	0.352	0.704	1	0.059 (0.123)	0.633	0.705	
HDL cholesterol	223	-0.127 (0.109)	0.243	0.336	0.068 (0.166)	0.684	0.947	0.103 (0.187)	0.583	0.930	0	-0.121 (0.109)	0.267	0.401	
Triglycerides	173	-0.058 (0.121)	0.635	0.735	0.009 (0.196)	0.964	0.991	-0.202 (0.203)	0.320	0.704	0	-0.052 (0.121)	0.666	0.705	
Blood pressure															
SBP	222	0.032 (0.016)	0.053	0.095	0.033 (0.023)	0.149	0.338	-0.013 (0.065)	0.840	0.930	1	0.038 (0.016)	0.015	0.036	
DBP	264	0.049 (0.024)	0.037	0.074	0.048 (0.034)	0.150	0.338	0.161 (0.084)	0.055	0.248	0	0.042 (0.023)	0.069	0.124	
CRP	298	0.349 (0.095)	2.49E-04	0.001	0.061 (0.150)	0.684	0.947	0.020 (0.132)	0.878	0.930	1	0.362 (0.093)	1.16E-04	4.18E-04	

Table 2. MR results of the associations between 18 modifiable factors and PhenoAgeAccel

PhenoAgeAccel represents epigenetic-age acceleration obtained using the PhenoAge clock; The q value represents the false discovery rate (FDR)-adjusted P value. β represents the associations with PhenoAgeAccel of respectively: 1-SD increase in years of educational attainment; Ever smoked regularly compared to never smoked; 1-SD increase in log-transformed alcoholic drinks per week; 1%-change in coffee consumption; 1-unit increase in napping category (never, sometimes, usually); 1-hour/day increase in sleep duration; 1-SD increase in MET-minutes/week of MVPA; 1-SD increase in BMI; 1-SD increase in waist circumference; 1-SD increase in BF%; 1-unit increase in log-transformed odds of childhood obesity; 1-unit increase in log-transformed odds of type 2 diabetes; 1-SD increase in LDL cholesterol; 1-SD increase in HDL cholesterol; 1-SD increase in triglycerides; 1-mmHg increase in SBP; 1-mmHg increase in DBP; 1-SD increase in serum CRP levels.

Abbreviations: BF%=body fat percentage; BMI=body mass index; CRP=C-reactive protein; DBP=diastolic blood pressure; HDL=high-density lipoprotein; IVW=Inverse variance weighted; LDL=low-density lipoprotein; MVPA=moderate-to vigorous physical activity; MR=Mendelian randomization; MR-PRESSO=Mendelian randomization pleiotropy residual sum and outlier; No=number; SBP=systolic blood pressure; SD=standard deviation; SNP=single nucleotide polymorphism

Figure 1. Study design and assumptions of the MR analysis

Assumption 1 indicates that the genetic variants proposed as instrumental variables should be robustly associated with the exposures; assumption 2 indicates that the used instrumental variables should not be associated with potential confounders of the relationship between exposures and outcomes; and assumption 3 indicates that the selected instrumental variables should influence the outcomes only through the exposures, not via alternative pathways.

Abbreviations: GrimAgeAccel=epigenetic-age acceleration obtained using the GrimAge clock; HDL=high-density lipoprotein; LDL=lowdensity lipoprotein; MR=mendelian randomization; PhenoAgeAccel=epigenetic-age acceleration obtained using the PhenoAge clock; PRESSO=pleiotropy residual sum and outlier.



Figure 2. Overview of the findings on associations of 18 modifiable factors with GrimAgeAccel and PhenoAgeAccel

a. Causal associations between 18 modifiable factors and GrimAgeAccel; b. Causal associations between 18 modifiable factors and PhenoAgeAccel.

GrimAgeAccel represents epigenetic-age acceleration obtained using the GrimAge clock; PhenoAgeAccel represents epigenetic-age acceleration obtained using the PhenoAge clock; β represents the effect of each modifiable factor on epigenetic-age acceleration. Red box indicates a strong association with p value <0.05 and FDR q value <0.05. Blue box indicates a suggestive association with p value <0.05 and FDR q value \geq 0.05. The IVW method was used for the main analysis. Sensitivity analyses included the MR-PRESSO, the MR-WM, and the MR-Egger methods.

Abbreviations: BF%=body fat percentage; BMI=body mass index; CRP=C-reactive protein; DBP=diastolic blood pressure; HDL=high-density lipoprotein; IVW=inverse-variance weighted; LDL=low-density lipoprotein; MR=mendelian randomization; MVPA=moderate to vigorous physical activity; PRESSO=pleiotropy residual sum and outlier; SBP=systolic blood pressure; SD=standard deviation; WM=weighted median.



Figure 3. Multivariable MR assessing the effects of the strongest protective factor and risk factor on GrimAgeAccel and PhenoAgeAccel

a. Effect of the strongest protective factor and risk factor on GrimAgeAccel; b. Effect of the strongest protective factor and risk factor on PhenoAgeAccel.

Causal estimates are β (95% CI) in years. Exposure factors with β coefficients >0.5 and P <0.05 in the main analysis as shown in Figure 2 were selected as covariates in the adjustment models.

Abbreviations: BF%=body fat percentage; BMI=body mass index; CI=confidence intervals; MR=mendelian randomization.

а	Modifiable factor	β (95% CI)	GrimAgeAccel	
	Education			
	Unadjusted	-1.143 (-1.380 to -0.906)		
	Adjusted for smoking initiation	-0.752 (-1.053 to -0.452)	-0-	
	Adjusted for alcohol intake	-1.141 (-1.411 to -0.872)	-0-	
	Adjusted for waist circumference	-1.050 (-1.365 to -0.735)	-0-	
	Adjusted for daytime napping	-1.152 (-1.422 to -0.881)	-0-	
	Adjusted for BF%	-0.947 (-1.228 to -0.666)	-0-	
	Adjusted for BMI	-1.012 (-1.302 to -0.722)	-0-	
	Smoking initiation			
	Unadjusted	1.299 (1.090 to 1.508)		-0-
	Adjusted for education	0.985 (0.688 to 1.282)		-0-
	Adjusted for alcohol intake	1.059 (0.759 to 1.358)		-0-
	Adjusted for waist circumference	1.103 (0.788 to 1.418)		-0-
	Adjusted for daytime napping	1.135 (0.851 to 1.420)		-0-
	Adjusted for BF%	1.017 (0.756 to 1.279)		-0-
	Adjusted for BMI	0.850 (0.604 to 1.096)		-0-
			-15 -10 -05 0	0 05 10 15
			← deceleration	acceleration→
h	Modifiable factor	β (95% CI)	PhenoAgeAccel	
~	Education			
	Unadjusted	-0.718 (-1.013 to -0.423)		
	Adjusted for waist circumference	-0.794 (-1.186 to -0.402)	— — —	
	Adjusted for BF%	-0.638 (-0.992 to -0.284)		
	Adjusted for BMI	-0.807 (-1.191 to -0.424)		
	Adjusted for smoking initiation	-0.691 (-1.070 to -0.313)	— — —	
	Waist circumference			
	Unadjusted	0.850 (0.323 to 1.377)		_
	Adjusted for education	0.582 (0.109 to 1.054)		
	Adjusted for BF%	0.733 (0.139 to 1.326)		
	Adjusted for BMI	0.649 (0.038 to 1.260)		_
	Adjusted for smoking initiation	0.767 (0.253 to 1.281)		_
			-15 -10 -05 0	0 05 10 15
			← deceleration	acceleration→