MERTIL FOR PARENTS: A STUDY PROTOCOL

- 1 **Full title:** My Early Relational Trust-Informed Learning (MERTIL) for Parents: A study
- protocol for a brief, universal, online, preventative parenting program to enhance relational
 health.
- 3 health.
- 4 **Short title:** MERTIL for Parents: A study protocol
- 5
- 6 Jessica Opie ^{a*#}, Leesa Hooker ^{b,c}, Tanudja Gibson ^a, Jennifer McIntosh ^a
- 7 ^a School of Psychology & Public Health, La Trobe University, Melbourne, Australia, 3056.
- ^b Judith Lumley Centre, School of Nursing and Midwifery, La Trobe University, Melbourne,
 Australia, 3056
- ^c La Trobe Rural Health School, La Trobe University, Bendigo, Australia, 3552.
- 12 * Correspondence for this article should be addressed to Jessica E. Opie:
- 13 j.opie@latrobe.edu.au
- 14 [#] Current address: The Bouverie Centre, School of Psychology & Public Health, La Trobe
- 15 University, 8 Gardiner Street, Brunswick 3056, Victoria Australia.
- 16

11

- 17 *Jessica E. Opie: j.opie@latrobe.edu.au
- 18 Leesa Hooker: <u>l.hooker@latrobe.edu.au</u>
- 19 Tanudja Gibson: <u>t.gibson@latrobe.edu.au</u>
- 20 Jennifer E McIntosh: jenn.mcintosh@latrobe.edu.au
- 21
 22 Jessica E. Opie: https://orcid.org/0000-0002-0559-0842
- 23 Leesa Hooker: https://orcid.org/0000-0002-4499-1139
- 24 Tanudja Gibson:
- 25 Jennifer E McIntosh: https://orcid.org/0000-0003-4709-5003
- 26

27 Ethics approval: This study has been approved by the Human Research Ethics Committee at

- 28 La Trobe University (HEC: 22096).
- 2930 Conflict of interest: No conflicts or competing interests to disclose.
- 31

Funding: JO, JM, and LH received funding from La Trobe University's (www.latrobe.edu.au)
 ABC Internal Investment Scheme. The funders had no role in study design, data collection

- 34 and analysis, decision to publish, or preparation of the manuscript
- 35
- 36 Author contributions:
- 37 Study conceptualisation: JM, JO, LH; Manuscript drafting: JO; Revision of manuscript and
- 38 final version approval: JO, JM, LH, TG
- 39
- 40 Note: This protocol was accepted and presented at the Family Relationship Services Australia
 41 (FRSA) conference in May 2022.
- 42
- 43
- 44

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

MERTIL FOR PARENTS: A STUDY PROTOCOL

45 **Background:** Early relational health is a key determinant of childhood development, while 46 relational trauma in the parent-infant dvad can instigate a cascading pattern of infant risk. Fortunately, early relational trauma is detectable and modifiable. In 2018, Australian 47 48 Maternal and Child Health (MCH) nurses participated in MERTIL (My Early Relational 49 Trauma-Informed Learning), a program to identify and prevent relational trauma. Program 50 evaluations revealed nurses felt competent and confident to identify and respond to relational 51 trauma; however, response capacity was inhibited by inadequate parent referral options. In 52 response, MERTIL for Parents (My Early Relational Trust-Informed Learning) was 53 developed, which is an online, evidence-based, self-paced parenting program that focuses on 54 enhancing parental knowledge of relational trust and its significance for infant development. 55 This low-cost, accessible prevention resource targets emerging relational concerns to reduce 56 later service system engagement. The potential for universal preventative online programs 57 that target parental and relational wellbeing remains under-explored. This paper reports on a 58 protocol for implementing a MERTIL for Parents pilot study describing nurses' and parents' 59 perspectives on program feasibility and efficacy. Methods: This study is a mixed-methods, 60 parallel-armed, uncontrolled, repeated measures design. We aim to recruit 48 Australian 61 MCH nurses from the states of Victoria and New South Wales. These nurses will in turn 62 recruit 480 parents with a child aged 0-5 years. All parents will receive MERTIL for Parents, 63 which entails a 40-minute video, tipsheets, worksheets, and support resources. Parent data 64 will be obtained at three periods: pre-program, program exit, and program follow-up. Nurse 65 data will be collected at two periods: parent recruitment completion and program follow-up. 66 Data collection will occur through surveys and focus groups. Primary parent outcomes will be socioemotional assessments of program efficacy. Nurses and parents will each report on 67 68 study program feasibility. Discussion: This protocol describes the feasibility and efficacy of a new online parenting program, MERTIL for Parents, with pilot field studies commencing 69

- in August 2022. We anticipate that this resource will be a valuable addition to various child
- 71 and family services, for use in individual support and group work.
- 72 *Keywords:* Relational trauma; online; parent program; feasibility; universal; maternal and
- 73 *child health; prevention; child.*

MERTIL FOR PARENTS: A STUDY PROTOCOL

74 Study Protocol Introduction

75 Early relational health is a key determinant of childhood development, while relational 76 trauma in the parent-infant dyad can instigate a cascading pattern of infant risk. Fortunately, 77 early relational trauma is detectable and modifiable. In 2018, Australian Maternal and Child Health (MCH) nurses based in the state of Victoria participated in MERTIL (My Early 78 79 Relational Trauma-Informed Learning), a program to identify and prevent relational trauma. 80 Program evaluations revealed nurses felt competent and confident to identify and respond to 81 relational trauma; however, response capacity was inhibited by inadequate parent referral 82 options for those with sub-clinical presentations. In response, MERTIL for Parents (My 83 Early Relational Trust-Informed Learning) was developed. MERTIL for Parents is an online, 84 evidence-based, self-paced parenting program that focuses on enhancing parental knowledge 85 of relational trust and its significance for infant development. This low cost, accessible 86 prevention resource targets emerging relational concerns to reduce later service system 87 engagement. The potential for universal preventative online programs that target parental and 88 relational wellbeing, as well as the development of early relational trust, remains under-89 explored.

90 This protocol describes the feasibility and efficacy of a new online parenting program, 91 MERTIL for Parents, describing MCH nurses' and parents' perspectives on program 92 feasibility and efficacy, with pilot field studies soon to commence. This study is a mixed-93 methods, parallel-armed (i.e., MCH nurses and parents), uncontrolled, repeated measures 94 design. We aim to recruit 48 MCH nurses who will in turn recruit 480 parents with a child 95 aged 0-5 years (inclusive) displaying early signs of relational trauma. All participants will 96 receive access to the MERTIL for Parents program, which entails a 40-minute video, tip 97 sheets, worksheets, and support resources. Parent and nurse data collection will occur through 98 surveys and focus groups. Primary parent outcomes will be socioemotional to examine

MERTIL FOR PARENTS: A STUDY PROTOCOL

99 program efficacy; such outcomes will include child-parent relationship quality and parental 100 reflective functioning. Nurses and parents will each report on study program feasibility. It is 101 anticipated that this resource will be a valuable addition to a wide range of child and family 102 services, for use in individual support and group work.

103 Background

104 Children develop in a relational context with everyday parent-child interactional patterns

105 contributing to their *relational health* (1). Relational health in the infant-parent dyad is

106 construed as parental security, trust, responsiveness, and sensitivity, which together form the

107 cornerstone of socioemotional development. Alternatively, relational trauma presents

108 ongoing and accruing disruptions to the infant's context of early care. Unsurprisingly,

109 relational health and relational trauma impact upon an array of child developmental domains,

110 instigating a cascading pattern of child resiliency or risk. Importantly, relational health

111 elements, including attachment organisation and parental sensitivity (2-4) are modifiable, for

112 better or worse. With this knowledge, promoting positive caregiving via psychoeducation and

113 increased parenting capacity is important, enabling parents to promote child development,

thereby affording their child the best start in life. Early parenting programs have been

115 developed as a result.

116 **Preventative Parenting Programs**

Preventative parenting programs can be universal, selective, or indicated. Universal programs are those available to all parents; selective programs target parents displaying above average risk: and indicated programs target parents showing signs or symptoms of emerging disorder

120 (5). Research has shown universal programs are well suited to mental health promotion,

121 offering significant preventative public health benefit (6).

122 Universal face-to-face parenting programs

MERTIL FOR PARENTS: A STUDY PROTOCOL

Universal parenting skills-based programs have historically been presented through varied inperson modes, including face-to-face lectures, workshops, and groups. Program examples
include CANparent (7) and *ACT Raising Safe Kids Program* (8). Such programs have been
shown to increase both child and parent mental health among an array of domains such as
parental self-efficacy, mental wellbeing, satisfaction, and child internalising and externalising
presentations (e.g., 7, 8, 9).

Despite the availability and evidence supporting these in-person parenting programs, 129 130 many families, particularly those experiencing numerous vulnerabilities, face multiple 131 barriers in accessing these programs (10, 11). For parents who live in regional centres and 132 rural or remote locations, attending in-person programs, typically city-based, is an accessibility obstacle. Transportation access is also an issue for in-person parent service 133 134 participation (12, 13). Such programs are additionally cost prohibitive to many due to their 135 high in-person contact hours with professionals (14). The inflexible structure of such 136 programs and rigid delivery approaches can be perceived as stigmatising by the individuals 137 these programs are intended to reach. Capped attendance due to physical room size, 138 occupational health and safety concerns, and the limited number of facilitators running 139 sessions provide further shortcomings. Parent logistical barriers, such as managing various 140 daily work and lifestyle demands have resulted in low in-person parent program enrolment 141 and poor program attendance for enrolees (e.g., 5, 15-17). Compounding these limitations, 142 the COVID-19 pandemic saw an abrupt stop to most in-person treatment programs due to 143 lockdowns and quarantine, severing preventative program resources at a time of heightened 144 psychological vulnerability for parents and children alike (18).

145 Universal online parenting programs

146 Online or app-based program delivery provides an avenue to address these limitations,

147 provided intended users have adequate technology access, reliable internet connection, and

MERTIL FOR PARENTS: A STUDY PROTOCOL

148 are digitally literate. Online universal parenting programs can be either entirely or partially 149 self-guided. In partially self-guided programs, clinician support is provided at varying 150 frequencies via online, phone, or in-person modalities (19). Prior meta-analytic research on 151 the efficacy of self-guided and partially self-guided online parenting programs has reported 152 benefits at the parent, child, and dyadic level. Specifically, at the parent level, programs have 153 been shown to enhance elements such as positive parenting perceptions, behaviours, 154 satisfaction, efficacy, and confidence, while lowering parental mental health problems (e.g., 155 anxiety, anger, depression, stress) and negative parenting practices (20-25). At the child level, 156 programs decrease emotional and behavioural problems (21-23, 25), while at the relational 157 level, they can reduce negative parent-child interactions (23), negative parental discipline 158 strategies, and parental conflict (24).

159 While both self-guided and partially self-guided online parenting programs are 160 efficacious, the formation and implementation of wholly self-guided, universal, digital 161 parenting programs has recently gained momentum (e.g., 15, 26). The rapid uptake of such 162 programs is particularly due to the ease of administration, scalability, sustainability, low-cost, 163 and program standardisation. Further rationale for the uptake of online program delivery is due to the ubiquitous nature of internet access, allowing widescale dissemination. For 164 165 example, 99% of Australians and 91% of Australian households have internet access (27), 166 making this a delivery method that can reach most of the population, thus making these 167 programs highly accessible. The self-directed nature of online programs may further allow 168 parents to flexibly complete these programs in the security of their homes, thereby 169 overcoming possible judgement and stigma felt from participating in in-person programs. 170 Such benefits may in turn reduce costs and barriers to program engagement and enhance 171 program reach.

MERTIL FOR PARENTS: A STUDY PROTOCOL

172 Due to their significant preventative potential, online parent resources may reduce the need for later intervention. This may occur by supporting and motivating parents with new 173 174 knowledge and behavioural skills to independently enhance their parent-child relationship, in 175 turn reducing later burden on service systems. Public health initiatives such as these may 176 empower parents to alter relational trajectories early on, within a preventative approach that 177 is parent-led and strengths-based. This may assist to disrupt emerging disadvantage before reaching pathological thresholds. Equitable access through online platforms has become more 178 179 important, particularly in the face of the COVID-19 global crisis. Further, evidence shows 180 that these programs promote help-seeking (28). Finally, from a research perspective, fully 181 online content allows for the collection and tracking of unique data on training progress. 182 completion times, and premature program cessation. Such data can be linked to outcomes and 183 inform program improvements.

184 Knowledge Gaps

While numerous clear benefits exist in relation to online, self-delivered, universal parenting 185 186 programs, gaps remain. Currently, there are few brief self-help, psychoeducational, program 187 options (in-person or online) suitable at the broad community level for parents where enhanced awareness of their role in promoting early relational trust may assist with 188 189 prevention of relational trauma, especially in the face of challenge. Second, most parenting 190 resources in this domain exist in the context of in-depth indicated parenting interventions 191 (e.g., Parent-Child Psychotherapy; 1). While these indicated programs are essential for dyads 192 in need of targeted treatment, there is a clear need for preventative education resources for 193 parents in the wider community wishing to enhance their understanding of the parent-child 194 relationship. This could act as an initial independent healthcare measure and reduce the need 195 for later targeted measures. Further, outcomes of online parenting programs tend of focus on 196 parents and, to a lesser extent, child outcomes (21, 24). Few studies have reported dyadic

197	outcomes and even fewer have reported familial outcomes associated with online parent
198	program participation (e.g., 23). Finally, despite MCH nurses completing MERTIL and
199	providing parenting support, they do not have a recommended suite of online parenting
200	programs within their toolbox to assist clients (29). MCH service practice guidelines focus on
201	health promotion, child development and to a lesser extend parent mental health; however,
202	nurses need evidence-based interventions to enhance dyadic relationships. This study aims to
203	address these resource and knowledge gaps.
204	Methods
205	Study aims
206	This paper reports on a protocol for the pilot implementation of a new self-directed, online
207	parenting program (MERTIL for Parents) within the Maternal and Child Health (MCH) and
208	allied Early Childhood sector. The aims of this study are twofold:
209	1) Determine the feasibility of MERTIL for Parents:
210	i. For MCH nurses and allied health professionals: Examine the programs
211	acceptability and utility for enhancing parent understanding of early relational
212	health; assess the practical dimension of implementing the program within a busy
213	service; and explore population reach.
214	ii. For parents: Examine program acceptability, efficacy, and utility.
215	2) Evaluate the short-term efficacy of the MERTIL <i>for Parents</i> program for parents across
216	selected parental caregiver knowledge, perceptions (e.g., parenting confidence &
217	competence), and behavioural domains (e.g., professional help-seeking amenability).
218	Intervention development
219	MERTIL for Parents is a new universally available, self-directed, online psychoeducation
220	resource for Australian parents focused on promoting parental knowledge and understanding
221	of relational trust and its significance for infant development. This program is informed by

222	evidence from attachment theory, developmental neuroscience, and psychoanalysis. As a
223	preventative early mental health resource, MERTIL for Parents aims to enhance infant-parent
224	relational trust and relational awareness, in turn modifying parents' attachment security-
225	oriented beliefs, attitudes, and behaviours. MERTIL for Parents builds upon prior research
226	by the authors. In 2018, over 1700 Australian MCH nurses from the state of Victoria
227	participated in MERTIL (My Early Relational Trauma-Informed Learning), a 14-hour self-
228	directed, interactive, professional development program for MCH nurses to identify and
229	prevent relational trauma. The MERTIL program evaluation from the 2018 Victorian cohort
230	revealed that MERTIL enhanced competence and confidence in nurses' identification of
231	relational trauma (30). However, capacity to respond to such identified early relational
232	trauma was inhibited by inadequate referral options for parents, particularly in rural and
233	remote settings. More broadly, it was identified that the potential for universal online
234	programs that target the development of early relational trust remains under-explored (30). In
235	response, MERTIL for Parents was developed, replacing the trauma in MERTIL for trust.
236	See Fig 1 for the MERTIL for Parents logo.
237	
238	Fig 1. MERTIL for Parents logo.
239	***Insert Fig 1 here***
240	The Intervention
241	Intervention Content
242	MERTIL for Parents is accessed via the internet and can be viewed on any online platform
243	(i.e., phone, computer, or tablet). The primary component of MERTIL for Parents comprises
244	a 40-minute pre-recorded video. Video content consists of four 'chapters', with each chapter
245	approximately nine-minutes in duration. See Table 1 for a description of the MERTIL for
246	Parents video content. Secondary program components include downloadable parent tip

247	sheets, parent worksheets, and professional support contacts. All secondary program elements
248	will be developed by the study authors. See Table 2 for additional downloadable program
249	resources. Participants progress through the varied program aspects at their own pace.
250	
251	Table 1. MERTIL for Parents video content.
252	***Insert Table 1***
253	
254	Table 2. Non-video MERTIL for Parents content
255	***Inser Table 2 here***
256	Program engagement
257	Numerous steps were taken to ensure MERTIL for Parents is highly engaging, in order to 1)
258	enhance program enrolment, 2) deliver memorable content, and 3) maximise successful
259	program completion. MERTIL for Parents was intentionally designed as a very-brief
260	parenting program, to counter the high dropout rates that have been observed in other online
261	parenting programs (31). The rationale for MERTIL for Parents' length was to introduce
262	parents to the pivotal elements of relational trust, in a manageable, user-friendly, and time-
263	efficient fashion that could be completed in one sitting or across several brief stints. A
264	production team and animator translated the content into an engaging, animated format. See
265	Fig 2 for screenshots of the MERTIL for Parents content.
266	
267	Fig 2. Screenshots from MERTIL for Parents video content.
268	***Insert Fig 2 here***
269	
270	Program content & delivery

MERTIL FOR PARENTS: A STUDY PROTOCOL

- 271 To ensure the evidence-based narrative content was engaging and well received, content
- included:
- 1. Occasional moments of humour to balance the program's occasional heavy content.
- Lay language to ensure the program's content was successfully communicated to a varied
 parent audience,
- 3. An emphasis on safety, relatability, and acceptance; to this end, the narrator's voice is
 warm, calm, non-blaming, and reassuring.
- 278 Co-designed nature of MERTIL for Parents

279 The MERTIL for Parents website (www.mertil.com.au) and all content within, was co-

280 designed and co-developed alongside consumers, child infant mental health clinicians, and

281 MCH nurses alike, from program commencement to completion. This collaborative process

enabled the voice of 'would be' end-user parents and professional referrers to be heard; prior

- 283 research has identified this co-design process as enhancing engagement, efficacy, and
- research applicability (32, 33). Researchers and clinicians engaged in a multiphase feedback

285 process, culminating in the resultant MERTIL for Parents program.

286 Design

287 MERTIL for Parents will be examined via a parallel armed (i.e., nurses & parents),

288 uncontrolled, mixed-methods, repeated measures study design. Nurses will complete two

- assessments (post-parent recruitment and three-month post-program) and parents will
- 290 complete three assessments (pre-program, program exit, and three-month post-program). All
- 291 nurse and parent data collection and completion will be conducted independently and in

292 parallel.

293 Participants

Eligible study participants will be twofold: 1) Australian MCH nurses, and 2) Australian

295 parents. See Fig 3 for an outline of all study participants.

MERTIL FOR PARENTS: A STUDY PROTOCOL

296 Eligibility criteria

297 Nurses

Nurses will i) have completed MERTIL, and ii) work at a participating pilot MCH site, in 298 299 metropolitan and regional/rural Victorian and New South Wales Local Government Areas 300 (LGAs) and iii) be able to make a follow-up phone call to each of the parents they refer to the 301 program, to gather feedback. Three programs sit within the MCH umbrella: 1) Universal 302 service, 2) Enhanced service and 3) 24-hour MCH Telephone Line (29). Participants will be 303 comprised of Universal and Enhanced services MCH nurses. The Universal MCH service is 304 available to all families, while the Enhanced MCH service supports families who meet strict 305 'at-risk' criteria, with limited places. Study participation invitations will be sent to MCH 306 nurses who work in both the Universal and Enhanced MCH services. For nurses who decide 307 to participate, free access to MERTIL (My Early Trauma-Informed Learning) will be 308 provided for the duration of the pilot study. This is the training program MCH nurses 309 completed in 2018 and will allow them to refresh their knowledge of and skills in identifying 310 early relational trauma, if necessary. Nurses will also have ongoing access to MERTIL for 311 Parents.

312 Parents

313 Consenting parent participants will: i) be the parent of a child 0-5 years (inclusive) or

314 currently pregnant, ii) be recommended to participate in the program by an MCH nurse from

- a participating pilot site, iii) aged 18 years and over, iv) currently reside in Australia, and v)
- 316 feel comfortable to complete questionnaires and program materials in English.
- 317 Sample size/power calculation

318 Power analysis was conducted using the G*Power 3.1 software to determine the required

- 319 sample size for this study (34). A single group within study design would require 199
- 320 participants to identify a small to moderate statistically significant effect (.20) at $p \le .05$, with

MERTIL FOR PARENTS: A STUDY PROTOCOL

- 321 80% power. To account for parent participant attrition, we aim to recruit approximately 480
- 322 parents.
- 323 Sample
- We aim to recruit 48 MCH nurses, consisting of 24 MCH nurses from the Universal MCH
- 325 service and 24 MCH nurses from the Enhanced service. These nurses will in turn recruit the
- 326 above-mentioned 480 parents.
- 327 Fig 3. MERTIL for Parents recruitment diagram.
- 328 ***Insert Fig 3***
- 329

330 Recruitment of study populations

331 Nurse

332 The authors will recruit through their existing MERTIL MCH nurse networks, ensuring a

- 333 representative sample of MCH nurses from differing Australian geographic locations.
- 334 Participating MCH nurse sites will include four Victorian and New South Wales metropolitan
- 335 LGAs and four Victorian and New South Wales regional/rural LGAs, based on pre-existing

336 nurse networks. To further ensure representativeness, recruitment will include MCH nurses

- 337 from the Universal and Enhanced MCH services. All participating MCH nurses will provide
- 338 written online consent to their study participation.
- 339 Parents

340 Parents who meet inclusion criteria will be informed about the study by their MCH nurse

341 during a visit to the Universal service (any of the scheduled visits from the initial post-birth

- home visit, consultations at 2, 4, & 8 weeks; 4, 8, 12, & 18 months; 2 & 3.5 years), or a visit
- 343 from the Enhanced service. The MCH nurse will explain MERTIL for Parents and the
- 344 associated research project, and if the parent verbally expresses interest, nurses will direct
- 345 them to the MERTIL *for Parents* website to register for the study and to access the materials

MERTIL FOR PARENTS: A STUDY PROTOCOL

346 (www.mertil.com.au). Nurses will provide parents with a printed flyer with MERTIL for 347 *Parents* details to aid this process. For parents who verbally agree to participate. MCH nurses 348 will follow-up with a phone call or email, to gauge parents' responses to the program. 349 During the study pilot phase, all individuals who arrive at the MERTIL for Parents 350 website landing page will be invited to join the research. Specifically, parents will be met 351 with a webpage popup with a brief research overview and the option to hear more about the 352 study. If they select to hear more about the research, they will be automatically prompted to 353 enter their first name and email address. This will trigger emailing of consent forms. Once the 354 completed forms are submitted, the pre-program parent survey will be automatically emailed 355 to the participant. Completing this survey will grant parents access to the MERTIL for 356 Parents program content.

357 **Procedure**

358 Nurses

Following informed consent, MCH nurses will attend a 120-minute web-based information and training session on MERTIL *for Parents*. This session will provide a study overview to promote program familiarisation, discuss parent program engagement, and explain the implementation options for parents accessing both the Universal and Enhanced MCH services. Immediately following this session, nurses will be emailed and asked to complete an online survey to report their first impressions of MERTIL *for Parents* regarding factors such as perceived program content, length, and suitability.

Nurses will be asked to make contact (phone call preferred) with parents they referred to the program, within two weeks of recommending MERTIL *for Parents*. On completion of all parent recommendations, a follow-up email survey will be conducted, assessing program use, perceived parent impacts, program engagement, ideas for program refinement, and wider implementation. Online focus group interviews will follow to elaborate on survey responses,

MERTIL FOR PARENTS: A STUDY PROTOCOL

371 assess nurses' views and experiences of the program, potential barriers to implementation,

- and to identify further uses and applications of MERTIL for Parents. These elements will
- 373 ensure MERTIL for Parents' co-designed nature endures. See Fig 4 for the MCH nurse data
- 374 collection flow diagram.

375 Fig 4. Flow diagram of MERTIL for Parents MCH nurse data collection timeline.

376

Insert Fig 4

377 Parents

378 During the study pilot phase, all individuals who arrive at the MERTIL for Parents website 379 landing page will be welcomed to the MERTIL for Parents pilot study and advised that the 380 program is open to parents who are willing to participate in the research (but will be open to 381 all following our test phase). They will be invited to read more about the study. If they 382 consent, parents will provide their contact details to be emailed the Participant Information 383 Statement and Participant Informed Consent Form. Once these forms are complete, the pre-384 program parent survey will be automatically electronically emailed to the parent participant. The brief 10-minute survey before, after, and at 3-months follow-up comprises two 385 386 sections: 1) close-ended questions, and 2) open-ended questions. Upon completion of Survey 387 1, participants will be able to access the MERTIL for Parents program. Participants have program access for 2 months from consent completion. Immediately upon MERTIL for 388 389 Parents main video completion, Survey 2 will be sent. As needed, survey reminders will be 390 sent via email at 7- and 14-days after the initial survey. If participants do not complete 391 surveys after these communication bids, no further contact will be made. Three-months after 392 completion of MERTIL for Parents, a follow-up survey will be sent to explore sustained 393 change; consenting parents will participate in a qualitative interview, to further identify lived 394 experience, program strengths and weaknesses. See Fig 5 for an outline of parent data 395 collection points.

MERTIL FOR PARENTS: A STUDY PROTOCOL

396	
397	Fig 5. Flow diagram of MERTIL for Parents parent data collection timeline.
398	***Insert Fig 5***
399	
400	During this research and piloting phase, MERTIL for Parents will be free of charge to
401	parents and caregivers who choose to participate in the research. Following this piloting
402	phase and program refinement, MERTIL for Parents will be made broadly available.
403	Measures
404	Our primary outcome will be to determine the efficacy of MERTIL for Parents in enhancing
405	socioemotional development. Secondary outcomes include evaluating the feasibility of
406	implementing MERTIL for Parents within the Victorian and New South Wales MCH
407	services.
408	Parents
409	Baseline and three-month post-program questionnaires
410	To evaluate program efficacy, participants will complete identical self-report measures pre-
411	program and three-month post-program (see Table 3); noting that demographic questions
412	would have been completed pre-program and will not be repeated. On both occasions parents
413	will receive surveys through a secure weblink sent to their email address. Program efficacy
414	questions will be both quantitative and qualitative in nature. Program efficacy will not be
415	assessed at program exit as not enough time would have elapsed for meaningful parent
416	change. All included measurement items were derived where possible from validated scales,
417	single items from validated scales, and/or items from established clinical cohort studies (i.e.,
418	Australian Temperament Project; Longitudinal Study of Australian Children). Table 3
419	displays outcomes and associated measures that will be used to evaluate MERTIL for Parents
420	pre- and post-program.

MERTIL FOR PARENTS: A STUDY PROTOCOL

421	1.	Parental distress (depression & anxiety): The Patient Health Questionnaire-4 (PHQ4;
422		35) will measure parent distress, with the two subscales measuring depression and
423		anxiety.
424	2.	Parenting practices: Parenting practices will be assessed by three parenting
425		dimensions: hostile parenting (4 items), parental warmth (4 items), and parental self-
426		efficacy (1 item; 36, 37).
427	3.	Social support: the Maternity Social Support Scale (MSSS; 38) will determine
428		parents' perceptions of support. Item wording will be modified so items are applicable
429		to all caregivers.
430	4.	Intimate partner safety: The Family Law DOORS (FL-DOORS; 39) will measure
431		partner safety via one item.
432	5.	Parenting experiences: Select items from the Perinatal Emotional Growth Index
433		(PEGI; 40) will assess parenting experiences. Seven items will assess parental
434		confidence (2 items), parenting satisfaction (1 item), parental agitation (2 items), and
435		general reflective capacity (2 items).
436	6.	Relational attunement: The Interest-Curiosity subscale from the Parental Reflecting
437		Functioning Questionnaire (PRFC; 41) will measure parents' interest and curiosity in
438		their child's mental states. Two further study-generated questions will assess a
439		parent's capacity to respond to their child's cues: 1) 'I have built trust with my child'
440		and 2) 'I know how to respond to my child's signals'. Response ranges from 1
441		(completely true) - 5 (completely untrue).
442	7.	Parent-child attachment: The Quality of Attachment subscale from the Maternal
443		Postnatal Attachment Scale (MPAS; 42) will measure the quality of parent-child
444		attachment. Item wording will be modified so items are applicable to all caregivers.

MERTIL FOR PARENTS: A STUDY PROTOCOL

445	8.	Professional help-seeking amenability: A study-specific item was developed by the
446		authors (JM & JO) to assess parent's help-seeking capacity: 'I feel able to find and
447		use support from other services in the community?' with ratings from 1 (not at all) -10
448		(definitely).
449		
450	Table	3. Pre- and post-MERTIL for Parents self-report outcomes, measures, & completion

451 time.

452

Insert Table 3

453 Parents' online program usage data will further be automatically collected via the MERTIL

454 for Parents website. This will record program engagement and completion elements such as

455 1) program completion rates, 2) number of access occasions, 3) program components

456 most/least accessed.

In addition to the above items, in the pre-program and three-months post-program questionnaires, parents will be asked open-ended free-text response questions relating to the program. Pre-program questions will cover hopes for learning, and three-month post-program feedback questions will include items on program components that parents found most useful.

Embedded within this study is an internal program evaluation consisting of multiplechoice questions and open-ended text-response questions. Specifically, upon program exit (data collection time two) parents will be asked to complete a five-minute survey relating to program usability and program satisfaction. Example questions asked include: 1) *How useful was the program for you, as a parent of a young child*? and 2) *What was most helpful about* MERTIL *for Parents*? This will be sent to parents via email with an attached secure link. *Nurses*

MERTIL FOR PARENTS: A STUDY PROTOCOL

After hearing from their parent participants, nurses will be asked to complete a brief mixedmethod online survey, allowing insight into nurse perceptions of the program. Questions will include: 1) *Would you recommend MERTIL for Parents to future clients/patients?* 2) *Would you recommend colleagues encourage suitable clients/patients to participant in MERTIL for Parents?* 3) *From your perspective, does MERTIL for Parents fill a current preventative health gap?* 4) *Are there any existing gaps in the MERTIL for Parents program?* For each question response options will be 'yes' or 'no', and then participants will be asked to expand

476 upon their answer with free text.

477 Focus groups

478 To further examine the feasibility and efficacy of MERTIL for Parents, focus groups will be

479 held upon program completion. Each group will consist of 4-6 parents or nurses. A semi-

480 structured interview schedule will guide focus group discussions. All focus groups will be

481 audio recorded to allow for later transcription and analysis.

482 *Nurse*

483 Example focus group questions will include: 1) Could you tell us about your MERTIL for

484 *Parents experience? 2) What was most relevant to you? 3) Can you see yourself in the future*

485 encouraging suitable parents to participate in MERTIL for Parents as a first intervention

486 step? (3) What are the long-term and short-term implications of programs such as MERTIL

487 *for Parents*?

488 *Parent*

- 489 Example focus group questions will include: 1) Could you tell us what it was like to
- 490 participate in MERTIL for Parents? 2) Since completing MERTIL for Parents, what, if
- 491 *anything, has changed in your thinking, or feeling, or in your behaviour as a parent?* 3) In
- 492 *MERTIL for Parents, what was most relevant to you?* 4) *Would you recommend MERTIL for*
- 493 *Parents to other parents?*

MERTIL FOR PARENTS: A STUDY PROTOCOL

494 **Program data management**

- 495 All MCH nurse and parent survey data will be disseminated through Moodle, an online
- 496 learning management system stored on a secure server, with data transferred securely for
- 497 analyses. All study data and participant information will be stored securely as per the criteria
- 498 required by the La Trobe University Human Ethics Committee.

499 Data analysis strategy

500 Quantitative

- 501 All statistical analyses will be conducted using the software IBM SSPS Statistics (43).
- 502 Descriptive statistics will be generated to identify nurse and parent population characteristics
- 503 and to evaluate parent and nurse program perceptions. To determine parenting change in dyad
- attachment, relational function, maternal mental health, from pre-program to three-months
- 505 post-program, inferential statistics will be used (i.e., univariate, bivariate, and multivariate
- analyses) such as t-tests, chi-squared analyses, ANOVA, and regression analyses.

507 Qualitative

508 Focus groups

- 509 All parent and nurse focus group interview data will be audio-recorded, transcribed verbatim,
- 510 and deidentified. Thematic and content analysis will be used to identify themes and sub-
- 511 themes from MCH nurse and parent transcript responses (44).
- 512 Free-text surveys responses
- 513 For any nurse or parent free-text survey responses, thematic analysis will also be used to
- 514 identify response themes and sub-themes.

515 **Data collection commencement**

- 516 Pilot field studies, including recruitment and data collection, will commence in August 2022
- 517 with data collection concluding in November 2022.
- 518

Discussion

MERTIL FOR PARENTS: A STUDY PROTOCOL

519 Limitations and challenges

520 While this study is innovative, we foresee several program content and delivery method 521 limitations: Firstly, it is possible that the brief self-guided content of MERTIL for Parents 522 may not provide parents with adequate content information and support (i.e., program dose 523 too low). Second, as a frontline service, MCH nurses may lack the time in session to explain 524 and recommend suitable parents to participate in MERTIL for Parents, due to competing 525 demands and scheduling pressures. Third, as the study population pertains only to nurses and 526 parents located in Victoria and New South Wales, Australia, generalisability concerns will be 527 present for inter-state Australian and international populations. Fourth, as all MERTIL for 528 Parents content is housed online, this may exclude those who lack the technology or skills to 529 participate, especially as on-demand technical support will not be available for live 530 participant trouble shooting. Further, because of this study's non-randomised sampling 531 strategy, cause-effect relationships cannot be drawn. That is, parents' socioemotional 532 outcome changes cannot be causally inferred by their MERTIL for Parents participation. 533 Finally, as with all research, there is a risk that attrition rates may be high. All parent data 534 will be self-report in nature, and interpretation of findings will therefore be subject to these 535 limitations.

536 Strengths and opportunities

There are multiple strengths to the present study: First, the mixed-methods design of this
study allows for greater understanding of the program's impact compared with either
qualitative- or quantitative-only designs. Second, diversity of parent demographics and needs
will be ensured through recruitment of participants from both the Universal and Enhanced
MCH services in both metropolitan and regional locations. Third, socioemotional-specific
content may fill an important gap in the available online parenting program market. Fourth,
the program's reach is expected to be wide due to the online delivery method. Finally, strong

MERTIL FOR PARENTS: A STUDY PROTOCOL

- 544 completion is expected due to 1) the program's brief nature, 2) the engaging, non-
- 545 judgemental, and supportive program content; and 3) the contribution of MCH nurses, infant
- 546 mental health experts, and parents in co-designing the program.
- 547 Conclusion
- 548 This study will examine implementation feasibility and efficacy of an evidence-based, online,
- 549 universal, self-directed parenting program, MERTIL for Parents. Multi-informant data from
- 550 parents and nurses will inform future iterations of MERTIL for Parents and universal online
- 551 relational programs.
- 552
- 553 Acknowledgements: We wish to thank the maternal child health nurses, early childhood

554 mental health clinicians, and parents who co-designed MERTIL for Parents.

555

556 Data availability statement: Not applicable as data collection has not commenced. Once all 557 study data is collected, the data supporting the findings of this study will be available from

- the corresponding author, JO, upon reasonable request.
- 559
- 560
- 561
- 562
- 563
- 564
- 565
- 566
- 567
- 568

569	
570	
571	References
572	1. Lieberman AF, Ippen CG, Van Horn P. Don't hit my mommy! A manual for child-
573	parent psychotherapy with young children exposed to violence and other trauma.
574	Washington, DC: Zero to Three; 2015.
575	2. Bakermans-Kranenburg MJ, van IJzendoorn MH, Juffer F. Less is more: Meta-
576	analyses of sensitivity and attachment interventions in early childhood. Psychological
577	Bulletin. 2003;129(2):195-215.
578	3. Opie JE, McIntosh JE, Esler TB, Duschinsky R, George C, Schore A, et al. Early
579	childhood attachment stability and change: A meta-analysis. Attachment & Human
580	Development. 2021;23(6):897-930.
581	4. Perrone L, Imrisek SD, Dash A, Rodriguez M, Monticciolo E, Bernard K. Changing
582	parental depression and sensitivity: Randomized clinical trial of ABC's effectiveness in the
583	community. Development and Psychopathology. 2021;33(3):1026-40.
584	5. Baker CN, Arnold DH, Meagher S. Enrollment and attendance in a parent training
585	prevention program for conduct problems. Prevention Science: The Official Journal of the
586	Society for Prevention Research. 2011;12(2):126-38.
587	6. Arango C, Díaz-Caneja CM, McGorry PD, Rapoport J, Sommer IE, Vorstman JA, et
588	al. Preventive strategies for mental health. The Lancet Psychiatry. 2018;5(7):591-604.
589	7. Lindsay G, Totsika V. The effectiveness of universal parenting programmes: The
590	CANparent trial. BMC Psychology. 2017;5(1):35.
591	8. Altafim ERP, McCoy DC, Linhares MBM. Unpacking the impacts of a universal
592	parenting program on child behavior. Child Development. 2021;92(2):626-37.

MERTIL FOR PARENTS: A STUDY PROTOCOL

- 593 9. Salari R, Enebrink P. Role of universal parenting programs in prevention. In: Sanders
- 594 MR, Morawska A, editors. Handbook of Parenting and Child Development Across the
- 595 Lifespan. Cham: Springer; 2018. p. 713-43.
- 596 10. Koerting J, Smith E, Knowles MM, Latter S, Elsey H, McCann DC, et al. Barriers to,
- 597 and facilitators of, parenting programmes for childhood behaviour problems: a qualitative
- 598 synthesis of studies of parents' and professionals' perceptions. European Child & Adolescent
- 599 Psychiatry. 2013;22(11):653-70.
- 600 11. Mytton J, Ingram J, Manns S, Thomas J. Facilitators and barriers to engagement in
- 601 parenting programs: A qualitative systematic review. Health Education & Behavior:
- 602 The official publication of the Society for Public Health
- 603 Education. 2014;41(2):127-37.
- 604 12. Berthelsen D, Williams K, Abad V, Vogel L, Nicholson J. The Parents at Playgroup
- 605 research report: Engaging families in supported playgroups 2012. Brisbane: Queensland

606 University of Technology; Playgroup Association of Queensland; 2012.

- 607 13. Oke N, Stanley J, Theobald J. The inclusive role of playgroups in Greater Dandenong.
 608 Melbourne, Australia: Brotherhood of St. Laurence; 2007.
- 609 14. Kazdin AE. Annual Research Review: Expanding mental health services through
- 610 novel models of intervention delivery. Journal of Child Psychology and Psychiatry.
- 611 2019;60(4):455-72.
- 612 15. Breitenstein SM, Gross D, Christophersen R. Digital delivery methods of parenting
- 613 training interventions: A systematic review. Worldviews on Evidence-Based Nursing.
- 614 2014;11(3):168-76.
- 615 16. Breitenstein SM, Gross D, Fogg L, Ridge A, Garvey C, Julion W, et al. The Chicago
- 616 Parent Program: Comparing 1-year outcomes for African American and Latino parents of
- 617 young children. Research in Nursing & Health. 2012;35(5):475-89.

MERTIL FOR PARENTS: A STUDY PROTOCOL

- 618 17. Thornton S, Calam R. Predicting intention to attend and actual attendance at a
- 619 universal parent-training programme: A comparison of social cognition models. Clinical
- 620 Child Psychology and Psychiatry. 2011;16(3):365-83.
- 621 18. Spinelli M, Lionetti F, Pastore M, Fasolo M. Parents' stress and children's
- 622 psychological problems in families facing the COVID-19 outbreak in Italy. Frontiers in
- 623 Psychology. 2020;11:1713.
- 19. Tully LA, Piotrowska PJ, Collins DAJ, Mairet KS, Black N, Kimonis ER, et al.

625 Optimising child outcomes from parenting interventions: fathers' experiences, preferences

and barriers to participation. BMC Public Health. 2017;17(1):550.

627 20. Baumel A, Pawar A, Kane JM, Correll CU. Digital parent training for children with

628 disruptive behaviors: Systematic review and meta-analysis of randomized trials. Journal of

629 Child and Adolescent Psychopharmacology. 2016;26(8):740-9.

- 630 21. Florean IS, Dobrean A, Păsărelu CR, Georgescu RD, Milea I. The efficacy of
- 631 internet-based parenting programs for children and adolescents with behavior problems: A
- 632 meta-analysis of randomized clinical trials. Clinical Child and Family Psychology Review.

633 2020;23(4):510-28.

634 22. Flujas-Contreras JM, García-Palacios A, Gómez I. Technology-based parenting

635 interventions for children's physical and psychological health: A systematic review and meta-

- analysis. Psychological Medicine. 2019;49(11):1787-98.
- 637 23. Opie JE, Clancy E, Painter F, Vuong A, Booth AT, Esler TB, et al. Universal digital

638 parent education programs for supporting early parent-child relational health: A systematic

review, meta-analysis, and qualitative examination. [Manuscript in preparation]. In press2022.

641 24. Spencer CM, Topham GL, King EL. Do online parenting programs create change?: A
642 meta-analysis. Journal of Family Psychology. 2020;34(3):364-74.

- 643 25. Thongseiratch T, Leijten P, Melendez-Torres GJ. Online parent programs for
- 644 children's behavioral problems: A meta-analytic review. European Child & Adolescent
- 645 Psychiatry. 2020;29(11):1555-68.
- 646 26. Nunes C, Martins C, Brás M, Carmo C, Gonçalves A, Pina A. Impact of an Online
- 647 Parenting Support Programme on Children's Quality of Life. Children (Basel). 2022;9(2):173.
- 648 27. Australian Communications and Media Authority. Communications and media in
- 649 Australia: How we use the internet. 2021.
- 650 28. Licheni SH, Devaraja L, Watson B, Simeonovic M, Cheng DR, Hoq M, et al. Impact
- of COVID-19 public health restrictions on hospital admissions for young infants in Victoria,
- Australia. Journal of Paediatrics and Child Health. 2022;58(6):1001-6.
- 65329.Department of Health and Human Services. Maternal and Child Health Service
- 654 Guidelines. 2019.
- 655 30. Clancy EM, McIntosh J, Booth AT, Sheen J, Johnson M, Gibson T, et al. Training
- 656 maternal and child health nurses in early relational trauma: An evaluation of the MERTIL
- workforce training. Nurse Education Today. 2020;89:104390.
- 658 31. Hall CM, Bierman KL. Technology-assisted interventions for parents of young
- 659 children: Emerging practices, current research, and future directions. Early Childhood
- 660 Research Quarterly. 2015;33:21-32.
- 661 32. Slattery P, Saeri AK, Bragge P. Research co-design in health: A rapid overview of
- reviews. Health Research Policy and Systems. 2020;18(1):1-13.
- 33. Thabrew H, Fleming T, Hetrick S, Merry S. Co-design of eHealth Interventions With
 Children and Young People. Frontiers in Psychiatry. 2018;9:481.
- 665 34. Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: A flexible statistical power
- analysis program for the social, behavioral, and biomedical sciences. Behavior Research
- 667 Methods. 2007;39(2):175-91.

MERTIL FOR PARENTS: A STUDY PROTOCOL

- 668 35. Kroenke KMD, Spitzer RLMD, Williams JBWDSW, Löwe BMDPD. An ultra-brief
- screening scale for anxiety and depression: The PHQ-4. Psychosomatics. 2011;50(6):613-21.
- 670 36. Paterson G, Sanson A. The association of behavioural adjustment to temperament,
- 671 parenting and family characteristics among 5-year-old children. Social Development.
- 672 1999;8(3):293-309.
- 673 37. Zubrick SR, Lucas N, Westrupp EM, Nicholson JM. Parenting measures in the
- 674 Longitudinal Study of Australian Children: Construct validity and measurement quality,
- 675 Waves 1 to 4. LSAC technical paper. 2014;12:1-110.
- 676 38. Webster J, Linnane JWJ, Dibley LM, Hinson JK, Starrenburg SE, Roberts JA.

677 Measuring social support in pregnancy: Can it be simple and meaningful? Birth (Berkeley,

678 Calif). 2000;27(2):97-101.

- McIntosh J. DOOR 1: Parent self-report form. In: McIntosh J, Ralfs C, editors. The
 family law DOORS handbook. Canbtment2011. p. 25-9.
- 681 40. McIntosh J, The Centre for Social and Early Emotional Development. The
- development of the Perinatal Emotional Growth Index (PEGI). [Manuscript in preparation.].In press 2022.
- 41. Luyten P, Mayes LC, Nijssens L, Fonagy P. The parental reflective functioning
- questionnaire: Development and preliminary validation. PLOS One. 2017;12(5):e0176218.
- 686 42. Condon JT, Corkindale CJ. The assessment of parent-to-infant attachment:
- 687 Development of a self-report questionnaire instrument. Journal of Reproductive and Infant
 688 Psychology. 1998;16(1):57-76.
- 689 43. IBM Corp. IBM SPSS Statistics for MacIntosh, Version 27.0. Armonk, NY: IBM
 690 Corp; 2020.
- 44. Braun V, Clarke V. Using thematic analysis in psychology. Qualitative Research in
 Psychology. 2006;3(2):77-101.



MY EARLY RELATIONAL TRUST INFORMED LEARNING







Note. LGAs = Local Government Areas; NSW = New south Wales. Fig 3 MERTIL for Parents

information session

Fig 4

Post-information session survey

Data collection time 1

Recommend MERTIL for Parents to parents, and follow-up phone call or email

Post-program recommendatio n survey Data collection

time 2

Qualitative interview Data collection time 3 Pre-program survey Data collection time 1 MERTIL for Parents program completion Program exit evaluation Data collection time 2

3-month postprogram survey Data collection time 3

Qualitative interview Data collection time 4

