OPERANT CONDITIONING OF RECTUS FEMORIS H-REFLEXES: A PROOF-OF-CONCEPT WITH IMPLICATIONS FOR POST-STROKE QUADRICEPS HYPERREFLEXIA

Δ
т

- 5 Kyoungsoon Kim¹, Tunc Akbas², Robert Lee³, Kathleen Manella⁴, and James Sulzer^{1,5}
- ⁶ ¹University of Texas at Austin, Austin, TX, USA, ²Intel Corporation, Hillsboro, OR, USA, ³St. David's Medical
- 7 Center, Austin, TX, USA, ⁴Nova Southeastern University, Clearwater, FL, USA, ⁵MetroHealth Hospital and Case
- 8 Western Reserve University, Cleveland, OH, USA
- 9 ⁵jss280@case.edu
- 10
- 11 Acknowledgements We would like to thank the participants for their efforts. We would also like to thank Dr.

12 Amir Eftekhar and Dr. Aiko Thompson for developing Evoked Potential Operant Conditioning System (EPOCS)

13 software and providing helpful guidance.

- 14
- 15 Author contributions KK: performed the experiment, recruited participants, and analyzed the data, TA: helped

16 with experimental setup, RL: helped with recruitment of post-stroke individuals, KM: helped with experimental

17 design, JS: designed the experiment and supervised data analysis. All authors helped edit the manuscript.

18

19 Funding This work was financially supported in part by the NIH/NICHD (P2CHD086844, Kautz), and JS is the

20 recipient. The contents are solely the responsibility of the authors and do not necessarily represent the official

- 21 views of the NIH or NICHD. The funders had no role in study design, data collection and analysis, decision to
- 22 publish, or preparation of the manuscript.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

23	Declaration
25	Deciaration

- 24 **Conflict of interest** The authors have declared that no competing interests exist.
- 25
- 26 Ethics approval This study was conducted according to the Declaration of Helsinki and had ethical approval
- 27 from University of Texas Institutional Review Board. All subjects gave written informed consent prior to data

28 collection.

29

- 30 Data availability The datasets generated during and/or analyzed during the current study are available from the
- 31 corresponding author on reasonable request.

32

Abstract

33 Hyperreflexia is common after neurological injury such as stroke, yet clinical interventions have had 34 mixed success. Our previous research has shown that hyperreflexia of the rectus femoris (RF) during pre-swing 35 is closely associated with reduced swing phase knee flexion in those with post-stroke Stiff-Knee gait (SKG). Thus, 36 reduction of RF hyperreflexia may improve walking function in those with SKG after stroke. A non-37 pharmacological and non-surgical procedure for reducing hyperreflexia has emerged based on operant 38 conditioning of spinal reflexes elicited using electrical stimulation of the peripheral nerve, known as an H-reflex. 39 Operant H-reflex conditioning training of the soleus has improved clinical gait function in those with spinal cord 40 injury. It is currently unknown whether operant conditioning can be applied to the RF or those after stroke. This 41 feasibility study trained 7 participants (5 neurologically intact, 2 post-stroke) over a period of 3 months, up to 3 42 times per week, to down-condition the RF H-reflex using visual feedback. We found an overall decrease in average 43 RF H-reflex amplitude among 7 participants (44% drop, p<0.001, paired t-test), of which the post-stroke 44 individuals contributed (49% drop). We observed a generalized training effect across quadriceps muscles. Post-45 stroke individuals exhibited improvements in peak knee-flexion velocity, reflex excitability during walking, and 46 clinical measures of spasticity. These aforementioned neuromuscular outcomes provide promising initial results 47 that operant RF H-reflex conditioning is feasible both in healthy and post-stroke individuals. This procedure could 48 provide a targeted alternative to spasticity management and avoid the drawbacks of drugs and surgery.

49

50 Keywords Stiff-knee gait · Hyperreflexia · H-reflex · Operant conditioning · Spasticity · Stroke

51

Introduction

52 Stiff-Knee gait (SKG) after stroke is a common disability often defined by decreased knee flexion angle 53 during the swing phase of walking (Perry and Burnfield 2010). Those with SKG suffer from joint pain, energy 54 inefficiency and increased risk of falls (Burpee and Lewek 2015). The causes of post-stroke SKG are unclear, 55 although it has been suggested that quadriceps spasticity contributes to SKG (Perry 1992). Our recent work 56 demonstrates this claim, showing an association between hyperreflexia, a component of spasticity, in the rectus 57 femoris (RF) and reduced knee flexion angle in those with post-stroke SKG (Akbas et al. 2020). Hyperreflexia is 58 defined as overactive or overresponsive reflex activity (Lance 1980; Young and Wiegner 1987), possibly resulting 59 from disinhibition of brainstem pathways such as the reticulospinal tract (Li and Francisco 2015; Li et al. 2018). 60 Thus, a reduction of RF hyperreflexia may help restore healthy gait through restoration of spinal inhibitory 61 pathways in those with post-stroke SKG.

62 Various pharmacological and surgical interventions have been used to treat quadriceps spasticity. 63 Botulinum neurotoxin (BoNT) injection blocks neurotransmitter release at the neuromuscular junction, resulting 64 in increased range of motion (ROM) in the knee joint angle (Robertson et al. 2009; Roche et al. 2015). BoNT 65 injections alleviate hyperreflexia (Stoquart et al. 2008; Robertson et al. 2009; Roche et al. 2015), but also weaken 66 the quadriceps throughout the gait cycle, resulting in limited clinical benefit in those with SKG (Stoquart et al. 67 2008; Wissel et al.), Surgical interventions such as tendon transfer surgery is also possible, but there is limited 68 evidence for clinical effectiveness (Platz 2021). Baclofen, a gamma-Aminobutyric acid (GABA) agonist, is the 69 most common oral treatment for spasticity, however its effectiveness is questionable and results in side effects 70 such as sedation (Dehnadi Moghadam et al. 2021; Navarrete-Opazo et al. 2016). Baclofen is more effective when 71 administered intrathecally in a targeted manner (Kofler et al. 2009; Dvorak et al. 2011), however it may negatively 72 affect compensatory motor systems and requires a surgical implant with regular maintenance. While 73 pharmacological and surgical interventions for hyperreflexia exist, they all have significant drawbacks.

A possible new non-pharmacological, non-surgical procedure could help target hyperreflexia using operant conditioning. Operant conditioning is a type of associative learning process through which the strength of a behavior is modified by reinforcement or punishment (Skinner 1951). Operant H-reflex conditioning allows the patient to self-modulate his/her own monosynaptic spinal reflex activity elicited via constant current electrical stimulation of a peripheral nerve (i.e. H-reflex) (Thompson et al. 2009). The typical procedure involves visual

79 feedback of H-reflex amplitude while the participant operantly learns how to control the feedback signal over 80 many training sessions. Initial work in animal models shows that operant H-reflex conditioning can modulate 81 descending activity of the corticospinal tract and thereby induce spinal cord plasticity that restores altered motor 82 function (Wolpaw et al. 1983; Wolpaw and Lee 1987; Chen and Wolpaw 1995). At present, operant H-reflex 83 conditioning has shown that individuals with spinal cord injury can learn to self-modulate soleus reflex 84 excitability, inducing plasticity at the spinal level (Manella et al. 2013; Thompson et al. 2013; Thompson and 85 Wolpaw 2021). Further, the ability to operantly condition soleus H-reflex correlates with improvements in 86 walking speed, locomotor symmetry, locomotor EMG activity, and H-reflex modulation during locomotion. For 87 instance, Thompson and colleagues (2013) found that some people with SCI can operantly down-condition the 88 soleus H-reflex to a degree comparable to neurologically intact individuals. Manella et al. (2013) found people 89 with motor-incomplete SCI can modulate reflex excitability through operant conditioning in both directions 90 (up/down) on agonist/antagonist pair (tibialis anterior/soleus). This training resulted in improved walking function 91 on a group level. While operant H-reflex conditioning shows promise for addressing hyperreflexia, it is unclear 92 whether quadriceps H-reflex can be successfully trained, and further, whether those post-strokes can down-93 condition reflex excitability given potential damage to the corticospinal tract (Chen et al. 2002).

94 In this study, our goal was to examine the feasibility of operant down-conditioning of the RF H-reflex in 95 a pilot cohort of both healthy and post-stroke individuals. We adapted an operant H-reflex conditioning protocol 96 from previous work which focused on regulating soleus H-reflex and targeted the femoral nerve instead of the 97 tibial nerve to innervate the quadriceps. We presented visual feedback of the electrically evoked RF H-reflex, 98 which appeared as a bar graph. Participants were asked to reduce the bar height below a given threshold and were 99 provided a running score of performance. The experiment consisted of 6 baseline sessions without feedback and 100 24 feedback training sessions, 2-3 sessions per week over a 3-month period. We expected down-regulation of RF 101 H-reflex would be feasible in both healthy and post-stroke individuals. This study represents a novel paradigm in 102 quadriceps spasticity management. This intervention could lead to a targeted, non-pharmacological and non-103 invasive treatment for hyperreflexia.

104

105

Methods

106 **SUBJECTS**

107 A total of 7 participants (5 unimpaired [H1-H5], 2 post-stroke [S1-S2]) were recruited for this study. The 108 participants were four men and three women aged 20-68, where two of the individuals had history of stroke (Table 109 1). Each participant was able to stand for 10-minute intervals unassisted, walk on a treadmill for 10-minutes, and 110 provide informed consent. Exclusion criteria included: lower limb musculoskeletal injury, functionally relevant 111 weight-bearing restrictions, cognitive impairment, vision impairment and must not have taken antispasmodic 112 medication one day prior to the session. All seven people participated in down-conditioning. The study was 113 approved by the University of Texas Institutional Review Board, and all subjects gave informed consent prior to 114 participation. To prevent diurnal variation in H-reflex magnitude (Carp et al. 2006), all training was performed at 115 the same time of the day for each individual. Both post-stroke individuals had right hemiparesis and the 116 experiment was conducted on the impaired side.

Subject		S1	S2
Age		50's	60's
Gender		М	F
Voare nost-strok	0	1	1
Type of stroke		Ischemic (Embolic)	Hemorrhagic
Location of stroke		Left Middle Cerebral Artery	Left Basal ganglia
		3 (Knee extensor)	1+ (Knee extensor)
	Pre	2 (Knee flexor)	0 (Knee flexor)
Modified		3 (Ankle plantarflexor)	1+ (Ankle plantarflexor)
Ashworth		2 (Knee extensor)	0 (Knee extensor)
Score (MAS)	Post	2 (Knee flexor)	0 (Knee flexor)
		1+ (Ankle plantarflexor)	1+ (Ankle plantarflexor)
Assistive device		- Customized AFO	- Cane, AFO, Wheelchair
		- Aphasia	
Other complicat	ions	- Mild cognitive impairment	

Table 1 Clinical information of post-stroke participants. Years post-stroke is the duration from the date of stroke to the beginning of the training. Modified Ashworth Score (Bohannon and Smith 1987) was measured by the clinician before and after training. Both participants received therapy outside their experimental participation. MAS Key: 0=no increase in tone, 1=slight increase in tone (catch/release at end ROM), 1+= slight increase in tone, (catch/release at ½ ROM with resistance to end ROM, 2=more marked increase in tone through ROM, passive movement easy, 3=considerable increase in tone, passive movement difficult, 4=affected part in rigid flexion or extension.

124 **DATA COLLECTION**

- 125 The experimental setup consisted of 8-channel surface electromyography (EMG) sensors (AMT-8,
- 126 Bortec, Calgary, AL), bipolar electrodes (Ag-AgCl, Noraxon, Scottsdale, AZ) a constant current electrical
- 127 stimulator (Digitimer DS8R, Hertfordshire, UK), a data acquisition board (NI-PCIe 6321, Austin, TX), a desktop
- 128 computer, and EPOCS software (Devetzoglou Toliou 2020) (Fig 1). Additionally, during assessment sessions,
- 129 stroke subjects walked on an instrumented split-belt force treadmill (Bertec, Columbus, OH), during which the
- 130 ground reaction forces (GRF) were recorded at 1kHz, and lower limb kinematic data were recorded via inertial
- 131 motion capture (IMU, Xsens, Enschede, Netherlands) at 60Hz.





133 Fig 1 Experimental setup and visual feedback of RF operant H-reflex conditioning.

(a) Participant with electrodes on the right leg is in the upright standing posture, facing visual feedback from a
computer monitor. (b) During the control session, only the background EMG level of the RF was provided on the
left bar graph. If the background EMG level was within the shaded region and medial hamstring (MH) activity
was within a predetermined window, the bar remained green and the electrical stimulation was delivered.
Otherwise, the bar turned red with no stimulation or trial progression. (c) During the training session, along with
control session's visual feedback, RF H-reflex size and cumulative success rate were provided. The bar turned
green if the RF H-reflex was below the threshold and turned red if larger than the threshold.

141 **EXPERIMENTAL PROTOCOL**

142 The experimental protocol consisted of the preparation stage, femoral nerve navigation, recruitment 143 stage, 6 baseline sessions, 24 training sessions, and an additional 2 assessment sessions (pre/post) only for the 144 post-stroke individuals. The experimental sessions occurred 3 times per week over a 12-week period for each 145 individual, where each session lasted 1-1.5 hrs. Each session consisted of preparation, navigation, recruitment, 146 and stimulation stages. The preparation stage started with drawing a grid on a surface of the femoral triangle on 147 the leg. The femoral nerve was identified using an anatomical landmark-based method known as the "four-three 148 finger technique" (Vloka et al. 2018). Surface EMG electrodes were applied over the muscle belly of the rectus 149 femoris (RF), medial hamstring (MH), vastus medialis (VM), and vastus lateralis (VL) based on commonly used 150 anatomical landmarks (Kasman and Wolf 2002; Bordlee and Wong 2015). RF was the primary focus for H-reflex 151 monitoring, MH was monitored for antagonist activity, and VM and VL were monitored to validate the training 152 specificity of the RF H-reflex conditioning. EMG activity was amplified, bandpass filtered (10-1000Hz), and 153 sampled at 1kHz.

154 During the femoral nerve navigation, the optimal stimulation location of the femoral nerve was 155 established via a monopolar probe electrode to each point on the grid drawn on the femoral triangle (Knikou 156 2006). Electrical stimulation was delivered as a square stimulus pulse, 1 ms in duration, intensity ranging between 157 10-20 mA or above depending on the individual, to elicit a reflex response with the subject in quiet standing. We 158 monitored the time-course of the reflex response. The H-reflex was determined as the peak response following 159 the initial stimulus artifact (<3 ms), and M-wave (~10-25 ms), typically within 30-40 ms post-stimulus. We used 160 the following criteria to identify the optimal H-reflex response: 1) a clear sinusoidal shape, 2) a clear distinction 161 from the M-wave or sinusoidal signal lies on the downward slope of the M-wave in case of contamination, and 3) 162 largest signal magnitude of all the candidate spots. After the optimal stimulation spot was located, the monopolar 163 probe was replaced with a disposable Ag/AgCl snap electrode (circular, 1cm diameter) and the anode electrode 164 (square 5x5cm) was placed over the surface of gluteus maximus (GMax) on the ipsilateral side. To avoid session-165 to-session variability in the electrodes' locations, the positions of all electrodes were marked using semi-166 permanent surgical skin marker. After electrodes were placed, RF maximum voluntary contraction (MVC) was 167 measured. Participants were asked to sit on a chair with their ankle fastened with a belt to the leg of the chair and 168 knee flexed to 90°. The investigator asked the participants to extend their knee with maximum effort for 5-7

seconds. The participant repeated this three times with 15 seconds intervals in between. The representative MVC
was determined as mean of three MVC trials and was used to determine the inactive window (15% MVC) for the
RF background EMG.

172 During the recruitment stage, the RF H-reflex recruitment curve was achieved while the subject was in 173 a quiet standing posture. Electrical stimulation with a fixed interval (7 sec) was delivered, only if the background EMG level of these two muscles were below a predetermined threshold of 15% MVC (Pierrot-Deseilligny and 174 175 Burke 2005). Stimulation intensity was gradually increased by 2mA increments from 10mA to the intensity when 176 the RF M-wave amplitude was the same for three successive trials (Palmieri et al. 2004). For each stimulus 177 intensity, two stimulations were recorded and the mean of two H-reflexes was taken as representative reflex size. 178 The corresponding intensity (I_{max}) that elicited maximal representative H-reflex (H_{max}) was used in the following 179 trials. Compared to a distal muscle, such as the soleus muscle, the RF H-reflex is often affected by overlap of the 180 M-wave. This overlap makes it difficult to isolate the H-reflex. To address this issue, we used the best decaying 181 exponential fit beginning at the peak of the concurrent M-wave and ending at the zero crossing, then subtracted 182 from the measured signal to obtain the H-reflex estimate (Akbas et al. 2020).

183 During the baseline/training sessions, surface electromyographic (EMG) activity was amplified, 184 bandpass filtered (3-3000Hz), and recorded from four muscles (RF, MH, VM, and VL), while the femoral nerve 185 was stimulated with the pre-determined intensity (I_{max}) at a frequency of 0.14Hz. Every 100ms post-stimulation, 186 peak-to-peak magnitude EMG activity was measured, and the result was provided as visual feedback to the 187 participant. As shown in Fig 1c, visual feedback consisted of 1) RF and MH background EMG (left bar), 2) RF 188 H-reflex (right bar), 3) trial number, and 4) cumulative success rate. The left bar was green if the background 189 EMG was below activation threshold but was red otherwise. The goal for the participant was to reduce the height 190 of the right bar, representing H-reflex amplitude, below a given threshold (shaded area in Fig 1c). If below the 191 performance threshold, the right bar turned green and cumulative success rate was adjusted accordingly. If above 192 the threshold, the bar turned red and success rate fell. Each trial was 7 s duration, with stimulation of 1ms pulse 193 width followed by feedback for the remainder of the trial. The 6 baseline sessions each consisted of 3 runs of 75 194 control trials (225 total), while each of 24 training sessions consisted of 20 control trials and 3 runs of 75 training 195 trials (245 total) in standing posture. Subjects were exposed to H-reflex feedback only during the training trials, 196 and no H-reflex feedback was given during the control trials. The participant's score for each run was cumulative

197 success rate ($Scr = \frac{N_{success}}{N_{trial}}$ %) and participants were rewarded in proportion to one's score (i.e., Reward(\$) =198 1.0 + 0.05 * (Scr - 50)). If the participant was successful in achieving three consecutive success rates that were 199 equal to or higher than 90% for all three runs, they were provided with an extra dollar as bonus, which made the 200 maximum additional compensation amount \$10.00. A rest period was provided between each run to minimize 201 fatigue, which could affect muscular responses to stimuli (Crone et al. 1999; Stutzig and Siebert 2017).

202 The two individuals with stroke participated in pre/post assessment sessions. Participants performed a 203 10 m walk test (10MWT) with instructions to walk with maximum effort for 10 m. This process was repeated for 204 4 times and the participant's gait kinematics were recorded using inertial motion capture. Next, participants 205 completed the quadriceps pendulum test, which is a non-invasive biomechanical method of evaluating spasticity 206 using gravity to provoke muscle stretch reflexes during passive swinging of the lower limb (Fowler et al. 2007). 207 Inertial measurement units were attached on top of thigh, shank, and foot to calculate the knee joint angle during 208 the process. This test was repeated 3 times with 30 second rest periods in between. The main outcome of this test 209 was quadriceps reflex threshold angle (QRTA), which was the angle difference from the maximum knee extension 210 position to the first swing excursion, the first transition period of knee flexion to knee extension. Third, the 211 participant was asked to walk on an instrumented treadmill while their gait kinematics, ground reaction forces 212 (GRF), and H-reflexes were obtained. During treadmill walking, four gait phases (heel-strike, toe-off, mid-stance, 213 and mid-swing) were detected in real-time using online GRF data. The H-reflex was elicited and measured only 214 when a desired gait phase was detected and occurred at least 7 seconds after the last H-reflex trial. Twenty H-215 reflexes were recorded for each gait phase. A 5-minute break after each of four runs was provided to minimize 216 the effect of fatigue on H-reflex. The main outcomes consisted of 1) knee flexion range of motion (RoM), 2) peak 217 knee flexion velocity, and 3) H-reflex amplitude during the four different gait phases.

218 **OUTCOME MEASURES**

219 We normalized the H-reflex measurements. For each session, the H-reflex was normalized with the 220 session's maximum motor response (M-wave, M_{max}), which is the most commonly used method of normalization 221 (Hopkins and Ingersoll 2000). To directly compare between session-to-session performances, each session's H-222 reflex trial was averaged and represented as the percentage of the mean of six baseline session's H-reflex. Main 223 outcome measures of each session were the average of 225 peak-to-peak H-reflex trials normalized to M_{max} and 224 baseline mean. Control trials, which were conducted without feedback in the beginning of each training session, 225 provide a window into long-term plasticity of H-reflex excitability. For each session, we compared the mean 226 amplitude of 20 control H-reflex trials, which were normalized to the M_{max} and baseline mean.

227 TRAINING SPECIFICITY OF OPERANT H-REFLEX CONDITIONING

228	Exploring training specificity of operant H-reflex conditioning can provide evidence that the training can
229	be targeted for a specific muscle, which in this study was the RF. During the experiment, only RF H-reflex visual
230	feedback was provided to the subject while two adjacent quadriceps muscles (VM and VL) were recorded without
231	being visualized by the participant. Since VM and VL share motor innervation with RF through a branch of
232	femoral nerve, we investigated whether the shared nerve root of the quadriceps would project a broader training
233	effect on the quadriceps. H-reflexes of VM and VL were processed in a similar way as RF; each sessions' H-
234	reflex trials were averaged, normalized by M_{max} , and represented as percentage (%) of 6 baseline sessions' mean.

235 STATISTICAL ANALYSIS

236	We investigated the hypothesis that RF H-reflex can be down-conditioned through operant H-reflex
237	conditioning both in healthy and post-stroke individuals. To validate the effect of training, a paired t-test ($a < 0.05$)
238	was used to compare the mean of the 225 normalized H-reflex trials during six baseline sessions to that of the last
239	six training sessions (Fig 2). We used a linear mixed model to evaluate any spontaneous change in H-reflex during
240	baseline, where session number was set as a fixed effect and subject number as random effect. Further, we
241	hypothesized RF operant H-reflex conditioning may be targeted to the RF muscle and would not affect other
242	adjacent muscles (VM and VL) using paired t-tests. We compared control H-reflex trials of conditioning sessions
243	to that of baseline sessions to examine the long-term effect of training (Thompson et al. 2009) using a paired t-
244	test.
245	For post-stroke individuals, we compared quadriceps reflex threshold angle (QRTA) as a quantitative
246	measure of spasticity, knee joint kinematics, and the H-reflex amplitude during four gait phases (heel-strike, toe-

247 off, mid-stance, and swing) before and after the training using a paired t-test.

248

Results

249 **RF OPERANT H-REFLEX CONDITIONING**

All participants exhibited the ability to down-regulate RF H-reflex through training. **Table 2** presents the mean response of the healthy participants and individual responses of the two post-stroke participants. Because performance of post-stroke individuals was not significantly different than healthy individuals, the data was pooled for subsequent analyses. There was no spontaneous decrease in H-reflex amplitude during baseline sessions (F = 0.53, p = 0.47), compared to the training sessions (F = 27.18, p < 0.0001). We observed a decrease in H-reflex amplitude when comparing the last 6 training sessions with the 6 baseline sessions ($43.82 \pm 4.81\%$ mean±SE, p < 0.0001). **Fig 2** illustrates the summary statistics of H-reflex amplitude of 7 participants over each

257 session.

Subject	Muscle	Conditioned H-reflex (% baseline)
H1 – H5	RF	58.37±2.91 (***)
	VM	52.46±3.44 (***)
	VL	71.74±6.87 (*)
S1	RF	46.87±5.59 (*)
	VM	74.62±10.09 (ns)
	VL	70.89±10.55 (*)
82	RF	54.56±3.88 (*)
	VM	17.34±9.27 (**)
	VL	14.58±0.92 (**)

258 Table 2 Summary of results based on neurological condition. Values represent mean±SE and are expressed as

259 the percentage of baseline H-reflex mean. The table shows the significant differences from the six baseline

260 sessions' H-reflex (* p<0.05, **p<0.01, ***p<0.001)



261

262 Fig 2 H-reflex magnitude change in RF over training.

263 Shaded area indicates the baseline sessions and blank area indicates training sessions. Each marker represents the

individuals' (H1-H5, S1-S2) RF H-reflex means for each session. (* p<0.05, **p<0.01, ***p<0.001)

265 LONG-TERM EFFECT OF CONDITIONING ON THE CONTROL H-REFLEX SIZE

Consolidation of H-reflex down-conditioning following training can be observed in the first 20 control (no feedback) trials of each session. **Fig 3** shows the time-course change of the control RF H-reflex for all participants. For the control trials (i.e., no feedback) of all 7 individuals, the change between baseline and the last 6 training sessions was (20.86 \pm 5.77%, mean \pm SE, *p*<0.01).

270







TRAINING SPECIFICITY OF RF OPERANT H-REFLEX CONDITIONING: COMPARISON WITH VM AND VL

- 278 The femoral nerve innervates all four quadriceps muscles, but feedback is only provided from one muscle, raising
- a question of muscle training specificity. Performance of healthy individuals separate from post-stroke individuals
- 280 can be seen in Table 2. For healthy individuals, H-reflex decreased significantly for both VM and VL, whereas
- 281 mixed result was monitored among post-stroke individuals. We pooled these results in Fig 4 for VM (49.16±
- 282 5.42% (SE) drop, p < 0.0001) and VL (37.24 ± 7.37% (SE) drop, p < 0.001), respectively.



VM H-reflex magnitude over training



284 Fig 4 Muscle training specificity of RF Operant H-reflex Conditioning in VM (a) and VL (b) in all

285 participants. Shaded area indicates the baseline sessions and blank area indicates training sessions. Each 286 marker represents the individuals' (H1-H5, S1-S2) VM and VL H-reflex means for each session. (* p<0.05, 287 **p<0.01, ***p<0.001, ****p<0.0001)

BEHAVIORAL EFFECT OF THE RF OPERANT H-REFLEX CONDITIONING IN POST STROKE INDIVIDUALS

We conducted a pre/post behavioral assessment on the two post-stroke individuals. Knee flexion RoM and peak knee flexion velocity significantly improved both during overground and treadmill walking for S1. However, for S2, the improvement in knee flexion kinematics was limited to RoM during overground walking. **Table 3** illustrates the results. Quadriceps spasticity, quantified as QRTA, improved in S1 by 10.55° (p=0.016), while S2 had a statistically insignificant change of 3.50° (p = 0.536). Reflex modulation (**Table 3**) for S1 improved significantly for all four gait phases, where the mean reduction was 19.21%. In S2, H-reflex excitability decreased significantly in three gait phases (heel-strike, toe-off, and mid-stance), with a mean decrease of 23.4%.

Subject	Measurement			
	QRTA (deg)		Pre-assess	Post-assess
			32.47±3.47	42.92±2.96 (**)
	Knee flexion			
		(Overground)	12.27±0.36	14.40±0.26 (****)
	RoM (deg)	(Treadmill)	6.57±0.40	10.10±0.39 (****)
		(Overground)	226.83±4.54	264.64±5.81 (****)
S 1	Pk vel. (deg/s)	(Treadmill)	138.01±6.73	190.59±5.95 (****)
	Gait pha	se H-reflex		
	Heel-strike		33.63±0.92	19.68±0.52 (****)
	Toe-off		47.44±2.94	17.41±0.76 (****)
	Mid-stance		31.69±2.38	18.97±0.42 (****)
	Mid-swing		37.05±2.87	16.89±0.60 (****)
	QRTA (deg)		Pre-assess	Post-assess
			41.81±0.7	45.31±5.18 (ns)
	Knee flexion			
	RoM (deg)	(Overground)	44.59±0.77	51.05±0.54 (****)
		(Treadmill)	46.15±1.13	46.53±1.20 (ns)
	Pk vel. (deg/s)	(Overground)	154.46±6.69	179.38±18.78 (ns)
S2		(Treadmill)	146.93±9.36	140.22±6.28 (ns)
	Gait phase H-reflex			
	Heel-strike		96.27±6.56	60.28±4.42 (****)
	Toe-off		102.65±5.58	68.63±1.11 (****)
	Mid-stance		80.43±2.45	54.71±2.76 (****)
	Mid-swing		78.95±11.09	81.66±3.95 (ns)

297 Table 3 Behavioral effect on gait kinematics

298 Values represent mean±SE of Quadriceps Reflex Threshold Angle (deg), knee flexion RoM (deg), and peak-

- velocity (deg/s) during overground (10MWT) walking and treadmill walking. The table shows the significant
- 300 differences between the pre-assessment and post-assessment session (* p<0.05, **p<0.01, ***p<0.001,
- 301 ****p<0.0001)

302

Discussion

The goal of this study was to investigate the feasibility of self-modulation of RF H-reflex excitability. We trained 7 individuals (5 healthy and 2 post-stroke), over 24 sessions. Our results provided the first evidence that RF H-reflex can be down-regulated, and we did not observe any decrement in ability to self-regulate in 2 post-stroke individuals. The down-regulation of RF generalized to other quadriceps muscles (VM and VL). The post-stroke individuals exhibited improvements in reflex modulation generalized over the gait cycle and gait performance. We conclude that RF H-reflex down-conditioning is feasible and its effects should be evaluated in a controlled clinical trial.

310 Our findings suggest that individuals were successful in down-regulating their RF H-reflex via operant 311 H-reflex conditioning. Comparing the H-reflex amplitude in the six baseline sessions to the last six training 312 sessions resulted in a 42% drop in healthy individuals, 49% drop for the two post-stroke individuals, and 44% 313 drop across all 7 individuals. Previous animal model research indicated that CST transmission was critical for 314 operant H-reflex learning (Chen et al. 2002) suggesting that those with reduced CST integrity resulting from 315 stroke may not be able to learn. Our findings indicate that damaged corticospinal pathways due to stroke do not 316 prevent successful learning. A controlled trial remains to be conducted that examines the effect of damaged 317 pathways on operant learning and performance.

318 We expected to observe successful down-conditioning of RF since previous work has shown successful 319 soleus H-reflex down-conditioning (Thompson et al. 2009; Manella et al. 2013). However, in neural operant 320 conditioning, even across modalities, approximately 30% of individuals are non-learners (Thompson et al. 2009; 321 Manella et al. 2013; Sitaram et al. 2017). Although only 7 individuals were included in our cohort, there were no 322 non-responders, suggesting the potential for a robust operant conditioning paradigm. Our analysis of 6 baseline 323 sessions showed no evidence of spontaneous decrease in H-reflex amplitude that could be mistaken as down-324 conditioning. Further, there is no evidence of spontaneous decrease in H-reflex amplitude over sessions in 325 previous work (Thompson et al. 2009). Thus, it is highly unlikely that RF H-reflex spontaneously decreased over 326 time.

Trials conducted without feedback, control trials, provide a window into long-term plasticity. Overall, we observed a decrease in all individuals $(21 \pm 6(SE) \%)$, where the drop in post-stroke individuals (S1 = 25%), S2 = 40%) was more evident than that of the healthy group (16 %). These results align with earlier findings, where

the drop of control H-reflex of the soleus was larger in participants with spinal cord injury (24%) than healthy
individuals (16%) (Thompson et al. 2013). This difference is attributed to the hypothesis of negotiated equilibrium
(Wolpaw 2018), the concept that neural circuitry already behaving well will not be affected by neuromodulation,
but neurologically affected circuitry will.

334 We found that operant conditioning based on femoral nerve stimulation generalized to other innervated 335 quadriceps muscles. Additional monitoring of adjacent quadriceps muscles (VM and VL) was performed to 336 validate the specificity of RF operant H-reflex conditioning. We found a significant drop in VM and VL in healthy 337 individuals between the first and last 6 sessions. Although not conclusive, these results suggest that operant 338 conditioning based on stimulation of a peripheral nerve that innervates multiple muscles may affect the entire 339 muscle group. Animal studies by Chen et al. (Chen et al. 2011) found that operant conditioning of the soleus 340 affected quadriceps H-reflex excitability, and as such the generalization of operant conditioning to multiple 341 muscles, especially innervated by the same nerve, was expected. The degree of generalization will indicate the 342 specificity of the effect of training and needs to be characterized further.

343 We found functional changes in gait in the post-stroke individuals following training. These results align 344 with earlier studies that have shown functional improvement following the operant H-reflex conditioning. Chen 345 et al. (2006) have shown rats with incomplete spinal cord injury (SCI) improved gait asymmetry following the 346 training. The studies by Thompson et al. and Manella et al. (Manella et al. 2013; Thompson et al. 2013) have 347 shown gait improvements (i.e., speed, endurance, less clonus, easier stepping etc.) in people with SCI. Most 348 recently, Tahayori and Koceja (2019) have shown 3 post-stroke individuals were able to down-regulate their 349 soleus H-reflex and the changes were correlated with movement improvements. In our study, S1 exhibited 350 improvements in knee ROM and knee flexion velocity in both overground and treadmill walking conditions, but 351 S2 only showed improvements in knee kinematics only in overground walking. Both participants exhibited better 352 improvements during overground walking than treadmill walking. Overground walking was measured during 10m 353 walk test, where the participants were asked to set their speed with the maximum effort. No wiring from EMG 354 sensors or the stimulator was attached, and a belt type harness was worn. In contrast, during treadmill walking, 355 participants had to walk in a fixed and self-selected speed that was predetermined, the wiring from EMG sensors 356 and stimulator was present, a full-body harness had to be worn, and no gait support devices (i.e., cane, rolling 357 walker) were allowed. Thus, it is possible that the differences between overground and treadmill walking could

be due to the experimental setup. Clinical and laboratory measurements of hyperreflexia showed improvements in both post-stroke individuals. S1 improved in QRTA, while we did not observe a change in QRTA of S2. Further, H-reflex amplitude during different gait cycle phases also decreased, suggesting generalization of training during standing to walking (Thompson and Wolpaw 2021). While no conclusions can be drawn across post-stroke individuals, these results suggest the efficacy of conducting a RF H-reflex down-conditioning clinical trial for individuals post-stroke with SKG and quadriceps hyperreflexia.

364 This study investigates the feasibility of RF H-reflex down-conditioning based on 7 individuals, 2 of 365 them post-stroke. One of the main limitations of this study is the small sample size (7). Despite the small sample, 366 the effect size (3.85) and statistical power (98.7%) were substantial and suggest that running more individuals 367 would not change our conclusions. Due to training only 2 post-stroke individuals, we cannot generalize the 368 feasibility in operant conditioning of RF H-reflex conditioning to a larger population. It is interesting that S2 369 demonstrated improvement in H-reflex during gait phases, except for mid-swing, yet did not show as great of a 370 functional improvement as S1. While the S2 mechanism (cortico-basal ganglia-thalamo-cortical loop pathway) 371 underlying this change was different than, S1 (corticospinal pathway), it is notable that both participants 372 successfully down-regulated RF H-reflex. Further investigations should be dedicated to understanding the role 373 between H-reflex excitability and clinical outcomes.

3	7	4
~		

Conclusions

375	Our observations suggest that RF H-reflex down-conditioning is feasible and generalizes to other
376	quadriceps muscles. The successful training of 2 post-stroke individuals provides promise that one's ability to
377	regulate H-reflex may extend to this neurologically impaired population, but a larger sample size is needed to
378	confirm feasibility, long-term plasticity and clinical effects. These initial results open the door for alternative
379	treatments of spasticity and may lead towards an effective intervention in post-stroke Stiff-Knee gait and other
380	spasticity-related disorders.

381 **REFERENCES**

- 382 Akbas T, Kim K, Doyle K, Manella K, Lee R, Spicer P, Knikou M, Sulzer J. 2020. Rectus
- femoris hyperreflexia contributes to Stiff-Knee gait after stroke. J NeuroEngineering Rehabil.
 17(1):117. doi:10.1186/s12984-020-00724-z.
- Bohannon RW, Smith MB. 1987. Interrater Reliability of a Modified Ashworth Scale of
 Muscle Spasticity. Physical Therapy. 67(2):206–207. doi:10.1093/ptj/67.2.206.
- 387 Bordlee R, Wong CK. 2015. Surface EMG: A how-to guide for practitioners | Lower
- Bordice R, Wong CR. 2015. Surface Live. A now-to guide for practitioners | Lower
 Extremity Review Magazine. [accessed 2021 Nov 2]. https://lermagazine.com/article/surface emg-a-how-to-guide-for-practitioners.
- Burpee J, Lewek M. 2015. Biomechanical gait characteristics of naturally occurring
- 391 unsuccessful foot clearance during swing in individuals with chronic stroke. Clinical
- 392 biomechanics (Bristol, Avon). 30(10). doi:10.1016/j.clinbiomech.2015.08.018. [accessed
- 393 2021 Nov 2]. https://pubmed.ncbi.nlm.nih.gov/26371855/.
- Carp JS, Tennissen AM, Chen XY, Wolpaw JR. 2006. Diurnal H-reflex variation in mice.
 Exp Brain Res. 168(4):517–528. doi:10.1007/s00221-005-0106-y.
- 396 Chen XY, Carp JS, Chen L, Wolpaw JR. 2002. Corticospinal tract transection prevents
- 397 operantly conditioned H-reflex increase in rats. Exp Brain Res. 144(1):88–94.
- 398 doi:10.1007/s00221-002-1026-8.
- Chen XY, Wolpaw JR. 1995. Operant conditioning of H-reflex in freely moving rats. Journal
 of Neurophysiology. 73(1):411–415. doi:10.1152/jn.1995.73.1.411.
- Chen Y, Chen L, Wang Y, Wolpaw JR, Chen XY. 2011. Operant Conditioning of Rat Soleus
 H-Reflex Oppositely Affects Another H-Reflex and Changes Locomotor Kinematics. Journal
- 403 of Neuroscience. 31(31):11370–11375. doi:10.1523/JNEUROSCI.1526-11.2011.
- 404 Chen Y, Chen XY, Jakeman LB, Chen L, Stokes BT, Wolpaw JR. 2006. Operant
- 405 Conditioning of H-Reflex Can Correct a Locomotor Abnormality after Spinal Cord Injury in
- 406 Rats. J Neurosci. 26(48):12537–12543. doi:10.1523/JNEUROSCI.2198-06.2006.
- 407 Crone C, Johnsen LL, Hultborn H, Ørsnes GB. 1999. Amplitude of the maximum motor
- response (M max) in human muscles typically decreases during the course of an experiment.
 Experimental Brain Research. 124(2):265–270. doi:10.1007/s002210050621.
- 410 Dehnadi Moghadam A, Hasanzadeh H, Dehnadi Moghadam F. 2021. Evaluation of the Effect
- 411 of Intranasal Lidocaine in the Treatment of Spasticity in Patients with Traumatic Brain
- 412 Injury. Anesth Pain Med. 11(4):e115849. doi:10.5812/aapm.115849.
- 413 Devetzoglou Toliou S. 2020. Brain-computer interface-based feedback to enhance motor
- 414 rehabilitation. [accessed 2022 Jul 14].
- 415 https://repository.tudelft.nl/islandora/object/uuid%3A58e681ae-9a2d-4046-b13f-
- 416 6d33e991af6e.

- 417 Dvorak EM, Ketchum NC, McGuire JR. 2011. The Underutilization of Intrathecal Baclofen
- 418 in Poststroke Spasticity. Topics in Stroke Rehabilitation. 18(3):195–202.
- 419 doi:10.1310/tsr1803-195.
- 420
- 421 Fowler EG, Nwigwe AI, Ho TW. 2007. Sensitivity of the pendulum test for assessing
- 422 spasticity in persons with cerebral palsy. Developmental Medicine & Child Neurology.
 423 42(3):182–189. doi:10.1111/j.1469-8749.2000.tb00067.x.
- Hopkins JT, Ingersoll CD. 2000. Arthrogenic Muscle inhibition: A Limiting Factor in Joint
 Rehabilitation. Journal of Sport Rehabilitation. 9(2):135–159. doi:10.1123/jsr.9.2.135.
- 426 Kasman G, Wolf S. 2002. SURFACE EMG MADE EASY: A Beginner's Guide for
- 427 Rehabilitation Clinicians. http://www.noraxon.com/wp-content/uploads/2015/01/nor-
- 428 19surface-emg-manual.pdf.
- 429 Knikou M. 2006. EFFECTS OF CHANGES IN HIP POSITION ON ACTIONS OF SPINAL
- 430 INHIBITORY INTERNEURONS IN HUMANS. Int J Neurosci. 116(8):945–961.
 431 doi:10.1080/00207450600675167.
- Kofler M, Quirbach E, Schauer R, Singer M, Saltuari L. 2009. Limitations of Intrathecal
 Baclofen for Spastic Hemiparesis Following Stroke. Neurorehabil Neural Repair. 23(1):26–
- 434 31. doi:10.1177/1545968308317700.
- 435 Lance J. 1980. The control of muscle tone, reflexes, and movement: Robert Wartenberg
- 436 Lecture. Neurology. 30(12). doi:10.1212/wnl.30.12.1303. [accessed 2021 Nov 1].
 437 https://pubmed.ncbi.nlm.nih.gov/7192811/.
- Li S, Francisco GE. 2015. New insights into the pathophysiology of post-stroke spasticity.
 Front Hum Neurosci. 9:192. doi:10.3389/fnhum.2015.00192.
- Li S, Francisco GE, Zhou P. 2018. Post-stroke Hemiplegic Gait: New Perspective and
 Insights. Front Physiol. 9:1021. doi:10.3389/fphys.2018.01021.
- 442 Manella KJ, Roach KE, Field-Fote EC. 2013. Operant conditioning to increase ankle control
- 443 or decrease reflex excitability improves reflex modulation and walking function in chronic
- 444 spinal cord injury. Journal of Neurophysiology. 109(11):2666–2679.
- 445 doi:10.1152/jn.01039.2011.
- 446 Navarrete-Opazo AA, Gonzalez W, Nahuelhual P. 2016. Effectiveness of Oral Baclofen in
 447 the Treatment of Spasticity in Children and Adolescents With Cerebral Palsy. Archives of
- 448 Physical Medicine and Rehabilitation. 97(4):604–618. doi:10.1016/j.apmr.2015.08.417.
- 449 Palmieri RM, Ingersoll CD, Hoffman MA. 2004. The Hoffmann Reflex: Methodologic
- 450 Considerations and Applications for Use in Sports Medicine and Athletic Training Research.
- 451 Journal of Athletic Training. 39(3):268.
- 452 Perry J. 1992. Gait analysis : normal and pathological function. Thorofare, N.J. : SLACK inc.
 453 [accessed 2022 Apr 12]. http://archive.org/details/gaitanalysisnorm0000perr.

- 454 Perry J, Burnfield J. 2010. Gait Analysis: Normal and Pathological Function. J Sports Sci
 455 Med. 9(2):353.
- 456 Pierrot-Deseilligny E, Burke D. 2005. The Circuitry of the Human Spinal Cord: Its Role in
- 457 Motor Control and Movement Disorders. Cambridge: Cambridge University Press.
- 458 https://www.cambridge.org/core/books/circuitry-of-the-human-spinal-
- 459 cord/BBDFC31B44453B61DD5FD801B34CFFFF.
- 460 Platz T, editor. 2021. Clinical Pathways in Stroke Rehabilitation: Evidence-based Clinical
- 461 Practice Recommendations. Cham: Springer International Publishing. [accessed 2021 Oct
- 462 29]. http://link.springer.com/10.1007/978-3-030-58505-1.
- 463 Robertson JVG, Pradon D, Bensmail D, Fermanian C, Bussel B, Roche N. 2009. Relevance
- 464 of botulinum toxin injection and nerve block of rectus femoris to kinematic and functional
- 465 parameters of stiff knee gait in hemiplegic adults. Gait & Posture. 29(1):108–112.
- 466 doi:10.1016/j.gaitpost.2008.07.005.
- 467 Roche N, Zory R, Sauthier A, Bonnyaud C, Pradon D, Bensmail D. 2015. Effect of
- 468 rehabilitation and botulinum toxin injection on gait in chronic stroke patients: A randomized
- 469 controlled study. J Rehabil Med. 47(1):31–37. doi:10.2340/16501977-1887.
- 470 Sitaram R, Ros T, Stoeckel L, Haller S, Scharnowski F, Lewis-Peacock J, Weiskopf N,
- 471 Blefari ML, Rana M, Oblak E, et al. 2017. Closed-loop brain training: the science of
- 472 neurofeedback. Nat Rev Neurosci. 18(2):86–100. doi:10.1038/nrn.2016.164.
- 473 Skinner BF. 1951. How to Teach Animals. Scientific American. 185(6):26–29.
- 474 Stoquart GG, Detrembleur C, Palumbo S, Deltombe T, Lejeune TM. 2008. Effect of
- 475 Botulinum Toxin Injection in the Rectus Femoris on Stiff-Knee Gait in People With Stroke:
- 476 A Prospective Observational Study. Archives of Physical Medicine and Rehabilitation.
- 477 89(1):56–61. doi:10.1016/j.apmr.2007.08.131.
- 478 Stutzig N, Siebert T. 2017. Assessment of the H-reflex at two contraction levels before and
- 479 after fatigue. Scandinavian Journal of Medicine & Science in Sports. 27(4):399–407.
- 480 doi:10.1111/sms.12663.
- 481 Tahayori B, Koceja D. 2019. Exercise induced operant conditioning of the H-reflex in stroke
- patients: Hopes for improving motor function through inducing plastic changes in the spinal
 pathways. J Neurol Neurol Sci Disord.:001–005. doi:10.17352/jnnsd.0000026.
- 465 patriways. J Neuror Neuror Sci Disord..001–005. doi:10.17552/jiiisd.0000020.
- 484 Thompson AK, Chen XY, Wolpaw JR. 2009. Acquisition of a Simple Motor Skill: Task-
- 485 Dependent Adaptation Plus Long-Term Change in the Human Soleus H-Reflex. Journal of
 486 Neuroscience. 29(18):5784–5792. doi:10.1523/JNEUROSCI.4326-08.2009.
- Thompson AK, Pomerantz FR, Wolpaw JR. 2013. Operant Conditioning of a Spinal Reflex
 Can Improve Locomotion after Spinal Cord Injury in Humans. Journal of Neuroscience.
- 488 Can Improve Locomotion after Spinal Cord Injury in Humans.
 489 33(6):2365–2375. doi:10.1523/JNEUROSCI.3968-12.2013.
- 490 Thompson AK, Wolpaw JR. 2021. H-reflex conditioning during locomotion in people with 491 spinal cord injury. J Physiol. 599(9):2453–2469. doi:10.1113/JP278173.

- 492 Vloka J, Hadzic A, Gautier P. 2018. Femoral Nerve Block Landmarks and Nerve Stimulator
- 493 Technique. NYSORA. [accessed 2021 Nov 2]. https://www.nysora.com/techniques/lower-
- 494 extremity/femoral/femoral-nerve-block/.
- 495 Wissel J, Ward A, Erztgaard P, Bensmail D, Hecht M. European consensus table on the use
- 496 of botulinum toxin type A in adult spasticity. doi:10.2340/16501977-0303. [accessed 2022
 497 Apr 18]. http://medicaljournals.se/jrm/content/abstract/10.2340/16501977-0303.
- Wolpaw JR. 2018. The negotiated equilibrium model of spinal cord function. The Journal of
 Physiology. 596(16):3469–3491. doi:10.1113/JP275532.
- 500 Wolpaw JR, Braitman DJ, Seegal RF. 1983. Adaptive plasticity in primate spinal stretch 501 reflex: Initial development. Journal of Neurophysiology. 50(6):1296–1311.
- 502 Wolpaw JR, Lee CL. 1987. Motoneuron response to dorsal root stimulation in anesthetized
- 503 monkeys after spinal cord transection. Exp Brain Res. 68(2). doi:10.1007/BF00248809.
- 504 [accessed 2021 Nov 2]. http://link.springer.com/10.1007/BF00248809.
- 505 Young RR, Wiegner AW. 1987. Spasticity. Clin Orthop Relat Res.(219):50–62.

506