

1 Model Evaluation of Secondary Chemistry due to Disinfection of Indoor Air with 2 Germicidal Ultraviolet Lamps

3
4 Zhe Peng,^{1*} Shelly L. Miller,² and Jose L. Jimenez^{1*}

5 ¹ Cooperative Institute for Research in Environmental Sciences and Department of Chemistry, University
6 of Colorado, Boulder, Colorado 80309, United States

7 ² Department of Mechanical Engineering, University of Colorado, Boulder, Colorado 80309, United States

8 9 * Corresponding authors:

10 Zhe Peng

11 Address: UCB 216, Boulder, Colorado 80309, United States

12 Phone number: +1 (303) 735-7258

13 Fax number: +1 (303) 492-1149

14 Email: zhe.peng@colorado.edu

15
16 Jose L. Jimenez

17 Address: UCB 216, Boulder, Colorado 80309, United States

18 Phone number: +1 (303) 492-3557

19 Fax number: +1 (303) 492-1149

20 Email: jose.jimenez@colorado.edu

21
22 **Abstract.** Air disinfection using Germicidal Ultraviolet light (GUV) has received increasing
23 attention during the COVID-19 pandemic. GUV uses UVC lamps to inactivate microorganisms,
24 but it also initiates photochemistry in air. However, GUV's indoor-air-quality impact has not been
25 investigated in detail. Here, we model the chemistry initiated by GUV at 254 ("GUV254") or 222
26 nm ("GUV222") in a typical indoor setting for different ventilation levels. Our analysis showed
27 that GUV254, usually installed in the upper room, can significantly photolyze O₃, generating OH
28 radicals that oxidize indoor volatile organic compounds (VOCs) into more oxidized VOCs.
29 Secondary organic aerosol (SOA) is also formed as a VOC-oxidation product. GUV254-induced
30 SOA formation is of the order of 0.1-1 µg/m³ for the cases studied here. GUV222 (described by
31 some as harmless to humans and thus applicable for the whole room) with the same effective
32 virus-removal rate makes a smaller indoor-air-quality impact at mid-to-high ventilation rates.
33 This is mainly because of the lower UV irradiance needed and also less efficient OH-generating
34 O₃ photolysis than GUV254. GUV222 has a higher impact than GUV254 under poor ventilation
35 due to a small but significant photochemical production of O₃ at 222 nm, which does not occur
36 with GUV254.

37

38 **Keywords:** ultraviolet germicidal irradiation; SARS-CoV-2; indoor air quality; photochemistry;
39 ventilation; airborne disease transmission

40

41 **Synopsis:** Germicidal ultraviolet light initiates indoor oxidation chemistry, potentially forming
42 indoor air pollutants. The amount is not negligible and depends on both the wavelength of light
43 and the ventilation level.


44

45 Introduction

46 Germicidal ultraviolet light (GUV) has been employed to disinfect air in indoor spaces since the
47 1930s.¹ It has been shown to effectively limit the airborne transmission of infectious diseases,
48 e.g., measles and tuberculosis.¹⁻³ This is due to photon-induced dimerization of pyrimidines in
49 the nucleic acids of airborne pathogens (and loss of their ability to replicate as a result). GUV
50 fixtures use lamps that emit in the UVC range, most commonly at 254 nm (hereinafter
51 “GUV254”).⁴ As 254 nm UV can cause skin and eye irritation,⁵ GUV254 is usually applied near
52 the ceiling (Figure 1a) or inside ventilation ducts. Recently, 222 nm UV has been shown to not
53 only have strong capability of inactivating airborne viruses,⁶ but also is claimed to be safe to
54 humans⁷ (although this is controversial),⁸ potentially allowing whole-room GUV applications
55 (“GUV222”) (Fig. 1b). Ground-resting GUV-based air cleaners have also been commercialized,
56 in which a fan continuously pulls air into a box and exposes it to UV light, from which the
57 occupants are shielded.⁹

58

59 (a)



60

61

62

63

64

65

66

67


68

69

70

71

(b)



72 (c)




Figure 1.

Schematics of a germicidal ultraviolet air disinfection system at (a) 254 nm and at (b) 222 nm in a room; (c) absorption cross sections of several important gas-phase species relevant to this study (a discontinuity in the spectrum of

87 *methacrolein is due to lack of data*).

88

89 During the COVID-19 pandemic, GUV has drawn renewed interest as a tool for airborne virus
90 inactivation. Inhalation of airborne virus is widely accepted as the main transmission route of
91 COVID-19,^{10–12} which explains the dominant indoor character of transmission.¹³ An important
92 component of the transmission is due to superspreading events,¹⁴ which have been shown to be
93 mostly explained by shared-room airborne transmission.¹⁵ Much transmission also happens in
94 close proximity due to short-range airborne transmission, but even in this situation a substantial
95 fraction of the inhaled virus may come from well-mixed room air.^{16,17} As COVID-19 remains
96 widespread, and with the possible appearance of new variants, there is a pressing need to
97 remove inhalable viruses from indoor environments.³ Similar measures would be beneficial for
98 other airborne diseases such as tuberculosis, measles, or a future pandemic virus.

99

100 Physical measures such as (natural and/or mechanical) ventilation and air filtration have been
101 proven safe and effective.¹⁸ Nevertheless, mechanical ventilation and air filtration usually can
102 remove airborne pathogens only at a few effective air changes per hour (ACH)¹⁹ and natural
103 ventilation can be highly variable and impractical depending on weather, or when pollution,
104 allergens or noise are present outdoors. When a high virus removal rate (e.g., >10 ACH) needs

105 to be ensured (e.g., in high-risk environments), GUV emerges as a practical and potentially
106 cost-effective way to achieve it.^{3,18}

107
108 UVC light is known to generate strong oxidants (e.g., OH radicals, and sometimes also O₃
109 depending on the wavelengths used),²⁰ which can subsequently oxidize volatile organic
110 compounds (VOCs) indoors and initiate organic radical chemistry in indoor air.^{21,22} Energetic
111 UVC photons can also directly photolyze many VOCs, such as peroxides^{23,24} and carbonyls,^{25,26}
112 and generate organic radicals. This radical chemistry is thought to lead to further oxidation of
113 indoor VOCs and the formation of oxygenated VOCs (OVOCs) and secondary organic aerosol
114 (SOA), both of which may have negative health effects.²⁷ Surveys of the concentration of total
115 VOCs in the indoor environments range from ~0.1-4 mg m⁻³.²⁸⁻³⁰ Thus there is always a
116 significant amount of VOC to react with any radicals and oxidants that are generated indoors,
117 and any “air cleaning” technique that can create radicals and/or oxidants indoors has the
118 potential to lead to secondary chemistry.²⁷ Very few studies on this topic have been conducted
119 with state-of-the-art measurements or models. Recently, air cleaning devices based on
120 chemistry induced by UV light (photocatalysis and OH generation, but not GUV), often also
121 marketed as suitable for air disinfection, have been experimentally shown to produce significant
122 amounts of OVOCs and SOA.^{31,32}

123
124 Despite the potential of GUV to cause secondary chemistry, to our knowledge this topic has not
125 been studied in detail to date. Some studies of GUV inactivation effectiveness have included
126 measurements of ozone, to assess whether any was generated.⁹ These studies report no
127 production of ozone when mercury vapor lamps coated to limit emission from wavelengths
128 nearer to the ozone generating wavelength of 185 nm are used, as expected. However, some
129 uncoated or improperly-coated lamps are commercially available, so ozone production can be a
130 problem in some cases. In this study, we perform a first modeling evaluation of the impacts of
131 GUV254 (assuming properly-coated lamps) and GUV222 on indoor air quality. The amounts of
132 OVOC and SOA that can be formed in typical indoor environments are investigated.

133

134

135 **Materials and Methods**

136 We include the photochemistry due to GUV and subsequent radical, oxidation, and SOA
137 formation chemistries. Given the complexity of the composition of indoor air, we simplify both
138 the chemical species present indoors and the reaction scheme, while keeping them consistent

139 with the state-of-the-art knowledge for indoor air. Surface reactions are neglected but could be
140 important, and should be investigated in future studies.

141
142 The chemical mechanism for this study is a combination of the inorganic radical chemistry in an
143 oxidation flow reactor (OFR) model^{20,33,34} and part of the Regional Atmospheric Chemistry
144 Mechanism (RACM)³⁵ relevant to this study. Section S1 (Supp. Info.) provides more details of
145 the mechanism. The mechanisms are run within the open-source KinSim chemical kinetics
146 simulator,³⁶ and are made available (See SI). We perform all simulations until a steady state is
147 reached.

148
149 We investigate a typical indoor space with representative indoor and urban outdoor air
150 concentrations from the literature (Table S2). The initial concentrations of most VOCs are
151 estimated based on McDonald et al,³⁷ with the total VOC concentration assumed to be 1.7 mg
152 m⁻³, a typical value for US indoor spaces.²⁸ See Section S2 for details on species lumping and
153 initial conditions.

154
155 The GUV254 fixture in our simulations is based on the AeroMed LEXUS L2.1 Open.³⁸ The
156 space irradiated by this device is 45 m³, and is placed in a room of 300 m³ (volume of a typical
157 classroom), thus the irradiated volume is 15% of the volume of the room, consistent with refs
158 39,40. The GUV254 model has two compartments, one for the irradiated zone and the other for
159 the rest of the room, while that for GUV222 has only one compartment. Based on the UV
160 inactivation rate constants at 222 nm for SARS-CoV-2,^{6,7} the UV intensity for GUV222 is
161 adjusted such that it provides the same whole-room effective virus-removal rate as GUV254
162 (see Section S3). Three levels of ventilation, i.e., a representative residential level (0.3 ACH,
163 “low ventilation”),⁴¹ a representative commercial level (3 ACH, “medium ventilation”),¹⁹ and a
164 representative medical level (9 ACH, “high ventilation”)³ are simulated in this study. Indoor VOC
165 emissions are set such that all VOC concentrations remain at their literature-constrained initial
166 values at low ventilation without chemistry occurring. To test COVID-19 infection risk in different
167 situations, we assume the presence of an infector shedding aerosolized SARS-CoV-2 at 16
168 quanta h⁻¹ (roughly for light exercise while speaking 50% time),⁴² which is consistent or lower
169 than values constrained for literature superspreading events.¹⁵ A quantum is an infectious dose,
170 that if inhaled by a susceptible person, will lead to a probability of infection of 1-1/e.^{15,43} The rate
171 of SARS-CoV-2 loss apart from ventilation and GUV (i.e., due to intrinsic loss of infectivity,
172 aerosol deposition etc.) is assumed to be 1 h⁻¹.


173

174

175 **Results and Discussion**

176 *Disinfection.* Figure 2 shows the amount of SARS-CoV-2 present in the room to be consistent
177 with the steady-state prediction. In the absence of GUV, the emission rate of SARS-CoV-2 is 16
178 quanta h^{-1} , and its total loss rate $1.3 h^{-1}$ ($0.3 h^{-1}$ from ventilation and $1 h^{-1}$ from decay and
179 deposition) for the low-ventilation case. The steady state SARS-CoV-2 quantity in the entire
180 room is 12.3 quanta. It is lowered to 4 and 1.6 quanta by increasing the ventilation rate to 3 and
181 9 ACH, respectively. When GUV with a whole-room virus-removal rate of $\sim 30 h^{-1}$ is applied,
182 SARS-CoV-2 decreases to ~ 0.6 , ~ 0.5 , and ~ 0.4 quanta in the low-, medium-, and high-
183 ventilation cases, respectively. The relative impact of GUV is higher at low ventilation, as
184 expected.

185



216
 217 **Figure 2.** Final quantity/concentration of the main (types of) species of interest in this study under different GUV and ventilation
 218 conditions. In the GUV254 cases, the volume-weighted average concentrations for the whole room are shown. The stable chemical
 219 species concentrations are similar between the irradiated and unirradiated zones, while the radical and SARS-CoV-2 concentrations
 220 in the unirradiated zone can be significantly lower and higher, respectively (Table S3). Note that some panels use log scale for
 221 concentrations while other panels use linear scale. SOA is assumed to have a molar weight of 200 g mol⁻¹.

222 The total quantity of SARS-CoV-2 does not directly reflect its infection risk, which also depends
223 on the volume of the room and the inhalation by susceptible occupants. For an occupant with a
224 breathing rate of $0.5 \text{ m}^3 \text{ h}^{-1}$ (typical for light physical activities)⁴⁴ present in the 300 m^3 room with
225 low ventilation and no GUV fixture for 1 h, ~ 0.02 quantum is inhaled. This corresponds to an
226 infection probability of $\sim 2\%$, since the infection probability is approximately equal to the inhaled
227 quanta if the latter is small.⁴⁵ For the cases studied in this work, infection risk is reduced by $x\sim 3$
228 by medium ventilation, and by a factor of $\sim 22/8/4$ when adding GUV to a low/medium/high
229 ventilation situation.

230
231 *Secondary Chemistry.* For chemical species in the room, ventilation alone (without GUV) can
232 make some difference (Fig. 2). The differences in O_3 , NO, and ketone concentrations are largely
233 due to these species being ventilated in or out. For other chemical species, secondary chemical
234 processes also play a role. OH radicals can form even without UV, i.e., from limonene
235 ozonolysis. As a result, OH radicals are higher at medium and high ventilation, which introduces
236 more O_3 from outdoors than at low ventilation. OH concentration at high ventilation is not higher
237 than medium ventilation because high ventilation also decreases indoor limonene, reducing the
238 overall limonene- O_3 reaction rate. HO_2 radicals are lower at higher ventilation because of higher
239 NO being ventilated into the room, which reacts with HO_2 . All other organic radicals and stable
240 products shown in Fig. 2 (including SOA) have higher concentrations in the low-ventilation case
241 due to higher VOC concentrations.

242
243 With GUV254, although the concentration of photolyzable O_3 remains relatively stable (due to
244 much stronger replenishment from outdoor air ventilation than photolytic destruction), the
245 chemistry is significantly altered by UV (Fig. 2). The fundamental cause of this change is OH
246 production from O_3 photolysis.²⁰ OH concentrations in the GUV254 cases are approximately a
247 factor of 1.2-5 and 3-20 times those in the corresponding no-UV cases for the whole room
248 average and for the irradiated zone, respectively. The difference in the higher-ventilation cases
249 is larger, due to more O_3 in the room from outside air (Fig. 2 and Table S3). OH in the higher-
250 ventilation cases is similar to daytime outdoor urban levels.⁴⁶ This OH level is high enough to
251 drive substantial oxidation of VOCs, production of other radicals (e.g., HO_2 and organic peroxy
252 radicals (RO_2)), and SOA formation. Organic peroxides (including hydroperoxides), carbonyls
253 (aldehydes (excluding formaldehyde) and ketones (excluding acetone)), and organic nitrates
254 (including peroxy nitrates) are among common VOC oxidation products and all have $\sim 10\%$ to
255 several-fold concentration increases (Fig. 2). The exceptions are ketones (excluding acetone),

256 whose production is dominated in the model by UV-independent limonene ozonolysis (Fig. 2).
257 Doubling RH significantly increases OH concentration, but changes of product species are
258 within 10% of the base RH results, likely due to non-linear buffering from the reaction scheme
259 (not shown). The exceptions are ketones, which are relatively unreactive and dominated in the
260 model by the chemistry-independent acetone emission and its dilution by ventilation (Fig. 2). In
261 addition to VOC oxidation by OH, radicals (OH, HO₂, and RO₂) are also produced by active
262 photolysis of carbonyls and peroxides at 254 nm, where both strongly absorb (Fig. 1c). Due to
263 higher peroxy radical concentrations, NO is lowered to ~30 ppt at low ventilation (Fig. 2). Such a
264 low NO concentration leads to reactions of RO₂ with both HO₂ and NO being important, as
265 estimated per ref 47 (Section S4).

266
267 SOA formation is estimated from the consumption of individual VOCs and SOA mass yields
268 from the literature (Table S4). Significant SOA production (~8 μg m⁻³ at low ventilation) occurs
269 even without GUV irradiation, through limonene ozonolysis, as this reaction has a high SOA
270 yield (20%). In the GUV254 cases, both limonene ozonolysis and VOC oxidation by OH
271 contribute to SOA. The overall SOA mass yield from total VOC is of the order of 0.1%, because
272 a large fraction of total organic molecules present are too small to form SOA through oxidation
273 (e.g., ethanol or acetone). Given that 1.5 mg m⁻³ of total VOCs (excluding limonene) are
274 present, that leads to ~1.2 μg m⁻³ due to GUV254. The enhancement of SOA formation by OH
275 oxidation relative to the no-UV cases decreases with increasing ventilation, as VOC
276 concentrations are lowered by ventilation (Fig. 2). The SOA precursors and mechanism used in
277 this work are likely incomplete, given the incomplete scientific understanding of this topic.⁴⁸

278
279 Due to the fast air exchange between the GUV254 irradiated and unirradiated spaces, the
280 concentrations of stable species are similar between these two spaces (Table S3). In contrast,
281 radicals are more rapidly consumed in the unirradiated space than supplied by the transport
282 from the irradiated space, and thus have much lower concentrations (up to >1 order of
283 magnitude for highly reactive ones such as OH and acylperoxy) in the unirradiated space.

284
285 The GUV222 cases assume irradiation of the entire room volume (Fig. 1b). 222 nm photons can
286 photolyze O₂ and produce O₃, albeit at a small rate, leading to higher O₃ in all cases relative to
287 GUV254. The amounts of organic products formed in the medium- and high-ventilation (low-
288 ventilation) cases are lower (higher) than in the unirradiated zone in the GUV254 cases (Fig. 2).

289

290 At medium and high ventilation rates, the main O₃ source in the GUV222 cases is still outdoor
291 O₃ through ventilation. Despite some O₃ production, the product enhancement is much smaller
292 than in the corresponding GUV254 cases (Fig. 2). These results indicate a weak OH-initiated
293 VOC oxidation on top of the VOC ozonolysis chemistry that is active in the no-UV cases.

294
295 This difference from the active photochemistry in the GUV254 cases can be attributed to several
296 factors. First, the UV irradiance of the 222 nm fixture is significantly lower, even in terms of the
297 number of photons emitted per unit time. 222 nm photons are ~40% more efficient in
298 inactivating SARS-CoV-2 than 254 nm photons.⁶ In addition the latter cannot be used in the
299 most efficient fashion. Due to the need to protect humans from irradiation, all photons are
300 concentrated in the small irradiated zone (15% of the room volume). Because of the limited rate
301 of transport of virus-containing aerosol to the irradiated zone, the steady-state infectious virus
302 concentration is ~70% lower than in the unirradiated zone, where the infector and the
303 susceptible individuals are present (Table S3). Even if the per-photon virus-inactivation
304 efficiency was the same, GUV254 would need about 3 times the photons for GUV222 to reach
305 the same effective GUV virus-removal rate for the occupied unirradiated space. Furthermore,
306 the first step of OH photochemical production is O₃ photolysis, whose corresponding absorption
307 at 222 nm is about ~5 times lower than at 254 nm (Fig. 1c). Simple carbonyl compounds, the
308 most abundant OVOCs in this study, also absorb much less efficiently at 222 nm than at 254 nm
309 (Fig. 1c), further reducing radical production. Although other products, such as peroxides and
310 conjugated carbonyl species, can have stronger absorption at 222 nm, their relatively low
311 concentrations (~1 ppb or lower vs. hundreds of ppb of ketones) limit their relative contributions
312 to the radical budget.

313
314 Because of the small direct production of O₃ by the GUV222 lights, O₃ in the GUV222 low-
315 ventilation case is substantially less depleted by limonene ozonolysis than for GUV254. As a
316 result, compared to GUV254 SOA formation through limonene ozonolysis is substantially
317 stronger and OH concentration is also higher (Fig. 2). Other gas-phase stable organic products
318 have comparable concentrations to those in the GUV254 case.

319
320 *Implications.* We have shown that GUV disinfection can induce active photochemistry producing
321 OVOCs and SOA in typical indoor environments. Under the conditions simulated here, these
322 products do not necessarily have significant negative effects on human health because of their
323 relatively low concentrations. Among the VOCs (including OVOCs) modeled in this study, only

324 formaldehyde has a concentration exceeding the Minimal Risk Level (MRL) recommended by
325 the CDC⁴⁹ due to strong indoor emissions. However, only a very limited number of species were
326 explicitly modeled in this study, particularly aldehydes, whose toxicity is generally high. Future
327 studies with higher chemical speciation are needed to better assess the toxicity of gas-phase
328 products. In polluted indoor spaces and/or outdoor atmospheric environments, the indoor
329 concentrations of the VOCs of interest can be much higher,²⁸ while the GUV-induced
330 photochemistry can still be active (Sections S5-S6). In this case, OVOC products might exceed
331 the MRLs and SOA formation might reach tens of $\mu\text{g m}^{-3}$.

332
333 The risk of GUV254 due to secondary photochemical products is not negligible but also not
334 dominant under typical indoor conditions. The risk for GUV222 appears to be substantially lower
335 when ventilation is not poor, but comparable or slightly higher in case of poor ventilation. We
336 note that many indoor environments, in particular homes and schools, have ventilation rates
337 similar to the definition of “poor” in this paper, even in high-income countries. If GUV222 is
338 confirmed to be safe for direct human exposure, it would have an advantage over GUV254 at
339 mid to high ventilation rates in terms of indoor chemistry, in addition to more efficient air
340 disinfection. When GUV254 is used, a strong air exchange between the irradiated and
341 unirradiated zones (e.g., by fans) is preferable, as recommended by the CDC/NIOSH.⁵⁰ It can
342 lower the UV irradiance needed for a given virus inactivation rate,⁵¹ and hence limit the induced
343 photochemistry. Good ventilation can not only remove airborne pathogens, but also limit the
344 production of secondary indoor pollutants, and is thus also recommended when outdoor air is
345 relatively clean.^{19,52} Similarly, particulate air filtration is also recommended as it removes both
346 virus-containing aerosol, indoor-formed SOA, and particulate pollution from other indoor and
347 outdoor sources. Gas filtration with sorbent materials such as activated carbon is also useful for
348 reducing VOC, NO_x and ozone levels indoors.^{53,54} The findings of this study are limited due to
349 modeling assumptions, e.g., the simplified chemical mechanism, limited data on photolysis
350 parameters in the UVC range, the assumed indoor and outdoor air pollutant compositions,
351 uncertainties over precursors and yields of SOA formation, and surface reactions. Experimental
352 studies in both simplified laboratory settings and real indoor conditions are needed to fully
353 constrain the impacts of GUV in indoor chemistry.

354

355

356 **Acknowledgements**

357 ZP and JLJ were supported by the CIRES Innovative Research Program and the Balvi
358 Filantropic Fund. We thank Vito Ilacqua, Zachary Finewax, Donald Milton, and Edward Nardell
359 for valuable discussions.

360

361 **Associated Content**

362 This paper has been previously submitted to medRxiv, a preprint server for health sciences. The
363 preprint can be cited as Peng, Z.; Miller, S. L.; Jimenez, J. L. Model Evaluation of Secondary
364 Chemistry due to Disinfection of Indoor Air with Germicidal Ultraviolet Lamps. 2022. medRxiv
365 10.1101/2022.08.25.22279238v3 (accessed November 16, 2022).

366

367 **Supporting Information**

368 The Supporting Information is available free of charge at
369 <https://pubs.acs.org/doi/10.1021/acs.estlett.xxxxxxx>.

370 Details about the model setup (reaction scheme, species lumping, indoor emission, outdoor air
371 composition, initial conditions, effective virus-removal rates of GUV), RO₂ fates in the GUV254
372 cases, concentrations in the irradiated and unirradiated zones in the GUV254 cases, and
373 sensitivity cases for highly polluted conditions.

374

375

376 **References**

- 377 (1) Wells, W. F.; Wells, M. W.; Wilder, T. S. The Environmental Control of Epidemic Contagion.
378 I. *An epidemiologic study of radiant disinfection of air in day schools* *Am J Hyg* **1942**, *35*,
379 97–121.
- 380 (2) Riley, R. L.; Mills, C. C.; O'grady, F.; Sultan, L. U.; Wittstadt, F.; Shivpuri, D. N.
381 Infectiousness of Air from a Tuberculosis Ward. Ultraviolet Irradiation of Infected Air:
382 Comparative Infectiousness of Different Patients. *Am. Rev. Respir. Dis.* **1962**, *85*, 511–525.
- 383 (3) Nardell, E. A. Air Disinfection for Airborne Infection Control with a Focus on COVID-19:
384 Why Germicidal UV Is Essential. *Photochem. Photobiol.* **2021**, *97* (3), 493–497.
- 385 (4) Riley, R. L.; Nardell, E. A. Clearing the Air: The Theory and Application of Ultraviolet Air
386 Disinfection. *American Review of Respiratory Disease*. 1989, pp 1832–1832.
387 <https://doi.org/10.1164/ajrccm/140.6.1832b>.
- 388 (5) Zaffina, S.; Camisa, V.; Lembo, M.; Vinci, M. R.; Tucci, M. G.; Borra, M.; Napolitano, A.;
389 Cannatà, V. Accidental Exposure to UV Radiation Produced by Germicidal Lamp: Case
390 Report and Risk Assessment. *Photochem. Photobiol.* **2012**, *88* (4), 1001–1004.
- 391 (6) Ma, B.; Gundy, P. M.; Gerba, C. P.; Sobsey, M. D.; Linden, K. G. UV Inactivation of SARS-
392 CoV-2 across the UVC Spectrum: KrCl* Excimer, Mercury-Vapor, and Light-Emitting-Diode
393 (LED) Sources. *Appl. Environ. Microbiol.* **2021**, *87* (22), e0153221.
- 394 (7) Buonanno, M.; Welch, D.; Shuryak, I.; Brenner, D. J. Far-UVC Light (222 Nm) Efficiently
395 and Safely Inactivates Airborne Human Coronaviruses. *Sci. Rep.* **2020**, *10* (1), 10285.
- 396 (8) Ong, Q.; Wee, W.; Cruz, J. D.; Ronnie Teo, J. W.; Han, W. 222-Nm Far UVC Exposure

- 397 Results in DNA Damage and Transcriptional Changes to Mammalian Cells. *bioRxiv*, 2022,
398 2022.02.22.481471. <https://doi.org/10.1101/2022.02.22.481471>.
- 399 (9) Kujundzic, E.; Matakah, F.; Howard, C. J.; Hernandez, M.; Miller, S. L. UV Air Cleaners
400 and Upper-Room Air Ultraviolet Germicidal Irradiation for Controlling Airborne Bacteria and
401 Fungal Spores. *J. Occup. Environ. Hyg.* **2006**, *3* (10), 536–546.
- 402 (10) Wang, C. C.; Prather, K. A.; Sznitman, J.; Jimenez, J. L.; Lakdawala, S. S.; Tufekci, Z.;
403 Marr, L. C. Airborne Transmission of Respiratory Viruses. *Science* **2021**, *373* (6558),
404 eabd9149.
- 405 (11) Greenhalgh, T.; Jimenez, J. L.; Prather, K. A.; Tufekci, Z.; Fisman, D.; Schooley, R. Ten
406 Scientific Reasons in Support of Airborne Transmission of SARS-CoV-2. *Lancet* **2021**.
407 [https://doi.org/10.1016/S0140-6736\(21\)00869-2](https://doi.org/10.1016/S0140-6736(21)00869-2).
- 408 (12) Klompas, M.; Milton, D. K.; Rhee, C.; Baker, M. A.; Leekha, S. Current Insights Into
409 Respiratory Virus Transmission and Potential Implications for Infection Control Programs :
410 A Narrative Review. *Ann. Intern. Med.* **2021**, *174* (12), 1710–1718.
- 411 (13) Qian, H.; Miao, T.; Liu, L.; Zheng, X.; Luo, D.; Li, Y. Indoor Transmission of SARS-CoV-
412 2. *Indoor Air* **2020**, in press.
- 413 (14) Adam, D. C.; Wu, P.; Wong, J. Y.; Lau, E. H. Y.; Tsang, T. K.; Cauchemez, S.; Leung,
414 G. M.; Cowling, B. J. Clustering and Superspreading Potential of SARS-CoV-2 Infections in
415 Hong Kong. *Nat. Med.* **2020**, *26* (11), 1714–1719.
- 416 (15) Peng, Z.; Rojas, A. L. P.; Kropff, E.; Bahnfleth, W.; Buonanno, G.; Dancer, S. J.;
417 Kurnitski, J.; Li, Y.; Loomans, M. G. L. C.; Marr, L. C.; Morawska, L.; Nazaroff, W.; Noakes,
418 C.; Querol, X.; Sekhar, C.; Tellier, R.; Greenhalgh, T.; Bourouiba, L.; Boerstra, A.; Tang, J.
419 W.; Miller, S. L.; Jimenez, J. L. Practical Indicators for Risk of Airborne Transmission in
420 Shared Indoor Environments and Their Application to COVID-19 Outbreaks. *Environ. Sci.*
421 *Technol.* **2022**, *56* (2), 1125–1137.
- 422 (16) Li, Y.; Cheng, P.; Jia, W. Poor Ventilation Worsens Short-Range Airborne Transmission
423 of Respiratory Infection. *Indoor Air* **2022**, *32* (1), e12946.
- 424 (17) Jimenez, J. L.; Peng, Z.; Pagonis, D. Systematic Way to Understand and Classify the
425 Shared-Room Airborne Transmission Risk of Indoor Spaces. *Indoor Air* **2022**, *32* (5),
426 e13025.
- 427 (18) Morawska, L.; Allen, J.; Bahnfleth, W.; Bluysen, P. M.; Boerstra, A.; Buonanno, G.;
428 Cao, J.; Dancer, S. J.; Floto, A.; Franchimon, F.; Greenhalgh, T.; Haworth, C.; Hogeling, J.;
429 Isaxon, C.; Jimenez, J. L.; Kurnitski, J.; Li, Y.; Loomans, M.; Marks, G.; Marr, L. C.;
430 Mazzarella, L.; Melikov, A. K.; Miller, S.; Milton, D. K.; Nazaroff, W.; Nielsen, P. V.; Noakes,
431 C.; Peccia, J.; Prather, K.; Querol, X.; Sekhar, C.; Seppänen, O.; Tanabe, S.-I.; Tang, J.
432 W.; Tellier, R.; Tham, K. W.; Wargocki, P.; Wierzbicka, A.; Yao, M. A Paradigm Shift to
433 Combat Indoor Respiratory Infection. *Science* **2021**, *372* (6543), 689–691.
- 434 (19) ASHRAE. *Ventilation for Acceptable Indoor Air Quality: ANSI/ASHRAE Standard 62.1-
435 2019*; ANSI/ASHRAE, 2019.
- 436 (20) Peng, Z.; Jimenez, J. L. Radical Chemistry in Oxidation Flow Reactors for Atmospheric
437 Chemistry Research. *Chem. Soc. Rev.* **2020**, *49* (9), 2570–2616.
- 438 (21) Atkinson, R.; Arey, J. Atmospheric Degradation of Volatile Organic Compounds. *Chem.*
439 *Rev.* **2003**, *103* (12), 4605–4638.
- 440 (22) Ziemann, P. J.; Atkinson, R. Kinetics, Products, and Mechanisms of Secondary Organic
441 Aerosol Formation. *Chem. Soc. Rev.* **2012**, *41* (19), 6582–6605.
- 442 (23) Blitz, M. A.; Heard, D. E.; Pilling, M. J. Wavelength Dependent Photodissociation of
443 CH₃OOH: Quantum Yields for CH₃O and OH, and Measurement of the OH+ CH₃OOH
444 Rate Coefficient. *J. Photochem. Photobiol. A Chem.* **2005**, *176* (1-3), 107–113.
- 445 (24) Vaghjiani, G. L.; Ravishankara, A. R. Photodissociation of H₂O₂ and CH₃OOH at 248
446 Nm and 298 K: Quantum Yields for OH, O(3P) and H(2S). *J. Chem. Phys.* **1990**, *92* (2),
447 996–1003.

- 448 (25) Link, M. F.; Farmer, D. K.; Berg, T.; Flocke, F.; Ravishankara, A. R. Measuring
449 Photodissociation Product Quantum Yields Using Chemical Ionization Mass Spectrometry:
450 A Case Study with Ketones. *J. Phys. Chem. A* **2021**, *125* (31), 6836–6844.
- 451 (26) Rajakumar, B.; Gierczak, T.; Flad, J. E.; Ravishankara, A. R.; Burkholder, J. B. The
452 CH₃CO Quantum Yield in the 248 Nm Photolysis of Acetone, Methyl Ethyl Ketone, and
453 Biacetyl. *J. Photochem. Photobiol. A Chem.* **2008**, *199* (2-3), 336–344.
- 454 (27) Collins, D. B.; Farmer, D. K. Unintended Consequences of Air Cleaning Chemistry.
455 *Environ. Sci. Technol.* **2021**. <https://doi.org/10.1021/acs.est.1c02582>.
- 456 (28) Logue, J. M.; McKone, T. E.; Sherman, M. H.; Singer, B. C. Hazard Assessment of
457 Chemical Air Contaminants Measured in Residences. *Indoor Air* **2011**, *21* (2), 92–109.
- 458 (29) Mattila, J. M.; Arata, C.; Abeleira, A.; Zhou, Y.; Wang, C.; Katz, E. F.; Goldstein, A. H.;
459 Abbatt, J. P. D.; DeCarlo, P. F.; Vance, M. E.; Farmer, D. K. Contrasting Chemical
460 Complexity and the Reactive Organic Carbon Budget of Indoor and Outdoor Air.
461 *Environmental Science & Technology*. 2022, pp 109–118.
462 <https://doi.org/10.1021/acs.est.1c03915>.
- 463 (30) Price, D. J.; Day, D. A.; Pagonis, D.; Stark, H.; Algrim, L. B.; Handschy, A. V.; Liu, S.;
464 Krechmer, J. E.; Miller, S. L.; Hunter, J. F.; de Gouw, J. A.; Ziemann, P. J.; Jimenez, J. L.
465 Budgets of Organic Carbon Composition and Oxidation in Indoor Air. *Environ. Sci. Technol.*
466 **2019**. <https://doi.org/10.1021/acs.est.9b04689>.
- 467 (31) Ye, Q.; Krechmer, J. E.; Shutter, J. D.; Barber, V. P.; Li, Y.; Helstrom, E.; Franco, L. J.;
468 Cox, J. L.; Hrdina, A. I. H.; Goss, M. B.; Tahsini, N.; Canagaratna, M.; Keutsch, F. N.; Kroll,
469 J. H. Real-Time Laboratory Measurements of VOC Emissions, Removal Rates, and
470 Byproduct Formation from Consumer-Grade Oxidation-Based Air Cleaners. *Environmental*
471 *Science & Technology Letters* **2021**. <https://doi.org/10.1021/acs.estlett.1c00773>.
- 472 (32) Joo, T.; Rivera-Rios, J. C.; Alvarado-Velez, D.; Westgate, S.; Lee Ng, N. Formation of
473 Oxidized Gases and Secondary Organic Aerosol from a Commercial Oxidant-Generating
474 Electronic Air Cleaner. *Environmental Science & Technology Letters* **2021**.
475 <https://doi.org/10.1021/acs.estlett.1c00416>.
- 476 (33) Peng, Z.; Day, D. A.; Stark, H.; Li, R.; Lee-Taylor, J.; Palm, B. B.; Brune, W. H.;
477 Jimenez, J. L. HO_x Radical Chemistry in Oxidation Flow Reactors with Low-Pressure
478 Mercury Lamps Systematically Examined by Modeling. *Atmospheric Measurement*
479 *Techniques* **2015**, *8* (11), 4863–4890.
- 480 (34) Peng, Z.; Jimenez, J. L. Modeling of the Chemistry in Oxidation Flow Reactors with High
481 Initial NO. *Atmos. Chem. Phys.* **2017**, *17* (19), 11991–12010.
- 482 (35) Stockwell, W. R.; Kirchner, F.; Kuhn, M.; Seefeld, S. A New Mechanism for Regional
483 Atmospheric Chemistry Modeling. *J. Geophys. Res.* **1997**, *102* (D22), 25847–25879.
- 484 (36) Peng, Z.; Jimenez, J. L. KinSim: A Research-Grade, User-Friendly, Visual Kinetics
485 Simulator for Chemical-Kinetics and Environmental-Chemistry Teaching. *J. Chem. Educ.*
486 **2019**, *96* (4), 806–811.
- 487 (37) McDonald, B. C.; de Gouw, J. A.; Gilman, J. B.; Jathar, S. H.; Akherati, A.; Cappa, C. D.;
488 Jimenez, J. L.; Lee-Taylor, J.; Hayes, P. L.; McKeen, S. A.; Cui, Y. Y.; Kim, S.-W.;
489 Gentner, D. R.; Isaacman-VanWertz, G.; Goldstein, A. H.; Harley, R. A.; Frost, G. J.;
490 Roberts, J. M.; Ryerson, T. B.; Trainer, M. Volatile Chemical Products Emerging as Largest
491 Petrochemical Source of Urban Organic Emissions. *Science* **2018**, *359* (6377), 760–764.
- 492 (38) *Upper room germicidal ultraviolet fixtures*. AeroMed Technologies.
493 <https://aeromed.com/product/upper-room-guv-fixtures/> (accessed 2022-02-21).
- 494 (39) Xu, P.; Kujundzic, E.; Peccia, J.; Schafer, M. P.; Moss, G.; Hernandez, M.; Miller, S. L.
495 Impact of Environmental Factors on Efficacy of Upper-Room Air Ultraviolet Germicidal
496 Irradiation for Inactivating Airborne Mycobacteria. *Environ. Sci. Technol.* **2005**, *39* (24),
497 9656–9664.
- 498 (40) Xu, P.; Peccia, J.; Fabian, P.; Martyny, J. W.; Fennelly, K. P.; Hernandez, M.; Miller, S.

- 499 L. Efficacy of Ultraviolet Germicidal Irradiation of Upper-Room Air in Inactivating Airborne
500 Bacterial Spores and Mycobacteria in Full-Scale Studies. *Atmos. Environ.* **2003**, 37 (3),
501 405–419.
- 502 (41) Daisey, J. M.; Angell, W. J.; Apte, M. G. Indoor Air Quality, Ventilation and Health
503 Symptoms in Schools: An Analysis of Existing Information. *Indoor Air* **2003**, 13 (1), 53–64.
- 504 (42) Buonanno, G.; Morawska, L.; Stabile, L. Quantitative Assessment of the Risk of Airborne
505 Transmission of SARS-CoV-2 Infection: Prospective and Retrospective Applications.
506 *Environ. Int.* **2020**, 145, 106112.
- 507 (43) Riley, E. C.; Murphy, G.; Riley, R. L. Airborne Spread of Measles in a Suburban
508 Elementary School. *Am. J. Epidemiol.* **1978**, 107 (5), 421–432.
- 509 (44) EPA. Chapter 6—Inhalation Rates. In *Exposure Factors Handbook*; U.S. Environmental
510 Protection Agency, 2011.
- 511 (45) Peng, Z.; Jimenez, J. L. Exhaled CO₂ as a COVID-19 Infection Risk Proxy for Different
512 Indoor Environments and Activities. *Environmental Science & Technology Letters* **2021**, 8
513 (5), 392–397.
- 514 (46) Ren, X.; Olson, J. R.; Crawford, J. H.; Brune, W. H.; Mao, J.; Long, R. B.; Chen, Z.;
515 Chen, G.; Avery, M. A.; Sachse, G. W.; Barrick, J. D.; Diskin, G. S.; Huey, L. G.; Fried, A.;
516 Cohen, R. C.; Heikes, B.; Wennberg, P. O.; Singh, H. B.; Blake, D. R.; Shetter, R. E.
517 HO_xchemistry during INTEX-A 2004: Observation, Model Calculation, and Comparison with
518 Previous Studies. *J. Geophys. Res.* **2008**, 113 (D5). <https://doi.org/10.1029/2007jd009166>.
- 519 (47) Peng, Z.; Lee-Taylor, J.; Orlando, J. J.; Tyndall, G. S.; Jimenez, J. L. Organic Peroxy
520 Radical Chemistry in Oxidation Flow Reactors and Environmental Chambers and Their
521 Atmospheric Relevance. *Atmos. Chem. Phys.* **2019**, 19 (2), 813–834.
- 522 (48) Shrivastava, M.; Cappa, C. D.; Fan, J.; Goldstein, A. H.; Guenther, A. B.; Jimenez, J. L.;
523 Kuang, C.; Laskin, A.; Martin, S. T.; Ng, N. L.; Petaja, T.; Pierce, J. R.; Rasch, P. J.; Roldin,
524 P.; Seinfeld, J. H.; Shilling, J.; Smith, J. N.; Thornton, J. A.; Volkamer, R.; Wang, J.;
525 Worsnop, D. R.; Zaveri, R. A.; Zelenyuk, A.; Zhang, Q. Recent Advances in Understanding
526 Secondary Organic Aerosol: Implications for Global Climate Forcing. *Rev. Geophys.* **2017**,
527 55 (2), 509–559.
- 528 (49) Hhs, U. S. Agency for Toxic Substances and Disease Registry Minimal Risk Levels
529 [WWW Document]. URL <https://www.atsdr.cdc.gov/about/index.html> (accessed 12. 18.
530 18) **2018**.
- 531 (50) Whalen, J. J. *Environmental Control for Tuberculosis; Basic Upper-Room Ultraviolet*
532 *Germicidal Irradiation Guidelines for Healthcare Settings Guide*; (NIOSH) 2009-105; 2009.
- 533 (51) Riley, R. L.; Permutt, S. Room Air Disinfection by Ultraviolet Irradiation of Upper Air. Air
534 Mixing and Germicidal Effectiveness. *Arch. Environ. Health* **1971**, 22 (2), 208–219.
- 535 (52) Morawska, L.; Tang, J. W.; Bahnfleth, W.; Bluysen, P. M.; Boerstra, A.; Buonanno, G.;
536 Cao, J.; Dancer, S.; Floto, A.; Franchimon, F.; Haworth, C.; Hogeling, J.; Isaxon, C.;
537 Jimenez, J. L.; Kurnitski, J.; Li, Y.; Loomans, M.; Marks, G.; Marr, L. C.; Mazzarella, L.;
538 Melikov, A. K.; Miller, S.; Milton, D. K.; Nazaroff, W.; Nielsen, P. V.; Noakes, C.; Peccia, J.;
539 Querol, X.; Sekhar, C.; Seppänen, O.; Tanabe, S.-I.; Tellier, R.; Tham, K. W.; Wargocki, P.;
540 Wierzbicka, A.; Yao, M. How Can Airborne Transmission of COVID-19 Indoors Be
541 Minimised? *Environ. Int.* **2020**, 142, 105832.
- 542 (53) Lee, P.; Davidson, J. Evaluation of Activated Carbon Filters for Removal of Ozone at the
543 PPB Level. *Am. Ind. Hyg. Assoc. J.* **1999**, 60 (5), 589–600.
- 544 (54) Mochida, I.; Korai, Y.; Shirahama, M.; Kawano, S.; Hada, T.; Seo, Y.; Yoshikawa, M.;
545 Yasutake, A. Removal of SO_x and NO_x over Activated Carbon Fibers. *Carbon N. Y.* **2000**,
546 38 (2), 227–239.