Effects of cardiovascular disease risk factors on cardiorespiratory fitness in

firefighters – a protocol for a systematic review and meta-analysis Jaron Ras, 1* Elpidoforos S. Soteriades, 2 Andre P. Kengne, 3 Denise Smith, 4 Lloyd Leach 1 ¹ Department of Sport, Recreation and Exercise Science, Faculty of Community and Health Sciences, University of the Western Cape, Cape Town, South Africa. ² School of Economics and Management, Healthcare Management Program. Open University of Cyprus, Nicosia, Cyprus. ³ Non-communicable Diseases Research Unit, South African Medical Research Council, Cape Town, South ⁴ Health and Human Physiological Sciences, Skidmore College, Saratoga Springs, New York. *Corresponding author: Department of Sport, Recreation and Exercise Science, Faculty of Community and Health Sciences, University of the Western Cape, Cape Town, South Africa. Jaron Ras Email: 3405618@myuwc.ac.za **Conflict of Interest:** The authors have no conflict of interest. Funding: The Ryoichi Sasakawa Young Leaders Fellowship Fund (Sylff) Program and the National research foundation (NRF) (grant number: 141282)

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

35 36

37 38

39

40

41

42

43

44

45

46

47

48 49

50

51

52

53

54

55

56

57

58

59 60

61

62 63

64

65

66

67

68 69

70

71

72

Abstract Introduction: Globally, firefighting is recognized as among the most physically demanding professions. In addition, because of the hazardous nature of firefighting, firefighters are required to wear heavy insulated personnel protective equipment that protects them from exposures to hazardous chemicals and fumes and severe temperatures. This requires that, throughout their careers, firefighters maintain their cardiovascular health and maintain a satisfactory cardiovascular fitness to manage these stressors and perform their duties sufficient vigour. The aim of this systematic is to determine the effects of cardiovascular disease risk factors on cardiorespiratory fitness of firefighters. **Methods:** Using the comprehensive search terms, a detailed literature search, with no limitation to publication year, will be conducted to identify relevant studies from PubMed/Medline, Web of Science, EBSCOHost, ScienceDirect and SCOPUS. Data-extraction will be extracted using a researcher-generated data extraction form. Data will be imported into Zotero® version 6.0.9, and duplicates removed. The article screening will be conducted using the Rayyan® intelligent systematic review tool. Thereafter, information from the included studies will be captured on the researcher-generated data extraction form. For the methodological assessment, the Appraisal Tool for Cross Sectional Studies (AXIS) checklist and the Critical Appraisal Skills Programme (CASP) toolkit will be used. For the meta-analysis, Review Manager 5.3 will be used to determine the exposure effects and MedCalc® statistical software Ltd. to determine the pooled correlation effects. Discussion: This systematic review arose from the global cardiovascular health concerns that firefighters face, and how the development and progression of CVD risk factors effect cardiorespiratory fitness in firefighters. Cardiorespiratory has been shown to be an essential factor in optimal occupational performance in firefighters but understudied related to CVD risk factors. It is anticipated that this review will make a considerable contribution to the international scientific literature on the effect that CVD risk factors will have on cardiorespiratory fitness. **Systematic review protocol registration:** PROSPERO (CRD42021258898) **Keywords:** Firefighters, cardiovascular disease, cardiorespiratory fitness, risk factors, Dyslipidaemia, obesity, hypertension, cigarette smoking.

INTRODUCTION

Globally, firefighting is recognized as among the most physically demanding professions [1–4]. In addition, because of the hazardous nature of firefighting, firefighters are required to wear heavy insulated personnel protective equipment that protects them from exposures to hazardous chemicals and fumes and severe temperatures [1–4]. The full PPE worn with the self-contained breathing apparatus (SCBA) may weigh up to 29.3 kg, which places significant strain on the cardiovascular system [1]. It has been shown that firefighters routinely exceed their age predicted maximal heart rates while performing work-related duties [5–7], which significantly predisposes firefighters to sudden cardiac events while on duty [3,4,8], which may account for the high mortality rates in this population.

Studies have shown that numerous firefighters have multiple cardiovascular disease (CVD) risk factors, with the most prevalent risk factors being obesity, cigarette smoking, dyslipidemia, and physical inactivity [9–15]. This is a concern given the physically strenuous occupation of firefighting [2–4,8]. Numerous studies have reported that firefighters have an inadmissibly high percentage of mortality due to cardiovascular related incidents, with the preponderance of these events related to underlying CVD risk factors [4,8,16]. An additional factor contributing toward the high prevalence of cardiovascular-related mortality is inadequate physical fitness, which results in increased cardiovascular strain and overexertion [17,18]. Firefighters are, therefore, required to remain in peak cardiorespiratory conditioning, while maintaining good CVD health throughout their careers [4,8,17]. Firefighters that have multiple CVD risk factors, and have suboptimal cardiorespiratory conditioning, are not only at risk for sustaining a SCE while on duty, but also at risk for endangering the public they serve to protect [4,8,16,17].

This study arose from the difficulties that firefighters routinely encounter, both globally and, in particular, locally in South Africa [12,19]. A disconcerting quantity of firefighters are at increased CVD risk and have poor cardiorespiratory fitness levels, which negatively impacts their overall health, well-being, and occupational performance [20–24]. The development of these CVD risk factors diminish firefighters ability to cope with the physical strain of firefighting, that has been described as being comparable to the demands placed on elite sportspersons [25]. It has been shown, in previous systematic reviews, that CVD risk factors are significantly related to cardiorespiratory fitness [26–29]. However, to the best of the authors knowledge, no systematic review and meta-analysis has investigated the effect of CVD risk factors on cardiorespiratory fitness in firefighters, which motivated the current systematic

review. One of the intentions of this review was to assist in informing policy change in South Africa, specifically on the need for remedial action and the development of strategies to enhance and maintain the cardiorespiratory fitness levels and cardiovascular wellbeing of firefighters.

METHODS AND ANALYSIS

Study Design

107

108

109

110

111

112

113

121

129

130

133

136

137

- 114 This protocol was registered in the PROSPERO (International prospective register of
- systematic reviews), with the registration number CRD42022330510. The systematic review
- protocol is reported in accordance with the Preferred Reporting Items for Systematic Review
- and Meta-Analysis Protocol (PRISMA-P). The systematic review methodology will be
- 118 conducted in accordance with the guidelines Meta-analysis of Observational Studies in
- Epidemiology studies (MOOSE) and Quality of Reporting of Meta-analysis (QUOROM)
- [30,31]. When considering studies for this review, the PRISMA guidelines will be followed.
- The authors have chosen to investigate the effect of CVD risk factors on cardiorespiratory
- fitness in adult, full-time firefighters. The focus of this review will be to determine the effect
- of CVD risk factors on cardiorespiratory fitness of firefighters. A quantitative systematic
- review is the preferred study design for this review [32].

126 Aim of the Review

- The aim of this systematic review is to determine the effects of cardiovascular disease risk
- factors on the cardiorespiratory fitness of firefighters.

Research Question

- What effects do cardiovascular disease risk factors have on the cardiorespiratory fitness of
- 132 firefighters?

Objectives of the Review

- The objective of the study is to investigate the effects of cardiovascular disease risk factors
- on the cardiorespiratory fitness of firefighters.

Participants

Full-time, part-time and volunteer firefighters 18 years and older.

Exposure Types

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

Cardiovascular disease risk factors in relation to the cardiorespiratory fitness of firefighters.

Outcome Types

Cardiovascular disease risk factors related to the cardiorespiratory fitness of firefighters.

Search Strategy for Identification of Studies

Using the comprehensive search terms, a detailed literature search will be conducted to identify studies investigating the effects of cardiovascular disease risk factors on the cardiorespiratory fitness of firefighters. Details on the inclusion and exclusion criteria for the systematic review is contained in Table 1:

The team of reviewers responsible for study identification will be made up of three main contributors:

- i. Reviewer I (JR) will be the primary investigator. JR will assume responsibility for all aspects of the review and will be one of the reviewers tasked with conducting the literature search, independently extracting the data, responsible for verifying the data collected, analysing the results, grading the quality of the data, and write up of the first draft of the systematic review.
- ii. Reviewer II (RN) will be the second reviewer and be tasked with independently conducting a literature search, extracting the data of studies, validating the data collected, and grading the quality of the data.
- iii. Reviewer III (LL) will serve as the adjudicator and resolve any disagreements between the two independent reviewers.

Table 1: Inclusion and exclusion criteria of studies.

Inclu	sion Criteria	Exclusion Criteria		
i.	Studies that included full-time, part-time	i. Studies not focusing on		
	and volunteer adult firefighters between	cardiorespiratory fitness as		
	the ages of 18 and 65 years, with no	the outcomes.		
	limitations to publication year.			

ii.	Observational studies, such as cross-	ii.	Studies not focusing on
	sectional, cohort and case controlled, and		cardiovascular disease risk
	experimental study designs.		factors as the main exposures.
iii.	Studies investigating the effects of	iii.	Systematic reviews or other
	cardiovascular disease risk factors on		types of reviews.
	cardiorespiratory fitness in firefighters.	iv.	Articles that are non-English.

Electronic Literature Search

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

Using the comprehensive search terms, an extensive literature search will be conducted to enable the capturing of as many relevant articles as possible to be included in the systematic review and meta-analysis, however, these studies will be limited to English papers only. The academic journal databases that will be searched will include PubMed/Medline, Web of Science, SCOPUS, ScienceDirect and EBSCOhost. The literature search will take place between August and September 2022. In addition, there will be no limitation to publication year to allow inclusivity. The comprehensive search strategy will include the arrangement of keywords and medical subject heading (MeSH) in various arrangements depending on the specific database searched. In Table 2, the various arrangements of each of the search terms for the population and the cardiovascular disease risk factors will be combined with each of the outcome terms. The search strategies for the other databases are presented in Appendix 1.

Table 2: Search strategy developed in PubMed.

Search #	Search terms
#1	"firefighter" OR "fire and rescue" OR "fire and rescue personnel" OR "fire
	fighters" OR "fire fighter" OR "firefight"
#2	("cardiovascular" [All Fields] AND "system" [All Fields])" OR cardiovascular
	system" [MeSH] OR "cardiovascular system" [All Fields] OR "cardiovascular*"
	[All Fields] OR "cardiovascular abnormalities" [MeSH] OR "heart rate
	variability" [All Fields] OR "HRV" [All Fields] OR "Heart Rate Interval" [All
	Fields] OR "cycle length variability" [All Fields] OR "RR variability" [All
	Fields] OR "heart period variability" [All Fields] OR "autonomic function" [All
	Fields] OR "vagal control" [All Fields] OR "lipid profile" [All Fields] OR
	"cholesterol" [MeSH] OR "dyslipidaemia" OR "dyslipidaemia" OR
	"hypercholesteremia" OR "diabetes" AND "mellitus" OR "blood glucose" OR
	"age" OR "aging" OR "age-group" OR "obesity" OR "central obesity" OR
	"waist circumference" OR "hypertension" OR "high blood pressure" OR

	"blood pressure" OR "metabolic syndrome" OR "hyperglycaemia" OR "cardiometabolic" OR "metabolic syndrome" OR "MetSyn"
#3	"physical fitness" [MeSH] OR "fitness" OR "exercise" [All Fields] OR "physical exertion" [All Fields] OR "muscular strength" OR "muscular endurance" OR "aerobic fitness" OR "cardiorespiratory fitness" OR "cardiorespiratory capacity" OR "VO ₂ max" OR "aerobic fitness" OR "power" OR "anaerobic power" OR "aerobic capacity"
#4	(#1 AND #2) OR (#1 AND #3) OR (#1 AND #2 AND #3)

Study Selection

Initially, the electronic database search will be performed to acquire all eligible research papers. All studies that meet the inclusion criteria will be selected for full-text screening. Authors will be contacted in the instance where data is missing or where a full-text is not available. Subsequently, the full-text articles will be assessed independently by the two principal reviewers (JR and RN), using the Rayyan® intelligent systematic review (RIS) tool [33]. When screening and categorizing the studies, three categories will be used, specifically, included, excluded and unsure. Where uncertainties in study inclusion or study grading, the dispute will be discussed between the two reviewers, and in the event of disagreement, the adjudicator (LL) will resolve the dispute.

Selection and Screening of Articles

Firstly, all relevant, pre-selected databases will be searched to identify and, initially, screen the titles and abstracts of potential studies for eligibility. Secondly, the search outputs will be downloaded and compiled in a reference software, namely, Zotero® version 6.0.9 and thirdly, the reference software will be used to remove duplicate studies. Fourthly, the studies remaining after the removal of duplicates, will be screened against the inclusion criteria to determine the final studies for inclusion in the review and meta-analysis. Next, data will be extracted from the included studies using a standard data extraction form. Lastly, studies included into the meta-analysis will be analysed and interpreted using Review Manager 5.3[34] and MedCalc® statistical software Ltd.

Data Management

All data that will be extracted from the initially selected studies will be captured using an initial electronic researcher-generated data extraction form (Appendix 2), by the two principal reviewers, in order to retrieve the key characteristics of each study. Thereafter, a detailed second data extraction form (Appendix 3) will be used to capture information of the included studies. The study data that will be extracted will include the general details of the study, such as the study title, authors, publication date, study design and the country of origin, the exposures evaluated, and the outcome measured. Thereafter, characteristics of the included studies will be collected, such as the sample size and method of sampling, participants details (age, height, weight, body mass index, cardiorespiratory fitness, gender). Lastly, the details of exposures and the outcome variables will be extracted, i.e., the research paper must report on at least one of the CVD risk factors under investigation in relation to cardiorespiratory fitness.

- Critical Appraisal of Included Studies
- Two critical appraisal tools will be used for this systematic review and meta-analysis, namely, 219
- The Critical Appraisal Skills Programme (CASP) toolkit (Middle Way, Oxford, UK) and the 220
- Appraisal Tool for Cross Sectional Studies (AXIS) checklist [35] (Table 2) (https://casp-221
- uk.net/casp-tools-checklists/) to conduct the methodological quality assessment of each study 222
- 223 that was included. The CASP toolkit and the AXIS toolkit was shown to be a reliable and valid
- tool for assessing the quality of cross-sectional studies [35,36]. For the grading of each study, 224
- where questions can be answered dichotomously, an article will be awarded a yes, denoted as 225
- a "\(\sigma\)", and when scored a no, an "\(\mathbf{x}\)" will be denoted. For questions that require written grading, 226
- all attempts will be made to adapt the question to a dichotomous scale, and if a rating is not 227
- possible, the question will be excluded from the checklist. For questions that are not relevant, 228
- an "na" will be given. 229

206

207

208

209

210

211

212

213

214

215

216

217

218

230

231

237

238

Data Synthesis and Meta-Analysis

- Data synthesis will commence once all relevant details of each study has been captured. 232
- Initially, a systematic synthesis of the results will be used to integrate the findings, narratively, 233
- from the various studies. This method will be used initially as it allows the researcher to 234
- identify, evaluate, and summarize study findings that are similar across all relevant studies [37– 235
- 39]. Thereafter, data will be captured, quantitatively, and analysed using a meta-analysis. 236
 - Meta-Analysis

Where there is sufficient reporting and the data permits, a meta-analysis will be conducted. For each study, quantitative data will be pooled to determine the pooled effect estimate [40]. For dichotomous data, the odds ratio (OR) and risk ratio (RR) will be generated, whereas for continuous data, the mean difference (MD) and standardized mean difference (SMD) of estimation will be used to estimate the effect of each cardiovascular disease risk factor on cardiorespiratory fitness of firefighters [41]. The data will be imported and analysed using the Review Manager 5.3 [34,39,41,42], using the inverse method of meta-analysis [40]. In addition, a meta-analysis of correlations will be performed to determine the relationship between continuous CVD risk factors and cardiorespiratory fitness of firefighters. To perform the analysis, the MedCalc® statistical software Ltd. (version 20.104) will be preferred [40]. The meta-analysis will involve using the original sample sizes and r values to generate the pooled r values between each cardiovascular risk factor in relation to cardiorespiratory fitness [40]. Using the Fisher's r to z transformation the original r values will be converted to a common test metric [40]:

$$Z_{ri} = \frac{1}{2} ln \left(\frac{1+ri}{1-ri} \right) s_z^2 = \frac{1}{n-3}$$

- The following will be used to indicate the strength of correlation [40]:
- 257 1. Very high correlation: 0.90 to 1.00 (-0.90 to -1.00).
- 258 2. High correlation: 0.70 to 0.90 (-0.70 to -0.90).
- 3. Moderate correlation: 0.50 to 0.70 (-0.50 to -0.70).
- 4. Low correlation: 0.30 to 0.50 (-0.30 to -0.50).
- 5. Negligible correlation: 0.00 to 0.30 (-0.00 to -0.30).
- 263 Assessment of Heterogeneity

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

255

262

269

- Heterogeneity will be evaluated, initially, through visual inspection of the forest plot, to
- judge the extent of overlap. In addition, to quantitively assess the heterogeneity, the Chi-
- square test and the I² test will be used [60]. The following will be used to explain I² statistics:
- 267 1) 0% to 30% not be important; 2) 31% to 60% moderate heterogeneity; 3) 61% to 80%
- substantial heterogeneity; and 4) 81% to 100% considerable heterogeneity [40,42].
- In the instance where substantial and considerable heterogeneity is found, potential reasons

will be determined by performing a subgroup analysis. A meta-analysis will be favoured by a low degree of heterogeneity [60]. Where significant heterogeneity is present, then a descriptive analysis of the results will be presented. If homogeneity is found between the studies, then a fixed-effects model will be preferred, and if heterogeneity is present, a random-effects model will be preferred [60].

- Subgroup Analysis and Investigation of Heterogeneity
- The following characteristics are expected to introduce significant clinical heterogeneity, i.e.,
- firefighters' age, gender, experience, methods of measuring CVD risk factors, type of testing
- procedure used to assess cardiorespiratory fitness, and plan to implement subgroup analysis on
- these variables, if possible. Although all exposures and outcomes will be measured using
- standardized instruments and protocols, the protocols preferred by each study may be different.
- This will necessitate the comparison and conversion of certain variables to produce analogous
- findings for comparison. The authors will use Review Manager for the subgroup analysis
- [39,41,42], and MedCalc® statistical software Ltd.[40] if adequate numbers of included studies
- are available.

271

272

273

274

275

276

277

287

292

298

299

- 288 Publication Bias
- Publication bias will be assessed visually using the Begg's funnel chart. Thereafter,
- 290 quantitative analysis will be conducted to assess the possibility of publication bias using the
- Egger's test and Begg's test [40].
- 293 Presenting and Reporting of Results
- The results generated will be presented using a combination of figures, graphs, and tables. The
- presentation will include the method involved in study selection, exclusion and inclusion, using
- the PRISMA guidelines [32]. In addition, forest plots, funnel charts and summary tables will
- be created to present the data.

DISCUSSION

- To the best of the authors' knowledge, no conclusive evidence currently exists on the effect of
- 301 CVD risk factors and cardiorespiratory fitness of firefighters. This systematic review arose
- from the global cardiovascular health concerns that firefighters face, and how the development
- and progression of CVD risk factors effect cardiorespiratory fitness in firefighters.
- Cardiorespiratory has been shown to be an important factor in optimal occupational

performance in firefighters, but understudied [21,43–45]. The results of this review is expected to make a significant contribution to the international scientific literature on the effect that CVD risk factors will have on cardiorespiratory fitness. In addition, the results of this review will assist policy makers in implementing policy changes, particularly in developing countries, such as South Africa, but also globally, to promote the wellbeing, health, and career longevity of firefighters. Moreover, the planned review will support researchers design original studies relating to this issue and, assist in identifying gaps in the research for further studies.

As with all research, there are inherent limitations in the current systematic review and metaanalysis protocol when interpreting the findings. One limitation of this study is that only English studies will be used, which may result in the exclusion of potentially relevant studies. Another limitation is that there may be considerable heterogeneity between studies, due to differences in study population age, gender and the method used to assess cardiorespiratory fitness. There are, however, several strengths of this study, one of which being that this review will be guided by reference methodologies to ensure that a high standard of quality is maintained throughout the review process. The exclusion of a publication date limit is a strength of this review, removing the possibility of high-quality studies being excluded due to falling outside the publication limit. Lastly, the inclusion of all study designs, using various study methodologies, in this review.

Ethics and Dissemination

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

This study has been registered onto PROSPERO (CRD42022330510) and has been granted ethical clearance by the University of the Western Cape (BM21/10/9). Because a systematic review and meta-analysis will be conducted, there will be no direct engagement with human participants. Freely accessible and published data will be used in the study and, therefore, no confidentiality or ethical procedures need to be considered for this review [46]. The information gathered will be presented to local firefighter organisations and health departments, at conferences and in webinars. Furthermore, the systematic review and metaanalysis will be published in an accredited peer-reviewed journal. Lastly, this protocol will form a chapter of a doctoral thesis.

AUTHOR CONTRIBUTIONS

- **Conceptualisation**: Jaron Ras 337
- **Data curation:** Jaron Ras 338

- 339 **Formal analysis:** Jaron Ras
- **Funding acquisition:** Jaron Ras
- 341 **Investigation:** Jaron Ras
- Methodology: Jaron Ras, Elpidofos Soteriades, Denise Smith, Andre Kengne and Lloyd Leach
- Project administration: Jaron Ras, Elpidofos Soteriades, Denise Smith, Andre Kengne and
- 344 Lloyd Leach
- Resources: Jaron Ras, Elpidofos Soteriades, Denise Smith, Andre Kengne and Lloyd Leach
- 346 Supervision: Elpidofos Soteriades, Denise Smith, Andre Kengne and Lloyd Leach
- Writing original draft preparation: Jaron Ras
- Writing review and editing: Elpidofos Soteriades, Denise Smith, Andre Kengne and Lloyd
- 349 Leach

350

351

355

356

357358

- **Protocol Registration**
- Details of the protocol for this systematic review were registered on PROSPERO
- 353 (CRD42022330510) and can be accessed at
- https://www.crd.york.ac.uk/prospero/display record.php?RecordID=330510

REFERENCES

- Li J, Wang Y, Jiang R, *et al.* Quantifying self-contained breathing apparatus on physiology and psychological responses during firefighting: a systematic review and meta-analysis. *International Journal of Occupational Safety and Ergonomics*Published Online First: 2022. doi:10.1080/10803548.2021.2024020
- Smith DL, Haller JM, Benedict R, *et al.* Cardiac Strain Associated with High-rise Firefighting. 2015;:213–21. doi:10.1080/15459624.2014.970272
- Smith DL, DeBlois JP, Kales SN, *et al.* Cardiovascular strain of firefighting and the risk of sudden cardiac events. *Exerc Sport Sci Rev* 2016;**44**:90–7. doi:10.1249/JES.00000000000001
- Smith DL, Barr DA, Kales SN. Extreme sacrifice: Sudden cardiac death in the US Fire Service. *Extrem Physiol Med* 2013;**2**:1–9. doi:10.1186/2046-7648-2-6
- 5 Smith DL, Haller JM, Benedict R, *et al.* Firefighter incident rehabilitation: Interpreting heart rate responses. *Prehospital Emergency Care* 2016;**20**:28–36. doi:10.3109/10903127.2015.1037477
- Marcel-Millet P, Ravier G, Grospretre S, *et al.* Physiological responses and parasympathetic reactivation in rescue interventions: The effect of the breathing apparatus. *Scand J Med Sci Sports* 2018;**28**:2710–22. doi:https://doi.org/10.1111/sms.13291
- Nazari G, Lu S, MacDermid JC. Quantifying physiological responses during simulated tasks among Canadian firefighters: A systematic review and meta-analysis. *J Mil Veteran Fam Health* 2021;7:55–75. doi:10.3138/JMVFH-2019-0063

- 8 Yang J, Teehan D, Farioli A, et al. Sudden cardiac death among firefighters ≤45 years 380 of age in the United States. American Journal of Cardiology 2013;112:1962–7. 381 doi:10.1016/j.amjcard.2013.08.029 382
- 9 Soares EMKVK, Smith D, Grossi Porto LG. Worldwide prevalence of obesity among 383 firefighters: A systematic review protocol. BMJ Open 2020;10:1–5. 384 doi:10.1136/bmjopen-2019-031282 385
- 10 Smith DL, Fehling PC, Frisch A, et al. The prevalence of cardiovascular disease risk 386 factors and obesity in firefighters. J Obes 2012;**2012**. doi:10.1155/2012/908267 387
- 11 Risavi BL, Staszko J. Prevalence of Risk Factors for Coronary Artery Disease in 388 Pennsylvania (USA) Firefighters. *Prehosp Disaster Med* 2015;**31**:102–7. 389 doi:10.1017/S1049023X15005415 390
- 12 Ras J, Leach L. Prevalence of coronary artery disease risk factors in firefighters in the 391 city of Cape Town fire and rescue service – A descriptive study. J Public Health Res 392 2021;**10**. doi:10.4081/jphr.2021.2000 393
- Damacena FC, Batista TJ, Ayres LR, et al. Obesity prevalence in Brazilian firefighters 13 394 and the association of central obesity with personal, occupational and cardiovascular 395 risk factors: a cross-sectional study. BMJ Open 2020;10:1–10. doi:10.1136/bmjopen-396 2019-032933 397
- 14 Savall A, Charles R, Bertholon A, et al. Volunteer and career French firefighters: 398 Cardiovascular risk factors and cardiovascular risk assessment. Eur J Prev Cardiol 399 2020;**27**:107–9. doi:10.1177/2047487319827463 400
- 15 Martin ZT, Schlaff RA, Hemenway JK, et al. Cardiovascular Disease Risk Factors and 401 Physical Fitness in Volunteer Firefighters. Int J Exerc Sci 2019;12:764– 402 76.http://www.ncbi.nlm.nih.gov/pubmed/31156744%0Ahttp://www.pubmedcentral.ni 403 h.gov/articlerender.fcgi?artid=PMC6533096 404
- 16 Soteriades ES, Smith DL, Tsismenakis AJ, et al. Cardiovascular disease in US 405 firefighters: A systematic review. Cardiol Rev 2011;19:202–15. 406 doi:10.1097/CRD.0b013e318215c105 407
- 17 Smith DL, Haller JM, Korre M, et al. The Relation of Emergency Duties to Cardiac 408 Death Among US Firefighters. American Journal of Cardiology 2019;123:736–41. 409 doi:10.1016/j.amjcard.2018.11.049 410
- Farioli A, Yang J, Teehan D, et al. Duty-related risk of sudden cardiac death among 18 411 young US firefighters. Occup Med (Chic Ill) 2014;64:428–35. 412 doi:10.1093/occmed/kgu102 413
- 19 Ras J, Mosie D, Strauss M, et al. Knowledge of and attitudes toward health and 414 415 cardiovascular disease risk factors among firefighters in Cape Town, South Africa. J Public Health Res 2021;11. doi:10.4081/jphr.2021.2307 416
- Phelps SM, Drew-Nord DC, Neitzel RL, et al. Characteristics and Predictors of 20 417 Occupational Injury Among Career Firefighters. Workplace Health Saf 2018;66:291– 418 301. doi:10.1177/2165079917740595 419
- Wynn P, Hawdon P. Cardiorespiratory fitness selection standard and occupational 21 420 outcomes in trainee firefighters. 2012;:123-8. doi:10.1093/occmed/kgr206 421
- 22 McDonough SL, Phillips JS, Twilbeck TJ. Determining Best Practices to Reduce 422 Occupational Health Risks in Firefighters. The Journal of Strength & Conditioning 423 Research 2015;29.https://journals.lww.com/nsca-424
- jscr/Fulltext/2015/07000/Determining Best Practices to Reduce Occupational.34.as 425 426
- 23 Nazari G, Osifeso TA, MacDermid JC. Distribution of Number, Location of Pain and 427 Comorbidities, and Determinants of Work Limitations among Firefighters. Rehabil 428 Res Pract 2020;**2020**:1942513. doi:10.1155/2020/1942513 429

- 24 Nazari G, MacDermid JC, Sinden KE, et al. The Relationship between Physical 430 Fitness and Simulated Firefighting Task Performance. Rehabil Res Pract 431 2018:**2018**:1–7. doi:10.1155/2018/3234176 432
- 25 Bucala M, Sweet E. Obesity in the fire service: An inside look at the perceptions of 433 firefighters towards obesity and other health issues. Res Sq Published Online First: 434 2019. doi:10.21203/rs.2.15518/v1 435
- 26 Chu DJ, al Rifai M, Virani SS, et al. The relationship between cardiorespiratory 436 fitness, cardiovascular risk factors and atherosclerosis. Atherosclerosis 2020;304:44– 437 52. doi:https://doi.org/10.1016/j.atherosclerosis.2020.04.019 438
- 27 Tarp J. Støle AP, Blond K, et al. Cardiorespiratory fitness, muscular strength and risk 439 of type 2 diabetes: a systematic review and meta-analysis. *Diabetologia* 440 2019;**62**:1129–42. doi:10.1007/s00125-019-4867-4 441
- 28 Vega GL, Grundy SM, Barlow CE, et al. Association of triglyceride-to-high density 442 lipoprotein cholesterol ratio to cardiorespiratory fitness in men. J Clin Lipidol 443 2016;**10**:1414-1422.e1. doi:https://doi.org/10.1016/j.jacl.2016.09.008 444
- 29 Parto P, Lavie CJ, Swift D, et al. The role of cardiorespiratory fitness on plasma lipid 445 levels. Expert Rev Cardiovasc Ther 2015;13:1177-83. 446 doi:10.1586/14779072.2015.1092384 447
- 30 Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of Observational Studies in 448 Epidemiology A Proposal for Reporting. *JAMA* 2000;**283**:2008–12. 449 doi:10.1001/jama.283.15.2008 450
- Brand RA. Editorial: Standards of Reporting: The CONSORT, QUORUM, and 31 451 STROBE Guidelines. Clin Orthop Relat Res 2009;467:1393-4. doi:10.1007/s11999-452 009-0786-x 453
- Moher D, Liberati A, Tetzlaff J, et al. Preferred Reporting Items for Systematic 32 454 Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 455 2009;**6**:e1000097.https://doi.org/10.1371/journal.pmed.1000097 456
- 33 Ouzzani M, Hammady H, Fedorowicz Z, et al. Rayyan-a web and mobile app for 457 systematic reviews. Syst Rev 2016;5:1–10. doi:10.1186/s13643-016-0384-4 458
- Collaboration C. Review manager (RevMan) version 5.3. Copenhagen: The Nordic 459 34 Cochrane Centre 2014. 460
- Downes MJ, Brennan ML, Williams HC, et al. Development of a critical appraisal tool 35 461 to assess the quality of cross-sectional studies (AXIS). BMJ Open 2016;6:e011458. 462 doi:10.1136/bmjopen-2016-011458 463
- Sanderson S, Tatt ID, Higgins JPT. Tools for assessing quality and susceptibility to 36 464 465 bias in observational studies in epidemiology: a systematic review and annotated bibliography. Int J Epidemiol 2007;**36**:666–76. doi:10.1093/ije/dym018 466
- 37 Harris JD, Quatman CE, Manring MM, et al. The American Journal of Sports 467 Medicine P < P How to Write a Systematic Review. Published Online First: 2013. 468 doi:10.1177/0363546513497567 469
- Metelli S, Chaimani A. Challenges in meta- analyses with observational studies. 38 470 2020::83-7. doi:10.1136/ebmental-2019-300129 471
- Gurevitch J, Koricheva J, Nakagawa S, et al. Meta-analysis and the science of research 39 472 synthesis. Nature 2018;555:175-82. doi:10.1038/nature25753 473
- 40 Akkaya-Hocagil T. Meta-analysis: Methods for health and experimental studies: 474 Shahjahan Khan Springer Nature Singapore Pte. Ltd. 2020. pp. 293. Biometrics 475 2022;78:414. doi:https://doi.org/10.1111/biom.13605
- 41 Haidich AB. Meta-analysis in medical research. *Hippokratia* 2010;14:29– 477 37.https://pubmed.ncbi.nlm.nih.gov/21487488 478

476

479	42	Lin L. Comparison of four heterogeneity measures for meta-analysis. <i>J Eval Clin Pract</i>
480		2020; 26 :376–84. doi:10.1111/jep.13159

- 481 43 Siddall AG, Stevenson RDM, Turner PJF, *et al.* Physical and Physiological Performance Determinants of a Firefighting Simulation Test. *J Occup Environ Med* 2018;**60**:637–43. doi:10.1097/JOM.000000000001313
- Heimburg E von, Ingulf Medbø J, Sandsund M, *et al.* Performance on a Work-Simulating Firefighter Test Versus Approved Laboratory Tests for Firefighters and Applicants. *International Journal of Occupational Safety and Ergonomics* 2013;19:227–43. doi:10.1080/10803548.2013.11076981
- Skinner TL, Kelly VG, Boytar AN, *et al.* Aviation Rescue Firefighters physical fitness and predictors of task performance. *J Sci Med Sport* 2020;**23**:1228–33. doi:10.1016/j.jsams.2020.05.013
- 491 46 Vergnes J-N, Marchal-Sixou C, Nabet C, *et al.* Ethics in systematic reviews. *J Med*492 *Ethics* 2010;**36**:771 LP 774. doi:10.1136/jme.2010.039941

493 494

495