1 Title

- 2 Cardiovascular risk of dementia is associated with brain–behaviour changes in cognitively
- 3 healthy, middle-aged individuals

4 Authors

- 5 Feng Deng ab, Maria-Eleni Dounavi c, Emanuele R. G. Plini a, Karen Ritchie d, Graciela
- 6 Muniz-Terrera ^{ef}, Siobhan Hutchinson ^g, Paresh Malhotra ^h, Craig W Ritchie ^e, Brian Lawlor
- 7 ab, Lorina Naci ab*

8 Affiliations

- ^a Trinity College Institute of Neuroscience, School of Psychology, Trinity College Dublin,
- 10 Dublin, Ireland.
- ^b Global Brain Health Institute, Trinity College Dublin, Dublin, Ireland
- ^c Department of Psychiatry, School of Clinical Medicine, University of Cambridge,
- 13 Cambridge CB2 0SP, UK
- d U1061 Neuropsychiatry, INSERM, University of Montpellier, Montpellier, France
- 15 ^e Edinburgh Dementia Prevention, University of Edinburgh, Edinburgh, UK
- 16 f Department of Social medicine, Ohio University, USA
- 17 g Department of Neurology, St James's Hospital, Dublin, Ireland
- ^h Department of Brain Science, Imperial College Healthcare NHS Trust, UK

19 *Corresponding author:

- 20 Lorina Naci
- 21 School of Psychology
- 22 Trinity College Institute of Neuroscience
- 23 Global Brain Health Institute
- 24 Trinity College Dublin
- 25 Dublin, Ireland
- 26 Email: <u>nacil@tcd.ie</u>

Abstract

27

45

Alzheimer's Disease (AD) neuropathology start decades before clinical manifestations, but 28 29 whether risk factors are associated with early cognitive and brain changes in midlife remains poorly understood. We examined whether AD risk factors were associated with cognition and 30 functional connectivity (FC) between the Locus Coeruleus (LC) and hippocampus – two key 31 32 brain structures in AD neuropathology – cross-sectionally and longitudinally in cognitively healthy midlife individuals. Neuropsychological assessments and functional Magnetic 33 Resonance Imaging were obtained at baseline (N=210), and two-years follow-up (N=188). 34 Associations of cognition and FC with apolipoprotein \(\epsilon 4 \) (APOE \(\epsilon 4 \)) genotype, family history 35 of dementia, and the Cardiovascular Risk Factors, Aging, and Incidence of Dementia 36 (CAIDE) score were investigated. Cross-sectionally, higher CAIDE scores were associated 37 with worse cognition. Menopausal status interacted with the CAIDE risk on cognition. 38 Furthermore, the CAIDE score significantly moderated the relationship between cognition 39 40 and LC-Hippocampus FC. Longitudinally, the LC-Hippocampus FC decreased significantly over 2 years. These results shed light on some of the earliest effects of AD risk on cognition 41 and the brain. 42 43 Keywords: Alzheimer's disease; Cognition; Locus Coeruleus; Hippocampus; Functional 44

connectivity; Midlife; Cardiovascular risk score

1. Introduction

- 47 Dementia, particularly Alzheimer's disease (AD), is a growing epidemic that presents
- 48 profound challenges to healthcare systems, families, and societies throughout the world
- 49 (World Health Organization, 2021). Midlife is a critical period for the beginning of AD
- 50 pathology (Jansen et al., 2015; Sperling et al., 2011) and potentially a unique disease-altering
- window prior to the manifestation of substantial brain damage. Therefore, there is an urgent
- need for risk reduction interventions focused on midlife (Barnes & Yaffe, 2011; Livingston et
- al., 2017; Ritchie et al., 2010). However, the indicators and brain mechanism of AD in
- midlife remain poorly understood (Irwin et al., 2018; Ritchie et al., 2017).
- To investigate early changes, studies in preclinical AD populations have used risk
- stratification approaches. Key risk factors include the Apolipoprotein E (APOE) & genotype
- 57 (Donix et al., 2012; Liu et al., 2013), the main genetic risk factor for sporadic late-onset AD
- in the Indo-European population (Lambert et al., 2013), and family history of dementia (FHD)
- (Berti et al., 2011; Donix et al., 2012; Scarabino et al., 2016). Additionally, there is growing
- evidence to suggest that up to 40% of all dementia cases are associated with known
- 61 modifiable risk factors (Livingston et al., 2020). Among dementia risk scores that
- 62 incorporate lifestyle risk factors (Barnes et al., 2014; Deckers et al., 2015; Kivipelto et al.,
- 63 2006), the Cardiovascular Risk Factors Aging and Dementia (CAIDE) score has been
- optimized for midlife (Kivipelto et al., 2006) and validated in a large US population followed
- longitudinally over 40 years (Exalto et al., 2014).
- AD is neuropathologically characterized by the accumulation of beta-amyloid $(A\beta)$ and
- 67 hyperphosphorylated tau (pTau) (Braak & Braak, 1991; Braak et al., 2011). Recent studies
- suggest that tau deposition is a key etiological factor that presages sporadic AD (Arnsten et
- al., 2021; Jacobs et al., 2021; Kametani & Hasegawa, 2018). Analyses of thousands of human

70 brains across the lifespan suggest that tau pathology may begin about a decade before formation of A β plaques (Braak & Del Tredici, 2015). Additionally, tau pathology correlates 71 with progressive grey matter loss to a greater extent than A β (La Joie et al., 2019) and 72 73 cognitive impairment (Giannakopoulos et al., 2003). Neuropathological findings show that the nucleus locus coeruleus (LC) is one of the subcortical regions where tau pathology starts 74 in late young adulthood and early midlife (30–40 years) (Gilvesy et al., 2022; Ehrenberg et al., 75 76 2017; Braak & Braak, 1991) then spreads to the transentorhinal and entorhinal cortex (Braak et al., 2011), to the hippocampal and neocortical association cortex, and finally throughout 77 78 the neocortex. It has been suggested that early tau pathology in the LC induces LC hyperactivity, thereby 79 promoting its own spread to interconnected brain regions and facilitating the progression of 80 the disease (Weinshenker, 2018). A recent review of the neuropathological literature across 81 82 the lifespan (Arnsten et al., 2021) suggests that the spread of tau pathology from the LC to 83 the medial temporal lobe (MTL) captures the earliest stages of incipient AD progression (but 84 see Kaufman et al. (2018) for a differing account). However, alterations of connectivity between the LC and the hippocampus, due to AD risk factors have not previously been 85 86 studied in midlife preclinical populations. LC is the earliest site of tau pathology accumulation, and, furthermore, the main brain site of 87 88 noradrenaline production (Amaral & Sinnamon 1977), and thus the originating node of the brain's arousal system (Sara, 2009; Aston-Jones & Cohen 2005). Therefore, early LC 89 pathology can critically impact arousal and cognitive processes throughout the brain (Mather 90 91 & Harley, 2016; Betts, Kirilina, et al., 2019). Previous in vivo studies of the LC have investigated healthy lifespan adult cohorts (Betts et al., 2017; Clewett et al., 2016; Lee et al., 92

2020; Liu et al., 2019; Porat et al., 2022), cognitively unimpaired older adults and preclinical

94 populations with risk for sporadic AD (Jacobs et al., 2021; Prokopiou et al., 2022; Van Egroo et al., 2021), individuals with preclinical autosomal dominant Alzheimer's disease (Jacobs et 95 al., 2022), mild cognitive impairment (Jacobs et al., 2015; Takahashi et al., 2015), and 96 Alzheimer's disease (Betts, Cardenas-Blanco, et al., 2019; Dahl et al., 2022; Olivieri et al., 97 2019; Takahashi et al., 2015) and found significant negative associations of LC integrity with 98 age, age-related cognitive decline (Calarco et al., 2022; Dahl et al., 2019; Liu et al., 2020), 99 100 AD pathology, such as amyloid (Betts, Cardenas-Blanco, et al., 2019) and tau deposition 101 (Dahl et al., 2022; Jacobs et al., 2022), as well as with symptom severity in AD (Sun et al., 102 2020). Prokopiu et al. (2022) also found that lower novelty-related LC activity and LC to hippocampus and parahippocampus functional connectivity were associated with steeper 103 104 beta-amyloid-related cognitive decline. However, by contrast to its key importance to 105 cognitive function and etiological role in AD progression, there is very limited in vivo understanding of LC function disruptions during the early preclinical period. One exception 106 is a recent study of asymptomatic midlife adults, offspring of late-onset AD (O-LOAD) 107 108 patients. Del Cerro and colleagues (2020) found that the connectivity of the LC and 109 cerebellum was significantly associated with delayed recall performance, and reduced in O-LOAD individuals relative to those with no family history. 110 The hippocampus is one of the first regions to demonstrate atrophy during prodromal AD. 111 Neurofibrillary tangle (NFT) accumulation within the MTL, including the hippocampus, is 112 one of the first hallmark pathological manifestations. Recent studies in cognitively healthy 113 114 individuals at risk for late-life AD have found that AD risk impacts the hippocampus in 115 midlife (Dounavi et al., 2022; 2020; Kerchner et al., 2014; for a review see Vilor-Tejedor et al. 2021). Dounavi et al. (2020) reported that the volume of the molecular layer of the 116 hippocampus was reduced in cognitively healthy midlife individuals with APOE & genotype. 117 118 A more recent study (Dounavi et al., 2022) reported that the hippocampal fissure was

enlarged in middle-aged individuals with high CAIDE dementia risk scores.

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

Critically, the connectivity between the LC and the hippocampus is important for cognition. Animal studies have demonstrated that the co-release of noradrenaline and dopamine from LC terminals in the hippocampus is critical for successful spatial learning and memory (James et al., 2021; Kaufman et al., 2020; Kempadoo et al., 2016). Human studies in healthy older adults are consistent with these data and show that higher LC-Hippocampus functional connectivity is significantly associated with better memory (Jacobs et al., 2015). While accumulating evidence indicates that, individually, the LC and hippocampus are affected in midlife by risk of late-life AD, it remains unknown whether their interactions and joint role in cognition are affected by AD risk at this stage. Understanding of changes in the interaction of LC and the hippocampus during this preclinical period in at-risk populations has the potential to not only shed light on the brain mechanisms of incipient AD, but also to provide urgently needed early disease biomarkers. To address this gap, we investigated the impact of three risk factors for late-life AD (APOE ε4 genotype, FHD and CAIDE score) on cognition, and its relationship to the LC-Hippocampus functional connectivity. We tested a cohort of cognitively healthy middle-aged individuals, at baseline (N=210), and two-years follow-up (N=188) with a detailed neuropsychological battery not restricted to functions usually implicated in dementia detection in the older adults. Structural and functional Magnetic Resonance Imaging (MRI) data was obtained at both research visits. Our hypotheses were that risk of late-life AD would be significantly associated with cognition, the LC-Hippocampus functional connectivity, and their brain-behaviour relationship in middle-aged and cognitively healthy individuals. Furthermore, we expected that risk factors would be associated with longitudinal change of cognition and functional connectivity in this cohort.

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

2. Methods

2.1 Participants

PREVENT-Dementia is an ongoing longitudinal multi-site research programme across the UK and Ireland, seeking to identify early biomarkers of AD and elaborate on risk-mechanism interactions for neurodegenerative diseases decades before the cardinal symptoms of dementia emerge. Its protocol has been described in detail elsewhere (Ritchie & Ritchie, 2012). In the first PREVENT programme phase, participants were recruited at a single site, via the dementia register database held at the West London National Health Service (NHS) Trust, of the UK National Health Service, the Join Dementia Research website (https://www.joindementiaresearch.nihr.ac.uk/), through public presentations, social media and word of mouth. Procedures involving experiments on human participants were carried out in accord with the ethical standards of the Institutional Review Board of Imperial College London and in accord with the Helsinki Declaration of 1975. Approval for the study was granted by the NHS Research Ethics Committee London Camberwell St Giles. Consented participants were seen at the West London NHS Trust, where they underwent a range of clinical and cognitive assessments (Ritchie & Ritchie, 2012). The cohort comprised cognitively healthy volunteers aged 40-59 years. Here we examined baseline and follow-up data from the West London dataset. 210 individuals (62 male; 148 female) were tested at baseline, with 188 (89.5%) (55 male; 133 female) retained at two-years follow-up (Table 1). Mild cognitive impairment and dementia were ruled out based on detailed clinical assessment on each visit. At baseline, 2 participants were missing APOE \(\varepsilon 4 \) genotype information, and therefore the CAIDE scores. At follow-up, 6 participants were missing information for calculating their

CAIDE scores, including the 2 missing APOE status. For cognitive tests, 2 participants were missing data at baseline, and 12 at follow-up. Therefore, the cognitive analyses cohort was N=208/N=176 at baseline/follow-up. 193 out of 210 participants finished MRI scanning at baseline, and 169 out of 188 at follow-up. Furthermore, at baseline, 6 were excluded due to incidental findings, and 58 due to not meeting the criterion of > 150 frames with minimal head motion (for details see the following section *2.4 MRI data acquisition and processing*). At follow-up, 3 were excluded due to incidental findings and 73 due to < 150 frames with small head motion. The final fMRI analyses cohort was N = 128/N = 93 at baseline/follow-up. Please see the Supporting Information (SI) file and STable 1–2 and SFigure 1 for a description of any missing data.

2.2 Biological sex

While the dichotomies of sex and gender are no longer considered to be sharply discrete, in this study 'sex' was defined as an individual's natal or biological sex and was self-reported in the Brain Injury Screening Questionnaire. The cohort is 70% female, reflecting higher female interest in the study, possibly due to the known female predisposition to developing Alzheimer's disease.

2.3 Menopausal Status

Menopausal status was classified based on answers from a pregnancy and menstruation survey administered during the clinical assessments. Self-reported menopausal status was determined via answers to the question "Are you postmenopausal?". Those who answered 'Yes' were categorized as postmenopausal, and those who answered 'No' were categorized

190 as premenopausal. Of the 148/210 female participants, 55 (37.16%) were postmenopausal, 73 were premenopausal, and 20 who answered 'don't know'. Those who answered 'Don't Know' 191 192 (20/210) were excluded from further analyses. 193 194 2.4 Assessments 195 2.4.1 Risk factors 196 APOE ε4 Genotyping The process of APOE \(\varepsilon 4 \) allele identification is outlined in detail in Ritchie et al. (2017). In 197 198 brief, genomic DNA was isolated from blood samples and APOE genotyping was performed. 199 All members of the research and clinical teams were blind to the result of APOE genotyping. In this study, APOE ε4 risk is determined by ≥1 APOE ε4 allele. 75/210 carried ≥1 APOE ε4 200 201 allele (See Table 1). The prevalence of APOE ε2ε4 in our cohort is 0.02% (see SFigure 2). Investigation of the effect of the APOE ε2ε4 allele can be found in the SI. The exclusion of 202 APOE £2£4 carriers did not affect our main results and we therefore report results including 203 APOE ε2ε4 carriers. 204 Family History of Dementia 205 FHD was determined by a 'yes'/'no' question during clinical visits, where participants were 206 asked whether a parent had a diagnosis of dementia. Participants were asked to include the 207 208 dementia subtype if known, but answering 'yes' alone categorized a participant as FHD+. The answer 'no' likely captured both participants with no FHD, and participants for whom 209 FHD was unknown. In summary, participants were defined as FHD+ if at least one parent 210

was diagnosed with dementia. Cases where the FHD was unknown or partially known were

not recorded outside of the binary yes/no scoring. 103/210 were FHD+.

Cardiovascular Risk Factors, Aging, and Incidence of Dementia (CAIDE) score

CAIDE is a composite scale of estimated future dementia risk based on mid-life cardiovascular measures (Fayosse et al., 2020; Sindi et al., 2015). It takes into consideration the individual's age, sex, educational attainment, APOE &4 genotype, activity level, BMI, cholesterol and systolic blood pressure (Kivipelto et al., 2006) and is scored on a range of 0 – 18. A higher score indicates greater risk. The CAIDE dementia risk score was calculated at

2.4.2 Cognitive testing

baseline and follow-up.

Cognitive function was assessed at baseline and follow up with the COGNITO neuropsychological battery (Ritchie et al., 2014), designed to examine information processing across a wide range of cognitive functions in adults of all ages and not restricted to those functions usually implicated in dementia detection in the older adults. Tests are administered using a tactile screen to capture information processing time as well as response accuracy and require about 40 minutes to complete. The tests, by order of presentation, are: reaction time; reading; comprehension of phonemes, phrases, and syntax; focused and divided attention in both visual and auditory modalities; visual working memory (visual tracking with auditory interference); the Stroop test; immediate, delayed, and recognition trials for verbal recall (name list); delayed recognition of spatial stimuli (faces); visuospatial associative learning; visuospatial span; form perception; denomination of common objects; spatial reasoning; copying of meaningful and meaningless figures; verbal fluency with se-mantic and phonetic prompts; immediate recall of a narrative; immediate recall of a description of the relative

position of objects; vocabulary; implicit memory (recognition of new and previously learned 235 material). 236 237 The COGNITO tests are designed to test several aspects of cognition, including attention (task: visual attention), memory (tasks: narrative recall, description recall, implicit memory, 238 name-face association, working memory), language (tasks: phoneme comprehension, verbal 239 fluency) and visuospatial abilities (task: geometric figure recognition) (Ritchie et al., 2014). 240 Based on previous studies (Ritchie et al., 2021; Ritchie et al., 2018; Ritchie et al., 2017), 11 241 summary variables from the COGNITO battery capturing the above functions were used here 242 (for a list see STable 2). 243 Additionally, we used the Visual Short-Term Memory Binding task (VSTMBT) (Parra et al., 244 245 2010), a computer-based task that assesses visual short-term memory binding of single features, e.g., complex shape or color combinations, or feature conjunctions, e.g., shape and 246 color combinations. In the single feature condition, participants must identify whether the test 247 stimuli (three random 6-sided polygons) are the "same" as or "different" to the studied 248 stimuli in terms of shape (shape only) or color (color only). In the binding condition, 249 250 participants are required to correctly identify if both the shape and color of the test stimuli match studied stimuli. The two summary variables from the VSTMBT were the percentage of 251 correctly recognized items from the two conditions. 252 The Addenbrooke's Cognitive Assessment (ACE) III (Hsieh et al., 2013) was recorded at the 253 follow-up session, but not at the baseline. Analyses that focused on the subset of assessments 254 255 performed at both testing sessions did not include the ACE III.

2.5 Behavioural data analyses

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

Rotated principal component analysis

Rotated principal component analysis (rPCA), a dimensionality reduction technique, was adopted to cluster the above-mentioned 13 cognitive measures into related domains, reduce the number of multiple comparisons between the cognitive tests (Jolliffe & Cadima, 2016). This step maximized power to investigate the impact of risk factors on cognition and brain behaviour relationships. rPCA was conducted by using the psych package (Version 2.0.12) in R software (https://www.r-project.org/), following steps as: (1) *Preprocessing*. Listwise deletion procedure was used to deal with the missing data. Only participants who had no missing observations were kept (N = 208 at baseline; N = 176 at follow-up). (2) Component extraction and estimation. Principal component analysis was used to extract a smaller number of components that capture the most variance of the initial variables, representing the common cognitive processes underlying the 13 original cognitive measures. Scree plots and parallel analysis were used to determine the number of components (Horn, 1965). Parallel analysis generated a set of random correlation matrices (n = 500) by using the same number of variables (n = 13) and participants (N = 208 at baseline; N = 176 at follow-up). Then, we compared the solution to the random simulated normal data. The eigenvalues obtained from the actual matrix were compared with those obtained from the randomly generated matrices, and the estimated number of components was determined by selecting components with eigenvalues larger than 95th percentile of the randomly generated components' eigenvalues (See also SI). (3) Rotation. To examine the component structure among the initial variables, an orthogonal rotation method Varimax that constrains the components to be uncorrelated, was applied to the coefficient matrix (Kaiser, 1958). (4) Component coefficients. Component coefficients (r), reflecting the correlation between each of the components and each of the

original variables, were thus obtained. Given the sample size in this study (N > 200), component coefficients that were > 0.4 were deemed practically significant (Hair Jr. et al., 1998; Stevens, 1992). (5) *Component scores*. Component scores were calculated by a regression method where the regression weights were found from the inverse of the correlation matrix times the coefficient of each variable on the corresponding components. The coefficients were used to interpret the components, and component scores were used for the following analyses. (6) *Component similarities*. Components were extracted at the baseline and follow-up respectively. To test the similarity of components extracted at each study session, to be capturing similar cognitive functions, Tucker's congruence coefficient was calculated (Lorenzo-Seva & ten Berge, 2006). In subsequent analyses, we used these cognitive components, rather than the individual tests to measure the impact of risk factors on cognition and brain–behaviour relationships.

2.6 MRI data acquisition and processing

Imaging data were obtained as part of a multimodal examinations in a 3T Siemens Verio MRI scanner and with 32-channel head coil (https://preventdementia.co.uk/for-researchers/). Resting-state fMRI data were acquired with T2*-weighted echo-planar imaging (EPI) sequence. Total acquisition time for the resting state scan was 11 minutes 06 sec. The eyes were closed during rest. 330 volumes were acquired, and each volume contained 35 slices (interleaved acquisition), with slice thickness of 3 mm (repetition time (TR) = 2000ms, echo time (TE) = 30ms, flip angle (FA) = 80°, Field of View (FOV) = 192 × 192mm², voxel size = 3 mm³ isotropic). A 3D T1-weighted magnetization prepared rapid gradient-echo image (MPRAGE, 160 slices, voxel size = 1 mm³ isotropic, TR = 2300ms, TE = 2.98ms, FOV = 240 × 256mm², FA = 9°) was also acquired. All scans were repeated after approximately two

years on the same scanner using the same protocol.

Standard preprocessing procedures for resting-state fMRI data were performed with SPM12 (Statistical Parametric Mapping, https://www.fil.ion.ucl.ac.uk/spm/software/spm12/) and the AA pipeline software (Cusack et al., 2015) implemented in MATLAB R2019a (The MathWorks, United States). We discarded the first 5 volumes to allow for adaptation to the environment and equilibrium of the MR signal. In this pipeline (Figure 1a), we performed slice timing correction, motion correction, co-registration of functional and structural images, normalization into standard MNI space, and spatial smoothing. Spatial normalization was performed using SPM12's segment-and-normalize procedure, whereby the T1 structural was segmented into grey matter (GM), white matter (WM) and cerebrospinal fluid (CSF) and normalized to a segmented MNI-152 template. These normalization parameters were then applied to all EPIs. The quality of spatial normalization was visually inspected for each participant and no participants showed a normalization failure. The data were then smoothed with a Gaussian kernel of 6mm full width at half maximum. We also applied a general linear model (GLM) that includes 24 head motion parameters – 6 rigid-body motion parameters, the first-order temporal derivative of them and quadratic terms of the original parameters and their derivatives (Satterthwaite et al., 2013) – and averaged signals from WM and CSF as covariates to reduce any residual effects of head motion and physiological noise. Lastly, all preprocessed data were temporally band-pass filtered (0.01-0.08 Hz) to remove lowfrequency drift and high-frequency physiological noise (Salvador et al., 2008; Zuo et al., 2010).

Figure 1 about here please

327

328

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

Because brain signal from the brainstem is susceptible to head movements, we performed

careful measurement and control for head motion, by using a "scrubbing" procedure (Power et al., 2012). We first calculated an index of data quality called framewise displacement (FD) (for details please also see SI), which measures how much the head has changed position from one frame to the next, and can therefore be used to flag frames of suspect quality based on the threshold of FD > 0.2 mm, in order to remove motion-contaminated data before calculating functional connectivity (Power et al., 2012). A participant inclusion criterion was then set to ensure a minimum amount of data remained after scrubbing, i.e., at least 150 frames (~5 min) of data (Figure 1b), based on wide agreement that ~5 min of fMRI data is an adequate for resting state functional connectivity analyses (Van Dijk et al., 2010). After completing the scrubbing procedure, the included participants had $74.0\% \pm 14.8\%$ frames at baseline and $73.5\% \pm 15.0\%$ at follow-up. Finally, we calculated the mean FD across the remaining good frames, which was further regressed in the group-level analyses (Power et al., 2014). The mean FD was 0.12 ± 0.01 at baseline and 0.12 ± 0.01 at follow-up. There were no significant associations of mean FD with any of the risk factors or the LC-Hippocampus functional connectivity (See SI). The fMRI-based exclusion criteria are also illustrated in Figure 1b. To ensure high accuracy in the localization of the LC, we created a binary LC mask of 3 mm³ isotropic voxel size that closely matched with two previously published LC maps (Dahl et al., 2022; Ye et al., 2021), which were developed in the standard MNI space with 0.5 mm³ isotropic voxel size. In order to achieve the highest precision, instead of resampling the LC maps to match the fMRI data's voxel size (3 mm³) using automatic scripts, we manually drew a 3 mm³ binary LC mask in MNI space with reference to the two LC maps. This is because the automatic scripts did not perform well when resampling the two referenced LC maps from high to low resolution due to the small size of these two LC regions of interest. The FSL software (https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/) was used for the drawing process, by

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

following the same procedure as in Plini et al. (2021). The two 0.5 mm³ LC maps were superimposed on a blank 3 mm³ template in standard MNI space. The drawing process was then performed on this blank template by adding or removing voxels to include different anatomical locations of the two referenced maps without crossing the dorsal and medial raphe nuclei, as defined by Beliveau et al. (2015), and the cerebellar vermis white matter. At the end of this procedure, the new bilateral LC mask was symmetrical and consisted of 16 3 mm³ voxels, was anatomically inclusive of the 7T LC map (Ye et al., 2021) and the LC meta-map (Dahl et al., 2022), and did not encroach on other pontine nuclei (Figure 1c). In an independent study (Dounavi et al., 2022), T1-weighted MPRAGE of this cohort were corrected for field inhomogeneities using the Advanced Normalisation Toolbox (ANTs) N4 algorithm (Tustison et al., 2010) and processed using Freesurfer version 7.1.0 (Desikan et al., 2006). The recon-all pipeline was run for every participant with standard settings. The brain masks and surfaces were inspected following recon-all and manual corrections were applied: (a) in the form of erosion of non-brain voxels from the brain mask or non-WM voxels from the WM mask, (b) in the form of filling of areas where the brain was not correctly identified, or (c) with the addition of control points in cases where white matter was not successfully identified. Following these steps, we generated individual masks of the hippocampus, which were then applied to the fMRI data to extract the time course of the region (Figure 1d). The functional connectivity (FC) between the LC and the hippocampus was derived using Pearson's correlation coefficient (r) on the denoised time courses (Figure 1d). The r value was then transformed to z value using Fisher's r-to-z transformation to improve normality for further statistical analysis. To avoid the formation of artificial anti-correlations (Anderson et al., 2011; Murphy et al., 2009), we did not perform global signal regression. In addition, volume of the hippocampus (HCV) was derived for each participant at baseline and followup as a covariate for functional connectivity analyses.

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

2.7 Statistical approach

All statistical analyses were performed in R software. The normality of the data was assessed by combining the visualization of a quantile-quantile plot and the Shapiro–Wilk test. Demographic and clinical information of the study cohort was analyzed across risk groups using chi-square (χ^2 tests) for categorical variables and Mann Whitney U tests for continuous (discrete) variables, given that they were not normally distributed in this cohort. To investigate the cross-sectional effects of risk factors (APOE ε4 genotype and FHD) on cognition and brain function we used multiple linear regression models. Age, years of education were included as covariates for cognition models, and mean FD values and HCV were additionally included as a covariate for models of functional connectivity. The effects of APOE ε4 and CAIDE risk factors were modelled independently to avoid modelling the variance associated with APOE genotype in the same model twice. Spearman correlation analyses were used to assess the associations between cognition and CAIDE score given the non-normal distribution of CAIDE in our cohort. For the CAIDE model, we did not control for the age, years of education, as they are considered in the CAIDE scoring. For any observed risk effect on cross-sectional changes of cognition or functional connectivity, interactions between risk factors and the LC-Hippocampus functional connectivity on cognition were assessed through multiple regression models with the cognition as the dependent variable, the LC-Hippocampus FC, risk factor and the risk × FC interaction as independent variables, and age, years of education, and HCV as covariates. Age, sex, and years of education were not included as covariates when CAIDE was included in the models, because CAIDE includes these variables in its scoring. Biological sex was investigated as variables of interest in these analyses especially given the higher proportion of females in the sample. If a sex effect was found, any interactions with risk factors, and the effect of menopausal status was also considered because the cohort age range [52+/- 11 years] cover menopause transition in females, which is known to affect cardiovascular health and cognition.

For any observed interactions, we plotted the regression of LC–Hippocampus FC on cognitive performance for each level/value of the risk factor, to interpret the effect (Aiken & West, 1991). For the continuous variable CAIDE, we applied the Johnson-Neyman technique (Bauer & Curran, 2005; Johnson & Fay, 1950), to test the significance of the slopes of the simple regression lines on each value of the CAIDE scores (ranged from 0-18). This examined the CAIDE interval at which the brain-cognition coupling was significant. All continuous variables in the interaction terms, i.e., CAIDE and FC, in the moderation models were mean-centered (subtracting each the mean from each value) to avoid multicollinearity. Longitudinal changes were first assessed using random intercept linear mixed effect models (Cnaan et al., 1997) for the entire cohort. Specifically, we evaluated longitudinal changes of cognition (Eq. 1) and functional connectivity (Eq. 2) over two years.

$$Cognition_{jt} = \beta_0 + \beta_1 (Time_{jt}) + u_j + r_{jt}$$
 (1)

$$FC_{jt} = \beta_0 + \beta_1 (Mean FD_{jt}) + \beta_2 (HCV_{jt}) + \beta_3 (Time_{jt}) + u_j + r_{jt}$$
 (2)

For each participant *j*, measures of cognition/functional connectivity at each time point *t* served as dependent variables, and the timepoint (a categorical variable with two levels: baseline and follow-up) served as the independent variable. Mean FD and HCV at each time

point t were included as covariates for the functional connectivity model only (Eq. 2). u_j represented individual variability in the intercepts, i.e., a random effect.

The presence of a longitudinal change for the entire cohort was then further assessed to determine whether it varied in relation to the risk factors, as assessed by the interaction term $(Risk_j \times Time_{jt})$. Age at baseline, sex, and years of education, were included as covariates for the cognition models that did not include CAIDE (Eq. 3). Mean FD and HCV were additionally included as covariates for the functional connectivity models (Eq. 4).

$$\begin{split} Cognition_{jt} \\ &= \beta_0 + \beta_1 (Age_j) + \beta_2 (Sex_j) + \beta_3 (Education_j) + \beta_4 (Risk_j) \\ &+ \beta_5 (Time_{jt}) + \beta_6 (Risk_j \times Time_{jt}) + u_j \\ &+ r_{jt} \end{split} \tag{3}$$

$$\begin{split} FC_{jt} &= \beta_0 + \beta_1 (Age_j) + \beta_2 (Sex_j) + \beta_3 (Education_j) + \beta_4 (Mean \, FD_{jt}) \\ &+ \beta_5 (HCV_{jt}) + \beta_6 (Risk_j) + \beta_7 (Time_{jt}) + \beta_8 (Risk_j \times Time_{jt}) \\ &+ u_j + r_{jt} \end{split}$$

Lastly, the presence of risk related longitudinal change in cognition or FC was further investigated to determine whether risk factors influence brain-cognition association longitudinally. Multiple linear regression modelling was used with a percentage change score [i.e. (follow-up - baseline)/baseline] in cognition as the dependent variable, and a risk factor, a percentage change in FC, and their interaction as independent variables. Baseline age, sex, years of education and changes in HCV were included as covariates. Only changes in HCV were controlled for when assessing CAIDE in the models.

Although the baseline and follow-up datasets were not independent, we considered each in its own right, in addition to testing for longitudinal changes in cognition and functional connectivity. During the two years, a proportion of our participants may have sustained brain health that are yet subthreshold to clinical manifestations, or that may not yet manifest as longitudinal change. Statistical results were corrected for multiple comparisons.

3. Results

3.1 Demographic characteristics

Demographic characteristics of the cohort at baseline and follow-up, stratified by APOE $\varepsilon 4$ genotype and FHD, are shown in Table 1. There were no significant differences in age, sex or years of education between the groups. APOE $\varepsilon 4$ allele genotype was more frequently found in the FHD+ than FHD- group at baseline (p=0.01) and follow up (p=0.02). CAIDE scores (including APOE status) were significantly higher for the FHD+ than FHD- group at baseline (p=0.03) and follow up (p=0.003). Naturally, CAIDE scores including APOE status were significantly higher for the APOE $\varepsilon 4$ + than APOE $\varepsilon 4$ - group at baseline (p=0.0003) and follow up (p=0.0002), but when APOE status was excluded the CAIDE scores did not differ between the two groups (Table 1).

Table 1 about here please

3.2 Cognitive domains

We first performed parallel analyses and plotted scree plots to determine the number of components that best represented the original 13 cognitive measures at baseline and follow-up, separately. Results showed that eigenvalues of the first three components were larger than 95^{th} percentile of the randomly generated eigenvalues at both study timepoints (Figure 2a), which indicated that the three component (C) solution best represented the data. The three components were then rotated to be uncorrelated with each other, and could cumulatively explain a total of 41% percentage of the variance, at baseline (C1 = 17%; C2 = 13%; C3 = 12%) and 40% at follow-up (C1 = 17%; C2 = 12%; C3 = 11%) (Figure 2b).

Figure 2 about here please

The loading values (weights) of the cognitive measures (Figure 3a) reflect the relationships between the original measures and the corresponding components. Measures with higher coefficients were more closely related to the components. Based on the cognitive functions that the highest loading measures tapped into (for details see STable 2), in subsequent analyses we refer to: C1 as 'episodic and relational memory'; C2 as 'working and short-term (single-feature) memory'; C3 as 'verbal, visuospatial functions, and short-term (conjunctive) memory.'

The highest loading measures and their coefficients for each component differed for the cross-sectional and longitudinal data. For example, verbal functions, visuospatial functions, and short-term (conjunctive) memory loaded most strongly on the third component at baseline. At follow-up, implicit memory rather than short-term memory loaded on the third

483 component (Figure 3a). We found high similarity between baseline and follow-up for C1 (ϕ = 0.90), and C3 (ϕ = 0.76), with relatively low similarly for C2 (ϕ = 0.40) (Figure 3b). 484 485 Figure 3 about here please 486 487 488 3.3 Impact of risk factors on cognition Episodic and relational memory 489 490 At baseline, the multiple linear regression model showed a significant positive association between episodic and relational memory with APOE $\varepsilon 4$ allele [β (SE) = 0.28 (0.13), p < 0.05], 491 independently of sex, age and years of education (Table 2). At follow-up, there was a trend 492 effect of APOE $\varepsilon 4$ allele [β (SE) = 0.27 (0.15), p = 0.07]. Education was significantly 493 positively associated with this domain at baseline $[\beta (SE) = 0.09 (0.02), p < 0.0001]$ and 494 follow-up [β (SE) = 0.10 (0.02), p < 0.0001]. Sex did not interact with the APOE $\varepsilon 4$ allele in 495 its effect on episodic and relational memory. The baseline effect of APEO &4 on episodic and 496 relational memory became a trend effect (SFigure 6) when corrected for multiple 497 498 comparisons and needs to be interpreted with caution. 499 *Table 2 about here please* 500 501 502 The multiple linear regression models showed no associations between FHD and episodic and

relational memory, at either timepoint. Spearman correlation analyses showed no significant

associations between the CAIDE score and episodic and relational memory, at either timepoints.

Working and short-term (single-feature) memory

We found no associations between APOE $\varepsilon 4$, FHD, or CAIDE score with working and short-term (single-feature) memory. Education was significantly positively associated with performance in this domain, independently of the other factors, at baseline [β (SE) = 0.04 (0.02), p < 0.05] only.

Verbal, visuospatial functions, and short-term (conjunctive) memory

We found no associations between APOE $\epsilon 4$ or FHD with verbal, visuospatial functions, and short-term (conjunctive) memory. Age was significantly negatively associated with performance in this domain, independently of the other factors, at baseline [β (SE) = -0.04 (0.01), p < 0.005]. The CAIDE score was significantly associated with verbal, visuospatial functions, and short-term (conjunctive) memory at baseline (rho = -0.17, p < 0.05, Figure 4a), and at follow-up (rho = -0.20, p < 0.01, Figure 4b). Higher CAIDE scores were significantly associated with worse performance.

As the cohort was 70% female, and the age range (40-60 years) covers the menopausal transition, which is known to affect cardiovascular health and cognition, the effect menopausal status on the relationship between CAIDE score and cognition was investigated. The CAIDE score includes sex in its calculation, with 'male' scored higher. Therefore, the effect of sex on relationship between the CAIDE risk and cognition not investigated.

However, there was a significant interaction between CAIDE and menopause on verbal, visuospatial functions and short-term (conjunctive) memory at baseline [β (SE) = -0.20 (0.09), p < 0.05] and at follow-up [β (SE) = -0.29 (0.13), p < 0.05] (Table 3). Increasing CAIDE was associated with worse cognitive performance in post-menopausal females, but not in premenopausal counterparts. This effect was controlled for age effects, as post-menopausal women were overall older than pre-menopausal counterparts (post-: median \pm IQR: 49.0 \pm 8.0 vs pre-: 56.0 ± 4.0).

Figure 4 about here please

Table 3 about here please

There was no significant longitudinal change of cognition in any of the three cognitive domains over two years.

3.4 Longitudinal change of LC-Hippocampus functional connectivity

We found a significant negative association between time and the LC–Hippocampus functional connectivity [β (SE) = -0.06 (0.03), p < 0.05], after controlling for mean FD and HCV (Table 4), indicating reduced connectivity at two-years follow-up relative to baseline across the whole cohort (Figure 5). Sex did not interact with time in affecting the LC–Hippocampus functional connectivity.

547	Table 4 about here please
548	Figure 5 about here please
549	
550	3.5 Impact of risk factors on the LC-Hippocampus functional connectivity and brain-
551	behavior relationships
552	There were no significant associations of <i>APOE</i> ε4, FHD, or CAIDE scores with the LC–
553	Hippocampus functional connectivity, cross-sectionally, or longitudinally.
554	At baseline, none of the risk factors moderated the relationship between functional
555	connectivity and cognition. At follow-up, the multiple linear regression model with CAIDE
556	LC-Hippocampus FC, and CAIDE × LC-Hippocampus FC interaction term as independent
557	variables, and verbal, visuospatial functions, and short-term (conjunctive) memory as the
558	dependent variable showed a significant negative association between the interaction term
559	(CAIDE \times LC-Hippocampus FC) and verbal, visuospatial functions, and short-term
560	(conjunctive) memory [β (SE) = -0.54 (0.22), $p < 0.05$] (Table 5). In individuals with
561	low/high CAIDE scores, higher functional connectivity was associated with better/worse
562	cognition (Figure 6). Menopausal status did not have an effect on this relationship.
563	
564	Table 5 about here please
565	Figure 6 about here please

4. Discussion

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

It is well acknowledged that AD neuropathology starts decades before clinical manifestations, but the brain mechanism of sporadic AD and its interactions with risk factors in midlife remain unclear. We examined whether risk factors for late-life AD were associated with cognition, functional connectivity between two of the key structures in AD neurogenesis – the LC and the hippocampus – and the brain-behaviour relationship between them, crosssectionally and longitudinally in a cohort of middle-aged, cognitively healthy individuals. We found that the CAIDE scores were significantly negatively associated with verbal, visuospatial functions and short-term (conjunctive) memory functions in this cohort. Menopausal status significantly interacted with CAIDE on cognition, independently of age. Post-menopausal women showed worse cognition with increasing CAIDE scores. Furthermore, the CAIDE score significantly moderated the relationship between cognition and the LC-Hippocampus functional connectivity. These results shed light on some of the earliest brain-behaviour alterations due to risk of future AD in cognitively healthy individuals cross-sectionally. We found a significant longitudinal effect of age on the LC-Hippocampus FC, with no longitudinal effects of risk factors on cognition, the LC-Hippocampus functional connectivity, or their relationship. The lack of prominent longitudinal effects of risk factors is likely due to the narrow follow-up time window (2 years) in this relatively young midlife cohort, an estimated 23 years from dementia onset (Dounavi et al., 2022). Higher CAIDE scores were significantly associated with worse cognition in verbal, visuospatial functions and short-term (conjunctive) memory, both at baseline and at followup. Previous studies from the same cohort have reported significant negative associations of CAIDE scores with visuospatial functions and navigational abilities (Ritchie et al., 2018;

Ritchie et al., 2017). We did not observe a longitudinal effect, similarly to other studies of this cohort (Dounavi et al., 2021; Low et al., 2021), possibly due to the relatively young age and the short follow-up window (Ritchie et al., 2018; Ritchie et al., 2017). Post-menopausal women only showed worse cognition with increasing CAIDE scores. The menopausal transition is a middle-age phenomenon with knock-on effects on endocrinal (Weber et al., 2014), cardiovascular (Yanes & Reckelhoff, 2011), and metabolic (Anagnostis et al., 2019; Rahman et al., 2020) systems that significantly impact brain health (Mosconi et al., 2018; Udeh-Momoh & Watermeyer, 2021). The rapid decline in estrogen, a key hormonal change during menopausal transition, is considered to increase vulnerability to AD in females (Scheyer et al., 2018; Udeh-Momoh & Watermeyer, 2021). Post-menopause has been associated with increased levels of Aβ, elevated tau burden, reduced white matter integrity, and glucose hypometabolism (Buckley et al., 2022; Mosconi et al., 2017; Rahman et al., 2020). Our results suggest that post-menopausal status exacerbates the effect of cardiovascular AD risk on cognition, rendering females particularly vulnerable to cognitive decline, and conversely, that addressing cardiovascular risk early may protect cognition in post-menopausal females. We found that the LC–Hippocampus functional connectivity decreased significantly over two years. The previously reported effect of age on the LC-Hippocampus functional connectivity is inconsistent, possibly due to different age ranges used in different studies. In healthy young and early midlife adults (age range: 18–49), LC functional connectivity to the parahippocampus was negatively associated with age (Zhang et al., 2016). However, another study of a lifespan adult cohort (age range: 19-74) found no association of age with the LC-Hippocampus/parahippocampus functional connectivity (Jacobs et al., 2018). In a broader age-range lifespan cohort (age range: 8–83), Song et al. (2021) found a positive quadratic age effect (u-shape) on LC-parahippocampal functional connectivity, suggesting an increased

591

592

593

594

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

interaction between these two regions in children and older adults. Therefore, our finding advances the literature, by showing that the LC-Hippocampus functional connectivity decreases with age in midlife. Another key question here was whether risk factors impact the LC-Hippocampus functional connectivity and its role in cognition. We found that the CAIDE score significantly moderated this brain-behaviour relationship. Stronger LC-Hippocampus functional connectivity was associated with better cognition for individuals with low CAIDE score, and with worse cognition in those with high scores. The presence of this effect in the follow-up, but not the baseline dataset is likely due to the older age of the cohort at follow-up. Previous studies in older adults found that stronger LC-Hippocampus functional connectivity was significantly associated with better cognition (Jacobs et al., 2015). Our finding advances understanding by suggesting that the connectivity between the LC and the hippocampus supports cognition in midlife individuals who have low dementia risk scores. Furthermore, our result suggests that, in the presence of high dementia risk scores, the LChippocampus functional connectivity is negatively associated with cognition. Previous studies have shown that a high CAIDE score, of more than 12, confers a probability of 16.4% for future dementia, with scores above 8 associated with a probability of more than 4% (Kivipelto et al., 2006). In a study of middle-aged individuals with a mean age of 46, participants with a score greater than 8 had a 29% 40-year risk for dementia (Exalto et al., 2014). These findings suggest that a proportion of our study's participants, who have a CAIDE score greater than 8, may develop dementia. Assuming that AD pathology is underway in this proportion of individuals with high dementia risk scores, our result lends indirect support to the hypothesis that LC hyperactivity and associated hyper-connectivity initiates the spread of pathological tau to MTL (Weinshenker, 2018), and particularly the

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

638

hippocampus, in the early stages of AD. The ensuing neurodegeneration may explain why hyper-connectivity is associated with worse performance in individuals with high dementia risk scores. However, the neural mechanisms of AD neuropathology genesis remain contested. Given the lack of tau biomarker status for this cohort, future studies that integrate tau and functional connectivity information are required to further test the proposal that tau spread from the LC to the hippocampus via functional connectivity pathways. Furthermore, future studies from the continuing longitudinal follow-up of this cohort will validate the role and nature of these changes in individuals who display AD neuropathology as they get older. We did not observe any significant effects of risk factors for AD on the LC–Hippocampus connectivity. Previous studies on MCI and AD patients have shown disrupted/lower LC-Hippocampus functional connectivity (MCI: Jacobs et al., 2015; age: 65.1 ± 4.5 years) (Liebe et al., 2022; age: 73.3 ± 7.5 years; AD patients: Zhao et al., 2017), compared to age-matched healthy controls. One previous study investigating cognitively healthy middle-aged adults with familial risk for late-onset dementia found no effect of AD risk on the LC-Hippocampus connectivity (Del Cerro et al., 2020; age: 50.4 ± 8.3 years). Consistent with this previous study, our null result suggests that AD risk factors are not associated with the LC-Hippocampus connectivity in middle-aged individuals who are cognitively healthy. We also observed a trend effect association of APOE ε4 allele with better cognition in episodic and relational memory, independently of sex, age and years of education. Previous research has suggested that the APOE & allele, although being associated with poorer health outcomes in old age, may have a positive effect earlier in life. For example, it has been associated with better visual perception (Ritchie et al., 2017), short-term memory (Zokaei et al., 2020), higher intelligence quotient and a more economic use of memory-related neural resources in young healthy humans (Mondadori et al., 2007). These results may also be

640

641

642

643

644

645

646

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

interpreted based on the antagonistic pleiotropy hypothesis of aging (William 1957), which proposes that deleterious genes, such as the APOE \(\epsilon 4 \) gene allele, have survived through evolution because they might confer an advantage early in life, when humans are reproductively fit. These results suggest an impact of genetic risk for late-life sporadic AD in cognitively healthy midlife individuals on episodic memory, the hallmark cognitive dysfunction in early clinical dementia and mild cognitive impairment (Irish et al., 2011). Thus, they highlight the importance of studying the origins of Alzheimer's disease in this age group, over 20 years before typical symptom onset. Given the weak effect however, further replication in larger samples is warranted to validate these findings. In summary, the PREVENT-Dementia study is a longitudinal multi-site study targeted at identifying early biomarkers for Alzheimer's disease. In this study, we provide early proof that the recruited cohort, with a mean age of 52/54 years at baseline/follow-up, showed a disrupted role of LC-Hippocampus connectivity in cognition with an increasing CAIDE score. APOE £4 genotype or FHD did not moderate this brain-behaviour relationship. In other words, when the genetic risk alone was considered, no alterations of brain-behaviour relationships were found. It was only when a risk score (CAIDE) incorporating genetic risk in combination with lifestyle factors, sex and age was considered, that such alterations were unravelled. Taken together, these results provide strong evidence that brain-behaviour alterations in individuals with higher CAIDE scores may be driven by lifestyle risk factors included in this dementia risk score (i.e., blood pressure, cholesterol, physical activity, body mass index, years of education). Hence, these findings highlight the importance of considering modifiable risk factors when stratifying risk populations, or potentially designing randomized control trials. Our previous work in this cohort has demonstrated that modifiable lifestyle factors affect cognition,

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

particularly in individuals at high risk for late-life AD (Heneghan et al., 2023) and lends further support to this idea. In fact, a randomized multi-domain control trial in the FINGER population (mean age 70 years old) over two years demonstrated that the applied lifestyle and vascular interventions had an impact on cognition (Stephen et al., 2019). Hence, further investigation of pathological alterations in relation to modifiable risk factors in midlife is warranted to unveil the sequalae of brain and behaviour alterations leading to dementia.

4.1 Strengths and Limitations

Strengths of this study are its large well-characterized middle-aged cohort. State-of-the-art analysis methods and a thorough quality control protocol were applied to the acquired data. Limitations include the short follow-up window of the study and the absence of further well-established preclinical biomarkers, such as amyloid and tau status. The interpretability of the effect of risk on individual cognitive abilities requires individuation in future studies. Standard MR functional sequences provide limited resolution of the diminutive LC volume, and thus, higher resolution imaging that specifically targets the LC region is required to further validate these findings in future studies. Like most other studies involving mid-life and older people in the UK and Ireland, the study population is mainly (95%) of white Caucasian ethnicity, not dissimilar to the historic ethnic mix of older people in the five study sites in this region. The inclusion of other ethnicities and representation from different world regions in future studies will ensure true diversity and findings that can be generalized to different populations across the globe.

708	Acknowledgement
709	F.D. was funded by the Provost PhD Award Scheme from Trinity College Dublin, to
710	L.N. L.N. was also funded by a L'Oréal-UNESCO for Women In Science International
711	Rising Talent Award, the Welcome Trust Institutional Strategic Support grant, and the Global
712	Brain Health Institute Project Grant.
713	The PREVENT-Dementia study is supported by the UK Alzheimer's Society (grant numbers
714	178 and 264), the US Alzheimer's Association (grant number TriBEKa-17-519007) and
715	philanthropic donations.
716	We thank all PREVENT-Dementia participants for their enthusiastic participation in this
717	study. We also thank the Research Delivery service at West London NHS Trust and the
718	Wolfson Clinical Imaging Facility at Imperial College London for their support in running
719	the study.
720	
721	Conflict of interest
722	The authors declare no conflict of interest.
723	

- 724 References:
- Aiken, L. S., & West, S. G. (1991). *Multiple regression: Testing and interpreting interactions*.
- Newbury Park, CA: Sage.
- Amaral, D. G., & Sinnamon, H. M. (1977). The locus coeruleus: neurobiology of a central
- noradrenergic nucleus. *Prog Neurobiol*, *9*(3), 147-196. https://doi.org/10.1016/0301-
- 729 <u>0082(77)90016-8</u>
- Anagnostis, P., Christou, K., Artzouchaltzi, A. M., Gkekas, N. K., Kosmidou, N., Siolos, P.,
- Paschou, S. A., Potoupnis, M., Kenanidis, E., Tsiridis, E., Lambrinoudaki, I.,
- Stevenson, J. C., & Goulis, D. G. (2019). Early menopause and premature ovarian
- insufficiency are associated with increased risk of type 2 diabetes: a systematic review
- and meta-analysis. Eur J Endocrinol, 180(1), 41-50. https://doi.org/10.1530/eje-18-
- 735 <u>0602</u>
- Anderson, J. S., Druzgal, T. J., Lopez-Larson, M., Jeong, E. K., Desai, K., & Yurgelun-Todd,
- D. (2011). Network anticorrelations, global regression, and phase-shifted soft tissue
- correction. *Hum Brain Mapp*, 32(6), 919-934. https://doi.org/10.1002/hbm.21079
- Arnsten, A. F. T., Datta, D., Del Tredici, K., & Braak, H. (2021). Hypothesis: Tau pathology
- is an initiating factor in sporadic Alzheimer's disease. *Alzheimer's & Dementia*, 17(1),
- 741 115-124. https://doi.org/10.1002/alz.12192
- Aston-Jones, G., & Cohen, J. D. (2005). An integrative theory of locus coeruleus-
- norepinephrine function: adaptive gain and optimal performance. *Annu Rev*
- 744 *Neurosci*, 28, 403-450. https://doi.org/10.1146/annurev.neuro.28.061604.135709
- Barnes, D. E., Beiser, A. S., Lee, A., Langa, K. M., Koyama, A., Preis, S. R., Neuhaus, J.,

- 746 McCammon, R. J., Yaffe, K., Seshadri, S., Haan, M. N., & Weir, D. R. (2014).
- Development and validation of a brief dementia screening indicator for primary care.
- 748 Alzheimers Dement, 10(6), 656-665.e651. https://doi.org/10.1016/j.jalz.2013.11.006
- Barnes, D. E., & Yaffe, K. (2011). The projected effect of risk factor reduction on
- 750 Alzheimer's disease prevalence. *Lancet Neurol*, 10(9), 819-828.
- 751 https://doi.org/10.1016/s1474-4422(11)70072-2
- Bauer, D. J., & Curran, P. J. (2005). Probing Interactions in Fixed and Multilevel Regression:
- 753 Inferential and Graphical Techniques. Multivariate behavioral research, 40(3), 373–
- 754 400. https://doi.org/10.1207/s15327906mbr4003 5
- Beliveau, V., Svarer, C., Frokjaer, V. G., Knudsen, G. M., Greve, D. N., & Fisher, P. M.
- 756 (2015). Functional connectivity of the dorsal and median raphe nuclei at rest.
- 757 *NeuroImage*, 116, 187–195. https://doi.org/10.1016/j.neuroimage.2015.04.065
- Betts, M. J., Cardenas-Blanco, A., Kanowski, M., Jessen, F., & Düzel, E. (2017). In vivo
- MRI assessment of the human locus coeruleus along its rostrocaudal extent in young
- and older adults. *NeuroImage*, 163, 150-159.
- 761 https://doi.org/10.1016/j.neuroimage.2017.09.042
- Betts, M. J., Cardenas-Blanco, A., Kanowski, M., Spottke, A., Teipel, S. J., Kilimann, I.,
- Jessen, F., & Düzel, E. (2019). Locus coeruleus MRI contrast is reduced in
- Alzheimer's disease dementia and correlates with CSF Aβ levels. *Alzheimers Dement*
- 765 (*Amst*), 11, 281-285. https://doi.org/10.1016/j.dadm.2019.02.001
- Betts, M. J., Kirilina, E., Otaduy, M. C. G., Ivanov, D., Acosta-Cabronero, J., Callaghan, M.
- F., Lambert, C., Cardenas-Blanco, A., Pine, K., Passamonti, L., Loane, C., Keuken, M.
- C., Trujillo, P., Lüsebrink, F., Mattern, H., Liu, K. Y., Priovoulos, N., Fliessbach, K.,

- Dahl, M. J., Maaß, A., Madelung, C. F., Meder, D., Ehrenberg, A. J., Speck, O.,
- Weiskopf, N., Dolan, R., Inglis, B., Tosun, D., Morawski, M., Zucca, F. A., Siebner,
- H. R., Mather, M., Uludag, K., Heinsen, H., Poser, B. A., Howard, R., Zecca, L.,
- Rowe, J. B., Grinberg, L. T., Jacobs, H. I. L., Düzel, E., & Hämmerer, D. (2019).
- Locus coeruleus imaging as a biomarker for noradrenergic dysfunction in
- neurodegenerative diseases. *Brain*, 142(9), 2558-2571.
- 775 https://doi.org/10.1093/brain/awz193
- Berti, V., Mosconi, L., Glodzik, L., Li, Y., Murray, J., De Santi, S., Pupi, A., Tsui, W., & De
- Leon, M. J. (2011). Structural brain changes in normal individuals with a maternal
- history of Alzheimer's. *Neurobiol Aging*, *32*(12), 2325.e2317-2326.
- https://doi.org/10.1016/j.neurobiolaging.2011.01.001
- 780 Braak, H., & Braak, E. (1991). Neuropathological stageing of Alzheimer-related changes.
- 781 *Acta Neuropathol*, 82(4), 239-259. https://doi.org/10.1007/bf00308809
- 782 Braak, H., & Del Tredici, K. (2015). Neuroanatomy and pathology of sporadic Alzheimer's
- 783 disease. Adv Anat Embryol Cell Biol, 215, 1-162.
- Braak, H., Thal, D. R., Ghebremedhin, E., & Del Tredici, K. (2011). Stages of the pathologic
- process in Alzheimer disease: age categories from 1 to 100 years. *J Neuropathol Exp*
- 786 Neurol, 70(11), 960-969. https://doi.org/10.1097/NEN.0b013e318232a379
- Buckley, R. F., O'Donnell, A., McGrath, E. R., Jacobs, H. I. L., Lois, C., Satizabal, C. L.,
- Ghosh, S., Rubinstein, Z. B., Murabito, J. M., Sperling, R. A., Johnson, K. A.,
- Seshadri, S., & Beiser, A. S. (2022). Menopause Status Moderates Sex Differences in
- Tau Burden: A Framingham PET Study. *Ann Neurol*, 92(1), 11-22.
- 791 <u>https://doi.org/10.1002/ana.26382</u>

792 Cacciaglia, R., Operto, G., Falcón, C., de Echavarri-Gómez, J. M. G., Sánchez-Benavides, G., Brugulat-Serrat, A., Milà-Alomà, M., Blennow, K., Zetterberg, H., Molinuevo, J. L., 793 Suárez-Calvet, M., & Gispert, J. D. (2022). Genotypic effects of APOE-\(\varepsilon\)4 on resting-794 state connectivity in cognitively intact individuals support functional brain 795 compensation. Cereb Cortex. https://doi.org/10.1093/cercor/bhac239 796 797 Calarco, N., Cassidy, C. M., Selby, B., Hawco, C., Voineskos, A. N., Diniz, B. S., & 798 Nikolova, Y. S. (2022). Associations between locus coeruleus integrity and diagnosis, age, and cognitive performance in older adults with and without late-life depression: 799 An exploratory study. *NeuroImage: Clinical*, 36, 103182. 800 https://doi.org/10.1016/j.nicl.2022.103182 801 802 Cnaan, A., Laird, N. M., & Slasor, P. (1997). Using the general linear mixed model to analyse unbalanced repeated measures and longitudinal data. Statistics in medicine, 803 16(20), 2349–2380. https://doi.org/10.1002/(sici)1097-804 0258(19971030)16:20<2349::aid-sim667>3.0.co;2-e 805 Cusack, R., Vicente-Grabovetsky, A., Mitchell, D. J., Wild, C. J., Auer, T., Linke, A. C., & 806 Peelle, J. E. (2015). Automatic analysis (aa): efficient neuroimaging workflows and 807 parallel processing using Matlab and XML. Front Neuroinform, 8, 90. 808 https://doi.org/10.3389/fninf.2014.00090 809 Clewett, D. V., Lee, T. H., Greening, S., Ponzio, A., Margalit, E., & Mather, M. (2016). 810 Neuromelanin marks the spot: identifying a locus coeruleus biomarker of cognitive 811 reserve in healthy aging. Neurobiol Aging, 37, 117-126. 812 https://doi.org/10.1016/j.neurobiolaging.2015.09.019 813

Dahl, M. J., Mather, M., Düzel, S., Bodammer, N. C., Lindenberger, U., Kühn, S., & Werkle-

Bergner, M. (2019). Rostral locus coeruleus integrity is associated with better 815 memory performance in older adults. Nat Hum Behav, 3(11), 1203-1214. 816 817 https://doi.org/10.1038/s41562-019-0715-2 Dahl, M. J., Mather, M., Werkle-Bergner, M., Kennedy, B. L., Guzman, S., Hurth, K., Miller, 818 C. A., Qiao, Y., Shi, Y., Chui, H. C., & Ringman, J. M. (2022). Locus coeruleus 819 integrity is related to tau burden and memory loss in autosomal-dominant Alzheimer's 820 disease. Neurobiol Aging, 112, 39-54. 821 https://doi.org/10.1016/j.neurobiolaging.2021.11.006 822 Deckers, K., van Boxtel, M. P., Schiepers, O. J., de Vugt, M., Muñoz Sánchez, J. L., Anstey, 823 K. J., Brayne, C., Dartigues, J. F., Engedal, K., Kivipelto, M., Ritchie, K., Starr, J. M., 824 825 Yaffe, K., Irving, K., Verhey, F. R., & Köhler, S. (2015). Target risk factors for dementia prevention: a systematic review and Delphi consensus study on the evidence 826 from observational studies. Int J Geriatr Psychiatry, 30(3), 234-246. 827 https://doi.org/10.1002/gps.4245 828 Del Cerro, I., Villarreal, M. F., Abulafia, C., Duarte-Abritta, B., Sánchez, S. M., Castro, M. 829 N., Bocaccio, H., Ferrer, I., Menchón, J. M., Sevlever, G., Nemeroff, C. B., Soriano-830 Mas, C., & Guinjoan, S. M. (2020). Disrupted functional connectivity of the locus 831 coeruleus in healthy adults with parental history of Alzheimer's disease. J Psychiatr 832 Res, 123, 81-88. https://doi.org/10.1016/j.jpsychires.2020.01.018 833 Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner, 834 R. L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., & Killiany, R. J. 835 (2006). An automated labeling system for subdividing the human cerebral cortex on 836 MRI scans into gyral based regions of interest. NeuroImage, 31(3), 968-980. 837 https://doi.org/10.1016/j.neuroimage.2006.01.021 838

839	Donix, M., Small, G. W., & Bookheimer, S. Y. (2012). Family history and APOE-4 genetic
840	risk in Alzheimer's disease. Neuropsychol Rev, 22(3), 298-309.
841	https://doi.org/10.1007/s11065-012-9193-2
842	Dounavi, ME., Low, A., Mckiernan, E. F., Mak, E., Muniz-Terrera, G., Ritchie, K., Ritchie,
843	C. W., Su, L., & O'Brien, J. T. (2021). Evidence of cerebral hemodynamic
844	dysregulation in middle-aged APOE ε4 carriers: The PREVENT-Dementia study.
845	Journal of Cerebral Blood Flow & Metabolism, 41(11), 2844-2855.
846	https://doi.org/10.1177/0271678x211020863
847	Dounavi, ME., Newton, C., Jenkins, N., Mak, E., Low, A., Muniz-Terrera, G., Williams, G.
848	B., Lawlor, B., Naci, L., Malhotra, P., Mackay, C. E., Koychev, I., Ritchie, K.,
849	Ritchie, C. W., Su, L., & O'Brien, J. T. (2022). Macrostructural brain alterations at
850	midlife are connected to cardiovascular and not inherited risk of future dementia: the
851	PREVENT-Dementia study. <i>Journal of Neurology</i> . https://doi.org/10.1007/s00415-
852	<u>022-11061-7</u>
853	Dounavi, M. E., Mak, E., Wells, K., Ritchie, K., Ritchie, C. W., Su, L., & JT, O. B. (2020).
854	Volumetric alterations in the hippocampal subfields of subjects at increased risk of
855	dementia. Neurobiol Aging, 91, 36-44.
856	https://doi.org/10.1016/j.neurobiolaging.2020.03.006
857	Ehrenberg, A. J., Nguy, A. K., Theofilas, P., Dunlop, S., Suemoto, C. K., Di Lorenzo Alho, A
858	T., Grinberg, L. T. (2017). Quantifying the accretion of hyperphosphorylated tau
859	in the locus coeruleus and dorsal raphe nucleus: the pathological building blocks of
860	early Alzheimer's disease. Neuropathol Appl Neurobiol, 43(5), 393-408.
861	https://doi.org/10.1111/nan.12387

Exalto, L. G., Quesenberry, C. P., Barnes, D., Kivipelto, M., Biessels, G. J., & Whitmer, R. A. 862 (2014). Midlife risk score for the prediction of dementia four decades later. 863 Alzheimer's & Dementia, 10(5), 562-570. https://doi.org/10.1016/j.jalz.2013.05.1772 864 Fayosse, A., Nguyen, D. P., Dugravot, A., Dumurgier, J., Tabak, A. G., Kivimäki, M., Sabia, 865 S., & Singh-Manoux, A. (2020). Risk prediction models for dementia: role of age and 866 cardiometabolic risk factors. BMC Med, 18(1), 107. https://doi.org/10.1186/s12916-867 020-01578-x 868 869 Giannakopoulos, P., Herrmann, F. R., Bussière, T., Bouras, C., Kövari, E., Perl, D. P., Morrison, J. H., Gold, G., & Hof, P. R. (2003). Tangle and neuron numbers, but not 870 amyloid load, predict cognitive status in Alzheimer's disease. Neurology, 60(9), 1495-871 1500. https://doi.org/10.1212/01.wnl.0000063311.58879.01 872 Gilvesy, A., Husen, E., Magloczky, Z., Mihaly, O., Hortobágyi, T., Kanatani, S., . . . Adori, C. 873 874 (2022). Spatiotemporal characterization of cellular tau pathology in the human locus coeruleus-pericoerulear complex by three-dimensional imaging. Acta Neuropathol. 875 https://doi.org/10.1007/s00401-022-02477-6 876 Hair Jr., J. F., Anderson, R. E., Tatham, R. L., & Black, W. C. (1998). Multivariate Data 877 Analysis (5th ed.). Prentice-Hall: London. 878 Heneghan, A., Deng, F., Wells, K., Ritchie, K., Muniz-Terrera, G., Ritchie, C. W., Lawlor, B., 879 & Naci, L. (2023). Modifiable Lifestyle Activities Affect Cognition in Cognitively 880 Healthy Middle-Aged Individuals at Risk for Late-Life Alzheimer's Disease. Journal 881 of Alzheimer's disease: JAD, 91(2), 833–846. https://doi.org/10.3233/JAD-220267 882 Horn, J. (1965). A rationale and test for the number of factors in factor analysis. 883 Psychometrika, 30, 179-185. https://doi.org/10.1007/BF02289447 884

Hsieh, S., Schubert, S., Hoon, C., Mioshi, E., & Hodges, J. R. (2013). Validation of the 885 Addenbrooke's Cognitive Examination III in frontotemporal dementia and 886 Alzheimer's disease. Dement Geriatr Cogn Disord, 36(3-4), 242-250. 887 https://doi.org/10.1159/000351671 888 Irish, M., Lawlor, B. A., Coen, R. F., & O'Mara, S. M. (2011). Everyday episodic memory in 889 890 amnestic mild cognitive impairment: a preliminary investigation. BMC neuroscience, 12(1), 1-13. https://doi.org/10.1186/1471-2202-12-80 891 892 Irwin, K., Sexton, C., Daniel, T., Lawlor, B., & Naci, L. (2018). Healthy Aging and Dementia: Two Roads Diverging in Midlife? Front Aging Neurosci, 10, 275. 893 https://doi.org/10.3389/fnagi.2018.00275 894 Jacobs, H. I., Wiese, S., van de Ven, V., Gronenschild, E. H., Verhey, F. R., & Matthews, P. 895 M. (2015). Relevance of parahippocampal-locus coeruleus connectivity to memory in 896 early dementia. Neurobiol Aging, 36(2), 618-626. 897 https://doi.org/10.1016/j.neurobiolaging.2014.10.041 898 899 Jacobs, H. I. L., Müller-Ehrenberg, L., Priovoulos, N., & Roebroeck, A. (2018). Curvilinear locus coeruleus functional connectivity trajectories over the adult lifespan: a 7T MRI 900 study. *Neurobiology of aging*, 69, 167–176. 901 https://doi.org/10.1016/j.neurobiolaging.2018.05.021 902 Jacobs, H. I. L., Becker, J. A., Kwong, K., Engels-Domínguez, N., Prokopiou, P. C., Papp, K. 903 V., Properzi, M., Hampton, O. L., d'Oleire Uquillas, F., Sanchez, J. S., Rentz, D. M., 904 El Fakhri, G., Normandin, M. D., Price, J. C., Bennett, D. A., Sperling, R. A., & 905 906 Johnson, K. A. (2021). In vivo and neuropathology data support locus coeruleus integrity as indicator of Alzheimer's disease pathology and cognitive decline. Sci 907

908	Transl Med, 13(612), eabj2511. https://doi.org/10.1126/scitranslmed.abj2511
909	Jacobs, H. I. L., Becker, J. A., Kwong, K., Munera, D., Ramirez-Gomez, L., Engels-
910	Domínguez, N., Sanchez, J. S., Vila-Castelar, C., Baena, A., Sperling, R. A., Johnson
911	K. A., Lopera, F., & Quiroz, Y. T. (2022). Waning locus coeruleus integrity precedes
912	cortical tau accrual in preclinical autosomal dominant Alzheimer's disease.
913	Alzheimers Dement. https://doi.org/10.1002/alz.12656
914	James, T., Kula, B., Choi, S., Khan, S. S., Bekar, L. K., & Smith, N. A. (2021). Locus
915	coeruleus in memory formation and Alzheimer's disease. European Journal of
916	Neuroscience, 54(8), 6948-6959. https://doi.org/10.1111/ejn.15045
917	Jansen, W. J., Ossenkoppele, R., Knol, D. L., Tijms, B. M., Scheltens, P., Verhey, F. R.,
918	Visser, P. J., Aalten, P., Aarsland, D., Alcolea, D., Alexander, M., Almdahl, I. S.,
919	Arnold, S. E., Baldeiras, I., Barthel, H., van Berckel, B. N., Bibeau, K., Blennow, K.,
920	Brooks, D. J., Zetterberg, H. (2015). Prevalence of cerebral amyloid pathology in
921	persons without dementia: a meta-analysis. JAMA, 313(19), 1924-1938.
922	https://doi.org/10.1001/jama.2015.4668
923	Johnson, P. O., & Fay, L. C. (1950). The Johnson-Neyman technique, its theory and
924	application. Psychometrika, 15(4), 349–367. https://doi.org/10.1007/BF02288864
925	Jolliffe, I. T., & Cadima, J. (2016). Principal component analysis: a review and recent
926	developments. Philos Trans A Math Phys Eng Sci, 374(2065), 20150202.
927	https://doi.org/10.1098/rsta.2015.0202.
928	Kaiser, H. F. (1958). The varimax criterion for analytic rotation in factor analysis.
929	Psychometrika, 23(3), 187-200. https://doi.org/10.1007/BF02289233

930	Kametani, F., & Hasegawa, M. (2018). Reconsideration of Amyloid Hypothesis and Tau
931	Hypothesis in Alzheimer's Disease. Front Neurosci, 12, 25.
932	https://doi.org/10.3389/fnins.2018.00025
933	Kaufman, S. K., Del Tredici, K., Thomas, T. L., Braak, H., & Diamond, M. I. (2018). Tau
934	seeding activity begins in the transentorhinal/entorhinal regions and anticipates
935	phospho-tau pathology in Alzheimer's disease and PART. Acta Neuropathol, 136(1),
936	57-67. https://doi.org/10.1007/s00401-018-1855-6
937	Kaufman, A. M., Geiller, T., & Losonczy, A. (2020). A Role for the Locus Coeruleus in
938	Hippocampal CA1 Place Cell Reorganization during Spatial Reward Learning.
939	Neuron, 105(6), 1018-1026.e1014. https://doi.org/10.1016/j.neuron.2019.12.029
940	Kempadoo, K. A., Mosharov, E. V., Choi, S. J., Sulzer, D., & Kandel, E. R. (2016).
941	Dopamine release from the locus coeruleus to the dorsal hippocampus promotes
942	spatial learning and memory. Proc Natl Acad Sci USA, 113(51), 14835-14840.
943	https://doi.org/10.1073/pnas.1616515114
944	Kerchner, G. A., Berdnik, D., Shen, J. C., Bernstein, J. D., Fenesy, M. C., Deutsch, G. K.,
945	Rutt, B. K. (2014). APOE ε4 worsens hippocampal CA1 apical neuropil atrophy and
946	episodic memory. Neurology, 82(8), 691-697.
947	https://doi.org/10.1212/wnl.00000000000154
948	Kivipelto, M., Ngandu, T., Laatikainen, T., Winblad, B., Soininen, H., & Tuomilehto, J.
949	(2006). Risk score for the prediction of dementia risk in 20 years among middle aged
950	people: a longitudinal, population-based study. The Lancet Neurology, 5(9), 735-741.
951	https://doi.org/10.1016/s1474-4422(06)70537-3
952	La Joie, R., Ayakta, N., Seeley, W. W., Borys, E., Boxer, A. L., DeCarli, C., Doré, V.,

- 953 Grinberg, L. T., Huang, E., Hwang, J. H., Ikonomovic, M. D., Jack, C., Jr., Jagust, W. J., Jin, L. W., Klunk, W. E., Kofler, J., Lesman-Segev, O. H., Lockhart, S. N., Lowe, 954 V. J., ... Rabinovici, G. D. (2019). Multisite study of the relationships between 955 antemortem [(11)C]PIB-PET Centiloid values and postmortem measures of 956 Alzheimer's disease neuropathology. *Alzheimers Dement*, 15(2), 205-216. 957 https://doi.org/10.1016/j.jalz.2018.09.001 958 959 Lambert, J. C., Ibrahim-Verbaas, C. A., Harold, D., Naj, A. C., Sims, R., Bellenguez, C., DeStafano, A. L., Bis, J. C., Beecham, G. W., Grenier-Boley, B., Russo, G., Thorton-960 Wells, T. A., Jones, N., Smith, A. V., Chouraki, V., Thomas, C., Ikram, M. A., 961 Zelenika, D., Vardarajan, B. N., . . . Amouyel, P. (2013). Meta-analysis of 74,046 962 individuals identifies 11 new susceptibility loci for Alzheimer's disease. Nat Genet, 963 45(12), 1452-1458. https://doi.org/10.1038/ng.2802 964 Lee, T. H., Kim, S. H., Katz, B., & Mather, M. (2020). The Decline in Intrinsic Connectivity 965 Between the Salience Network and Locus Coeruleus in Older Adults: Implications for 966 Distractibility. Front Aging Neurosci, 12, 2. https://doi.org/10.3389/fnagi.2020.00002 967 Liebe, T., Dordevic, M., Kaufmann, J., Avetisyan, A., Skalej, M., & Müller, N. (2022). 968 Investigation of the functional pathogenesis of mild cognitive impairment by 969 localisation-based locus coeruleus resting-state fMRI. Human brain mapping, 43(18), 970 5630–5642. https://doi.org/10.1002/hbm.26039 971 972 Liu, C. C., Liu, C. C., Kanekiyo, T., Xu, H., & Bu, G. (2013). Apolipoprotein E and Alzheimer disease: risk, mechanisms and therapy. Nat Rev Neurol, 9(2), 106-118. 973 https://doi.org/10.1038/nrneurol.2012.263 974
- Liu, K. Y., Acosta-Cabronero, J., Cardenas-Blanco, A., Loane, C., Berry, A. J., Betts, M. J.,

976	Kievit, R. A., Henson, R. N., Düzel, E., Howard, R., & Hämmerer, D. (2019). In vivo
977	visualization of age-related differences in the locus coeruleus. Neurobiol Aging, 74,
978	101-111. https://doi.org/10.1016/j.neurobiolaging.2018.10.014
979	Liu, K. Y., Kievit, R. A., Tsvetanov, K. A., Betts, M. J., Düzel, E., Rowe, J. B., Cam-CAN,
980	Howard, R., & Hämmerer, D. (2020). Noradrenergic-dependent functions are
981	associated with age-related locus coeruleus signal intensity differences. Nature
982	communications, 11(1), 1712. https://doi.org/10.1038/s41467-020-15410-w
983	Livingston, G., Huntley, J., Sommerlad, A., Ames, D., Ballard, C., Banerjee, S., Brayne, C.,
984	Burns, A., Cohen-Mansfield, J., Cooper, C., Costafreda, S. G., Dias, A., Fox, N.,
985	Gitlin, L. N., Howard, R., Kales, H. C., Kivimäki, M., Larson, E. B., Ogunniyi, A.,
986	Mukadam, N. (2020). Dementia prevention, intervention, and care: 2020 report of the
987	Lancet Commission. The Lancet, 396(10248), 413-446.
988	https://doi.org/10.1016/s0140-6736(20)30367-6
989	Livingston, G., Sommerlad, A., Orgeta, V., Costafreda, S. G., Huntley, J., Ames, D., Ballard,
990	C., Banerjee, S., Burns, A., Cohen-Mansfield, J., Cooper, C., Fox, N., Gitlin, L. N.,
991	Howard, R., Kales, H. C., Larson, E. B., Ritchie, K., Rockwood, K., Sampson, E.
992	L., Mukadam, N. (2017). Dementia prevention, intervention, and care. Lancet,
993	390(10113), 2673-2734. https://doi.org/10.1016/s0140-6736(17)31363-6
994	Lorenzo-Seva, U., & ten Berge, J. M. F. (2006). Tucker's congruence coefficient as a
995	meaningful index of factor similarity. Methodology: European Journal of Research
996	Methods for the Behavioral and Social Sciences, 2(2), 57-64.
997	https://doi.org/10.1027/1614-2241.2.2.57

Low, A., Su, L., Stefaniak, J. D., Mak, E., Dounavi, M. E., Muniz-Terrera, G., Ritchie, K.,

999	Ritchie, C. W., Markus, H. S., & O'Brien, J. T. (2021). Inherited risk of dementia and
1000	the progression of cerebral small vessel disease and inflammatory markers in
1001	cognitively healthy midlife adults: the PREVENT-Dementia study. Neurobiol Aging,
1002	98, 124-133. https://doi.org/10.1016/j.neurobiolaging.2020.10.029
1003	Mather, M., & Harley, C. W. (2016). The Locus Coeruleus: Essential for Maintaining
1004	Cognitive Function and the Aging Brain. Trends Cogn Sci, 20(3), 214-226.
1005	https://doi.org/10.1016/j.tics.2016.01.001
1006	Mondadori, C. R. A., De Quervain, D. JF., Buchmann, A., Mustovic, H., Wollmer, M. A.,
1007	Schmidt, C. F., Boesiger, P., Hock, C., Nitsch, R. M., Papassotiropoulos, A., & Henke,
1008	K. (2007). Better Memory and Neural Efficiency in Young Apolipoprotein E 4
1009	Carriers. Cerebral Cortex, 17(8), 1934-1947. https://doi.org/10.1093/cercor/bhl103
1010	Mosconi, L., Berti, V., Quinn, C., McHugh, P., Petrongolo, G., Varsavsky, I., Osorio, R. S.,
1011	Pupi, A., Vallabhajosula, S., Isaacson, R. S., de Leon, M. J., & Brinton, R. D. (2017).
1012	Sex differences in Alzheimer risk: Brain imaging of endocrine vs chronologic aging.
1013	Neurology, 89(13), 1382-1390. https://doi.org/10.1212/wnl.0000000000004425
1014	Mosconi, L., Rahman, A., Diaz, I., Wu, X., Scheyer, O., Hristov, H. W., Vallabhajosula, S.,
1015	Isaacson, R. S., de Leon, M. J., & Brinton, R. D. (2018). Increased Alzheimer's risk
1016	during the menopause transition: A 3-year longitudinal brain imaging study. PLoS
1017	One, 13(12), e0207885. https://doi.org/10.1371/journal.pone.0207885
1018	Murphy, K., Birn, R. M., Handwerker, D. A., Jones, T. B., & Bandettini, P. A. (2009). The
1019	impact of global signal regression on resting state correlations: are anti-correlated
1020	networks introduced? NeuroImage, 44(3), 893-905.
1021	https://doi.org/10.1016/j.neuroimage.2008.09.036

1022 Olivieri, P., Lagarde, J., Lehericy, S., Valabrègue, R., Michel, A., Macé, P., Caillé, F., Gervais, P., Bottlaender, M., & Sarazin, M. (2019). Early alteration of the locus 1023 coeruleus in phenotypic variants of Alzheimer's disease. Ann Clin Transl Neurol, 6(7), 1024 1345-1351. https://doi.org/10.1002/acn3.50818 1025 1026 Parra, M. A., Abrahams, S., Logie, R. H., Méndez, L. G., Lopera, F., & Della Sala, S. (2010). 1027 Visual short-term memory binding deficits in familial Alzheimer's disease. *Brain*, 1028 133(9), 2702-2713. https://doi.org/10.1093/brain/awq148 1029 Plini, E. R. G., O'Hanlon, E., Boyle, R., Sibilia, F., Rikhye, G., Kenney, J., Whelan, R., Melnychuk, M. C., Robertson, I. H., & Dockree, P. M. (2021). Examining the Role of 1030 1031 the Noradrenergic Locus Coeruleus for Predicting Attention and Brain Maintenance 1032 in Healthy Old Age and Disease: An MRI Structural Study for the Alzheimer's 1033 Disease Neuroimaging Initiative. *Cells*, 10(7), 1829. https://doi.org/10.3390/cells10071829 1034 1035 Porat, S., Sibilia, F., Yoon, J., Shi, Y., Dahl, M. J., Werkle-Bergner, M., Düzel, S., Bodammer, N., Lindenberger, U., Kühn, S., & Mather, M. (2022). Age differences in 1036 1037 diffusivity in the locus coeruleus and its ascending noradrenergic tract. NeuroImage, 251, 119022. https://doi.org/10.1016/j.neuroimage.2022.119022 1038 Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2012). 1039 Spurious but systematic correlations in functional connectivity MRI networks arise 1040 1041 from subject motion. NeuroImage, 59(3), 2142-2154. https://doi.org/10.1016/j.neuroimage.2011.10.018 1042 Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. 1043 1044 (2014). Methods to detect, characterize, and remove motion artifact in resting state

1045	fMRI. NeuroImage, 84, 320-341. https://doi.org/10.1016/j.neuroimage.2013.08.048
1046	Prokopiou, P. C., Engels-Domínguez, N., Papp, K. V., Scott, M. R., Schultz, A. P., Schneider
1047	C., Farrell, M. E., Buckley, R. F., Quiroz, Y. T., El Fakhri, G., Rentz, D. M., Sperling,
1048	R. A., Johnson, K. A., & Jacobs, H. I. L. (2022). Lower novelty-related locus
1049	coeruleus function is associated with Aβ-related cognitive decline in clinically healthy
1050	individuals. Nature Communications, 13(1). https://doi.org/10.1038/s41467-022-
1051	<u>28986-2</u>
1052	Rahman, A., Schelbaum, E., Hoffman, K., Diaz, I., Hristov, H., Andrews, R., Jett, S., Jackson
1053	H., Lee, A., Sarva, H., Pahlajani, S., Matthews, D., Dyke, J., de Leon, M. J., Isaacson,
1054	R. S., Brinton, R. D., & Mosconi, L. (2020). Sex-driven modifiers of Alzheimer risk:
1055	A multimodality brain imaging study. <i>Neurology</i> , 95(2), e166-e178.
1056	https://doi.org/10.1212/wnl.0000000000009781
1057	Ritchie, C. W., & Ritchie, K. (2012). The PREVENT study: a prospective cohort study to
1058	identify mid-life biomarkers of late-onset Alzheimer's disease. BMJ Open, 2(6),
1059	e001893. https://doi.org/10.1136/bmjopen-2012-001893
1060	Ritchie, K., Carrière, I., Gregory, S., Watermeyer, T., Danso, S., Su, L., Ritchie, C. W., &
1061	O'Brien, J. T. (2021). Trauma and depressive symptomatology in middle-aged
1062	persons at high risk of dementia: the PREVENT Dementia Study. Journal of
1063	Neurology, Neurosurgery & Psychiatry, 92(1), 16-21. https://doi.org/10.1136/jnnp-
1064	<u>2020-323823</u>
1065	Ritchie, K., Carrière, I., Howett, D., Su, L., Hornberger, M., O'Brien, J. T., Ritchie, C. W., &
1066	Chan, D. (2018). Allocentric and Egocentric Spatial Processing in Middle-Aged
1067	Adults at High Risk of Late-Onset Alzheimer's Disease: The PREVENT Dementia

1068 Study. Journal of Alzheimer's Disease, 65(3), 885-896. https://doi.org/10.3233/jad-180432 1069 1070 Ritchie, K., Carrière, I., Ritchie, C. W., Berr, C., Artero, S., & Ancelin, M. L. (2010). 1071 Designing prevention programmes to reduce incidence of dementia: prospective 1072 cohort study of modifiable risk factors. Bmj, 341, c3885. 1073 https://doi.org/10.1136/bmj.c3885 Ritchie, K., Carrière, I., Su, L., O'Brien, J. T., Lovestone, S., Wells, K., & Ritchie, C. W. 1074 (2017). The midlife cognitive profiles of adults at high risk of late-onset Alzheimer's 1075 1076 disease: The PREVENT study. Alzheimer's & Dementia, 13(10), 1089-1097. 1077 https://doi.org/10.1016/j.jalz.2017.02.008 Ritchie, K., de Roquefeuil, G., Ritchie, C., Besset, A., Poulain, V., Artero, S., Ancelin, M. L. 1078 (2014). COGNITO: computerized assessment of information processing. Journal of 1079 1080 Psychology & Psychotherapy, 4(2). https://doi.org/10.4172/2161-0487.1000136 Salvador, R., Martínez, A., Pomarol-Clotet, E., Gomar, J., Vila, F., Sarró, S., Capdevila, A., 1081 1082 & Bullmore, E. (2008). A simple view of the brain through a frequency-specific functional connectivity measure. NeuroImage, 39(1), 279-289. 1083 https://doi.org/10.1016/j.neuroimage.2007.08.018 1084 1085 Sara, S. J. (2009). The locus coeruleus and noradrenergic modulation of cognition. *Nat Rev* 1086 *Neurosci*, 10(3), 211-223. https://doi.org/10.1038/nrn2573 1087 Satterthwaite, T. D., Elliott, M. A., Gerraty, R. T., Ruparel, K., Loughead, J., Calkins, M. E., Eickhoff, S. B., Hakonarson, H., Gur, R. C., Gur, R. E., & Wolf, D. H. (2013). An 1088 1089 improved framework for confound regression and filtering for control of motion artifact in the preprocessing of resting-state functional connectivity data. NeuroImage, 1090

1091	64, 240-256. https://doi.org/10.1016/j.neuroimage.2012.08.052
1092	Scarabino, D., Gambina, G., Broggio, E., Pelliccia, F., & Corbo, R. M. (2016). Influence of
1093	family history of dementia in the development and progression of late-onset
1094	Alzheimer's disease. Am J Med Genet B Neuropsychiatr Genet, 171b(2), 250-256.
1095	https://doi.org/10.1002/ajmg.b.32399
1096	Scheyer, O., Rahman, A., Hristov, H., Berkowitz, C., Isaacson, R. S., Diaz Brinton, R., &
1097	Mosconi, L. (2018). Female Sex and Alzheimer's Risk: The Menopause Connection. J
1098	Prev Alzheimers Dis, 5(4), 225-230. https://doi.org/10.14283/jpad.2018.34
1099	Sindi, S., Calov, E., Fokkens, J., Ngandu, T., Soininen, H., Tuomilehto, J., & Kivipelto, M.
1100	(2015). The CAIDE Dementia Risk Score App: The development of an evidence-
1101	based mobile application to predict the risk of dementia. Alzheimers Dement (Amst),
1102	1(3), 328-333. https://doi.org/10.1016/j.dadm.2015.06.005
1103	Song, I., Neal, J., & Lee, T. H. (2021). Age-Related Intrinsic Functional Connectivity
1104	Changes of Locus Coeruleus from Childhood to Older Adults. Brain sciences, 11(11),
1105	1485. https://doi.org/10.3390/brainsci111111485
1106	Sperling, R. A., Aisen, P. S., Beckett, L. A., Bennett, D. A., Craft, S., Fagan, A. M., Iwatsubo
1107	T., Jack, C. R., Kaye, J., Montine, T. J., Park, D. C., Reiman, E. M., Rowe, C. C.,
1108	Siemers, E., Stern, Y., Yaffe, K., Carrillo, M. C., Thies, B., Morrison-Bogorad, M.,
1109	Phelps, C. H. (2011). Toward defining the preclinical stages of Alzheimer's disease:
1110	Recommendations from the National Institute on Aging-Alzheimer's Association
1111	workgroups on diagnostic guidelines for Alzheimer's disease. Alzheimer's &
1112	Dementia, 7(3), 280-292. https://doi.org/10.1016/j.jalz.2011.03.003
1113	Stephen R. Liu V. Ngandu T. Antikainen R. Hulkkonen I. Koikkalainen I.

1114	Kemppainen, N., Lötjönen, J., Levälahti, E., Parkkola, R., Pippola, P., Rinne, J.,
1115	Strandberg, T., Tuomilehto, J., Vanninen, R., Kivipelto, M., Soininen, H., & Solomon,
1116	A. (2019). Brain volumes and cortical thickness on MRI in the Finnish Geriatric
1117	Intervention Study to Prevent Cognitive Impairment and Disability (FINGER).
1118	Alzheimers Res Ther, 11(1), 53. https://doi.org/10.1186/s13195-019-0506-z
1119	Stevens, J. P. (1992). Applied multivariate statistics for the social sciences (2nd ed.).
1120	Hillsdale, NJ: Erlbaum.
1121	Sun, W., Tang, Y., Qiao, Y., Ge, X., Mather, M., Ringman, J. M., & Shi, Y. (2020). A
1122	probabilistic atlas of locus coeruleus pathways to transentorhinal cortex for
1123	connectome imaging in Alzheimer's disease. NeuroImage, 223, 117301.
1124	https://doi.org/10.1016/j.neuroimage.2020.117301
1125	Takahashi, J., Shibata, T., Sasaki, M., Kudo, M., Yanezawa, H., Obara, S., Kudo, K., Ito, K.,
1126	Yamashita, F., & Terayama, Y. (2015). Detection of changes in the locus coeruleus in
1127	patients with mild cognitive impairment and Alzheimer's disease: high-resolution fast
1128	spin-echo T1-weighted imaging. Geriatr Gerontol Int, 15(3), 334-340.
1129	https://doi.org/10.1111/ggi.12280
1130	Tustison, N. J., Avants, B. B., Cook, P. A., Zheng, Y., Egan, A., Yushkevich, P. A., & Gee, J.
1131	C. (2010). N4ITK: improved N3 bias correction. IEEE transactions on medical
1132	imaging, 29(6), 1310–1320. https://doi.org/10.1109/TMI.2010.2046908
1133	Udeh-Momoh, C., & Watermeyer, T. (2021). Female specific risk factors for the
1134	development of Alzheimer's disease neuropathology and cognitive impairment: Call
1135	for a precision medicine approach. Ageing Res Rev, 71, 101459.
1136	https://doi.org/10.1016/j.arr.2021.101459

Van Dijk, K. R., Hedden, T., Venkataraman, A., Evans, K. C., Lazar, S. W., & Buckner, R. L. 1137 (2010). Intrinsic functional connectivity as a tool for human connectomics: theory, 1138 properties, and optimization. *Journal of neurophysiology*, 103(1), 297–321. 1139 https://doi.org/10.1152/jn.00783.2009 1140 1141 Van Egroo, M., van Hooren, R. W. E., & Jacobs, H. I. L. (2021). Associations between locus 1142 coeruleus integrity and nocturnal awakenings in the context of Alzheimer's disease 1143 plasma biomarkers: a 7T MRI study. Alzheimers Res Ther, 13(1), 159. https://doi.org/10.1186/s13195-021-00902-8 1144 Vilor-Tejedor, N., Evans, T. E., Adams, H. H., González-de-Echávarri, J. M., Molinuevo, J. 1145 L., Guigo, R., ... Operto, G. (2021). Genetic Influences on Hippocampal Subfields: 1146 1147 An Emerging Area of Neuroscience Research. Neurol Genet, 7(3), e591. https://doi.org/10.1212/nxg.0000000000000591 1148 1149 Weber, M. T., Maki, P. M., & McDermott, M. P. (2014). Cognition and mood in perimenopause: a systematic review and meta-analysis. J Steroid Biochem Mol Biol, 1150 142, 90-98. https://doi.org/10.1016/j.jsbmb.2013.06.001 1151 Weinshenker, D. (2018). Long Road to Ruin: Noradrenergic Dysfunction in 1152 Neurodegenerative Disease. Trends Neurosci, 41(4), 211-223. 1153 https://doi.org/10.1016/j.tins.2018.01.010 1154 Williams GC. Pleiotropy, Natural Selection, and the Evolution of Senescence. Evolution. 1155 1957;11:398-411. https://doi.org/10.1111/j.1558-5646.1957.tb02911.x 1156 World Health Organization. (2021). Global status report on the public health response to 1157 dementia. 1158

1159 Yanes, L. L., & Reckelhoff, J. F. (2011). Postmenopausal hypertension. Am J Hypertens, 24(7), 740-749. https://doi.org/10.1038/ajh.2011.71 1160 1161 Ye, R., Rua, C., O'Callaghan, C., Jones, P. S., Hezemans, F. H., Kaalund, S. S., Tsvetanov, K. A., Rodgers, C. T., Williams, G., Passamonti, L., & Rowe, J. B. (2021). An in vivo 1162 1163 probabilistic atlas of the human locus coeruleus at ultra-high field. NeuroImage, 225, 117487. https://doi.org/10.1016/j.neuroimage.2020.117487 1164 Zhang, S., Hu, S., Chao, H. H., & Li, C.-S. R. (2016). Resting-state functional connectivity of 1165 1166 the locus coeruleus in humans: in comparison with the ventral tegmental area/substantia nigra pars compacta and the effects of age. Cerebral Cortex, 26(8), 1167 3413-3427. https://doi.org/10.1093/cercor/bhv172 1168 1169 Zhao, S., Rangaprakash, D., Venkataraman, A., Liang, P., & Deshpande, G. (2017). 1170 Investigating Focal Connectivity Deficits in Alzheimer's Disease Using Directional 1171 Brain Networks Derived from Resting-State fMRI. Frontiers in aging neuroscience, 9, 211. https://doi.org/10.3389/fnagi.2017.00211 1172 1173 Zokaei, N., Grogan, J., Fallon, S. J., Slavkova, E., Hadida, J., Manohar, S., . . . Husain, M. (2020). Short-term memory advantage for brief durations in human APOE ε4 carriers. 1174 Sci Rep, 10(1), 9503. https://doi.org/10.1038/s41598-020-66114-6 1175 Zuo, X. N., Di Martino, A., Kelly, C., Shehzad, Z. E., Gee, D. G., Klein, D. F., Castellanos, F. 1176 X., Biswal, B. B., & Milham, M. P. (2010). The oscillating brain: complex and 1177 reliable. NeuroImage, 49(2), 1432-1445. 1178 https://doi.org/10.1016/j.neuroimage.2009.09.037 1179

Figure Legends

Figure 1. The flowchart of functional magnetic resonance imaging (fMRI) data processing. (a) The standard preprocessing pipeline for fMRI data. (b) Three exclusion criteria were used to exclude participants. A 'scrubbing' procedure was used to account for the effect of head movement on functional connectivity analyses (Power et al., 2012). (c) The top panels shows the study-specific 3mm locus coeruleus (LC) mask (in blue), which closely matched the 7T probabilistic atlas developed by Ye et al. (2021) and the LC meta-mask developed by Dahl et al. (2022). The bottom panel shows the study-specific mask superimposed on a participant's T1w structural data in MNI space. (d) Functional connectivity was calculated between the bilateral LC (in blue) and the hippocampus (in red). The hippocampus was defined after *recon-all* pipeline using Freesurfer version 7.1.0.

Figure 2. Data reduction of cognitive information. (a) Scree plots and parallel analysis for the baseline and follow-up datasets. Eigenvalues of the principal components obtained from the actual data (in blue line) were compared to those of random simulated data (mean values in grey line and 95th percentile in red line). Eigenvalues larger than 95th percentile of the randomly generated 500 eigenvalues (simulation) determined the number of components. (b) The three extracted components were rotated to be uncorrelated with each other. The proportion of variance explained by each component, and the cumulative variance explained by the three are presented for the baseline and follow-up datasets. Abbreviations: C1, component 1; C2, component 2; C3, component 3.

Figure 3. Interpretation of data-derived cognitive domains. Component (*C*) coefficients and Tucker's congruence coefficients between (a) baseline and (b) follow-up, are shown. (a) The coefficients for 13 original cognitive measures (in rows) on the three components (in columns). A larger absolute coefficient (darker colors and larger solid circles) represents a closer relationship between the cognitive measure and corresponding component. Cool/warm colors represent the positive/negative relationships between cognitive measures and components. Only cognitive measures with a coefficient > 0.4 on each component, highlighted with purple rectangles, were used to interpret the cognitive functions driving each component. (b) Similarities among the components across baseline (in rows) and follow-up (in columns) were measured by Tucker's congruence coefficients. The diagonal values represented the similarities for each of the components with itself across time, and off diagonal values indicated the similarities for each of the components with the other two components across time. Cool/warm colors represent the positive/negative relationships with darker colors representing correlation magnitude, as shown in the color-bar scale.

Abbreviations: VSTMBT, visual short-term memory binding test; diff, difference.

Figure 4. The associations between the CAIDE dementia risk score and verbal, visuospatial functions, and short-term (conjunctive) memory at (a) baseline and (b) follow-up.

Abbreviation: CAIDE, Cardiovascular risk factor, Aging, and Incidence of Dementia.

Figure 5. Longitudinal decline of the LC–Hippocampus functional connectivity over 2 years in cognitively healthy middle-aged adults. * = p < 0.05. Abbreviations: LC, Locus Coeruleus.

Figure 6. The CAIDE score significantly moderated the brain–behaviour relationship. The relationship was plotted at the fixed values of CAIDE (mean and \pm 1 SD) to show the effect of CAIDE at (a) baseline and (b) follow-up. The LC–Hippocampus functional connectivity and the verbal, visuospatial functions, and short-term (conjunctive) memory were positively associated for individuals with low CAIDE score, and negatively associated for those with high scores, at (b) follow-up, but not baseline (a). Johnson-Neyman plots show the specific range of CAIDE scores where FC was significantly associated with cognition at (c) baseline and (d) follow-up. The bolded horizontal line shows the range of CAIDE scores of our cohort. Two blue dashed vertical lines indicate the interval of significant (blue area) and non-significant (pink area) brain-behaviour relationships. At follow-up, there was a significant positive relationship between FC and cognition for individuals with a CAIDE score < 3.25 and a significant negative relationship for individuals with a CAIDE score > 9.65. Abbreviations: LC, locus coeruleus; CAIDE, Cardiovascular risk factor, Aging, and Incidence of Dementia; FC, functional connectivity; SD, standard deviation; n.s., not significant.

Table 1. Demographic characteristic of the cohort at baseline and follow-up based on the dementia family history and on APOE genotype

	Baseline				Follow-up			
	FHD- (n=107)	FHD+ (n=103)	p (Mann-Whitney U)	FHD- (n=89)	FHD+ (n=99)	p (Mann-Whitney U)		
Age (y)	52.0 ± 11.0	53.0 ± 7.0	0.52	54.0 ± 10.0 55.0 ± 6.0 0		0.22		
Years of Education	16.0 ± 5.0	16.0 ± 5.0	0.34	17.0 ± 5.0	16.0 ± 5.0	0.21		
CAIDE (incl. APOE status)	5.0 ± 4.0 (n=107)	6.0 ± 2.0 (n=101)	<0.05	6.0 ± 4.0 (n=89)	7.0 ± 3.0 (n=93)	< 0.005		
			p (Chi-Square)			p (Chi-Square)		
Sex (%female)	69.2%	71.8%	0.67	69.7%	71.7%	0.76		
APOE ε4 (% Carriers)	28.0% (n=107)	44.6% (n=101)	<0.05	28.1% (n=89)	44.3% (n=97)	<0.05		
	APOE ε4- (n=133)	APOE ε4+ (n=75)	p (Mann-Whitney U)	APOE ε4- (n=118)	APOE ε4+ (n=68)	p (Mann-Whitney U)		
Age (y)	53.0 ± 8.0	52.0 ± 8.0	0.06	55.0 ± 8.0	54.5 ± 8.0	0.08		
Years of Education	16.0 ± 5.0	17.0 ± 4.5	0.51	16.0 ± 4.8	17.0 ± 5.0	0.47		
CAIDE (incl. APOE status)	5.0 ± 3.0 (n=133)	7.0 ± 3.0 (n=75)	<0.0005	6.0 ± 3.0 (n=118)	7.0 ± 4.0 (n=64)	< 0.0005		
CAIDE (excl. APOE status)	5.0 ± 2.0 (n=133)	4.0 ± 3.0 (n=75)	0.12	5.0 ± 2.0 (n=118)	4.5 ± 3.0 (n=64)	0.25		
			p (Chi-Square)			p (Chi-Square)		
Sex (%female)	69.9%	70.7%	0.91	71.2%	69.1%	0.77		

Note: median ± interquartile range (IQR) was reported for continuous variables. Abbreviations: FHD-, Negative family history of dementia;

FHD+, positive family history of dementia; APOE ε4+, Apolipoprotein ε4 genotype positive; APOE ε4-, Apolipoprotein ε4 genotype negative;

CAIDE, Cardiovascular Risk Factors, Aging and Dementia.

Table 2. Statistical summary of the multiple regression model testing the relationship between APOE and episodic and relational memory

			Baseline			Follow-up			
Model summary -		R^2	$F_{(4,201)}$	p	R^2	$F_{(4, 169)}$	p		
		0.16	9.55	< 0.0001	0.17	8.48	< 0.0001		
DV	IV	β	se	p	β	se	p		
	ΑΡΟΕ ε4	0.28	0.13	< 0.05	0.27	0.15	0.07		
Episodic and	Age	-0.02	0.01	0.17	-0.008	0.01	0.55		
relational	Sex	0.27	0.14	0.06	0.25	0.16	0.11		
memory	Years of education	0.09	0.02	<0.0001	0.10	0.02	<0.0001		

Note: unstandardized coefficient (β) and standard error (se) are reported. DV, dependent variable; IV, independent variable; APOE ϵ 4, Apolipoprotein ϵ 4 genotype.

1255

Table 3. Statistical summary of the multiple regression model testing the interaction between CAIDE and menopausal status

	_	Baseline			Follow-up			
Modal s	R^2	$F_{(4, 120)}$	p	R^2	$F_{(4,96)}$	p		
Model summary –		0.07	2.34	0.06	0.06	1.45	0.22	
DV	IV	β	se	p	β	se	p	
Verbal, visuospatial	CAIDE	0.09	0.06	0.18	0.07	0.07	0.35	
functions, and short-	Menopause	1.12	0.58	0.05	1.81	0.87	< 0.05	
term (conjunctive)	CAIDE*Menopause	-0.20	0.09	< 0.05	-0.29	0.13	< 0.05	
memory	Age	-0.04	0.03	0.10	-0.02	0.03	0.52	

Note: unstandardized coefficient (β) and standard error (se) are reported. DV, dependent variable; IV, independent variable; APOE ϵ 4, Apolipoprotein ϵ 4 genotype.

1263

Table 4. Statistical summary of longitudinal linear mixed effect model testing for longitudinal change in the LC–Hippocampus functional connectivity

DV	IV	β	se	p	
rem.	Time	-0.06	0.03	< 0.05	
LC–Hippocampus connectivity	mean FD	0.34	1.42	0.81	
	Hippocampus volume	0.01	0.06	0.87	


Note: unstandardized coefficient (β) and standard error (se) are reported. DV, dependent variable; IV, independent variable; FD, framewise displacement; LC, Locus Coeruleus

1264

Table 5. Regression coefficients of the model testing the relationship between CAIDE, LC–Hippocampus and verbal, visuospatial functions, and short-term (conjunctive) memory

Model summary		Baseline			Follow-up		
		R^2	$F_{(4, 121)}$	p	R^2	$F_{(4,77)}$	p
		0.07	2.22	0.07	0.07	1.55	0.20
DV	IV	β	se	p	β	se	p
Verbal, visuospatial functions, and short- term (conjunctive) memory	CAIDE	-0.09	0.03	< 0.01	-0.04	0.05	0.45
	LC-HC	-0.10	0.34	0.78	0.48	0.53	0.37
	CAIDE × LC–HC	-0.10	0.14	0.48	-0.54	0.22	< 0.05
	Hippocampus volume	-1.97E-04	2.29E-04	0.39	5.39E-06	3.26E-04	0.99

Note: unstandardized coefficient (β) and standard error (se) are reported. CAIDE and LC-HC have been centred in this model. DV, dependent variable; IV, independent variable; CAIDE, Cardiovascular Risk Factors, Aging and Dementia; LC-HC, Locus Coeruleus–Hippocampus functional connectivity.



Participants

Bandpass filtering

