

1 **New primer sets to detect recent human adenovirus F41 variants in wastewater: Is it**  
2 **linked to the new acute hepatitis?**

3 Mariel Perez-Zabaleta<sup>a\*</sup>, Cecilia Williams<sup>b</sup>, Zeynep Cetecioglu<sup>a</sup>

4 a Department of Industrial Biotechnology, KTH Royal Institute of Technology, AlbaNova  
5 University Center, SE-10691, Stockholm, Sweden.

6 b Department of Protein Science, KTH Royal Institute of Technology, Science for Life  
7 Laboratory, Solna, Sweden.

8

9 \*Corresponding author: [mnpz@kth.se](mailto:mnpz@kth.se) (M. Perez-Zabaleta)

10

11 **Abstract**

12 Human adenovirus type F-41 has been pursued as one of the potential reasons for the  
13 new acute hepatitis cases of unknown cause in young children. Tracking the spread of this  
14 virus in the population using wastewater-based epidemiology tools can help clinical  
15 investigations to determine its relation to this new hepatitis outbreak.

16 In this study, methods to detect human adenovirus type F (40 and 41) and specifically  
17 type F41 were designed and implemented to quantify the amount of these pathogens in  
18 wastewater samples from Stockholm, Sweden. An assay based on reverse transcriptase  
19 quantitative polymerase chain reaction using TaqMan technology and primers targeting the  
20 three main capsid genes of adenoviruses: hexon, penton and fiber, was designed. The hexon  
21 primers were specific to adenovirus F41, while fiber primers could quantify both  
22 adenoviruses, F40 and F41. Wastewater samples from Stockholm were used to validate the  
23 designed assay and, in addition, pepper mild mottle virus (PMMoV) levels were quantified to  
24 study the data normalization.

25 Our results can help link the occurrence of the virus variant with new cases of acute  
26 hepatitis and contribute to a better understanding of the possible causes. It can also provide  
27 valuable information that can be used in future investigations on the monitoring of human  
28 adenovirus type F in wastewater.

29 **Keywords:**

30 Human adenovirus F41 (HAdV-F41), human adenovirus F40 (HAdV-F40), wastewater-based  
31 epidemiology (WBE), acute hepatitis, pepper mild mottle virus (PMMoV).

## 32        **1 Introduction**

33        Human adenoviruses are nonenveloped DNA viruses belonging to the family  
34 Adenoviridae. These viruses are classified into seven species (A to G) and at least 104  
35 genotypes (HAdV Working Group, 2022). Adenoviruses can cause respiratory illness,  
36 gastroenteritis, conjunctivitis, or cystitis in young children (Ikner and Gerba, 2017). There is  
37 no specific treatment for adenovirus infections. Adenovirus type F (40 and 41) usually causes  
38 acute gastroenteritis in children. However, cases of hepatitis have been reported in  
39 immunocompromised children associated with adenovirus type 41 (HAdV-F41) infection  
40 (Ikner and Gerba, 2017).

41        A sudden increase in acute hepatitis cases of unknown cause was reported to the World  
42 Health Organization (WHO) in 2022. Between April 5 and July 8, 2022, 1 010 probable cases  
43 and 22 deaths were reported by 35 countries, including five pediatric patients with severe  
44 hepatitis of unknown origin identified at a children’s hospital in Alabama, U.S., in early  
45 October 2021 (WHO, 2022). Of the 1 010 probable cases, 5% of the children required a liver  
46 transplant, and almost half of the cases (48%) were reported from European countries with  
47 the UK accounting for the highest number. The children were under 16 years old, presented  
48 with liver inflammation (including some with liver failure), and tested negative for hepatitis  
49 A to E (WHO, 2022). The cases were not associated with travel, and most children tested  
50 positive for adenovirus, in which type 41 was predominant. Of 251 cases tested for  
51 adenovirus in the UK, 68% gave positive results against this pathogen (UK Health Security  
52 Agency, 2022). In Alabama, HAdV-F41 was detected in all five infected patients (Baker et  
53 al., 2021). A recent investigation has pointed to the synergism between HAdV-F41 and  
54 SARS-CoV-2 as responsible to activate higher levels of inflammation, which eventually can  
55 cause acute hepatitis in children (Luan et al., 2022).

56 Wastewater-based epidemiology (WBE) is a useful tool to detect the occurrence of  
57 viruses in a population, as viruses do not possess the ability to grow outside their host and  
58 their concentration in wastewater can be representative of the infection rate (Xagorarakis and  
59 O’Brien, 2020). The wastewater-based epidemiology concept was initially perceived in 2001  
60 (Daughton C.G. and Jones-Lepp T.L., 2001) and applied to grasp the circulation of viruses in  
61 previous outbreaks such as poliovirus in the Netherlands or hepatitis A and norovirus in  
62 Sweden (Hellmér et al., 2014). Simultaneous detection of different adenoviruses genotypes  
63 was previously investigated to determine the removal efficiency of “generic” adenoviruses  
64 through the wastewater treatment process (Carducci et al., 2008; Hata et al., 2012; Quintão et  
65 al., 2021). Few studies were able to specifically identify HAdV-F41 (Heim et al., 2003; Kuo  
66 et al., 2009; Tiemessen and Nel’, 1996), and previously designed primers to detect HAdV-  
67 F41 can be less specific for emergent variants. According to Götting and colleagues (2022),  
68 HAdV-F41 has evolved significantly over the past decade, which possibly has resulted in  
69 altered virulence of this pathogen. The emergence of a novel strain of HAdV-F41 could be  
70 suspected, which may be causing hepatitis. Unfortunately, recovery of high-quality HAdV-  
71 F41 genomes from hepatitis-affected children has failed due to low virus loads (Kajon and St  
72 George, 2022).

73 In relation to the genetic diversity of adenoviruses, sequencing from patient samples  
74 revealed that the genes of hexon, fiber and penton include hypervariable regions that can be  
75 used for genotyping (Risso-Ballester et al., 2016). Fiber, penton, and hexon as the main  
76 capsid proteins of adenoviruses are responsible for binding to the host cell. (Larson et al.,  
77 2015).

78 In this study, the gene sequences of these three major capsid genes, hexon, penton and  
79 fiber, were used to design the primers and probes for the quantification of adenovirus type F  
80 by RT-qPCR. The difference between the specificity of the primers for different variants was

81 tested. Wastewater samples from Stockholm were used to validate the designed methods.  
82 Contents of human adenovirus F40 (HAdV-F40) and HAdV-F41 were detected in the  
83 wastewater samples. In addition, pepper mild mottle virus (PMMoV) was also determined in  
84 the samples and studied for data normalization.

## 85 2 Materials and Methods

### 86 2.1 Design of primers to detect human adenovirus type F and in silico validation

87 Primers and probes targeting the main capsid genes of adenovirus: hexon, penton and  
88 fiber, were designed using recently published genome sequences of HAdV-F41 variants  
89 current in 2021 and 2022: MW567962.1 (March 2021), ON532826.1 (May 2022) and  
90 OP174922.1 (August 2022). The primers and probes are listed in Table 1.

91 **Table 1. List of primer sequences used to quantify Human Adenovirus type F**

Region	Primer	Sequence 5'--> 3'
Hexon	HVR_fw	CCAAATTACTGCTTTCCGCTG
	HVR_Probe	CAGTCCAGTTTCCGCTGTTTGCTTT
	HVR_rev	ACTCAATTTCTGCCCTCTG
Penton	Penton_fw	TGAGTTTACCTTGCCCGAAG
	Penton_Probe	CGCCGTTTTGTCTTCCCCTTCC
	Penton_rev	TCAAACCTTCACCCCTATGTCG
Fiber	Fiber_fw	TCTACCCCTATGACACCTTCTC
	Fiber_Probe	CCTATGTAGCTCCGCCCTTCGTTT
	Fiber_rev	TTTCCTGTAACCCGTCAGAAG

92  
93 The primers were analyzed in silico using Basic Local Alignment Tool (BLAST)  
94 from the National Center for Biotechnology Information (NCBI) and the sequences of the  
95 three primer sets were confirmed to match HAdV-F41. Then, the primer sequences were  
96 aligned against twenty HAdV-F41 species and two HAdV-F40 species from NCBI GenBank.  
97 The alignments were performed using DNASTAR Bioinformatics Software. The twenty  
98 HAdV-F41 genomes were selected according to a recent phylogenetic study, which indicated

99 that HAdV-F41 evolved with three lineages (Götting et al., 2022). This recent investigation  
100 analyzed 65 genomes from which Lineage 1 included HAdV-F41 prototype strain Tak from  
101 1973 (GenBank DQ315364) and six isolates from 2007 to 2017. Lineage 2 included 53  
102 isolates of the 65 genomes studied and strains from 2000 to 2022. Lineage 3 included 6  
103 isolates and this group was closer to the prototype strain HAdV-F40 Dugan, isolated in 1979.  
104 The selected HAdV-F41 genomes were from lineage 1 and 2. In lineage 1, we included the  
105 Tak strain, while lineage 2 included HAdV-F41 genomes used in this investigation  
106 (MW567962.1, ON532826.1, and OP174922.1) to design the primers/probes.

## 107 **2.2 In vitro validation of primers to detect human adenovirus type F**

108 The hexon, penton, and fiber primers were evaluated against thirteen genotypes of  
109 human adenovirus for species A, B, C, D and E (Table 3). In addition, three positive controls  
110 of adenovirus type F were tested. Two variants of HAdV-F41, one prototype strain Tak  
111 (GenBank: DQ315364.2) and one strain isolated from a Swedish patient (GenBank:  
112 KX868523.2) (Lá szló Kajá et al., 2018), and a positive control for HAd-V-F40 (the  
113 prototype strain HoviX, GenBank: KU162869.1) were also tested. The adenovirus species  
114 were provided by the Norrlands Universitetssjukhus (Klinisk Mikrobiologi, Umeå, Sweden).  
115 Adenovirus controls were heat inactivated at 56°C for 30 min and then, 200 µL of each  
116 culture was used for DNA extraction using Maxwell RSC Enviro TNA Promega Kit and  
117 Maxwell RSC Instrument (Promega Biotech AB, Sweden) following the manufacturer's  
118 instructions.

## 119 **2.3 Reverse transcriptase quantitative polymerase chain reaction (RT-qPCR)** 120 **method using the designed primer sets**

121 RT-qPCR assays were performed using 5 µL of TaqMan Reliance One-Step  
122 Multiplex Supermix (BioRad, Cat # 12010221). 1 µL of 20 mg/ml Bovine Serum Albumin

123 (BSA) (Thermo Scientific, Cat # B14) was added to reduce PCR inhibitors and enhance  
124 efficacy, for a final reaction volume of 20  $\mu$ L. For each reaction, 5  $\mu$ L of DNA template was  
125 used. Nuclease-free water and DNA extracted from tap water were included as negative  
126 controls for all qPCR reactions. Thermal cycling (50 °C 10 min, 95 °C 10 s, followed by  
127 45 cycles of 95 °C 3 s, 60 °C for 30 s) was performed on an Applied Biosystems QuantStudio  
128 3 RT-qPCR machine (ThermoFisher Scientific). Reactions were considered positive if the  
129 cycle threshold (Ct) was below 42 cycles. Amplifications were carried out on the FAM  
130 channel.

131 PMMoV was also quantified in the samples to study data normalization. The PMMoV  
132 primers used were forward: 5'-GAGTGGTTTGACCTTAACGTTTGA-3' reverse: 5'-  
133 TTGTCGGTTGCAATGCAAGT-3' and probe 5'-CCTACCGAAGCAAATG-3' (Zhang et  
134 al., 2006).

135 Standard curves were created using constructed plasmids containing appropriate  
136 regions for hexon, penton, fiber, and PMMoV (IDT, Custom MiniGene 25-500 bp).

## 137 **2.4 Validation of primer sets in wastewater samples**

### 138 ***2.4.1 Wastewater samples***

139 The validation of the primer sets was performed in wastewater samples from Stockholm,  
140 which is the capital and largest city of Sweden and accounts for a population of approx. 2.4  
141 million people in the metropolitan area. The wastewater samples were collected from the  
142 three main municipal wastewater treatment plants (WWTP): Stockholm Vatten och Avfall  
143 Bromma, Stockholm Vatten och Avfall Henriksdal, and Käppala WWTP. Bromma WWTP  
144 treats wastewater from approximately 377 500 inhabitants and samples were obtained from  
145 three inlets (corresponding to three regions: Hässelby, Riksby, and Järva). The second  
146 analyzed WWTP, Henriksdal, treats wastewater from 862 100 inhabitants and has two inlets,

147 Sickla and Henriksdal. Käppala WWTP covers a region of 700 000 inhabitants and has one  
148 inlet.

149 Approximately 500 ml of wastewater samples from each inlet were taken over a period of  
150 24 hours from Monday to Tuesday, using a flow-proportional composite sampler.  
151 Wastewater samples were transported to the laboratory on ice and kept at 4°C until further  
152 analyses. After receiving the samples, two representative samples for the whole Stockholm  
153 region were attained by mixing the different inlets according to the percentage of flow rates  
154 of each region (Table 2). The representative samples from Stockholm were further  
155 concentrated and analyzed.

156 **Table 2. Flow rates and inhabitants of the WWTPs from Stockholm. Average flow rates**  
157 **were calculated using historical data of flow rates from April 2020 to June 2022.**

WWTP	Inlet	Average flow rate (m <sup>3</sup> /s)	Inhabitants	Flow rate (%)
<b>Henriksdal</b>	Henriksdal	1.52	371800	23%
	Sickla	1.84	490200	28%
<b>Bromma</b>	Hässelby	0.44	125800	7%
	Järva	0.78	197200	12%
	Riksby	0.24	37500	4%
<b>Käppala</b>	Käppala	1.76	700000	27%

158

#### 159 ***2.4.2 Wastewater concentration and DNA extraction***

160 Wastewater viral concentration and total nucleic acid (TNA) extraction were usually  
161 performed the same or the next day after receiving the samples. Concentration and TNA  
162 extraction were performed using Maxwell RSC Enviro TNA Promega Kit and Maxwell RSC  
163 Instrument (Promega Biotech AB, Sweden). High concentrations and yields were reported  
164 when this kit has been used in wastewater samples (Perez-Zabaleta et al., 2022), (Isaksson et  
165 al., 2022). Briefly, 40 ml of wastewater was treated with a protease solution and centrifuged

166 at 3000 g for 10 min to remove proteins and solids present in the wastewater. Then, the  
167 supernatant was filtrated, eluted to 500 L using a column-based system, and then loaded into  
168 a cartridge provided by the kit. Maxwell RSC Pure Food GMO program was selected in the  
169 instrument software for automatic TNA extraction and eluted in 80 µL nuclease-free water.  
170 Two independent replicates were analyzed for each sample and tap water was used as a  
171 negative control for viral concentration and DNA extraction steps.

#### 172 *2.4.3 Sanger sequencing of PCR amplicons from wastewater samples*

173 The PCR amplicons from the wastewater samples obtained after RT-qPCR were purified  
174 using GeneJET PCR Purification Kit (ThermoFisher Scientific, Cat # K0701) following the  
175 manufacturer's instructions. Two pure PCR products per sample were sent for sequencing,  
176 one PCR product was sequenced in the forward direction and the other one in the reverse  
177 direction. Sanger sequencing was performed by Eurofins using Mix2Seq Kit.

178 To determine the HAdV genotypes in the wastewater samples, hexon, penton and fiber  
179 sequences obtained by Sanger sequencing were compared with sequences in the NCBI  
180 databases by using BLAST.

### 181 **3 Results**

#### 182 **3.1 Primers and probes do not amplify a panel of related human adenoviruses**

183 The primers and probes were designed to be specific to adenovirus type F. To test  
184 their specificity, different adenovirus species (A, B, C, D and E) were used as negative  
185 controls. In total, thirteen adenovirus genotypes were analyzed (Table 3). The three sets of  
186 primers (hexon, penton and fiber) gave negative results for species A to E. Thus, they should  
187 not detect these species.

188

189

190

191

192

193

194

195 **Table 3. Specificity test of hexon, penton and fiber primer-sets toward various human**  
196 **adenovirus species.**

Adenovirus Control	Species	Classification	RT-qPCR results		
			Hexon	Penton	Fiber
Adenovirus 31	A	A31	Negative	Negative	Negative
Adenovirus 3	B1	B3	Negative	Negative	Negative
Adenovirus 7	B1	B7	Negative	Negative	Negative
Adenovirus 11	B2	B11	Negative	Negative	Negative
Adenovirus 14	B2	B14	Negative	Negative	Negative
Adenovirus 34	B2	B34	Negative	Negative	Negative
Adenovirus 1	C	C1	Negative	Negative	Negative
Adenovirus 2	C	C2	Negative	Negative	Negative
Adenovirus 5	C	C5	Negative	Negative	Negative
Adenovirus 9	D	D9	Negative	Negative	Negative
Adenovirus 13	D	D13	Negative	Negative	Negative
Adenovirus 20	D	D20	Negative	Negative	Negative
Adenovirus 4	E	E4	Negative	Negative	Negative

197


### 198 **3.2 Primers and probes amplify adenovirus type F**

199 Next, the primers and probes were tested for adenovirus type F. One positive control  
200 for HAdV-F40 (strain HoviX) and two variants of HAdV-F41 (strain Tak and strain isolated  
201 from a Swedish patient) were analyzed and used to validate the primer sets in vitro.

202 Two primer sets (fiber and penton) detected the HAdV-F40 (strain HoviX). The  
203 amount detected by the fiber primer-set was almost 8000 times higher than the detected by

204 the penton primers. Hexon primers gave negative results, showing no specificity towards  
205 HAdV-F40 (Figure 1a). The amount of gene copy numbers of HAdV-F40 (strain HoviX) per  
206  $\mu\text{L}$  was significantly different ( $P$ -value = 0.0007) among the three primer sets. For adenovirus  
207 F41 (strain Tak), the penton or fiber primer sets detected similar amounts and were not  
208 significantly different ( $P$ -value = 0.30, Figure 1b). The primers targeting the hexon region  
209 could not detect this strain either. Alignments of the genome of HAdV-F41 strain Tak and  
210 hexon primers showed that forward and reverse primers could bind the hexon region but the  
211 designed probe (HVR\_probe) had in total seven mismatches in the nucleotides, in which four  
212 of them were consecutive. However, the HAdV-F41 variant previously isolated from a  
213 Swedish patient was detected by all three primer sets, which quantified similar amounts of  
214 the virus, with no significant difference ( $P$ -value = 0.35, Figure 1c).

215



216

217 **Figure 1. Analysis of positive controls for human adenovirus F40 and F41 using**  
218 **primers-sets for hexon, penton and fiber regions.** Detection and gene copy numbers (per  
219  $\mu\text{L}$ ) of HAdV-F40 and F41 were determined by qPCR analysis of the hexon, penton or fiber  
220 region in three controls (a) Prototype HAdV-F40 HoviX strain (b) Prototype HAdV-F41 Tak  
221 strain (c) and an HAdV-F41 strain isolated from a patient in Sweden.

222 The validation of primers in silico and in vitro showed that hexon primers are  
223 selective for recent HAdV-41 strains evolved in lineage 2. Fiber and penton primers can be  
224 used to detect both HAdV-40 and HAdV-41 variants and detect the HAdV-41 Swedish

225 variant at the same sensitivity as hexon primers (but are not selective for this variant).  
226 Comparing penton and fiber primers, penton is less sensitive to HAdV-40 than fiber primers,  
227 however, both primer sets can detect similar amounts of both HAdV-41 strains (Figure 1b  
228 and Figure 1c).

### 229 **3.3 Validation of primer sets using wastewater samples**


#### 230 ***3.3.1 Quantification of Human Adenovirus type F***

231 Next, the primer sets were tested for the detection of human adenovirus F in  
232 Stockholm wastewater. Wastewater samples from six inlets (three WWTPs) were mixed  
233 proportionally to the flow rates to obtain a representative Stockholm sample. Two biological  
234 replicates were analyzed by RT-qPCR with the three primer sets (hexon, penton and fiber) as  
235 well as with PMMoV primer set, and quantified. The results obtained from the wastewater  
236 samples showed that primers targeting the hexon region detected lower amounts of  
237 adenovirus compared to penton or fiber primers (Figure 2). These results are in line with the  
238 outcomes obtained in the specificity study (section 3.2) since hexon primers were more  
239 specific to HAdV-41 lineage 2 and did not detect HAdV-F40 or HAdV-F41 Tak strain. Fiber  
240 primers detected higher levels of adenovirus compared to penton primers in all the samples  
241 (Figure 2b). In all the analyzed samples, the fiber and penton assays detected the same trends  
242 of increasing or decreasing amounts of virus, whereas the hexon primers detect a different  
243 trend (Figure 2b). The fiber and penton assays also both detected about 100 times more viral  
244 DNA copies (around  $10^{17}$ ) than the hexon primer ( $10^{15}$ ). This suggests that the wastewater  
245 samples contain a mixture of HAdV-F strains (detected by fiber and penton) and strains  
246 similar to the HAdV-F41 Swedish isolate (detected selectively by the hexon assay) make up  
247 1% or less of the circulating HAdV-F40 and 41 strains. However, this estimation is uncertain

248 as it cannot account for potential variations in the sequences of the primer and probe sites

249 among developing variants which would impact amplification efficacy or detection.

250



251

252 **Figure 2. Quantification of HAdV- F40 and F41 using primers-sets for hexon, penton**

253 **and fiber regions in wastewater samples.** Gene copy numbers of HAdV-F41 and F40 per

254  $\mu\text{L}$  were determined by targeting (a) hexon region (purple), (b) penton (pink) and fiber

255 (green) regions. Two biological replicates were analyzed for each sampling point.

256

### 3.3.2 Sanger sequencing of PCR products from wastewater samples


257 To determine the specific HAdV genotypes detected by the assays in the wastewater  
258 samples, the amplified hexon, penton and fiber PCR products were analyzed by Sanger  
259 sequencing. The obtained sequences were compared in the NCBI databases by BLAST. The  
260 DNA sequences from the PCR products analyzed by hexon primers were specific for HAdV-  
261 F41, which confirms that the hexon method only detects adenovirus F41. In the case of  
262 penton and fiber, the DNA sequences were not specific to HAdV-F41 and a double detection  
263 of HAdV-F40 and HAdV-F41 can be suspected.

### 264 *3.3.3 Normalization of human adenovirus type F detection to populations*

265 The size of a population in a specific area is a variable factor difficult to predict. The  
266 PMMoV is an RNA virus that is abundant in human faeces and has been reported to be a  
267 good population indicator in wastewater samples (Kitajima et al., 2018; Zhang et al., 2006).  
268 Furthermore, in our previous investigations of SARS-CoV-2 in wastewater, PMMoV was  
269 found to be an effective normalization factor to correct for fluctuations in the wastewater  
270 content (Perez-Zabaleta et al., 2022). However, the normalization of adenovirus to PMMoV  
271 has not yet been explored.

272 Comparing our data for HAdV-F with and without normalization to PMMoV, it can  
273 be observed that the trends of increasing or decreasing levels are different (Figure 3  
274 compared to Figure 2). The PMMoV-normalized data show more constant values among the  
275 analyzed weeks for the three assays (hexon, penton and fiber primer sets, Figure 3). While  
276 non-normalized data indicated a sudden peak of HAdV-F41 (Stockholm variant) in week 21,  
277 the normalized data show a higher level during spring (week 14-21) that decreases toward the  
278 end of May. Similarly, the overall HAdV-F40 and 41 variants appear to decrease during May.  
279 As adenoviruses are more common in the Scandinavian regions during winter and early  
280 spring, this may suggest that PMMoV normalization can help to decrease fluctuation in the

281 measurement due to population size. However, to corroborate this, the trends should be  
282 correlated to the actual number of clinical cases. Unfortunately, a such comparison was not  
283 possible since the number of infected persons by adenovirus type F per week in Stockholm is  
284 unknown.



285

286 **Figure 3. Normalization of HAdV-F40 and F41 in wastewater samples with respect to**  
287 **PMMoV.** Samples were analyzed using primers sets for hexon, penton and fiber regions.  
288 PMMoV content was determined and used to normalize the data. The amount of adenovirus  
289 is presented as (a) hexon (purple) gene copy numbers per PMMoV gene copy number \*100,

290 (b) penton (pink) or fiber (green) gene copy numbers per PMMoV gene copy number \*100.

291 Two biological replicates were analyzed for each sampling point.

## 292 **Conclusion**

- 293 • Hexon primer sets are more specific to HAdV-41, especially from lineage 2.
- 294 • Fiber primer sets can detect both HAdV-40 and HAdV-41.
- 295 • PMMoV may be appropriate for the normalization of HAdV measurements in  
296 wastewater but further studies are needed.
- 297 • Routine sequencing of PCR products from wastewater could provide detailed  
298 information on HAdV variant spread.

## 299 **Author approval**

300 All authors interpreted the data, revised the manuscript, and approved the final version of  
301 the manuscript.

## 302 **Competing interests**

303 All authors declare no competing interests.

## 304 **Funding**

305 This research was supported by SciLifeLab, Pandemic Laboratory Preparedness (VC-2021-  
306 0033), and the Knut and Alice Wallenberg Foundation (KAW 2020.0241, V-2020-0699).

## 307 **Acknowledgements**

308 The authors would like to thank the Science for Life Laboratories Environmental Virus  
309 Profile Platform, Swedish Environmental Epidemiology Center (SEEC), the Kappala  
310 Association, and Stockholm Vatten och Avfall. Additionally, the authors acknowledge SEEC  
311 collaborators for project management, and Annika Alland from Norrlands

312 Universitetssjukhus, Klinisk Mikrobiologi, Umeå, Sweden for providing positive controls of  
313 different adenovirus genotypes and scientific discussions.

## 314 **References**

- 315 Baker, J.M., Buchfellner, M., Britt, W., Sanchez, V., Potter, J.L., Ingram, ; L Amanda, Shiau, H.,  
316 Luz, ;, Gutierrez Sanchez, H., Saaybi, S., Kelly, D., Lu, X., Vega, E.M., Ayers-Millsap, S.,  
317 Willeford, W.G., Rassaei, N., Bullock, H., Reagan-Steiner, S., Martin, A., Moulton, E.A.,  
318 Lamson, D.M., Kirsten, ;, George, S., Umesh, ;, Parashar, D., Hall, A.J., Macneil, A., Tate,  
319 J.E., Kirking, H.L., 2021. MMWR, Acute Hepatitis and Adenovirus Infection Among  
320 Children — Alabama, October 2021–February 2022.  
321 <https://doi.org/10.1128/CMR.5.3.262>
- 322 Carducci, A., Morici, P., Pizzi, F., Battistini, R., Rovini, E., Verani, M., Carducci, A.A., 2008.  
323 Study of the viral removal efficiency in a urban wastewater treatment plant. *Water*  
324 *Science & Technology*. <https://doi.org/10.2166/wst.2008.437>
- 325 Daughton C.G., Jones-Lepp T.L., 2001. Pharmaceuticals and care products in the  
326 environment: Scientific and regulatory issues. American Chemical Society.
- 327 Götting, J., Cordes, A.K., Steinbrück, L., Heim, A., 2022. Molecular Phylogeny of human 1  
328 adenovirus type 41 lineages 2 3. <https://doi.org/10.1101/2022.05.30.493978>
- 329 HAdV Working Group, 2022. Human Adenovirus genotypes [WWW Document]. URL  
330 <http://hadvwg.gmu.edu/> (accessed 9.7.22).
- 331 Hata, A., Kitajima, M., Katayama, H., 2012. Occurrence and reduction of human viruses, F-  
332 specific RNA coliphage genogroups and microbial indicators at a full-scale wastewater  
333 treatment plant in Japan. <https://doi.org/10.1111/jam.12051>
- 334 Heim, A., Ebnet, C., Harste, G., Pring-Åkerblom, P., 2003. Rapid and quantitative detection of  
335 human adenovirus DNA by real-time PCR. *J Med Virol* 70, 228–239.  
336 <https://doi.org/10.1002/jmv.10382>
- 337 Hellmér, M., Paxéus, N., Magnius, L., Enache, L., Arnholm, B., Johansson, A., Bergström, T.,  
338 Norder, H., 2014. Detection of Pathogenic Viruses in Sewage Provided Early Warnings  
339 of Hepatitis A Virus and Norovirus Outbreaks. *Appl Environ Microbiol* 80, 6771–6781.  
340 <https://doi.org/10.1128/AEM.01981-14>
- 341 Ikner, L.A., Gerba, C.P., 2017. Adenoviruses, in: Quah, S.R. (Ed.), *International Encyclopedia*  
342 *of Public Health (Second Edition)*. Academic Press, Oxford, pp. 5–9.  
343 <https://doi.org/https://doi.org/10.1016/B978-0-12-803678-5.00004-7>
- 344 Isaksson, F., Lundy, L., Hedström, A., Székely, A.J., Mohamed, N., 2022. Evaluating the use  
345 of alternative normalization approaches on SARS-CoV-2 concentrations in wastewater:  
346 experiences from two catchments in northern Sweden.  
347 <https://doi.org/10.3390/environments9030039>
- 348 Kajon, A.E., St George, K., 2022. Mysterious cases of acute hepatitis in children: is  
349 adenovirus still a lead suspect? <https://doi.org/10.1080/22221751.2022.2095933>
- 350 Kitajima, M., Sassi, H.P., Torrey, J.R., 2018. Pepper mild mottle virus as a water quality  
351 indicator 1, 19. <https://doi.org/10.1038/s41545-018-0019-5>
- 352 Kuo, D.H.W., Simmons, F., Xagorarakis, I., 2009. A new set of PCR assays for the identification  
353 of multiple human adenovirus species in environmental samples. *J Appl Microbiol* 107,  
354 1219–1229. <https://doi.org/10.1111/j.1365-2672.2009.04300.x>

- 355 Lá szló Kajá, G.I., Lipiec, A., niel Bartha, D., Allard, A.I., Arnberg, N., 2018. A multigene typing  
356 system for human adenoviruses reveals a new genotype in a collection of Swedish  
357 clinical isolates. <https://doi.org/10.1371/journal.pone.0209038>
- 358 Larson, C., Oronsky, B., Scicinski, J., Fanger, G.R., Stirn, M., Oronsky, A., Reid, T.R., 2015.  
359 Going viral: a review of replication-selective oncolytic adenoviruses.  
360 <https://doi.org/10.18632/oncotarget.5116>
- 361 Luan, J., Zhou, Y., Wang, X., Ming, A., Liu, Y., Yuan, Y., Liang, J., Zhang, L., 2022. Single-cell  
362 RNA sequencing analysis of liver reveals the enhanced entry and release abilities of  
363 human adenovirus F41, partially explaining acute hepatitis in children. *Journal of*  
364 *Infection* 85, 334–363. <https://doi.org/10.1016/j.jinf.2022.06.007>
- 365 Perez-Zabaleta, M., Archer, A., Khatami, K., Jafferli, M.H., Nandy, P., Atasoy, M., Birgersson,  
366 M., Williams, C., Cetecioglu, Z., 2022. Long-term SARS-CoV-2 surveillance in the  
367 wastewater of Stockholm: what lessons can be learned from the Swedish perspective?  
368 Quintão, T.S.C., Silva, F G, Pereira, A L, Araújo, W N, Oliveira, P M, Souza, M B L D,  
369 Lamounier, T.A., Haddad, R, 2021. Detection and molecular characterization of enteric  
370 adenovirus in treated wastewater in the Brazilian Federal District 3, 691.  
371 <https://doi.org/10.1007/s42452-021-04678-2>
- 372 Risso-Ballester, J., Cuevas, J.M., Sanjuá, R., 2016. Genome-Wide Estimation of the  
373 Spontaneous Mutation Rate of Human Adenovirus 5 by High-Fidelity Deep Sequencing.  
374 <https://doi.org/10.1371/journal.ppat.1006013>
- 375 Tiemessen, C.T., Nel, M.J., 1996. Detection and typing of subgroup F adenoviruses using the  
376 polymerase chain reaction, *Journal of Virological Methods ELSEVIER Journal of*  
377 *Virological Methods*.
- 378 UK Health Security Agency, 2022. Investigation into acute hepatitis of unknown aetiology in  
379 children in England.
- 380 WHO, 2022. Severe acute hepatitis of unknown aetiology in children - Multi-country [WWW  
381 Document]. URL [https://www.who.int/emergencies/disease-outbreak-](https://www.who.int/emergencies/disease-outbreak-news/item/2022-DON400)  
382 [news/item/2022-DON400](https://www.who.int/emergencies/disease-outbreak-news/item/2022-DON400) (accessed 9.7.22).
- 383 Xagorarakis, I., O'Brien, E., 2020. Wastewater-Based Epidemiology for Early Detection of Viral  
384 Outbreaks. [https://doi.org/10.1007/978-3-030-17819-2\\_5](https://doi.org/10.1007/978-3-030-17819-2_5)
- 385 Zhang, T., Breitbart, M., Heng Lee, W., Run, J.-Q., Lin Wei, C., Wee Ling Soh, S., Hibberd,  
386 M.L., Liu, E.T., Rohwer, F., Ruan, Y., 2006. RNA Viral Community in Human Feces:  
387 Prevalence of Plant Pathogenic Viruses. *Plosbiology* 4.  
388 <https://doi.org/10.1371/journal.pbio.0040003>  
389