- Title: Additional Evidence Implicating GPD1L in the Pathogenesis of Brugada Syndrome in A Large Multi-1
- generational Family 2
- 3 Brief title: GPD1L Mutation and Brugada Syndrome 4
- 5 Authors: 6
- 1. Greiner, Alexander M., B.A.^{a,b,c}
- 2. Mehdi, Haider, Ph.D.^{a,b}
- 3. Cevan, Chloe^a
- 4. Gutmann, Rebecca, B.S.N.^a
- 10 5. London, Barry, M.D., Ph.D.^{a,b,c}
- 11 12

14

15

16

17

18 19

20

7

8

9

- Department and institutional affiliations: 13
 - A Division of Cardiovascular Medicine, Department of Internal Medicine, University of Iowa Carver College of Medicine, Iowa City, IA, United States of America
 - B Abboud Cardiovascular Research Center, University of Iowa Carver College of Medicine, Iowa City, IA, United States of America
 - C Interdisciplinary Graduate Program in Genetics, University of Iowa, Iowa City, IA, United States of America
- Funding: This work was supported by R01HL062300, R01HL077398, and R01HL115955. Mr. Greiner is 21 also supported by an NIH Ruth L. Kirschstein F30 fellowship (F30HL143908) and by a T32 training grant 22 held by the University of Iowa Medical Scientist Training Program (T32GM007337). 23
- 24
- Conflicts of Interest: The authors have nothing to disclose. 25
- 26
- Word count: 3379 27
- Address for correspondence: 28
- Barry London, M.D., Ph.D. 29
- 285 Newton Road 30
- 2283 Carver Biomedical Research Building 31
- Iowa City, Iowa 52242 32
- United States of America 33
- Fax: 319-353-6343 34
- Telephone: 319-356-2750 35
- e-mail address: barry-london@uiowa.edu 36

37

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

38 Abstract

39 Background

⁴⁰ Brugada Syndrome (BrS) is an inherited arrhythmia syndrome in which mutations in SCN5A account for

 $_{41}$ 20% of cases. Mutations in other ion channels or channel-modifying genes may account for an additional

 $_{42}$ 10% of cases, though recent analysis has suggested that SCN5A should be regarded as the sole monogenic

⁴³ cause of BrS.

44 Objective

⁴⁵ We sought to re-assess the genetic underpinnings of BrS in a large mutligenerational family with a putative ⁴⁶ GPD1L-A280V mutation.

47 Methods

Fine linkage mapping was performed in the family using the Illumina Global Screening array. Whole exome sequencing of the proband was performed to identify rare variants and mutations, and Sanger sequencing

⁵⁰ was used to assay previously-reported risk single nucleotide polymorphsims (SNPs) for BrS.

51 Results

Linkage analysis decreased the size of the previously-reported microsatellite linkage region to ~3 megabases. GPD1L-A280V was the only rare coding non-synonymous variation present at <1% allele frequency in the proband within the linkage region. Other variants were either synonymous, or in genes not known to play a role in BrS and that failed to co-segregate with BrS in the large family. Risk SNPs known to predispose to BrS were overrepresented in affected members of the family.

57 Conclusion

Together, our linkage and sequencing data suggest GPD1L-A280V remains the most likely cause of BrS in

⁵⁹ this large mutligenerational family. While care should be taken in interpreting variant pathogenicity given

⁶⁰ the genetic uncertainty of BrS, our data support inclusion of other putative BrS in clinical genetic panels.

⁶¹ Keywords: inherited, arrhythmia, genetics, exome sequencing, sudden death

62

⁶³ Abbreviations: BrS: Brugada Syndrome, SNP: Single nucleotide polymorphism, CM: centimorgan, LOD:
⁶⁴ logarithm of the odds, Mb: megabase, ICD: implantable cardioverter defibrillator, RBBB: right bundle
⁶⁵ branch block, VT: ventricular tachycardia

Introduction 1 67

Brugada Syndrome (BrS) is an inherited arrhythmia syndrome characterized by ST-segment elevation in the 68 right precordial leads (V_1 through V_3) and sudden cardiac death.¹ The molecular mechanism underpinning 69 BrS was initially described as autosomal-dominant loss-of-function mutations in the main cardiac sodium 70 channel, $Na_V 1.5$, encoded on chromosome 3 by SCN5A.² These loss-of-function mutations result in decreased 71 inward depolarizing sodium current, which can result in premature repolarization of the epicardium in the 72 right ventricle, slowed conduction, ventricular tachyarrhythmias and sudden cardiac death.³ However, only 73 around 20% of patients with Brugada Syndrome have mutations in SCN5A. The advent of massively parallel 74 sequencing has allowed many groups to investigate the genetic underpinnings of BrS over the past two 75 decades. These genetic data, in combination with molecular studies, have expanded the number of putative 76 disease-causing genes in BrS from one gene, SCN5A, to over 20 genes today. Genes reported to have mutations 77 which cause BrS can be broadly classified as those which encode 1) sodium channels, 2) potassium channels, 78 3) channel interacting proteins, and 4) metabolic proteins.⁴ Recent reports have also suggested BrS may be 79 polygenic in nature, with single nucleotide polymorphisms (SNPs) at the SCN5A-SCN10A and HEY2 loci 80 predisposing individuals to BrS.⁵ 81 Our lab first reported a large, multigenerational family with BrS.⁶ Initial linkage mapping using mi-82 crosatellites identified an ~15 centimorgan (CM) region on chromosome 3 with a LOD score >4 that did 83 not contain SCN5A or SCN10A.⁷ Positional cloning and subsequent molecular analysis identified a variant 84 in glycerol 3-phosphate dehydrogenase 1-like (GPD1L) as the putative cause of this family's disease. Peak

sodium current in HEK293 cells transfected with $Na_V 1.5$ decreased if cells were co-transfected with GPD1L-86

A280V compared to those co-transfected with GPD1L-WT. Additional studies suggested GPD1L-A280V 87

decreases Na_V1.5 membrane expression. Together, a decrease in peak sodium current and a decrease in 88

membrane localization suggest that GPD1L-A280V is pathogenic for BrS.

The explosion of genetic data, especially variants of uncertain significance, from clinical genetic testing 90 has led to a plethora of putative mutations but a dearth of scientific evidence supporting the pathogenicity of these variants. Concerns about variants of uncertain significance being classified as pathogenic without 92 data to support the claim has been reviewed in many publications.^{8,9} This concern sparked a review of the 93 pathogenicity of reported BrS genes using the Clinical Genome Resources (ClinGen) framework. This review determined that SCN5A was the only gene with sufficient scientific evidence to definitively be regarded as a causative gene for BrS.¹⁰ GPD1L was excluded due to a large linkage region in which other genes had 96 not been sequenced and because the allele frequency of GPD1L-A280V was relatively high (gnomAD V2.1.1 97 allele frequency = 1.29E-4).¹¹ 98

In the present study, we provide high-depth whole exome sequencing data in the proband of our originally-99 reported large multigenerational family, SNP-based linkage analysis of affected individuals, and sequencing 100 data of previously defined BrS risk SNPs to support our initial reports that GPD1L-A280V is pathogenic 101 for BrS. 102

$\mathbf{2}$ Methods 103

85

Patient enrollment and phenotype validation $\mathbf{2.1}$ 104

Patients were enrolled under protocols approved through the University of Pittsburgh and University of Iowa 105 Institutional Review Boards (IRBs). All individuals in the study provided informed consent for participation 106 in the study. Type 1, Type 2, or Type 3 Brugada Syndrome, or clinically unaffected status, was determined 107

through clinical history, electrocardiogram analysis, and clinical provocative testing using procainamide as

¹⁰⁹ previously described.⁶

110 2.2 Genomic DNA isolation and RNAse A treatment

Genomic DNA was isolated from peripheral blood using commercially-available kits or automated machine as previously described.⁶ Isolated DNA was treated with RNAse A and re-isolated by ethanol precipitation prior to sequencing. DNA integrity was assessed by agarose gel electrophoresis and quantified using Qubit High Sensitivity DNA assays (ThermoFisher Scientific).

115 2.3 SNP calling and Linkage Analysis

Ten affected and five unaffected individuals from the family had genomic DNA prepared as described above. After passing quality control and subsequent library preparation, SNPs were called using the Infinium Global Screening Array (Illumina) at CD Genomics (Shirley, NY, USA) which sequences approximately 654,027 SNPs. Call rates were greater than 98% for all individuals. (Supplemental Table 1) Chromosomes were phased using ShapeIt2 with DuoHMM enabled and the reference HapMap Phase II data.^{12,13} The resultant phased haplotypes were investigated manually for recombination using our previously-reported microstaellite data as the starting point. Logarithm of odds scores were calculated using Superlink-Online SNP.¹⁴

¹²³ 2.4 High-depth whole exome sequencing and analysis

Whole exome sequencing was performed at the Iowa Institute of Human Genetics (IIHG) on genomic DNA 124 prepared as described above. Library preparation was performed using standard protocols for the Agilent 125 SureSelect V6 + UTR library preparation kit (Agilent Technologies). The resulting 150 base pair paired-126 end library was sequenced on an Illumina HiSeq 4000. Resultant .fastq files underwent quality control using 127 FastQC.¹⁵ Alignment, variant calling, and variant annotation were performed using BWA-MEM (alignment), 128 GATK4 (variant calling), and SnpEff (variant annotation) using an implementation of BCBIO and the 129 GRCh37 reference genome.^{16–21}. A second variant calling tool, Freebayes, was used in a near-identical 130 pipeline to ensure consensus variant calls were achieved²². Greater than 80% of targeted bases had greater 131 than 50x coverage. (Supplemental Figure 1) A brief summary of sequencing results is shown in Supplemental 132 Table 2. 133

Two approaches were used to identify candidate variants 1) a linkage-region focused analyses and 2) 134 a linkage-naive approach. The linkage-region focused analysis was limited to the newly-defined linkage 135 region on chromosome 3, while the linkage-naive approach entertained variants on any autosome. For 136 both approaches, the resulting annotated variant call file was a used to generate a database compatible with 137 Gemini.²³ Variants present at less than 1% minor allele frequency, a lenient threshold for autosomal dominant 138 disease, were included. Variants were then segregated according to impact (nonsense, splicing, missense, 139 synonymous) with synonymous variants removed from consideration. Variant location information, either 140 exonic, UTR, intronic, or intergenic, was subsequently used in prioritization of variants. 141

¹⁴² 2.5 Sanger Sequencing of Risk SNPs

Sanger sequencing of BrS risk SNPs was performed at the Iowa Institute of Human Genetics using custom primers designed in-lab and synthesized by Integrated DNA Technologies. (Supplemental Table 3) The resulting chromatograms were analyzed using NCBI Blast, Finch, and SnapGene from which wild-type, heterozygous, or homozygous SNP calls were made and manually validated.

¹⁴⁷ 2.6 Protein and variant modeling

A single unpublished protein model of GPD1L is present on the RCSB Protein Databank.^{23,24}. AlphaFold computational models for GPD1L are also publicly available from EMBL. The AlphaFold GPD1L model AF-Q8N335-F1-model_v1 was downloaded and imported into PyMol.²⁵ The mutagenesis function of PyMol was used to visualize the impact of the three predicted GPD1L-A280V rotamers. Strain scores were calculated for all rotamers within PyMol. (Supplemental Figure 2) Variant data for *GPD1L* was downloaded from ClinVar.²⁶

$_{154}$ 3 Results

¹⁵⁵ 3.1 Linkage analysis reaffirms a region on chromosome 3 co-segregates with ¹⁵⁶ Brugada Syndrome

To understand the linked genomic regions within this family we performed linkage analysis using the Illumina 157 Global Screening Array with ten affected and five unaffected family members. The previously-identified 15 158 CM linkage region in this family which we reported was used as a starting point for our investigation.⁶ 159 Analysis of phased haplotypes identified a narrowed linkage region defined by the SNPs rs13059657 and 160 rs7651953 (GRCh37, chr3:29,899,567-32,970,737). (Figure 1) Individuals III-2 and III-8 defined the 5' and 161 3' breakpoints, respectively. Calculation of LOD scores using individuals having a Type 1 EKG pattern or 162 positive procainamide test, in addition to I-I and I-II from Figure 1, using Superlink-Online SNP results in 163 a LOD score of 2.02 for this region.¹⁴ The addition of individuals with Type II EKG patterns results in a LOD score of 3.18. The region of interest contains 14 genes, including GPD1L, but not SCN5A or SCN10A 165 which are approximately 6 Mb downstream. (Figure 1) 166

¹⁶⁷ 3.2 Whole-exome sequencing analysis affirms GPD1L-A280V as the only rare ¹⁶⁸ exonic variant in the linked region

High-depth (100x) whole exome sequencing of the proband was performed using the Agilent SureSelect 169 V6+UTR capture kit. Limiting the analysis to the newly defined linkage region identified GPD1L-A280V 170 as the only rare, exonic coding variation. No other exonic or splice variants occurred at an allele fre-171 quency of less than 1%. The proband carried missense variants in arrhythmia and cardiomyopathy genes 172 not usually associated with BrS: one missense variant in ANK2 (p.R2416G;c.7246C>G, unreported), one 173 in FHL2 (p.V187M;c.559G>A, f = 6.78E-5) and two in TTN (p.A31503T;c94507G>A, f = 5.04E-5 and 174 p.F14410C;c.43229T>G, unreported) (Table 1). Linkage analysis and Sanger sequencing showed that none 175 of these variants cosegregate with BrS in this family, and no other variants are annotated as pathogenic for 176 cardiovascular diseases in the ClinVar database. 177

The SCN5A and SCN10A BrS risk SNPs are linked to GPD1L-A280V in this family

Three published risk SNPs for BrS [rs10428132 (SCN10A; total minor allele frequency, dbSNP (MAF) = 0.40), rs11708996 (SCN5A; MAF=0.14), and rs9388451 (HEY2; MAF=0.42)] identified by Bezzina et al. were assessed in family members with sufficient DNA, showing 8/15 affected subjects (those demonstrating a Type I, II, or III BrS pattern) in generations II and III are homozygous for the SCN10A risk allele and 12/15 subjects are heterozygous for the SCN5A risk allele (Figure 1). SCN5A and SCN10A are near GPD1L on

 $chromosome \ 3; in most affected individuals the \ GPD1L-A280V \ mutation \ co-segregated \ with \ one \ SCN5A \ and \ Not \ Not$ one SCN10A risk allele. The HEY2 risk SNP on chromosome 6 segregated in Mendelian fashion. Overall, 186 individuals affected with BrS (n=15) carried 3.3 ± 0.2 risk SNPs, while unaffected individuals (n=12) carried 187 2.1 ± 0.4 risk SNPs (p=0.012). (Figure 2) The SCN5A and SCN10A risk SNPs carried along with GPD1L-188 A280V in this family may potentiate the pathogenicity of GPD1L-A280V. 189

$\mathbf{3.4}$ GPD1L-A280V may introduce protein instability in computationally-predicted 190 models 191

In silico mutagenesis of the A280 residue to V280 in the computational model of GPD1L demonstrates 192 increased strain within the alpha helix in which residue 280 resides. (Figure 3 and Supplemental Figure 2) 193 Thus, in silico mutagenesis results in increased computationally-prediced strain stores indicating GPD1L-194 A280V may introduce steric hinderance in the GPD1L tertiary structure.

4 Discussion 196

195

4.1 The genetic architecture of BrS in one large, multigenerational family 197

Our analysis of this large, multigenerational family narrows the linkage region to approximately 3.1 megabases 198 which includes GPD1L but excludes SCN5A and SCN10A. Whole exome sequencing confirmed GPD1L-199 A280V as the only rare exonic variant at an allele frequency <1% within this region. The affected-only 200 LOD score of 3.18 for individuals displaying Type I and Type II BrS patterns (Type 1 BrS alone = 2.02), suggests this region is linked to BrS in this family and is consistent with the previous reportd LOD score of 202 >3 which included unaffected individuals. While not proving pathogenicity of the GPD1L-A280V mutation, these data, along with previously published experiment data should be considered when assessing the role 20/ of GPD1L in the pathogenesis of BrS. 205

While the allele frequency of GPD1L-A280V is high relative to published allele frequency cutoffs, GPD1L-206 A280V is not the lone reported *GPD1L* putative mutation.²⁷ Four other putative mutations, p.I124V, 207 p.E174K, p.R231C, and p.Q345H, have been reported in ClinVar. (Figure 4)²⁸ Nevertheless, we believe 208 that mutations in GPD1L, in isolation, are likely unable to cause sufficient decreases in sodium current to 209 precipitate BrS. It is likely that GPD1L mutations and risk polymorphisms act in concert to precipitate 210 BrS, as is the case of this large multigenerational family. We believe that SCN5A is the only gene in which 211 mutations are sufficient to cause enough inward sodium current decrease to cause BrS, while GPD1L ---and 212 possibly other reported BrS genes —precipitate BrS in a polygenic manner or alongside risk SNPs. 213

The genetic architecture of BrS: past, present, and future 4.2214

Mutations in SCN5A are certainly the predominant genetically-identifiable cause of BrS.¹ Future studies of 215 GPD1L and other putative BrS mutations in heterologous systems and in mouse models will continue provide 216 information on the role these genes play in cardiac physiology. Surprisingly, few reports exist of the use of 217 whole-exome or whole-genome sequencing to identify the genetic underpinnings of BrS, and the majority of 218 these studies identify genetic variation in previously reported disease genes. Future large genomic datasets 219 (e.g. UK Biobank, gnomAD) have and will continue to provide further information on allele frequency of genes reported to cause BrS. In all, these data and future studies will further our understanding of the still 221 uncertain genetic architecture of BrS. 222

223 4.3 Study limitations

Our study uses linkage and exome sequencing data from a single family or single proband. Linkage data provide an incomplete picture of the genetic architecture for a disease, though we believe the addition of exome sequencing partly addresses this limitation. Notably, exome sequencing libraries are not targeted to capture intronic sequences. Deep intronic variants within *GPD1L* may contribute to disease in this family.

228 4.4 Clinical implications

Recent recommendations to clinical genetic testing panels exclude all genes except *SCN5A* in their testing panels. While well-intentioned to avoid undue harm from provider misinterpretation, this may reduce identification of rare variation contributing to BrS in sodium channel modifying genes, among others. While concerns for generating undue harm from misinterpretation of a variant of uncertain significance in a disputed gene included on a clinical genetic testing panel are valid, we believe these concerns highlight the necessity of thorough understanding of genetic testing results and the importance both of cardiologists trained in genetics and the use of genetic counselors.^{29,30}

Our study demonstrates that mutation in GPD1L likely contributes to BrS in this multigenerational family. Given this, inclusion of *GDP1L* on genetic testing panels should be reconsidered.

238 5 Conclusion

We identified GPD1L-A280V as the only rare non-synonymous coding variation in the refined linkage region in a large, multigenerational family. Associated BrS risk SNPs may be permissive for the BrS phenotype in affected family members. In addition, the A280V mutation may alter the GPD1L tertiary structure. While mutation in *SCN5A* is the most common cause genetically-identified cause of BrS, our study suggests that GPD1L should still be considered when evaluating the genetic underpinnings of BrS.

244

245 **References**

- ²⁴⁶ [1] J Brugada, P Brugada, and R Brugada. "The Syndrome of Right Bundle Branch Block ST Segment
- Elevation in V1 to V3 and Sudden Death-the Brugada Syndrome." In: *Europace : European pacing*,
- arrhythmias, and cardiac electrophysiology : journal of the working groups on cardiac pacing, arrhyth-
- mias, and cardiac cellular electrophysiology of the European Society of Cardiology 1.3 (1999), pp. 156–
- ²⁵⁰ 66. DOI: 10.1053/eupc.1999.0033. pmid: 11225790.
- [2] Qiuyun Chen et al. "Genetic Basis and Molecular Mechanism for Idiopathic Ventricular Fibrillation".
 In: Nature 392.6673 (6673 Mar. 1998), pp. 293–296. DOI: 10.1038/32675.
- [3] Charles Antzelevitch et al. "Brugada Syndrome: 1992–2002". In: Journal of the American College of
 Cardiology 41.10 (2003).
- In: Josep Brugada et al. "Present Status of Brugada Syndrome: JACC State-of-the-Art Review". In: Journal of the American College of Cardiology 72.9 (Aug. 28, 2018), pp. 1046–1059. DOI: 10.1016/j.
 jacc.2018.06.037. pmid: 30139433.
- [5] Connie R Bezzina et al. "Common Variants at SCN5A-SCN10A and HEY2 Are Associated with
 Brugada Syndrome, a Rare Disease with High Risk of Sudden Cardiac Death". In: *Nature Genetics*45.9 (Sept. 21, 2013), pp. 1044–1049. DOI: 10.1038/ng.2712. pmid: 23872634.
- [6] Barry London et al. "Mutation in Glycerol-3-Phosphate Dehydrogenase 1 like Gene (GPD1-L) De creases Cardiac Na+ Current and Causes Inherited Arrhythmias." In: *Circulation* 116.20 (Nov. 13, 2007), pp. 2260–8. DOI: 10.1161/CIRCULATIONAHA.107.703330. pmid: 17967977.
- [7] Raul Weiss et al. "Clinical and Molecular Heterogeneity in the Brugada Syndrome". In: *Circulation* 105.6 (Feb. 12, 2002), pp. 707–713. DOI: 10.1161/hc0602.103618.
- [8] Michael J. Ackerman. "Genetic Purgatory and the Cardiac Channelopathies: Exposing the Variants of
 Uncertain/Unknown Significance Issue". In: *Heart Rhythm* 12.11 (Nov. 1, 2015), pp. 2325–2331. DOI:
 10.1016/j.hrthm.2015.07.002. pmid: 26144349.
- [9] Jamie D. Kapplinger et al. "Enhanced Classification of Brugada Syndrome–Associated and Long-QT
 Syndrome–Associated Genetic Variants in the SCN5A -Encoded Na v 1.5 Cardiac Sodium Channel".
 In: Circulation: Cardiovascular Genetics 8.4 (Aug. 2015), pp. 582–595. DOI: 10.1161/CIRCGENETICS.
 114.000831. pmid: 25904541.
- [10] S. Mohsen Hosseini et al. "Reappraisal of Reported Genes for Sudden Arrhythmic Death: Evidence based Evaluation of Gene Validity for Brugada Syndrome". In: *Circulation* 138.12 (June 29, 2018),
 pp. 1195–1205. DOI: 10.1161/CIRCULATIONAHA.118.035070. pmid: 29959160.
- [11] Konrad J. Karczewski et al. "The Mutational Constraint Spectrum Quantified from Variation in
 141,456 Humans". In: *Nature* 581.7809 (May 28, 2020), pp. 434–443. DOI: 10.1038/s41586-020 2308-7. pmid: 32461654.
- [12] Olivier Delaneau, Jean-Francois Zagury, and Jonathan Marchini. "Improved Whole-Chromosome Phas ing for Disease and Population Genetic Studies". In: *Nat Methods* 10.1 (1 Jan. 2013), pp. 5–6. DOI:
 10.1038/nmeth.2307.
- [13] "A Second Generation Human Haplotype Map of over 3.1 Million SNPs". In: *Nature* 449.7164 (Oct. 18, 2007), pp. 851–861. DOI: 10.1038/nature06258. pmid: 17943122.

- ²⁸⁴ [14] M. Silberstein et al. "A System for Exact and Approximate Genetic Linkage Analysis of SNP Data in
- Large Pedigrees". In: *Bioinformatics* 29.2 (Jan. 15, 2013), pp. 197–205. DOI: 10.1093/bioinformatics/ bts658. pmid: 23162081.
- [15] S Andrews. FastQC: A Quality Control Tool for High Throuhgput Sequence Data. URL: http://www.
 bioinformatics.babraham.ac.uk/projects/fastqc.
- [16] Heng Li. Aligning Sequence Reads, Clone Sequences and Assembly Contigs with BWA-MEM. May 26,
 2013. arXiv: 1303.3997 [q-bio]. URL: http://arxiv.org/abs/1303.3997 (visited on 11/22/2021).
- ²⁹¹ [17] Mark A DePristo et al. "A Framework for Variation Discovery and Genotyping Using Next-Generation
- ²⁹² DNA Sequencing Data". In: *Nature Genetics* 43.5 (May 10, 2011), pp. 491–498. DOI: 10.1038/ng.806.
- [18] Ryan Poplin et al. "Scaling Accurate Genetic Variant Discovery to Tens of Thousands of Samples".
 In: *bioRxiv* (July 24, 2018), p. 201178. DOI: 10.1101/201178.
- [19] Pablo Cingolani et al. "A Program for Annotating and Predicting the Effects of Single Nucleotide
 Polymorphisms, SnpEff: SNPs in the Genome of Drosophila Melanogaster Strain W1118; Iso-2; Iso-3".
 In: Fly (Austin) 6.2 (2012 Apr-Jun), pp. 80–92. DOI: 10.4161/fly.19695. pmid: 22728672.
- ²⁹⁸ [20] Brad Chapman et al. *Bcbio/Bcbio-Nextgen:* Zenodo, Apr. 14, 2021. DOI: 10.5281/zenodo.4686097.
- [21] Deanna M. Church et al. "Modernizing Reference Genome Assemblies". In: *PLoS Biol* 9.7 (July 2011),
 e1001091. DOI: 10.1371/journal.pbio.1001091. pmid: 21750661.
- [22] Erik Garrison and Mark Marth. "Haplotype-Based Variant Detection from Short-Read Sequencing
 Erik Garrison and Gabor Marth January 12, 2016 Abstract". In: (2016), pp. 1–20. DOI: 10.1093/
 bioinformatics/btv440. pmid: 24136966.
- ³⁰⁴ [23] Umadevi Paila et al. "GEMINI: Integrative Exploration of Genetic Variation and Genome Annota ³⁰⁵ tions." In: *PLoS computational biology* 9.7 (2013), e1003153. DOI: 10.1371/journal.pcbi.1003153.
 ³⁰⁶ pmid: 23874191.
- RCSB Protein Data Bank. RCSB PDB 2PLA: Crystal Structure of Human Glycerol-3-Phosphate
 Dehydrogenase 1-like Protein. URL: https://www.rcsb.org/structure/2PLA (visited on 11/22/2021).
- ³⁰⁹ [25] John Jumper et al. "Highly Accurate Protein Structure Prediction with AlphaFold". In: *Nature* ³¹⁰ 596.7873 (7873 Aug. 2021), pp. 583–589. DOI: 10.1038/s41586-021-03819-2.
- ³¹¹ [26] Melissa J. Landrum et al. "ClinVar: Improving Access to Variant Interpretations and Supporting
 ³¹² Evidence". In: Nucleic Acids Res 46.D1 (Jan. 4, 2018), pp. D1062–D1067. DOI: 10.1093/nar/gkx1153.
 ³¹³ pmid: 29165669.
- ³¹⁴ [27] Nicola Whiffin et al. "Using High-Resolution Variant Frequencies to Empower Clinical Genome Inter pretation". In: *Genetics in Medicine* 19.10 (Oct. 18, 2017), pp. 1151–1158. DOI: 10.1038/gim.2017.26.
 pmid: 28518168.
- ³¹⁷ [28] David W. Van Norstrand et al. "Molecular and Functional Characterization of Novel Glycerol-3 ³¹⁸ Phosphate Dehydrogenase 1 like Gene (GPD1-L) Mutations in Sudden Infant Death Syndrome". In:
 ³¹⁹ Circulation 116.20 (Nov. 13, 2007), pp. 2253–2259. DOI: 10.1161/CIRCULATIONAHA.107.704627. pmid:
 ³²⁰ 17967976.
- [29] Ferhaan Ahmad et al. "Establishment of Specialized Clinical Cardiovascular Genetics Programs: Rec ognizing the Need and Meeting Standards: A Scientific Statement From the American Heart Associ ation". In: *Circulation: Genomic and Precision Medicine* 12.6 (June 2019), e000054. DOI: 10.1161/
- ³²⁴ HCG.0000000000054. pmid: 31117808.

- 325 [30] Eric D. Green et al. "Strategic Vision for Improving Human Health at The Forefront of Genomics".
- In: Nature 586.7831 (7831 Oct. 2020), pp. 683–692. DOI: 10.1038/s41586-020-2817-4.

327 Figures and Tables



329

³³⁰ Figure legend:

Title: Representative pedigree of the family, linkage analysis, and assessment of Brugada Syndrome (BrS)
 risk SNPs.

Caption: A) Linkage analysis demonstrates a small region on chromosome 3 (inset map and green bracketed region) which has undergone recombination in the third generation of the pedigree. Previously reported
risk SNPs (SCN5A and SCN10A, red) for the BrS phenotype are associated with the GPD1L-A280V. B)
The 3 megabase region on chromosome 3 identified by linkage analysis contains 14 genes, including *GPD1L*. *SCN5A* and *SCN10A* lie upstream of GPD1L (ideogram, red text) and outside of the refined linkage region
(ideogram, red box). ICD: implantable cardioverter defibrillator. +Proc: positive procainamide challenge.

³³⁹ RBBB: right bundle branch block. VT: ventricular tachycardia.



342

Figure legend: 343

Title: Distribution of Brugada Syndrome Risk Polymoprhisms in the large mutli-generational family 344

Caption: Individuals with Brguada Syndrome in our family carry more risk polymorphisms on average 345

(3.3) than unaffected individuals (2.1) in our large mutli-generational family. The distribution of risk poly-346

moprhisms is similar to that which was reported by Bezzina et al.⁵ 347





350

- ³⁵¹ Figure legend:
- ³⁵² Title: Computational modeling of GPD1L suggests A280V introduces steric clashing
- 353 Caption:

Left: The wild-type GPD1L structure as calculated by AlphaFold (AF-Q8N335-F1-model_v1) with residue

A280 highlighted in blue. A280 resides in an alpha helix. Right: One of three putative GPD1L-A280V

³⁵⁶ rotamers as calculated by PyMol. All three predicted rotamers introduce strain in the alpha helix.



- ³⁶⁰ Figure legend:
- ${}_{361} \quad {\bf Title: Reported GPD1L variants in ClinVar show GPD1L-A280V is not the sole reported putative mutation.}$
- 362 Caption: Missense variants with conflicting interpretation (yellow circles, with GPD1L-A280V highlighted
- $_{\tt 363}$ $\,$ as red) and variants of uncertain significance (black bars, 83 total) were downloaded from ClinVar (accessed
- November 12, 2021) and plotted along the full length of GPD1L. All five variants with conflicting interpre-
- $_{365}$ tation, including GPD1L-A280V, were submitted to ClinVar by their respective submitters with Brugada
- 366 Syndrome as the clinical condition.
- 367

Table 1. Variants in reported Brugada Syndrome or Inherited Cardiac Condition (ICC) List Genes

Gene List	Position	Consequence	Gene	Frequency
Linkage region	chr3:32200588C>T	p.Ala280Val	GPD1L	1.29E-04
BrS	chr12:2788732C>T	p.Gly1738Gly	CACNA1C	1.21E-05
	chr2:105979871C>T	p.Val187Met	FHL2	6.78E-05
	chr2:179411745C>T	p.Ala31503Thr	TTN	5.04E-05
ICC	chr2:179497504A>C	p.Phe14410Cys	TTN	Unreported
	chr4:114277020C>G	p.Arg2416Gly	ANK2	Unreported
	chr16:15844004G>A	p.His690His	MYH11	5.25E-04