

1 **Explanatory factors for Ethnic inequalities in Multimorbidity; findings from**
2 **pooled Health Survey for England 2011-2018**

3 **Authors:** Linda Ng Fat¹, Jennifer S Mindell¹, Logan Manikam¹, Shaun Scholes¹

4 * **Corresponding author:** Linda Ng Fat, l.ngfat@ucl.ac.uk, Health and Social Surveys
5 Group, Department of Epidemiology and Public Health, UCL, 1-19 Torrington Place,
6 London, WC1E 7HB

7

8

9 ¹ Health and Social Surveys Group, Research Department of Epidemiology and
10 Public Health, University College London (UCL), 1-19 Torrington Place, London,
11 WC1E 7HB

12

13

14 **Key words:** Ethnic inequalities, Multimorbidity, Cardiovascular multimorbidity,
15 England

16

17

18 **Abstract**

19 **Background**

20 Social-economic factors and health behaviours may be driving variation in ethnic health
21 inequalities in multimorbidity including among distinct ethnic groups.

22 **Methods**

23 Using the cross-sectional nationally-representative Health Surveys for England 2011-2018
24 (N=54,438, aged 16+), we carried out multivariable logistic regression on the odds of having
25 general multimorbidity (≥ 2 longstanding conditions) by ethnicity (British White (reference
26 group), White Irish, Other White, Indian, Pakistani, Bangladeshi, Chinese, African,
27 Caribbean, White Mixed, Other Mixed, Other), adjusting for age, sex, education, area-
28 deprivation, smoking status, and survey year. This was repeated for cardiovascular
29 multimorbidity (N=37,148, aged 40+: having ≥ 2 of the following: doctor-diagnosed diabetes
30 or hypertension, heart attack or stroke) and multiple cardiometabolic risk biomarkers (HbA1c
31 $\geq 6.5\%$, raised blood pressure, total cholesterol $\geq 5\text{mmol/L}$).

32 **Results**

33 20% of adults had general multimorbidity. In fully-adjusted models, compared with the White
34 British majority, Other White (Odds Ratio (OR)=0.63 (95% confidence interval=(0.53-0.74)),
35 Chinese (OR=0.58 (0.36-0.93)), and African adults (OR=0.54 (0.42-0.69)), had lower odds of
36 general multimorbidity. Among adults aged 40+, Pakistani (OR=1.27 (0.97-1.66) $p=0.080$)
37 and Bangladeshi (OR=1.75 (1.16-2.65)) had increased odds, and African adults had
38 decreased odds (OR=0.63 (0.47-0.83)) of general multimorbidity. Risk of cardiovascular
39 multimorbidity was higher among Indian (OR=3.31 (2.56-4.28)), Pakistani (OR=3.48 (2.52-
40 4.80)), Bangladeshi (OR=3.67 (1.98-6.78)), African (OR=1.61 (1.05-2.47)), Caribbean
41 (OR=2.18 (1.59-2.99)) and White Mixed (OR=1.98 (1.14-3.44)) adults. Indian adults were
42 also at risk of having multiple cardiometabolic risk biomarkers.

43 **Conclusion**

44 Ethnic inequalities in multimorbidity are independent of socioeconomic factors. Ethnic
45 minority groups are particularly at risk of cardiovascular multimorbidity, which may be
46 exacerbated by poorer management of cardiometabolic risk factors.

47

48

49

50 **What is already known on this topic**

- 51 • Studies have found ethnic inequalities in multimorbidity among broad ethnic groups;
52 however, it is not known whether lower socioeconomic status or health behaviours
53 are driving this inequality in England, nor whether there are differences between
54 distinct ethnic groups.

55

56 **What this study adds**

- 57 • Ethnic inequalities in multimorbidity in a nationally-representative sample in England
58 vary between older and younger adults, distinct ethnic groups, and are independent
59 of socioeconomic status, smoking and obesity.
- 60 • African, Caribbean, Bangladeshi, Indian, Pakistani and White mixed adults aged 40
61 and over had higher risk of self-reported cardiovascular multimorbidity compared with
62 British White adults. Indian adults were also at risk of multiple uncontrolled
63 cardiometabolic risk biomarkers.
- 64 • African, Chinese, Other White and Indian adults (<40 years) had lower risk of general
65 multimorbidity compared with the British White majority, suggesting a healthy migrant
66 effect or differences in interpretation of chronic conditions.

67

68 **How this study might affect research, practice or policy**

- 69 • Further research is needed to explore whether underdiagnosis and/or poorer
70 management of cardiometabolic risk factors among ethnic groups may be driving
71 factors behind inequalities in self-reported cardiovascular multimorbidity.

72

73

74 **Explanatory factors for variations in multimorbidity among ethnic groups; findings**
75 **from pooled nationally representative Health Survey for England 2011-2018**

76

77 **Background**

78

79 Multimorbidity is increasing, given an ageing population, affecting individuals' quality of life
80 and increasing mortality,(1,2). Understanding the factors related to different levels of
81 multimorbidity in ethnic groups could be used to prevent harm and reduce ethnic
82 inequalities. Non-white ethnic groups were disproportionately affected with Covid-19, with
83 multimorbidity potentially being one contributing factor. It was found that compared with the
84 white population, non-white ethnicities with multimorbidity had almost triple the risk of having
85 Covid-19(3).

86

87 Ethnic health inequalities in multimorbidity, including cardiovascular multimorbidity, exist(4–
88 9). A study in East London found the Black and South Asian population had higher rates of
89 cardiovascular multimorbidity than the white majority(4). Studies of ethnic health inequalities
90 in multimorbidity in England often use GP records, which may not reflect differences in a
91 national population sample,(4,5). Furthermore, studies in England have not comprehensively
92 explored what may be accounting for some of the health differences, such as socioeconomic
93 differences. Deprivation and lower socioeconomic status could increase the risk of
94 multimorbidity,(6–8,10–12). Behavioural risk-factors could also confound the relationship.
95 Smoking and obesity were found to be risk-factors for multimorbidity,(7,10,13), and may
96 have a stronger effect than deprivation and ethnicity,(7,13). Smoking and obesity were found
97 to vary in The Health Survey for England 2011-2019,(14).

98

99 Furthermore, broad ethnic groups such as 'South Asian' and 'Black' may mask heterogeneity
100 within groups, for example among Indian versus Pakistani/Bangladeshi, and Black African
101 versus Black Caribbean groups,(15). The aims of this study were to examine how risk of
102 multimorbidity including cardiovascular multimorbidity varied among distinct ethnic groups,
103 and whether this relationship was explained by difference in socioeconomic status or
104 behavioural risk factors for cardiovascular diseases, namely smoking and obesity.

105

106 **Methods**

107

108 **Data Sources**

109

110 The Health Survey for England (HSE) is a cross-sectional, nationally-representative survey
111 of the population occurring annually. Data is collected via new multistage stratified
112 probability sampling each year, as described elsewhere(16). This study uses data spanning
113 from 2011 to 2018, to maximise the number of participants from different ethnic groups. Self-
114 reported data were collected from an initial face-to-face interview conducted by trained
115 interviewers using computer-assisted personal interviewing (CAPI). The interviewers also
116 measured height and weight. At the second stage, further biophysical measurements were
117 collected at a nurse visit using standard protocols,(17). All data collection took place within
118 participants' homes. Response rates have gradually declined over the period, e.g., from 59%
119 in 2011 to 54% in 2018 for the face-to-face interview.

120

121 **General and cardiovascular multimorbidity**

122

123 This study examines ethnic group differences in two types of multimorbidity: general
124 multimorbidity and cardiovascular multimorbidity, using information provided by participants
125 at the interview stage. General multimorbidity was derived through self-reported responses
126 to a question on longstanding conditions: whether the participant had any longstanding
127 physical or mental health conditions or illnesses expected to last over a period of time (2011)
128 or 12 months or more (2012-2018). Participants were then asked to list up to six conditions
129 that affected them. Responses were recoded to indicate the number of longstanding
130 conditions (none/one/two or more). Participants having two or more longstanding conditions
131 were classified as having general multimorbidity.

132

133 Self-reported doctor-diagnosed cardiovascular multimorbidity was defined as having two or
134 more of the following conditions: reporting having doctor-diagnosed diabetes or high blood
135 pressure (in response to specific questions on whether participants were ever told by a
136 doctor they had diabetes/high blood pressure) or reporting stroke or heart disease when
137 responding to the aforementioned follow-up question to the longstanding illness question.
138 We included doctor-diagnosed hypertension following the example of Mathur et al.(4),
139 although other definitions do not include hypertension,(18).

140

141

142 **Multiple cardiometabolic risk biomarkers**

143

144 Using objective health examination measures, we also examined differences by ethnicity in
145 multiple cardiometabolic risk biomarkers. Cardiometabolic biomarkers were treated
146 separately from doctor-diagnosed conditions (cardiovascular multimorbidity) to identify
147 uncontrolled risk factors. Having multiple cardiometabolic risk biomarkers was defined as
148 two or more of the following: raised glycated haemoglobin (HbA_{1c} levels $\geq 6.5\%$); raised blood
149 pressure (systolic blood pressure ≥ 140 mmHg and/or diastolic blood pressure ≥ 90 mmHg);
150 and/or raised total cholesterol level (≥ 5 mmol/L), suggesting uncontrolled risk-factors or
151 disease such as diabetes or hypertension. Further details on data collection during the nurse
152 visit have been published elsewhere,(15).

153

154 **Ethnicity and other covariates**

155

156 Ethnicity was self-reported at the main interview and grouped in the following way: White
157 British, White Irish, Other White, Indian, Pakistani, Bangladeshi, Chinese, African,
158 Caribbean, White Mixed, Other Mixed, Other.

159

160 Other covariates included the Index of Multiple Deprivation, a standard measure of area
161 deprivation covering different social and economic dimensions, assigned based on
162 participants' residence,(17), grouped into quintiles. Highest educational qualification was
163 self-reported and grouped into degree or higher; lower than degree; no qualifications. Self-
164 reported smoking status was grouped into never; former; current smoker. BMI categories
165 were based on height and weight measurements of the participants performed by the trained
166 interviewer at the main interview. Detailed information on collection can be found
167 elsewhere,(17).

168

169

170 **Statistical analyses**

171

172 Complete case analyses were used and the sample was limited to (i) adults aged 16+ for
173 general multimorbidity (N=54,438), (ii) adults aged 40+ for self-reported cardiovascular
174 multimorbidity (N= 37,148), and (iii) adults aged 16+ with biological measurements for
175 multiple cardiometabolic biological risk factors (N=24,203). The complete case approach
176 resulted in a reduction in sample size of around 17.5%; this was predominately driven by
177 missing data for BMI, which required physical measurements of height and weight (17.0%).
178 Few participants (<0.5%) had missing responses on key variables: general multimorbidity
179 (n=46), cardiovascular multimorbidity (n=7), and ethnicity (n=220). Missing data on

180 biomarkers was high due to not all the sample providing blood samples (59.8%). We
181 accounted for non-response weighting, specific to non-response at the interview or blood
182 samples for biological risk factors were applied to account for this, and the complex survey
183 design (geographical clustering of participants) in all analyses.

184

185 Characteristics of ethnic groups were explored and chi-squared used to test for bivariate
186 associations. Logistic regression was carried out on the risk of having multimorbidity for each
187 of the three definitions of multimorbidity. Since longstanding conditions are broader, and
188 multimorbidity is not restricted to older participants,(8,19), analyses of general multimorbidity
189 focused on all adults. However, we also stratify results by aged under 40 and 40 and over. A
190 cut-off of age 40 has been used in other large scale studies,(20), and sample sizes among
191 some of the ethnic groups were too small if limiting to an older sample (e.g. <100, if based
192 on adults aged 65 and over).

193

194 To explore whether the relationship between ethnicity and multimorbidity may be
195 confounded by socioeconomic status or behavioural risk factors, variables were added in the
196 following way, with changes in odds ratios examined to explore the effect of adjusting for
197 each additional potential confounder.

198

199 Model 1: Sex + Age in years

200 Model 2: Model 1 + Area Deprivation

201 Model 3: Model 1 + Educational qualifications

202 Model 4: Model 1 + Smoking status

203 Model 5: Model 1 + BMI category

204 Model 6: All covariates + survey year

205

206 As a sensitivity analyses, we replicated the final models on a sample including participants
207 without BMI measurements (without adjusting for BMI), resulting in fewer missing cases
208 (1%). Analyses were conducted in Stata 17,(21).

209

210 **Results**

211

212 Chinese, Indian and Other White adults had the highest proportion of adults with the highest
213 educational qualifications (53%-44%), whereas Bangladeshi, White Irish, Pakistani, and
214 Caribbean had the highest proportions with no qualifications (29%-25%) (Table 1). Over two-

215 fifths of adults in the Bangladeshi, Pakistani, African and Caribbean groups lived in the most
216 deprived areas; this proportion was lowest for White British adults (15%).
217

Table 1 Characteristics of ethnic groups (column %), Health Survey for England 2011-2018

	White British	White Irish	Other White	Indian	Pakistani	Bangladeshi	Chinese	African	Caribbean	White Mixed	Other Mixed	Other	Total	p-value
Sample size (N)	44,922	440	2,876	1,382	859	368	278	804	496	477	268	1,268	54,438	
Mean age (years)	48.8	53.4	37.6	40.6	36.8	36.5	36.2	37.3	48.6	34.1	38.8	39.8	46.8	
<i>standard deviation</i>	19	17.2	12.5	14.1	13	12.4	13.9	13.1	17.4	13.4	14	14.6	18.5	
Female	50	47	52	48	50	49	53	52	56	49	51	51	50	0.103
Highest Educational qualification														
Degree or higher	25	39	44	45	29	26	53	37	23	31	40	41	28	
Lower than degree	57	33	32	38	45	45	32	48	52	54	51	43	53	
No qualifications	18	28	25	17	26	29	14	15	25	15	10	17	19	<0.001
Index of Multiple Deprivation														
Least deprived	23	18	13	15	5	3	19	5	8	12	11	13	20	
2 nd	23	15	16	17	7	7	15	9	7	16	18	16	21	
3 rd	22	20	20	20	11	8	24	12	14	17	24	17	21	
4 th	18	26	27	25	29	26	27	31	28	25	27	25	20	
Most deprived	15	21	24	23	49	56	15	43	43	30	20	29	19	<0.001
Smoking status														
Current smoker	19	20	27	8	15	18	10	7	21	29	17	12	19	
Ex-smoker	27	34	22	8	8	8	7	8	18	17	17	12	25	
Never smoker	54	47	52	84	77	74	83	85	61	54	67	76	57	<0.001
Body Mass Index (BMI) category														
Not overweight	36	38	48	44	36	44	74	36	29	46	44	43	38	
Overweight	37	35	33	38	37	42	22	33	32	31	35	35	36	
Obese	27	27	20	18	27	15	3	31	39	23	21	22	26	<0.001
Number of longstanding conditions														
None	58	57	76	70	71	67	86	77	55	68	72	71	61	
One	21	17	14	17	15	16	7	14	22	17	15	16	20	
Two or more	22	25	10	13	15	17	7	9	24	15	13	13	20	<0.001
Number of longstanding conditions (aged 40 and over, N=37,135)														
None	49	50	63	55	53	49	79	67	46	51	64	56	51	
One	23	20	17	22	19	18	10	17	24	24	16	23	22	
Two or more	28	31	20	23	29	33	12	16	30	25	20	21	27	<0.001
Number of cardiovascular conditions (aged 40 and over, N= 37,148)¹														
None	62	59	75	62	62	64	81	64	51	69	67	65	63	
One	31	32	21	26	23	22	15	28	33	22	29	26	30	
Two or more	7	9	4	12	15	15	4	7	16	9	4	9	7	<0.001
Cardiometabolic risk biomarkers (aged 16 and over with blood samples (aged 16 and over, N=24,203)²														
None	38	31	53	44	44	45	58	53	36	56	54	47	40	
One	47	52	40	42	48	47	35	38	45	39	39	43	46	
Two or more	14	17	7	15	8	9	7	10	19	6	7	10	14	<0.001

1 Doctor-diagnosed hypertension, diabetes or self-reported stroke or heart attack; 2 Raised blood pressure, raised glycated haemoglobin or raised total cholesterol;

Table 2 Characteristics of those with single and multiple limiting longstanding illness conditions (column %), HSE 2011-2018

	None	One	Two or more	Total	p-value
Sample size (N)	31,233	11,087	12,118	54,438	
Mean age (years)	41.9	50	58.3	46.8	
<i>Standard deviation</i>	16.5	18.5	18.2	18.5	
Female	49	52	54	50	<0.001
Highest educational qualification					
Degree or higher	31	27	19	28	
Lower than degree	55	54	50	53	
No qualifications	14	20	32	19	<0.001
Index of multiple deprivation					
Least deprived quintile	21	21	17	20	
2 nd quintile	21	21	20	21	
3 rd quintile	21	22	21	21	
4 th quintile	19	19	21	20	
Most deprived quintile	18	17	22	19	<0.001
Smoking status					
Current	18	19	19	19	
Ex-regular	21	28	34	25	
Never regular	61	53	47	57	<0.001
Body Mass Index (BMI) category					
Not overweight	43	34	25	38	
Overweight	37	37	35	36	
Obese	21	29	40	26	<0.001
Number of cardiovascular conditions (aged 40+, row %)¹					
None	63	21	17	100	
One condition	36	26	39	100	
Two or more conditions	9	18	73	100	<0.001
Year (row %)					
2011	61	21	18	100	
2012	63	19	18	100	
2013	62	19	20	100	
2014	61	19	20	100	
2015	62	20	19	100	
2016	60	20	21	100	
2017	58	20	22	100	
2018	59	19	22	100	
2011-2018	61	20	20	100	<0.001

¹ Doctor-diagnosed hypertension, diabetes or self-reported stroke or heart attack

220 Obesity was highest among Caribbean and African adults (39%-31%); and was lowest
221 among Chinese, Bangladeshi and Indian adults (3%-18%). The proportion of never smoking
222 was highest among African, Indian, Chinese, Pakistani and Bangladeshi adults (85%-74%)
223 and was lowest among White Irish, White and White mixed adults (47%-54%).

224

225 20% of all adults had general multimorbidity (27% of adults aged 40+); this was higher
226 among White Irish and Caribbean adults (24% and 25%, respectively), and lowest among
227 Chinese and African adults (7% and 9%) (Table 2). 7% of adults aged 40+ had
228 cardiovascular multimorbidity. This was highest among Caribbean, Pakistani and
229 Bangladeshi adults (15%) and lowest among Chinese and Other White adults (4%).

230

231 Among those aged 40+ with cardiovascular multimorbidity, the combination of diabetes and
232 hypertension was the most common dyad (70%) followed by hypertension and heart attack
233 (12%) (Figure S1). 14% of adults aged 16+ had multiple cardiometabolic risk biomarkers;
234 this varied by ethnicity, being highest among White Irish, Caribbean and Indian adults (15%-
235 19%). This largely comprised the combination of raised blood pressure and raised
236 cholesterol (76%), followed by raised cholesterol and raised glycated haemoglobin (9%)

237

238 Those with general multimorbidity were more likely to have no qualifications than those with
239 no longstanding conditions, lived in the most deprived areas, were obese or were a current
240 or former smoker ($p < 0.001$) (Table 2). Among those classified as having cardiovascular
241 multimorbidity (aged 40+), 73% were classified as having general multimorbidity ($p < 0.001$).
242 The prevalence of general multimorbidity increased steadily over the period, from 18% in
243 2011 to 22% in 2018 ($p < 0.001$).

244

245

Figure 1 General multimorbidity

246

247 In age-sex adjusted models, compared with White British adults, Other White (OR=0.66 95%
248 Confidence Interval (0.57-0.77)), Indian (0.78, (0.65-0.93)), Chinese (0.42 (0.26-0.68)),
249 African (0.63 (0.50-0.80)) and Other adults (0.76, (0.63-0.91)) had lower odds of general
250 multimorbidity. The odds for other ethnic groups were not statistically significant although
251 Bangladeshi adults had higher odds (1.34, (0.98, 1.84), $p = 0.069$) (Figure 1, Table S1). In
252 fully adjusted models, the results remained significant only for Other White (0.63 (0.53-
253 0.74)), Chinese (0.58, (0.36-0.93)) and African adults (0.54 (0.42-0.69)).

254

255 In models adjusting for single additional covariates, significant effect sizes of reduced risk
256 were strengthened after adjusting for area deprivation, whilst adjusting for current smoking
257 status reduced effect sizes. Bangladeshi adults had higher odds of general multimorbidity
258 after adjusting for smoking status or BMI status, however, this association was not significant
259 after fully adjusting for all covariates.

260

261 Among adults aged under 40, in fully adjusted models, African (0.37 (0.21-0.62)), Indian
262 (0.37 (0.24-0.57)), Other White (0.41 (0.29-0.58)), Pakistani (0.51 (0.32-0.78)) and Other
263 adults (0.50 (0.34-0.75)), had reduced odds of general multimorbidity compared with White
264 British adults (Table S2). Among adults aged 40 and over in fully adjusted models (Table
265 S3), compared with White British adults, Bangladeshi (1.27 (0.97-1.66) $p=0.080$) and
266 Pakistani adults (1.75 (1.16-2.65)) had higher odds of having general multimorbidity,
267 whereas Other White (0.81, (0.68-0.96)), Chinese adults (0.62 (0.36-1.07, $p=0.086$)) and
268 African adults had lower odds (0.63, (0.47-0.83)).

269

270 *Figure 2 Cardiovascular multimorbidity*

271

272 Figure 2 (Table S4) presents odds ratios for cardiovascular multimorbidity among adults
273 aged 40+. In age-sex adjusted models, compared with White British adults, Pakistani (4.01
274 (2.94-5.46)), Bangladeshi (3.73 (2.15-6.47)), Caribbean (2.93 (2.15-3.98)), Indian (2.52
275 (1.97-3.24)), White mixed (2.23 (1.32-3.76)), Other (1.93 (1.46-2.54)), and African adults
276 (1.86 (1.23-2.81)) had higher odds of having cardiovascular multimorbidity. In fully adjusted
277 models, the relationship remained significant for all ethnicities. The effect sizes were
278 attenuated for Caribbean (2.18 (1.58-2.99)), African (1.61 (1.05-2.47)), Pakistani (3.48 (2.52-
279 4.80)), White Mixed (1.98 (1.14-3.44)), and Bangladeshi (3.67 (1.98-6.78)) adults and
280 strengthened for Indian (3.31 (2.56-4.28)) and Other adults (2.12 (1.60-2.82)).

281

282 Broadly, introducing area-deprivation reduced effect sizes for all significant effects, whereas
283 adjusting for smoking strengthened the effects (with the exception of White Mixed adults).
284 Adjusting for BMI status had a differential effect, slightly attenuating effect sizes for African
285 and Caribbean adults whilst strengthening effect sizes for Bangladeshi, Indian, Other Mixed
286 and Pakistani adults.

287

288 *Figure 3 Multiple cardiometabolic risk biomarkers*

289

290 In sex-age adjusted models, among adults aged 16+, compared with White British adults,
291 Indian (1.54 (1.17-2.03)) and Caribbean adults (1.46 (0.98- 2.19), $p=0.064$) had greater odds
292 of multiple risk biomarkers (Figure 3). In fully adjusted models, the results were strengthened
293 for Indian adults (1.81 (1.36- 2.41)) and were no longer significant for Caribbean adults.

294

295 When based on a larger sample (not restricted to those with BMI measurements),
296 statistically significant results were apparent for similar ethnic groups in the same direction
297 (Table S5). Additionally, Indian adults had lower odds of general multimorbidity (0.78 (0.66-
298 0.91) and Caribbean adults had increased odds of multiple risk biomarkers (1.47 (1.01-
299 2.14)).

300

301

302 **Discussion**

303

304 General multimorbidity varied with age among ethnic groups, with the risk being greater
305 among the British White population aged under 40, and greater among ethnic minorities
306 (Pakistani and Bangladeshi) aged 40 and over. Age-related differences in patterns of
307 multimorbidity by ethnicity, was also found in a recent study,(22). This correlates well with
308 the increased risk of cardiovascular multimorbidity among Pakistani and Bangladeshi adults
309 aged 40 and over; however, differences between other ethnic groups and greater risk of
310 cardiovascular multimorbidity but not general multimorbidity suggest the general white
311 population has higher prevalence of non-cardiovascular longstanding conditions and/or there
312 are ethnic differences in reporting or in being diagnosed.

313

314 Chinese and Asian Indian adults were less likely to report having multimorbidity in the US,
315 however, they were more likely to have a combination of high cholesterol and hypertension,
316 risk factors for CVD consistent with our study,(9). Cultural barriers or differences in
317 considering what constitutes a long-term condition could be a possible explanation for the
318 discrepancy, which requires further investigation. In this study, among those who had
319 cardiovascular multimorbidity, around one in four did not report having general
320 multimorbidity, indicating a discrepancy between definitions, supporting this theory. Healthier
321 ethnic minorities compared with the British White majority, in terms of multimorbidity, could
322 also be due to a 'healthy migrant effect',(23). In our study, African, Chinese, Other White and
323 Indian adults (<40 years) had a lower risk of general multimorbidity. Likewise similar groups
324 were found to be less likely than the majority white population to report poorer self-rated

325 health in a different study using the HSE,(15), further highlighting the need to separate these
326 distinct ethnic groups.

327

328 Despite the reduced risk of general multimorbidity for some ethnic groups, Indian and African
329 adults and other ethnic groups (Pakistani, Bangladeshi, Caribbean and White Mixed)), were
330 at increased risk of cardiovascular multimorbidity, consistent with a study which found the
331 South Asian and Black populations to have the highest rates of cardiovascular multimorbidity
332 in East London,(4). It is established that ethnic minorities have greater cardiometabolic risk
333 factors,(24). In particular, South Asian and Black adults are more likely to have hypertension
334 and Type 2 Diabetes,(23–25). Moreover, for the South Asian population, the risks of Type 2
335 diabetes and dyslipidaemia are apparent at lower levels of BMI compared with the white
336 population,(24). We too found Indian and Bangladeshi adults to be at greater risk of
337 cardiometabolic multimorbidity despite having lower levels of obesity than the average in this
338 study. The exact mechanisms behind the differences in risk among ethnicities are not fully
339 understood but are likely due to a complex interplay between genetics and environmental
340 factors,(24).

341 The association between ethnicity and multimorbidity remained after adjusting for
342 socioeconomic factors, obesity and smoking, suggesting ethnicity to be an independent risk
343 factor for general and cardiovascular multimorbidity. Other studies which adjust for
344 socioeconomic factors have found the greater risk of multimorbidity for ethnic minorities
345 remain,(6,8) or even strengthen, when based on a younger sample,(8). In our models,
346 adjusting for area-deprivation and education tended to reduce the effect, whereas
347 adjustment for smoking and obesity tended to strengthen the effect. In general, after full
348 adjustment, the net effect was a slight reduction in risk, although the Indian population was
349 an exception with a slightly stronger effect after full adjustment, suggesting that the risk for
350 the Indian population is greater despite lower levels of smoking and obesity, and higher
351 levels of educational attainment, which requires further investigation.

352

353 Multiple cardiometabolic risk biomarkers exceeding recommended targets, particularly
354 among Indian and Caribbean adults, could be one explanation behind the increase in
355 cardiovascular multimorbidity. Exceeding targets could be exacerbated by undiagnosed
356 diseases or poorer management, such as lack of treatment or adherence to treatment.
357 Ethnic minority groups may face barriers to treatment or health-seeking behaviours.
358 Difficulties with healthcare access are well-recognised,(5). Discrimination, which we were
359 unable to account for, was found to be a risk-factor for multimorbidity in the US,(28) and
360 metabolic syndrome in the Netherlands,(29).

361

362 **Strengths and limitations**

363

364 Strengths of this study include the use of a nationally representative sample in England,
365 where studies based on GP records may only capture people in contact with health
366 services,(5) and studies using a selective sample of volunteers may generate conservative
367 prevalence estimates of multimorbidity,(20). Furthermore, we were able to separate out
368 distinct ethnic groups where differences were found especially in terms of general
369 multimorbidity.

370

371 There are many limitations to this study. Even though eight years of data was pooled,
372 sample sizes within distinct ethnic minorities were still relatively small. Consequently, except
373 for cardiovascular multimorbidity, combinations of conditions could not be explored in detail
374 and we were unable to stratify analyses by smaller age groups. We could not adjust for
375 important covariates, such as foreign-born status, which could have suggested a healthy
376 migrant effect, as the question was not asked in the survey years assessed. Furthermore,
377 diet, physical activity, and mental health were not accounted for due to a lack of consistency
378 of data being collected in all years. The extent to which culture, including influences on diet,
379 affect multimorbidity risk could also not be assessed

380

381 There is no agreed definition of multimorbidity, and the conditions listed by participants did
382 not include conditions that may be counted in the definition elsewhere, such as alcohol
383 problems,(12) which makes comparing prevalence rates across studies challenging. Finally,
384 our study was cross-sectional therefore we cannot determine the temporal order of events
385 between variables such as deprivation and multimorbidity. However, we believe the use of a
386 nationally representative sample and the ability to assess distinct ethnic groups offset this
387 limitation, where research on ethnicity and multimorbidity is lacking.

388

389 **Conclusion**

390 Ethnic inequalities in multimorbidity vary between older and younger adults, distinct ethnic
391 groups, and are independent of social–economic status, smoking and obesity. Differences in
392 findings between general multimorbidity and cardiovascular multimorbidity suggests a
393 possible healthy migrant effect, different susceptibility to different diseases, differential use of
394 healthcare for diagnosis and management, or differences in interpretation of chronic
395 conditions. Several ethnic minority groups among the South Asian and Black ethnic groups

396 are at risk of cardiovascular multimorbidity, which may be exacerbated by underdiagnosis
397 and/or poorer management of cardiometabolic risk factors.

398

399 **Acknowledgements**

400 We thank NHS Digital for funding The Health Survey for England, participants, fieldwork staff
401 and colleagues at Natcen, for collecting, processing the data and creating the dataset

402 **Contributors**

403 LNF designed and conceived the idea of the study, conducted analyses and drafted the
404 manuscript. JM, LM, SS advised on analyses, and contributed to revisions of the manuscript
405 and approved the final version.

406 **Role of the funding source**

407 The HSE is funded by NHS Digital. The authors are funded to produce HSE reports but this
408 study received no specific funding. NHS Digital had no role in study design, interpretation or
409 production of the manuscript.

410 **Competing interests**

411 The authors have no competing interests to declare.

412 **Ethical approval**

413 Ethical approval was obtained from the following Research Ethics Committees (REC): HSE
414 2011 and 2012: Oxford A REC: 10/H0604/56; HSE 2013 and 2014: Oxford A REC:
415 12/sc/0317; HSE 2015: West London NRES Committee: 14/LO/0862; HSE 2016:
416 Nottingham REC: 15/EE/0299, HSE 2017: East of England Research Ethics Committee
417 (Reference no 15/EE/0229), HSE 2018: East Midlands Nottingham 2 Research Ethics
418 Committee (Reference no. 15/EM/0254).. Ethical approval was obtained ahead of data
419 collection, no further ethical approval is needed for secondary analyses.

420 **Patient consent**

421 Verbal or written consent was obtained from each participant for their involvement ahead of
422 data collection for the various stages of the HSE

423 **Data availability statement**

424 The Health Survey for England is available to UK Academic Institutions from the UK Data
425 Archive subject to their end-user license. This study utilised a dataset used for the Health
426 Survey for England report on Ethnicity, which the authors were funded to produce.

427 **Transparency statement**

428 The lead author affirms that this manuscript is an honest, accurate, and transparent account
429 of the study being reported; no important aspects of the study have been omitted; and that
430 any discrepancies from the study as planned have been explained.

431

432

433 **References**

- 434 1. Pearson-Stuttard J, Ezzati M, Gregg EW. Multimorbidity—a defining challenge for
435 health systems. *The Lancet Public Health*. 2019 Dec 1;4(12):e599–600.
- 436 2. Jani BD, Hanlon P, Nicholl BI, McQueenie R, Gallacher KI, Lee D, et al. Relationship
437 between multimorbidity, demographic factors and mortality: findings from the UK
438 Biobank cohort. *BMC Medicine*. 2019 Apr 10;17(1):74.
- 439 3. McQueenie R, Foster HME, Jani BD, Katikireddi SV, Sattar N, Pell JP, et al.
440 Multimorbidity, polypharmacy, and COVID-19 infection within the UK Biobank cohort.
441 *PLOS ONE*. 2020 Aug 20;15(8):e0238091.
- 442 4. Mathur R, Hull SA, Badrick E, Robson J. Cardiovascular multimorbidity: the effect of
443 ethnicity on prevalence and risk factor management. *Br J Gen Pract*. 2011
444 May;61(586):e262-270.
- 445 5. Hayanga B, Stafford M, Bécarea L. Ethnic Inequalities in Healthcare Use and Care
446 Quality among People with Multiple Long-Term Health Conditions Living in the United
447 Kingdom: A Systematic Review and Narrative Synthesis. *International Journal of
448 Environmental Research and Public Health* [Internet]. 2021;18(23). Available from:
449 <https://www.mdpi.com/1660-4601/18/23/12599>
- 450 6. Verest WJGM, Galenkamp H, Spek B, Snijder MB, Stronks K, van Valkengoed IGM.
451 Do ethnic inequalities in multimorbidity reflect ethnic differences in socioeconomic
452 status? The HELIUS study. *European Journal of Public Health*. 2019 Aug 1;29(4):687–
453 93.
- 454 7. Ashworth M, Durbaba S, Whitney D, Crompton J, Wright M, Dodhia H. Journey to
455 multimorbidity: longitudinal analysis exploring cardiovascular risk factors and
456 sociodemographic determinants in an urban setting. *BMJ Open*. 2019 Dec
457 1;9(12):e031649.
- 458 8. Johnson-Lawrence V, Zajacova A, Sneed R. Education, race/ethnicity, and
459 multimorbidity among adults aged 30-64 in the National Health Interview Survey. *SSM
460 Popul Health*. 2017 Dec;3:366–72.
- 461 9. Zhang Y, Misra R, Sambamoorthi U. Prevalence of Multimorbidity among Asian Indian,
462 Chinese, and Non-Hispanic White Adults in the United States. *Int J Environ Res Public
463 Health*. 2020 May 11;17(9).

- 464 10. Gallacher KI, McQueenie R, Nicholl B, Jani BD, Lee D, Mair FS. Risk factors and
465 mortality associated with multimorbidity in people with stroke or transient ischaemic
466 attack: a study of 8,751 UK Biobank participants. *J Comorb*. 2018 Feb 19;8(1):1–8.
- 467 11. Ingram E, Ledden S, Beardon S, Gomes M, Hogarth S, McDonald H, et al. Household
468 and area-level social determinants of multimorbidity: a systematic review. *J Epidemiol
469 Community Health*. 2021 Mar 1;75(3):232.
- 470 12. Knies G, Kumari M. Multimorbidity is associated with the income, education,
471 employment and health domains of area-level deprivation in adult residents in the UK.
472 *Sci Rep*. 2022 May 4;12(1):7280.
- 473 13. Blümel JE, Carrillo-Larco RM, Vallejo MS, Chedraui P. Multimorbidity in a cohort of
474 middle-aged women: Risk factors and disease clustering. *Maturitas*. 2020/05/04 ed.
475 2020 Jul;137:45–9.
- 476 14. NHS Digital. Health Survey for England Additional Analyses, Ethnicity and Health,
477 2011-2019 Experimental statistics [Internet]. [cited 2022 Feb 8]. Available from:
478 [https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-
479 england-additional-analyses/ethnicity-and-health-2011-2019-experimental-statistics](https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-england-additional-analyses/ethnicity-and-health-2011-2019-experimental-statistics)
- 480 15. Mindell JS, Knott CS, Ng Fat LS, Roth MA, Manor O, Soskolne V, et al. Explanatory
481 factors for health inequalities across different ethnic and gender groups: data from a
482 national survey in England. *J Epidemiol Community Health*. 2014 Dec 1;68(12):1133.
- 483 16. Mindell J, Biddulph JP, Hirani V, Stamatakis E, Craig R, Nunn S, et al. Cohort Profile:
484 The Health Survey for England. *International Journal of Epidemiology*. 2012 Dec
485 1;41(6):1585–93.
- 486 17. NatCen Social Research, UCL. Methods [Internet]. NHS Digital; 2019 Mar. (Health
487 Survey for England 2018). Available from:
488 <https://files.digital.nhs.uk/CA/2393EF/HSE18-Methods-rep.pdf>
- 489 18. The Emerging Risk Factors Collaboration. Association of Cardiometabolic
490 Multimorbidity With Mortality. *JAMA*. 2015 Jul 7;314(1):52–60.
- 491 19. Whitty CJM, MacEwen C, Goddard A, Alderson D, Marshall M, Calderwood C, et al.
492 Rising to the challenge of multimorbidity. *BMJ*. 2020 Jan 6;368:l6964.
- 493 20. Hanlon P, Jani BD, Nicholl B, Lewsey J, McAllister DA, Mair FS. Associations between
494 multimorbidity and adverse health outcomes in UK Biobank and the SAIL Databank: A
495 comparison of longitudinal cohort studies. *PLOS Medicine*. 2022 Mar
496 7;19(3):e1003931.
- 497 21. StataCorp. 2021. Stata Statistical Software: Release 17. College Station, TX:
498 StataCorp LLC.;
- 499 22. Hayanga B, Stafford M, Saunders CL, Bécares L. Ethnic inequalities in age-related
500 patterns of multiple long-term conditions in England: analysis of primary care and
501 nationally representative survey data. *medRxiv*. 2022 Jan 1;2022.08.05.22278462.
- 502 23. Moullan Y, Jusot F. Why is the ‘healthy immigrant effect’ different between European
503 countries? *Eur J Public Health*. 2014 Aug;24 Suppl 1:80–6.

- 504 24. Dal Canto E, Farukh B, Faconti L. Why are there ethnic differences in cardio-metabolic
505 risk factors and cardiovascular diseases? *JRSM Cardiovascular Disease*. 2018 Jan
506 1;7:2048004018818923.
- 507 25. Cappuccio FP. Ethnicity and cardiovascular risk: variations in people of African
508 ancestry and South Asian origin. *J Hum Hypertens*. 1997 Sep;11(9):571–6.
- 509 26. Pham TM, Carpenter JR, Morris TP, Sharma M, Petersen I. Ethnic Differences in the
510 Prevalence of Type 2 Diabetes Diagnoses in the UK: Cross-Sectional Analysis of the
511 Health Improvement Network Primary Care Database. *Clin Epidemiol*. 2019;11:1081–8.
- 512 27. Schofield P, Saka O, Ashworth M. Ethnic differences in blood pressure monitoring and
513 control in south east London. *Br J Gen Pract*. 2011 Apr;61(585):190–6.
- 514 28. Oh H, Glass J, Narita Z, Koyanagi A, Sinha S, Jacob L. Discrimination and
515 Multimorbidity Among Black Americans: Findings from the National Survey of American
516 Life. *Journal of Racial and Ethnic Health Disparities*. 2021 Feb 1;8(1):210–9.
- 517 29. Ikram UZ, Snijder MB, Agyemang C, Schene AH, Peters RJG, Stronks K, et al.
518 Perceived Ethnic Discrimination and the Metabolic Syndrome in Ethnic Minority
519 Groups: The Healthy Life in an Urban Setting Study. *Psychosomatic Medicine*
520 [Internet]. 2017;79(1). Available from:
521 [https://journals.lww.com/psychosomaticmedicine/Fulltext/2017/01000/Perceived_Ethnic](https://journals.lww.com/psychosomaticmedicine/Fulltext/2017/01000/Perceived_Ethnic_Discrimination_and_the_Metabolic.12.aspx)
522 [_Discrimination_and_the_Metabolic.12.aspx](https://journals.lww.com/psychosomaticmedicine/Fulltext/2017/01000/Perceived_Ethnic_Discrimination_and_the_Metabolic.12.aspx)
- 523
- 524
- 525

Odds Ratio for General Multimorbidity

Model (ref:
White British)

Odds
ratio (95% CI)

White Irish

Age-sex only	1.04 (0.82, 1.31)
+ Education	0.95 (0.74, 1.20)
+ Area-deprivation	1.01 (0.80, 1.28)
+ Smoking	1.05 (0.83, 1.34)
+ BMI	0.96 (0.75, 1.22)
Fully-adjusted	0.96 (0.75, 1.22)

Other White

Age-sex only	0.66 (0.57, 0.77)
+ Education	0.66 (0.57, 0.78)
+ Area-deprivation	0.61 (0.52, 0.72)
+ Smoking	0.65 (0.56, 0.76)
+ BMI	0.69 (0.59, 0.81)
Fully-adjusted	0.63 (0.53, 0.74)

Indian

Age-sex only	0.78 (0.65, 0.93)
+ Education	0.79 (0.66, 0.95)
+ Area-deprivation	0.72 (0.60, 0.86)
+ Smoking	0.88 (0.74, 1.06)
+ BMI	0.83 (0.69, 0.99)
Fully-adjusted	0.86 (0.72, 1.04)

Pakistani

Age-sex only	1.08 (0.85, 1.36)
+ Education	1.02 (0.81, 1.28)
+ Area-deprivation	0.84 (0.67, 1.05)
+ Smoking	1.18 (0.93, 1.49)
+ BMI	1.04 (0.82, 1.32)
Fully-adjusted	0.91 (0.72, 1.15)

Bangladeshi

Age-sex only	1.34 (0.98, 1.84)
+ Education	1.24 (0.90, 1.71)
+ Area-deprivation	1.01 (0.73, 1.40)
+ Smoking	1.46 (1.06, 2.00)
+ BMI	1.47 (1.06, 2.02)
Fully-adjusted	1.21 (0.87, 1.68)

Chinese

Age-sex only	0.42 (0.26, 0.68)
+ Education	0.43 (0.27, 0.70)
+ Area-deprivation	0.42 (0.26, 0.67)
+ Smoking	0.48 (0.30, 0.77)
+ BMI	0.53 (0.33, 0.85)
Fully-adjusted	0.58 (0.36, 0.93)

African

Age-sex only	0.63 (0.50, 0.80)
+ Education	0.64 (0.50, 0.81)
+ Area-deprivation	0.49 (0.39, 0.63)
+ Smoking	0.72 (0.57, 0.91)
+ BMI	0.58 (0.46, 0.73)
Fully-adjusted	0.54 (0.42, 0.69)

Caribbean

Age-sex only	1.15 (0.93, 1.43)
+ Education	1.12 (0.90, 1.39)
+ Area-deprivation	0.90 (0.72, 1.12)
+ Smoking	1.17 (0.95, 1.46)
+ BMI	1.05 (0.85, 1.30)
Fully-adjusted	0.88 (0.70, 1.10)

White mixed

Age-sex only	1.24 (0.94, 1.63)
+ Education	1.21 (0.92, 1.61)
+ Area-deprivation	1.10 (0.83, 1.46)
+ Smoking	1.21 (0.91, 1.60)
+ BMI	1.25 (0.94, 1.67)
Fully-adjusted	1.11 (0.83, 1.48)

Other mixed

Age-sex only	0.81 (0.55, 1.19)
+ Education	0.85 (0.57, 1.26)
+ Area-deprivation	0.75 (0.50, 1.11)
+ Smoking	0.84 (0.57, 1.24)
+ BMI	0.84 (0.57, 1.26)
Fully-adjusted	0.84 (0.56, 1.26)

.25 .5 1 2 4

medRxiv preprint doi: <https://doi.org/10.1101/2022.10.03.22280637>; this version posted October 4, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Odds Ratio for Cardiovascular Multimorbidity

Model (ref:
White British)

Odds
ratio (95% CI)

bioRxiv preprint doi: <https://doi.org/10.1101/2022.10.03.22280637>; this version posted October 4, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Model	Odds ratio (95% CI)
White Irish	
Age-sex only	1.07 (0.72, 1.60)
+ Education	1.01 (0.68, 1.50)
+ Area-deprivation	1.10 (0.74, 1.63)
+ Smoking	1.10 (0.74, 1.63)
+ BMI	1.15 (0.77, 1.71)
Fully-adjusted	0.98 (0.65, 1.48)
Other White	
Age-sex only	0.80 (0.57, 1.13)
+ Education	0.82 (0.58, 1.15)
+ Area-deprivation	0.76 (0.54, 1.07)
+ Smoking	0.80 (0.57, 1.13)
+ BMI	0.84 (0.60, 1.19)
Fully-adjusted	0.82 (0.58, 1.15)
Indian	
Age-sex only	2.52 (1.97, 3.24)
+ Education	2.55 (1.98, 3.28)
+ Area-deprivation	2.37 (1.84, 3.04)
+ Smoking	2.84 (2.21, 3.66)
+ BMI	3.16 (2.46, 4.06)
Fully-adjusted	3.31 (2.56, 4.28)
Pakistani	
Age-sex only	4.01 (2.94, 5.46)
+ Education	3.64 (2.65, 5.00)
+ Area-deprivation	3.07 (2.24, 4.20)
+ Smoking	4.40 (3.22, 6.01)
+ BMI	4.11 (2.99, 5.65)
Fully-adjusted	3.48 (2.52, 4.80)
Bangladeshi	
Age-sex only	3.73 (2.15, 6.47)
+ Education	3.14 (1.79, 5.51)
+ Area-deprivation	2.66 (1.48, 4.78)
+ Smoking	4.07 (2.33, 7.10)
+ BMI	4.92 (2.77, 8.75)
Fully-adjusted	3.67 (1.98, 6.78)
Chinese	
Age-sex only	0.83 (0.29, 2.40)
+ Education	0.85 (0.29, 2.51)
+ Area-deprivation	0.86 (0.30, 2.50)
+ Smoking	0.94 (0.32, 2.71)
+ BMI	1.48 (0.52, 4.21)
Fully-adjusted	1.69 (0.59, 4.83)
African	
Age-sex only	1.86 (1.23, 2.81)
+ Education	1.95 (1.29, 2.96)
+ Area-deprivation	1.42 (0.94, 2.16)
+ Smoking	2.08 (1.37, 3.14)
+ BMI	1.70 (1.11, 2.60)
Fully-adjusted	1.61 (1.05, 2.47)
Caribbean	
Age-sex only	2.93 (2.15, 3.98)
+ Education	2.71 (1.97, 3.73)
+ Area-deprivation	2.19 (1.59, 3.01)
+ Smoking	3.02 (2.22, 4.13)
+ BMI	2.67 (1.97, 3.62)
Fully-adjusted	2.18 (1.58, 2.99)
White mixed	
Age-sex only	2.23 (1.32, 3.76)
+ Education	2.15 (1.26, 3.66)
+ Area-deprivation	1.93 (1.14, 3.27)
+ Smoking	2.22 (1.30, 3.78)
+ BMI	2.25 (1.32, 3.82)
Fully-adjusted	1.98 (1.14, 3.44)
Other mixed	
Age-sex only	0.92 (0.40, 2.12)
+ Education	1.02 (0.45, 2.33)
+ Area-deprivation	0.85 (0.37, 1.95)
+ Smoking	0.94 (0.41, 2.18)
+ BMI	1.04 (0.45, 2.39)
Fully-adjusted	1.06 (0.47, 2.39)

.25 .5 1 2 4 8 16

Odds Ratio for Multiple Biological Risk Factors

Model (ref:
White British)

Odds
ratio (95% CI)

White Irish

bioRxiv preprint doi: <https://doi.org/10.1101/2022.10.03.22280637>; this version posted October 4, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Model	Odds ratio (95% CI)
Age-sex only	0.92 (0.69, 1.43)
+ Education	0.89 (0.57, 1.39)
+ Area-deprivation	0.90 (0.58, 1.39)
+ Smoking	0.90 (0.57, 1.43)
+ BMI	0.90 (0.57, 1.43)
Fully-adjusted	0.92 (0.58, 1.46)

Other White

Age-sex only	0.84 (0.66, 1.07)
+ Education	0.88 (0.69, 1.12)
+ Area-deprivation	0.83 (0.66, 1.06)
+ Smoking	0.84 (0.66, 1.07)
+ BMI	0.92 (0.72, 1.17)
Fully-adjusted	0.95 (0.74, 1.21)

Indian

Age-sex only	1.54 (1.17, 2.03)
+ Education	1.60 (1.22, 2.11)
+ Area-deprivation	1.53 (1.16, 2.02)
+ Smoking	1.58 (1.20, 2.09)
+ BMI	1.68 (1.27, 2.24)
Fully-adjusted	1.81 (1.36, 2.41)

Pakistani

Age-sex only	0.95 (0.61, 1.48)
+ Education	0.96 (0.62, 1.50)
+ Area-deprivation	0.92 (0.59, 1.44)
+ Smoking	0.96 (0.62, 1.50)
+ BMI	0.89 (0.56, 1.40)
Fully-adjusted	0.93 (0.58, 1.47)

Bangladeshi

Age-sex only	1.20 (0.63, 2.28)
+ Education	1.19 (0.63, 2.27)
+ Area-deprivation	1.16 (0.61, 2.20)
+ Smoking	1.21 (0.64, 2.29)
+ BMI	1.36 (0.71, 2.61)
Fully-adjusted	1.42 (0.73, 2.76)

Chinese

Age-sex only	0.77 (0.41, 1.46)
+ Education	0.80 (0.42, 1.52)
+ Area-deprivation	0.77 (0.40, 1.46)
+ Smoking	0.79 (0.42, 1.50)
+ BMI	1.12 (0.59, 2.12)
Fully-adjusted	1.19 (0.63, 2.25)

African

Age-sex only	1.15 (0.72, 1.85)
+ Education	1.18 (0.73, 1.89)
+ Area-deprivation	1.12 (0.69, 1.80)
+ Smoking	1.19 (0.74, 1.91)
+ BMI	1.04 (0.64, 1.70)
Fully-adjusted	1.10 (0.67, 1.81)

Caribbean

Age-sex only	1.46 (0.98, 2.19)
+ Education	1.47 (0.98, 2.20)
+ Area-deprivation	1.42 (0.94, 2.13)
+ Smoking	1.46 (0.97, 2.18)
+ BMI	1.31 (0.87, 1.95)
Fully-adjusted	1.33 (0.89, 2.00)

White mixed

Age-sex only	0.73 (0.41, 1.28)
+ Education	0.73 (0.41, 1.30)
+ Area-deprivation	0.72 (0.41, 1.27)
+ Smoking	0.72 (0.40, 1.27)
+ BMI	0.74 (0.41, 1.31)
Fully-adjusted	0.74 (0.41, 1.32)

Other mixed

Age-sex only	0.66 (0.32, 1.35)
+ Education	0.69 (0.34, 1.42)
+ Area-deprivation	0.65 (0.32, 1.34)
+ Smoking	0.67 (0.32, 1.37)
+ BMI	0.74 (0.36, 1.53)
Fully-adjusted	0.77 (0.37, 1.58)

.125 .25 .5 1 2 4