

Title:

Low daytime light and bright night-time light are associated with psychiatric disorders: an objective light study in >85,000 UK Biobank participants

Proposed Author List:

Angus C. Burns^{1,2,3}, Daniel P. Windred¹, Martin K. Rutter^{4,5}, Patrick Olivier⁶, Céline Vetter⁷, Richa Saxena^{2,3,8}, Jacqueline M. Lane^{2,3,9}, Andrew J. K. Phillips^{1*}, Sean W. Cain^{1*}

Affiliations:

¹Turner Institute for Brain and Mental Health, School of Psychological Sciences, Monash University, Melbourne, VIC, Australia;

²Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, 02114, USA;

³Broad Institute, Cambridge, MA, USA;

⁴Division of Endocrinology, Diabetes & Gastroenterology, School of Medical Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester, UK;

⁵Diabetes, Endocrinology and Metabolism Centre, Manchester University NHS Foundation Trust, Manchester Academic Health Science Centre, Manchester, UK;

⁶Action Lab, Faculty of Information Technology, Monash University, Clayton, VIC, Australia.

⁷Department of Integrative Physiology, University of Colorado Boulder, Boulder, CO, USA;

⁸Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, MA, 02114, USA;

⁹Division of Sleep and Circadian Disorders, Brigham and Women's Hospital, Boston, MA, 02115, USA

*Equal contribution

Corresponding Authors:

Angus C. Burns (angus.burns@monash.edu), Sean W. Cain (sean.cain@monash.edu)

Abstract.

Circadian rhythm disturbance is a common feature of many psychiatric disorders. Light is the primary input to the circadian clock, with daytime light strengthening rhythms and night light disrupting them. Therefore, habitual light exposure may represent an environmental risk factor for susceptibility to psychiatric disorders. We performed the largest to-date cross-sectional analysis of light, sleep, physical activity, and mental health ($n = 86,772$ adults; aged 62.4 ± 7.4 years; 57% women). We examined the independent association of day and night light exposure with covariate-adjusted risk for psychiatric disorders and self-harm. Greater night light exposure was associated with increased risk for major depressive disorder, generalized anxiety disorder, PTSD, psychosis, bipolar disorder, and self-harm behavior. Independent of night light, greater day light exposure was associated with reduced risk for major depressive disorder, PTSD, psychosis, and self-harm behavior. These findings were robust to adjustment for sociodemographics, photoperiod, physical activity, and sleep quality. Avoiding light at night and seeking light during the day may be a simple and effective, non-pharmacological means of broadly improving mental health.

Introduction.

Healthy circadian rhythms are essential for mental health and wellbeing¹. Many psychiatric disorders are characterized by disrupted circadian rhythms and sleep^{2,3}. In humans, a central circadian (~24-hour) clock in the suprachiasmatic nuclei (SCN) of the hypothalamus regulates the timing of basic cellular functions⁴, physiology, cognition, and behavior^{5,6}. Rhythms within the SCN are regulated by daily light exposure patterns. This biological system evolved under predictable conditions of bright light during the day and darkness at night to ensure stable, robust rhythms^{7,8}. Humans in modern, industrialized societies challenge this biology, spending ~90% of the day indoors under electric lighting⁹ which is dim during the day and bright at night compared to natural light/dark cycles¹⁰. Deviations from our natural light/dark cycle lead to disrupted circadian rhythms and therefore could contribute to adverse psychiatric outcomes.

Here, we report the largest sample to-date ($n = 86,772$) of objective recordings of individual, 24-hour light exposure data. We investigated the association of day and night light with a constellation of psychiatric disorders that feature circadian rhythm disturbance, while controlling for confounding variables in multivariable models. We tested two primary hypotheses: (i) greater light exposure in the day is associated with lower risk for psychiatric disorders and better mood; and (ii) greater light exposure at night is associated with higher risk for psychiatric disorders and poorer mood. These hypotheses were motivated by the known effects of day and night light exposure on the human circadian system and the well-established links between circadian disruption and psychiatric disorders.

Results.

Of the 103,720 UK Biobank participants that completed the actigraphy assessment, we excluded those with poor quality or unreliable accelerometry, sleep, and light data (detailed in Supplementary Methods and Supplementary Figure 1), leaving 86,772 participants. Of these, 86,631 had complete data for day and night light as well as the covariates in our fully adjusted Model 3. Characteristics of the sample and their missingness are presented in Table 1 for the lowest (Q1 & Q2) and highest (Q3 & Q4) quartiles of day and night light exposure. The light measurements showed good reliability, intra-class correlation (ICC) = 0.82 (95% CI 0.81-0.83; see Supplementary Methods). The analytical sample comprised participants who completed both the MHQ and actigraphy assessment and sample sizes varied depending on the outcome variable assessed, ranging from 26,956 to 61,442 for Model 1 and 26,824 to 61,147 in Model 3. Summary statistics for outcome variables derived from the MHQ are detailed in Supplementary Tables 1 and 2.

In fully adjusted regression models (Model 3, adjusted for age, sex, ethnicity, photoperiod, employment, and physical activity; Figure 1 & Table 2), higher night light exposure was associated with higher odds of major depressive disorder ($\chi^2 P < 0.0001$; fourth quartile odds ratio (OR_{Q4}) = 1.30, 95% CI = 1.23-1.38), self-harm ($\chi^2 P < 0.0001$; OR_{Q4} = 1.27, 95% CI = 1.14-1.42), generalized anxiety disorder ($\chi^2 P = 0.0001$; OR_{Q4} = 1.23, 95% CI = 1.11-1.36), PTSD ($\chi^2 P < 0.0001$; OR_{Q4} = 1.34, 95% CI = 1.22-1.48), and psychosis ($\chi^2 P = 0.0009$; OR_{Q4} = 1.21, 95% CI = 1.09-1.34). There was no overall association with bipolar disorder ($\chi^2 P = 0.18$), however those in the brightest night light quartile had 1.20 (95% CI = 1.02-1.42) times higher risk. Greater night light exposure was associated with higher scores on the PHQ-9 ($\chi^2 P < 0.0001$; Standardized $\beta_{Q4} = 0.13$, 95% CI = 0.11-0.15), GAD-7 ($\chi^2 P < 0.0001$; $\beta_{Q4} = 0.07$, 95% CI = 0.05-0.09), PCL-6 ($\chi^2 P < 0.0001$; $\beta_{Q4} = 0.11$, 95% CI = 0.08-0.14), and

lower wellbeing scores ($\chi^2 P < 0.0001$; $\beta_{Q4} = -0.11$, 95% CI = -0.13--0.08; Figure 2, Supplementary Table 3). The results for Model 1 (unadjusted) and Model 2 (adjusted for age, sex, ethnicity, and photoperiod) were similar in terms of direction, strength, and significance to Model 3.

Higher day light exposure (Model 3; Figure 1 & Table 3), was associated with lower odds of major depressive disorder ($\chi^2 P < 0.0001$; $OR_{Q4} = 0.81$, 95% CI = 0.76-0.87), self-harm ($\chi^2 P = 0.0001$; $OR_{Q4} = 0.76$, 95% CI = 0.67-0.87), PTSD ($\chi^2 P = 0.01$; $OR_{Q4} = 0.82$, 95% CI = 0.73-0.92), and psychosis ($\chi^2 P < 0.0001$; $OR_{Q4} = 0.69$, 95% CI = 0.61-0.79). There was no association of day light exposure with generalized anxiety disorder ($\chi^2 P = 0.61$; $OR_{Q4} = 0.96$, 95% CI = 0.85-1.09) or lifetime bipolar disorder ($\chi^2 P = 0.22$; $OR_{Q4} = 0.86$, 95% CI = 0.70-1.05). Figure 2 and Supplementary Table 4 show that greater day light exposure was associated with lower scores on the PHQ-9 ($\chi^2 P < 0.0001$; $\beta_{Q4} = -0.09$, 95% CI = -0.12--0.07), GAD-7 ($\chi^2 P < 0.0001$; $\beta_{Q4} = -0.05$, 95% CI = -0.08--0.03), PCL-6 ($\chi^2 P = 0.002$; $\beta_{Q4} = -0.08$, 95% CI = -0.12--0.04), and higher wellbeing scores ($\chi^2 P < 0.0001$; $\beta_{Q4} = 0.08$, 95% CI = 0.05-0.11). The results for Model 1 (unadjusted) and Model 2 (adjusted for age, sex, ethnicity, and photoperiod) were similar in terms of direction and significance to Model 3. However, ORs and betas tended to be smaller in Model 3 after the addition of physical activity and employment covariates.

A series of sensitivity analyses were completed. First, we re-ran Model 3 regressions excluding participants who reported doing shift work ($n = 6,840$, 7.9%). The results for both psychiatric disorders and symptom severity scores were unchanged in this subsample (Supplementary Tables 5 and 6). The second sensitivity analysis examined whether observed associations of day and night light with symptom severity scales and wellbeing were driven by the presence of clinical subgroups. We re-ran these analyses excluding participants with relevant disorders (see Methods), adjusting for the same covariates in Model 3. The associations were unchanged, excepting small variations in strength (Supplementary Table 7). The third sensitivity analysis examined whether the associations between light and psychiatric outcomes were independent of sleep characteristics. After adjustment for actigraphically derived sleep duration and sleep efficiency in addition to the Model 3 covariates (Model 4) the association of night light exposure with psychosis was no longer significant ($\chi^2 P < 0.09$; $OR_{Q4} = 1.12$, 95% CI = 1.01-1.25). All other associations of day and night light with psychiatric outcomes and symptom severity scales were unchanged, though associations tended to be weaker (Supplementary Tables 8 and 9). Finally, to assess whether residential density as a measure of urbanicity could explain the light and mood relationship, we conducted an additional sensitivity analysis (Model 5) adjusting for residence type (urban vs. rural) as well as the covariates in Model 3. Adjustment for urbanicity did not change any of the associations of day and night light with psychiatric outcomes and symptom severity scales in strength or significance (Supplementary Tables 10 and 11).

Discussion.

We found that objectively measured light exposure patterns under free-living conditions were associated with the risk for psychiatric disorders and the severity of mood symptoms. Brighter light at night was associated with a greater risk for major depressive disorder, self-harm behavior, PTSD, psychosis, generalized anxiety disorder, and bipolar disorder, as well as poorer self-reported mood and wellbeing. Conversely, brighter light in the day was associated with lower odds of major depressive disorder, self-harm behavior, PTSD, and psychosis, as well as better self-reported mood and wellbeing. Remarkably, these associations

were independent and additive. For example, greater night light exposure was associated with increased odds of major depressive disorder even for those in the brightest day light quartile. These associations were also independent of demographic, physical activity, photoperiod, and employment covariates. Sensitivity analyses showed these findings to be consistent when accounting for shift work, sleep, and urbanicity.

For individuals in the brightest night light quartile, we observed ~30% higher risk of MDD and self-harm, while individuals in the brightest day light quartile had ~20% lower risk of MDD and self-harm. Night light exposure was also associated with poorer self-reported mood and wellbeing, while day light exposure was associated with better mood and wellbeing. Previous studies have reported an association of night light exposure with low mood¹¹ and one study has linked night light to MDD risk, though this study examined group-level, outdoor light at night which may not be a good proxy for individual, ambient light levels¹². Other limitations of these studies were generally small sample sizes, poor control for confounders such as physical activity and sleep quality, and no studies considered the independent effects of day and night light. Conversely, day light therapy has long been shown to be efficacious in treating depression¹³ and has been shown to significantly enhance treatment efficacy when combined with an SSRI¹⁴. Fewer studies have linked free-living day light exposure to MDD risk, though one study linked self-reported time spent in outdoor light with lower risk¹⁵. Depression has long been associated with circadian disruption. Patients with depression have both delayed and low amplitude circadian rhythms^{16,17} which are reversed in recovery¹⁸. The severity of mood symptoms and the duration of depressive episodes are greater in those who experience circadian rhythm disturbance^{16,19} and the presence of circadian rhythm disturbance in depression is predictive of recurrence²⁰. Our findings are consistent with the known time-dependent effects of light on the properties of the circadian system, such that light at night tends to delay rhythms and reduce circadian amplitude, whereas early morning and day light tends to advance rhythms and boost circadian amplitude^{8,21,22}. Therefore, the euthymic effect of bright day light and dim night light exposure may occur by boosting the amplitude and advancing the timing of circadian rhythms, correcting the delayed and blunted rhythms seen in depression^{17,23}. Seeking greater day light and minimizing night light exposure could be a simple means of improving depression trajectories by treating underlying circadian disturbance.

In people with high levels of light exposure at night, we observed ~20% higher risk for bipolar disorder. Bipolar disorder has long been associated with dampened amplitude of behavioral rhythms and more variable circadian timing^{3,24}. A recent study found that brighter night light exposure predicted manic/hypomanic episodes in bipolar patients²⁵ and outdoor light at night has been associated with bipolar disorder¹². Hypersensitivity of the circadian system to light at night has been proposed to be a trait marker of bipolar disorder²⁶. Drugs used to treat bipolar disorder reduce the sensitivity of the circadian system to light²⁷, suggesting that reducing the effects of light at night on the circadian system may play a role in recovery. Consistent with this, nighttime dark therapy and wearing blue light blocking glasses at night are effective at reducing mania in patients^{28,29}. We did not see an association of bipolar disorder with day light exposure. This finding is novel, as the association of day light with manic symptoms in bipolar disorder and the efficacy of day light therapy on manic episodes has not been examined³⁰. The avoidance of light at night specifically may be beneficial in mitigating risk for bipolar disorder.

We found both an adverse association of night light exposure and a beneficial association of day light exposure with PTSD risk and symptom severity. To our knowledge, no studies have

examined the association of free-living light exposure patterns in the day or night to PTSD risk. There is some evidence of disturbed circadian rhythms in PTSD. Delayed activity rhythms are associated with more severe PTSD³¹ symptoms while lower urinary melatonin rhythm amplitude after a trauma exposure predicts a higher risk for PTSD³², a finding replicated in military personnel³³. Our results suggest bright night and dim day light exposure may be antecedent factors leading to blunted and delayed rhythms. This is supported by evidence that day light therapy may be an effective treatment for PTSD symptoms³⁴. Avoidance of light at night and seeking bright day light after trauma could reduce the risk of developing PTSD or the severity of symptoms in those with the disorder.

Finally, bright night light exposure was associated with ~20% increased risk for psychosis, while bright day light exposure was associated with ~30% reduced risk for psychosis. There is also little data linking free-living light exposure in the day or night to psychosis and psychotic disorders, despite sleep and circadian rhythm disruption being common features of patients on and off medication¹. One small study found that patients with schizophrenia had lower daytime light exposure and in a natural experiment observed that boosting daytime light exposure could normalize the sleep and circadian disruption seen in the disorder³⁵. Studies of day light therapy and schizophrenia have reported mixed results, though these studies have been small and further, more rigorous trials are needed^{36,37}. In addition to day light, our findings point to night light as a novel therapeutic target for psychosis.

Beyond effects on the circadian clock, non-visual photoreception is appreciated to have a direct effect on mood via projections to brain areas implicated in mood regulation. Light exposure acutely enhances both mood and alertness^{38,39}. Intrinsically photosensitive retinal ganglion cells (ipRGCs) expressing the photopigment melanopsin are the primary input of light information to the circadian clock in the SCN⁴⁰. These cells also project to the medial amygdala and lateral habenula, brain areas implicated in depression, and these projections mediate the acute euthymic effect of light exposure^{41,42}. The direct effects of light may partially explain the association of day light with lower MDD and self-harm risk, though a mechanism for direct effects of day light on other disorders such as PTSD, psychosis, and bipolar disorder is unclear. As mood is generally poorer in the night/early morning hours, people may seek out the acute euthymic effects of light at night. Though this may immediately improve mood, it would lead to circadian disruption in the long term and could perpetuate mental illness. This represents a potential challenge for promoting healthy light behaviors.

Taken together, our findings are consistent with bright day light and low night light strengthening circadian rhythms as an antecedent to more robust mental health. Patterns of bright day light and low night light serve to enhance the amplitude and stability of the circadian clock as well as align its activity appropriately with daily activities^{7,8,21}. As modern humans spend ~90% of the day indoors⁹, our light exposure patterns are typically less bright in the day and more bright at night than naturalistic patterns across our evolutionary history¹⁰. Addressing this deviation from our natural light/dark cycles may improve the general mental health of people in industrialized societies.

This study has a number of important limitations. First, the findings we report are cross-sectional. While there are well-supported causal mechanisms linking bright night light and dim day light with circadian disruption, and linking circadian disruption with mental health, we acknowledge the possibility of reverse causation and that longitudinal studies will be needed to establish the temporality of the associations we observed. However, the robustness

of our findings to adjustment for confounders including physical activity and sleep provides support for our interpretation. Second, light monitoring was performed using a wrist-worn device, which is not designed to measure light at the ocular level. The data therefore provide a coarse estimate of the actual effects of light on the circadian system. Third, the actigraphy data and outcome variables were not measured simultaneously, with the latter measured an average of 1.86 years later. We note that we did observe good reliability of within-individual repeat light exposure assessments completed over the course of a year, suggesting that light measurements at one timepoint are a good proxy for other timepoints.

This study is the first to report an analysis of objectively measured light exposure data from the UK Biobank and is the largest examination of objective light exposure and mental health to date. Our findings demonstrate a consistent association of healthy light exposure patterns with better psychiatric outcomes. These results suggest that brighter days and darker nights may be a simple, freely available, non-pharmacological intervention to enhance mental health.

Conflicts of interest

AJKP and SWC have received research funding from Delos and Versalux, and they are co-founders and co-directors of Circadian Health Innovations PTY LTD. SWC has also received research funding from Beacon Lighting and has consulted for Dyson. PO was a co-founder of Axivity Ltd and a Director until 2015. CV is a board member of the Working Time Society and a research committee member for DiME. ACB: none; DPW: none; MKR: none; RS: none; JML: none.

Data availability statement

The data used in this study are available in the UK Biobank resource. Derived data supporting the findings of this study are available from the corresponding author (ACB) on request at angus.burns@monash.edu.

Methods.

Study design and participants

In this cross-sectional study we drew on the UK Biobank prospective general population cohort which contains more than 502,000 UK residents recruited via National Health Service (NHS) patient registers from 2006 to 2010. The study population is described in detail elsewhere^{43,44}. Accelerometry and light data were measured in a subset of 103,720 participants in 2013-2015 and a separate subset of 157,366 participants completed an online Mental Health Questionnaire (MHQ) in 2016-2017. Participants who accepted the invitation to join the UK Biobank cohort provided written, informed consent and the UK Biobank has generic ethical approval from the North West Multi-Center Research Ethics Committee (ref 11/NW/03820).

Measures

Light exposure

In 2013, 236,519 UK Biobank participants were invited to participate in a seven day physical activity and light monitoring study. Of these participants, 103,720 (43.9%) accepted, and returned the accelerometer to the UK Biobank. Participants who accepted the invitation received a wrist-worn AX3 triaxial accelerometer (Axivity, Newcastle upon Tyne, UK) with in-built light sensor (APDS9007 silicon photodiode sensor; spectral sensitivity $\lambda = 470\text{-}650\text{nm}$) and were asked to wear the device on their dominant wrist for seven days under free-living conditions.

Quality control and definition of light exposure predictors was completed with custom R (version 4.1.0) scripts detailed in the Supplementary Methods. Briefly, the daily light profiles of participants meeting quality control criteria were entered into a factor analysis to identify independent patterns in light exposure in the sample. Factor analysis supported the extraction of day (7.30am-8.30pm) and night (12.30am-6am) factors. Light exposure during the day had a small positive correlation with light exposure at night ($r_s = 0.10$, $p < 0.0001$). Due to large positive skew in both day and night light variables (skewness coefficient > 1 for both), both variables were converted into categorical predictors for analysis by dividing them into four equal-sized quartiles in ascending brightness.

Psychiatric outcomes

A total of 339,092 UK Biobank participants were invited to complete the online Mental Health Questionnaire (MHQ) in 2016 and 157,366 completed the questionnaire. Measurement of psychiatric outcomes as part of the MHQ took place an average of 1.86 years (SD = 0.66) after the actigraphy assessment. Of the 86,772 participants with complete light data, 61,466 (70.8%) completed the UK Biobank MHQ. Definition of case-control psychiatric disorder outcomes from the MHQ are based on Composite International Diagnostic Interview (CIDI) and Diagnostic and Statistical Manual IV (DSM-IV) criteria and followed guidelines established by Davis, Coleman⁴⁵. Case/control outcomes were major depressive disorder (MDD), generalized anxiety disorder (GAD), bipolar disorder, post-traumatic stress disorder (PTSD), psychosis, and self-harm. Detailed definitions are given in the Supplementary Methods. Continuous outcomes were symptom severity scales for depression (Patient Health Questionnaire, PHQ-9), anxiety (GAD-7), PTSD (PTSD Checklist-6, PCL-6), and an overall wellbeing score indexing euthymia and eudaemonia (see Supplementary Methods).

Statistical analysis

The association between day and night light and case/control outcomes were examined with multiple logistic regression. Odds ratios (OR) and their 95% confidence intervals are reported. Multiple linear regression was used for the continuous symptom severity scales and wellbeing, with standardized beta values reported. These associations were tested hierarchically in three models with increasing adjustment for potential confounders. Each model included both day and night light categorical predictors to examine their independent effects. Likelihood-ratio χ^2 tests were used as an omnibus test of significance for the day and night light factors. Model 1 examined the unadjusted association between day and night light and psychiatric outcomes. Model 2 adjusted for age (at the time of actigraphy), sex, ethnicity (white vs. non-white; data-field 21000), and photoperiod (as a measure of seasonality, defined as the duration between sunrise and sunset at the beginning of the actigraphy assessment). Finally, model 3 additionally adjusted for employment (employed vs. unemployed; data-field 6142) and physical activity (defined as overall acceleration average

over the actigraphy period in milligravity units; data-field 90012). The physical activity variable was previously defined by the UK Biobank accelerometer expert working group⁴⁶.

A series of sensitivity analyses were completed to examine the robustness of the primary findings. The first sensitivity analysis examined whether the presence of shift workers in the sample ($n = 6,840$, 7.9%) was driving the observed associations. Model 3 regressions were re-run excluding those who reported doing shiftwork (data-field 826) to any degree (sometimes, usually, or always). The second sensitivity analysis assessed whether observed associations of light with symptom severity scales and wellbeing were driven by clinical subgroups. We re-ran these analyses excluding participants with MDD for the PHQ-9 model; with GAD for the GAD-7 model; with PTSD for the PCL-6 model; and with MDD, GAD, PTSD, bipolar disorder, or psychosis for the wellbeing model. The third sensitivity analysis examined whether the relationship between light and psychiatric outcomes was independent of objectively measured sleep duration and efficiency (Model 4; see Supplementary Methods). The fourth sensitivity analysis examined whether participants' residential population density as a measure of urbanicity could explain the light and mood relationship (Model 5). Urbanicity (data-field 20118; urban vs. rural) was defined according to the UK Office for National Statistics population density classification of participant residential postcodes where urban postcodes have a population of 10,000 or more and rural postcodes have a population of less than 10,000. Reporting of statistical analyses and results followed the STROBE guidelines.

References.

1. Walker WH, Walton JC, DeVries AC, Nelson RJ. Circadian rhythm disruption and mental health. *Translational Psychiatry* 2020; **10**(1): 28.
2. Freeman D, Sheaves B, Waite F, Harvey AG, Harrison PJ. Sleep disturbance and psychiatric disorders. *The Lancet Psychiatry* 2020; **7**(7): 628-37.
3. Lyall LM, Wyse CA, Graham N, et al. Association of disrupted circadian rhythmicity with mood disorders, subjective wellbeing, and cognitive function: a cross-sectional study of 91 105 participants from the UK Biobank. *The Lancet Psychiatry* 2018; **5**(6): 507-14.
4. Mure LS, Le HD, Benegiamo G, et al. Diurnal transcriptome atlas of a primate across major neural and peripheral tissues. *Science* 2018; **359**(6381).
5. Schmidt C, Collette F, Cajochen C, Peigneux P. A time to think: circadian rhythms in human cognition. *Cognitive neuropsychology* 2007; **24**(7): 755-89.
6. Reppert SM, Weaver DR. Coordination of circadian timing in mammals. *Nature* 2002; **418**(6901): 935-41.
7. Winfree AT. *The Geometry of Biological Time*: Springer New York; 1980.
8. Bano-Otalora B, Martial F, Harding C, et al. Bright daytime light enhances circadian amplitude in a diurnal mammal. *Proceedings of the National Academy of Sciences* 2021; **118**(22): e2100094118.
9. Klepeis NE, Nelson WC, Ott WR, et al. The National Human Activity Pattern Survey (NHAPS): a resource for assessing exposure to environmental pollutants. *Journal of Exposure Science & Environmental Epidemiology* 2001; **11**(3): 231-52.
10. Wright Kenneth P, McHill Andrew W, Birks Brian R, Griffin Brandon R, Rusterholz T, Chinoy Evan D. Entrainment of the Human Circadian Clock to the Natural Light-Dark Cycle. *Current Biology* 2013; **23**(16): 1554-8.
11. Tancredi S, Urbano T, Vinceti M, Filippini T. Artificial light at night and risk of mental disorders: A systematic review. *Science of The Total Environment* 2022; **833**: 155185.
12. Paksarian D, Rudolph KE, Stapp EK, et al. Association of Outdoor Artificial Light at Night With Mental Disorders and Sleep Patterns Among US Adolescents. *JAMA Psychiatry* 2020.
13. Golden RN, Gaynes BN, Ekstrom RD, et al. The efficacy of light therapy in the treatment of mood disorders: a review and meta-analysis of the evidence. *Am J Psychiatry* 2005; **162**(4): 656-62.
14. Lam RW, Levitt AJ, Levitan RD, et al. Efficacy of Bright Light Treatment, Fluoxetine, and the Combination in Patients With Nonseasonal Major Depressive Disorder: A Randomized Clinical Trial. *JAMA Psychiatry* 2016; **73**(1): 56-63.
15. Burns AC, Saxena R, Vetter C, Phillips AJK, Lane JM, Cain SW. Time spent in outdoor light is associated with mood, sleep, and circadian rhythm-related outcomes: A cross-sectional and longitudinal study in over 400,000 UK Biobank participants. *J Affect Disord* 2021; **295**: 347-52.
16. Emens J, Lewy A, Kinzie JM, Arntz D, Rough J. Circadian misalignment in major depressive disorder. *Psychiatry research* 2009; **168**(3): 259-61.

17. Sou etre E, Salvati E, Belugou J-L, et al. Circadian rhythms in depression and recovery: evidence for blunted amplitude as the main chronobiological abnormality. *Psychiatry Research* 1989; **28**(3): 263-78.
18. Avery DH, Wildschijdtz G, Rafaelsen OJ. Nocturnal temperature in affective disorder. *J Affect Disord* 1982; **4**(1): 61-71.
19. Hasler BP, Buysse DJ, Kupfer DJ, Germain A. Phase relationships between core body temperature, melatonin, and sleep are associated with depression severity: further evidence for circadian misalignment in non-seasonal depression. *Psychiatry Res* 2010; **178**(1): 205-7.
20. Harvey A. Sleep and circadian functioning: critical mechanisms in the mood disorders? *Annual review of clinical psychology* 2011; **7**: 297-319.
21. Jewett ME, Kronauer RE, Czeisler CA. Phase-Amplitude Resetting of the Human Circadian Pacemaker via Bright Light: A Further Analysis. *Journal of Biological Rhythms* 1994; **9**(3-4): 295-314.
22. Khalsa SBS, Jewett ME, Cajochen C, Czeisler CA. A phase response curve to single bright light pulses in human subjects. *The Journal of physiology* 2003; **549**(3): 945-52.
23. Emens J, Lewy A, Kinzie JM, Arntz D, Rough J. Circadian misalignment in major depressive disorder. *Psychiatry Res* 2009; **168**(3): 259-61.
24. McCarthy MJ, Gottlieb JF, Gonzalez R, et al. Neurobiological and behavioral mechanisms of circadian rhythm disruption in bipolar disorder: A critical multi-disciplinary literature review and agenda for future research from the ISBD task force on chronobiology. *Bipolar disorders* 2022; **24**(3): 232-63.
25. Esaki Y, Obayashi K, Saeki K, Fujita K, Iwata N, Kitajima T. Effect of nighttime bedroom light exposure on mood episode relapses in bipolar disorder. *Acta Psychiatrica Scandinavica* 2022; **146**(1): 64-73.
26. Lewy AJ, Wehr TA, Goodwin FK, Newsome DA, Rosenthal NE. Manic-depressive patients may be supersensitive to light. *Lancet (London, England)* 1981; **1**(8216): 383-4.
27. Hallam KT, Olver JS, Horgan JE, McGrath C, Norman TR. Low doses of lithium carbonate reduce melatonin light sensitivity in healthy volunteers. *The international journal of neuropsychopharmacology* 2005; **8**(2): 255-9.
28. Henriksen TE, Skrede S, Fasmer OB, et al. Blue-blocking glasses as additive treatment for mania: a randomized placebo-controlled trial. *Bipolar disorders* 2016; **18**(3): 221-32.
29. Barbini B, Benedetti F, Colombo C, et al. Dark therapy for mania: a pilot study. *Bipolar disorders* 2005; **7**(1): 98-101.
30. Takeshima M, Utsumi T, Aoki Y, et al. Efficacy and safety of bright light therapy for manic and depressive symptoms in patients with bipolar disorder: A systematic review and meta-analysis. *Psychiatry and Clinical Neurosciences* 2020; **74**(4): 247-56.
31. Sandahl H, Baandrup L, Vindbjerg E, Jennum P, Carlsson J. Social zeitgebers and circadian dysrhythmia are associated with severity of symptoms of PTSD and depression in trauma-affected refugees. *European Archives of Psychiatry and Clinical Neuroscience* 2021; **271**(7): 1319-29.
32. McFarlane AC, Barton CA, Briggs N, Kennaway DJ. The relationship between urinary melatonin metabolite excretion and posttraumatic symptoms following traumatic injury. *J Affect Disord* 2010; **127**(1-3): 365-9.
33. Paul MA, Love RJ, Jetly R, et al. Blunted Nocturnal Salivary Melatonin Secretion Profiles in Military-Related Posttraumatic Stress Disorder. *Frontiers in psychiatry* 2019; **10**: 882.
34. Youngstedt SD, Kline CE, Reynolds AM, et al. Bright Light Treatment of Combat-related PTSD: A Randomized Controlled Trial. *Military Medicine* 2022; **187**(3-4): e435-e44.
35. Skeldon AC, Dijk D-J, Meyer N, Wulff K. Extracting Circadian and Sleep Parameters from Longitudinal Data in Schizophrenia for the Design of Pragmatic Light Interventions. *Schizophrenia Bulletin* 2021; **48**(2): 447-56.

36. Aichhorn W, Stelzig-Schoeler R, Geretsegger C, Stuppaeck C, Kemmler G. Bright light therapy for negative symptoms in schizophrenia: a pilot study. *The Journal of clinical psychiatry* 2007; **68**(7): 2142.
37. Roopram SM, Burger AM, van Dijk DA, Enterman J, Haffmans J. A pilot study of bright light therapy in schizophrenia. *Psychiatry Research* 2016; **245**: 317-20.
38. Cajochen C, Zeitzer JM, Czeisler CA, Dijk D-J. Dose-response relationship for light intensity and ocular and electroencephalographic correlates of human alertness. *Behavioural Brain Research* 2000; **115**(1): 75-83.
39. Vandewalle G, Schwartz S, Grandjean D, et al. Spectral quality of light modulates emotional brain responses in humans. *Proceedings of the National Academy of Sciences* 2010; **107**(45): 19549-54.
40. Hattar S, Liao H-W, Takao M, Berson DM, Yau K-W. Melanopsin-containing retinal ganglion cells: architecture, projections, and intrinsic photosensitivity. *Science* 2002; **295**(5557): 1065-70.
41. Fernandez DC, Fogerson PM, Lazzerini Ospri L, et al. Light Affects Mood and Learning through Distinct Retina-Brain Pathways. *Cell* 2018; **175**(1): 71-84.e18.
42. LeGates TA, Fernandez DC, Hattar S. Light as a central modulator of circadian rhythms, sleep and affect. *Nature Reviews Neuroscience* 2014; **15**(7): 443-54.
43. Collins R. What makes UK Biobank special? *The Lancet* 2012; **9822**(379): 1173-4.
44. Sudlow C, Gallacher J, Allen N, et al. UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLOS Medicine* 2015; **12**(3): e1001779.
45. Davis KAS, Coleman JRI, Adams M, et al. Mental health in UK Biobank - development, implementation and results from an online questionnaire completed by 157 366 participants: a reanalysis. *BJPsych open* 2020; **6**(2): e18.
46. Doherty A, Jackson D, Hammerla N, et al. Large Scale Population Assessment of Physical Activity Using Wrist Worn Accelerometers: The UK Biobank Study. *PLOS ONE* 2017; **12**(2): e0169649.

Figures




Figure 1. Associations of day and night light exposure with psychiatric disorders and self-harm. Coefficients plot for day and night-time light associations with (A) major depressive disorder, (B) self-harm behavior, (C) generalized anxiety disorder, (D) PTSD, (E) bipolar disorder and (F) psychosis. Coefficients represent the odds ratios \pm the SEM (inner error bar) and 95% CI (outer error bar) for each quartile of day and night light exposure relative to the low light (Q1) referent. Three models are presented with increasing adjustment for confounders: Model 1 (green) is unadjusted, Model 2 (blue) adjusts for age, sex, ethnicity and photoperiod and Model 3 (red) additionally adjusts for employment and physical activity.




Figure 2. Associations of day and night light exposure with symptom severity scales and wellbeing. Coefficients plot for day and night-time light associations with (A) PHQ-9 score, (B) wellbeing score, (C) GAD-7 score and (D) PCL-6 score. Coefficients represent the standardized betas \pm the SEM (inner error bar) and 95% CI (outer error bar) for each quartile of day and night light exposure relative to the low light (Q1) referent. Three models are presented with increasing adjustment for confounders: Model 1 (green) is unadjusted, Model 2 (blue) adjusts for age, sex, ethnicity and photoperiod and Model 3 (red) additionally adjusts for employment and physical activity.

Tables

Table 1. Demographic characteristics of participants by low and high day and night light exposure

	Day light			Night light		
	Low Q1 & Q2 N=43386	High Q3 & Q4 N=43386	<i>p</i> value	Low Q1 & Q2 N=43387	High Q3 & Q4 N=43385	<i>p</i> value
Age						
Mean (SD)	62.0 (7.94)	62.7 (7.71)	<0.0001	62.6 (7.86)	62.1 (7.79)	<0.0001
Median [IQR]	62.9 [55.7, 68.4]	64.0 [57.0, 68.7]		63.8 [56.6, 68.8]	63.2 [56.1, 68.3]	
Sex						
Female	25159 (58.0%)	24321 (56.1%)	<0.0001	25352 (58.4%)	24128 (55.6%)	<0.0001
Male	18227 (42.0%)	19065 (43.9%)		18035 (41.6%)	19257 (44.4%)	
BMI						
Mean (SD)	26.7 (4.57)	26.7 (4.47)	0.21	26.3 (4.31)	27.1 (4.68)	<0.0001
Median [IQR]	26.0 [23.6, 29.1]	26.1 [23.6, 29.0]		25.7 [23.3, 28.6]	26.4 [23.9, 29.5]	
Missing	102 (0.2%)	83 (0.2%)		83 (0.2%)	102 (0.2%)	
White ethnicity						
	41679 (96.1%)	42226 (97.3%)	<0.0001	42233 (97.3%)	41672 (96.1%)	<0.0001
Missing	150 (0.3%)	141 (0.3%)		136 (0.3%)	155 (0.4%)	
Employment						
No	15912 (36.7%)	17412 (40.1%)	<0.0001	17353 (40.0%)	15971 (36.8%)	<0.0001
Yes	27474 (63.3%)	25974 (59.9%)		26034 (60.0%)	27414 (63.2%)	
Shiftwork						
No	39780 (91.7%)	40152 (92.5%)	<0.0001	40545 (93.4%)	39387 (90.8%)	<0.0001
Yes	3606 (8.3%)	3234 (7.5%)		2842 (6.6%)	3998 (9.2%)	
Physical activity						
Mean (SD)	27.3 (7.89)	29.0 (8.16)	<0.0001	28.1 (7.97)	28.3 (8.17)	0.0003
Median [IQR]	26.4 [21.9, 31.7]	28.1 [23.4, 33.6]		27.1 [22.6, 32.5]	27.3 [22.6, 32.8]	
Missing	39 (0.1%)	94 (0.2%)		60 (0.1%)	73 (0.2%)	
Photoperiod						
Mean (SD)	10.9 (2.87)	14.1 (2.58)	<0.0001	12.4 (3.06)	12.6 (3.28)	<0.0001
Median [IQR]	10.2 [8.47, 12.8]	14.8 [12.3, 16.4]		12.4 [9.58, 15.2]	12.8 [9.61, 15.9]	
Sleep duration						
Mean (SD)	7.56 (1.06)	7.58 (1.02)	0.004	7.79 (0.95)	7.34 (1.07)	<0.0001
Median [IQR]	7.60 [6.95, 8.21]	7.61 [6.98, 8.22]		7.81 [7.23, 8.38]	7.37 [6.72, 8.00]	
Sleep efficiency						
Mean (SD)	0.88 (0.06)	0.88 (0.06)	<0.0001	0.89 (0.06)	0.88 (0.07)	<0.0001
Median [IQR]	0.89 [0.85, 0.92]	0.90 [0.86, 0.92]		0.90 [0.86, 0.93]	0.89 [0.85, 0.92]	
Urban residence						
Yes	36703 (84.6%)	35533 (81.9%)	<0.0001	35687 (82.3%)	36549 (84.2%)	<0.0001
Missing	430 (1.0%)	404 (0.9%)		423 (1.0%)	411 (0.9%)	

IQR = interquartile range, SD = standard deviation. *p* values correspond to *t* tests for continuous variables and χ^2 tests for categorical variables.

Table 2. Associations between night light exposure and psychiatric outcomes

Night light	Model 1 OR (95% CI)				Model 2 aOR (95% CI)*				Model 3 aOR (95% CI) [†]			
	2 nd Quartile	3 rd Quartile	4 th Quartile	LRT <i>p</i> [‡]	2 nd Quartile	3 rd Quartile	4 th Quartile	LRT <i>p</i> [‡]	2 nd Quartile	3 rd Quartile	4 th Quartile	LRT <i>p</i> [‡]
Major depressive disorder	1.05 (1.00 – 1.11)	1.17 (1.11 – 1.24)	1.28 (1.22 – 1.36)	<0.0001	1.04 (0.98 – 1.09)	1.17 (1.11 – 1.24)	1.32 (1.25 – 1.39)	<0.0001	1.04 (0.98 – 1.10)	1.18 (1.12 – 1.25)	1.30 (1.23 – 1.38)	<0.0001
Self-harm	1.02 (0.91 – 1.14)	1.03 (0.92 – 1.15)	1.28 (1.15 – 1.42)	<0.0001	0.99 (0.88 – 1.10)	0.99 (0.88 – 1.11)	1.28 (1.15 – 1.43)	<0.0001	0.99 (0.88 – 1.11)	1.00 (0.89 – 1.12)	1.27 (1.14 – 1.42)	<0.0001
Generalized anxiety disorder	1.10 (0.99 – 1.21)	1.22 (1.10 – 1.35)	1.23 (1.11 – 1.36)	<0.0001	1.06 (0.96 – 1.18)	1.18 (1.07 – 1.31)	1.23 (1.11 – 1.36)	0.0002	1.07 (0.97 – 1.19)	1.20 (1.08 – 1.33)	1.23 (1.11 – 1.36)	0.0001
PTSD	1.03 (0.93 – 1.13)	1.19 (1.08 – 1.31)	1.35 (1.23 – 1.48)	<0.0001	1.00 (0.90 – 1.11)	1.15 (1.04 – 1.27)	1.35 (1.22 – 1.48)	<0.0001	1.00 (0.91 – 1.11)	1.16 (1.05 – 1.29)	1.34 (1.22 – 1.48)	<0.0001
Bipolar disorder	1.15 (0.98 – 1.36)	1.17 (0.99 – 1.38)	1.25 (1.06 – 1.47)	0.06	1.11 (0.94 – 1.31)	1.11 (0.94 – 1.31)	1.21 (1.02 – 1.43)	0.18	1.11 (0.94 – 1.31)	1.12 (0.95 – 1.32)	1.20 (1.02 – 1.42)	0.18
Psychosis	1.05 (0.95 – 1.17)	1.17 (1.05 – 1.29)	1.21 (1.10 – 1.35)	0.0005	1.03 (0.93 – 1.15)	1.13 (1.02 – 1.26)	1.22 (1.10 – 1.35)	0.0004	1.04 (0.93 – 1.15)	1.14 (1.03 – 1.27)	1.21 (1.09 – 1.34)	0.0009

Note: logistic regression was used for all outcomes and odds ratios (ORs) are presented with their 95% confidence intervals for each ascending night light quartile relative to the low light (Q1) referent. Model 1 is unadjusted. *Model 2 adjusted for age, sex, ethnicity, and photoperiod as a measure of seasonality. [†]Model 3 is additionally adjusted for employment and physical activity. Bolded coefficients indicate significance at the .05 level.

Table 3. Associations between day light exposure and psychiatric outcomes

Day light	Model 1 OR (95% CI)				Model 2 aOR (95% CI)*				Model 3 aOR (95% CI)†			
	2 nd Quartile	3 rd Quartile	4 th Quartile	LRT <i>p</i> ‡	2 nd Quartile	3 rd Quartile	4 th Quartile	LRT <i>p</i> ‡	2 nd Quartile	3 rd Quartile	4 th Quartile	LRT <i>p</i> ‡
Major depressive disorder	0.94 (0.90 – 1.00)	0.91 (0.86 – 0.96)	0.80 (0.76 – 0.85)	<0.0001	0.91 (0.86 – 0.97)	0.85 (0.80 – 0.90)	0.75 (0.70 – 0.80)	<0.0001	0.93 (0.88 – 0.99)	0.88 (0.83 – 0.93)	0.81 (0.76 – 0.87)	<0.0001
Self-harm	0.87 (0.78 – 0.97)	0.81 (0.73 – 0.90)	0.77 (0.69 – 0.85)	<0.0001	0.83 (0.75 – 0.93)	0.76 (0.67 – 0.85)	0.71 (0.63 – 0.81)	<0.0001	0.85 (0.76 – 0.94)	0.78 (0.70 – 0.88)	0.76 (0.67 – 0.87)	0.0001
Generalized anxiety disorder	1.00 (0.90 – 1.10)	0.94 (0.85 – 1.03)	0.91 (0.82 – 1.00)	0.14	0.98 (0.89 – 1.09)	0.91 (0.82 – 1.02)	0.91 (0.81 – 1.03)	0.28	0.99 (0.90 – 1.10)	0.93 (0.84 – 1.04)	0.96 (0.85 – 1.09)	0.61
PTSD	0.94 (0.86 – 1.04)	0.92 (0.83 – 1.01)	0.80 (0.73 – 0.89)	0.0001	0.91 (0.83 – 1.00)	0.86 (0.77 – 0.95)	0.75 (0.67 – 0.84)	<0.0001	0.93 (0.85 – 1.03)	0.89 (0.81 – 0.99)	0.82 (0.73 – 0.92)	0.01
Bipolar disorder	0.88 (0.75 – 1.04)	0.95 (0.81 – 1.12)	0.81 (0.69 – 0.95)	0.06	0.86 (0.73 – 1.02)	0.94 (0.80 – 1.12)	0.81 (0.67 – 0.99)	0.10	0.87 (0.74 – 1.03)	0.97 (0.81 – 1.15)	0.86 (0.70 – 1.05)	0.22
Psychosis	1.00 (0.91 – 1.10)	0.88 (0.79 – 0.97)	0.76 (0.69 – 0.85)	<0.0001	0.95 (0.86 – 1.06)	0.79 (0.71 – 0.88)	0.65 (0.57 – 0.74)	<0.0001	0.97 (0.88 – 1.07)	0.81 (0.73 – 0.91)	0.69 (0.61 – 0.79)	<0.0001

Note: logistic regression was used for all outcomes and odds ratios (ORs) are presented with their 95% confidence intervals for each ascending day light quartile relative to the low light (Q1) referent. Model 1 is unadjusted. *Model 2 adjusted for age, sex, ethnicity, and photoperiod as a measure of seasonality. †Model 3 is additionally adjusted for employment and physical activity. Bolded coefficients indicate significance at the .05 level.