

1 **Epidemiological surveillance of respiratory pathogens in wastewater in** 2 **Belgium**

3 Annabel Rector¹, Mandy Bloemen¹, Marijn Thijssen¹, Bram Pussig², Kurt Beuselinck³, Marc
4 Van Ranst^{1,3} and Elke Wollants^{1*}

5

6 ¹KU Leuven, Rega Institute, Department of Microbiology, Immunology and Transplantation,
7 Laboratory of Clinical and Epidemiological Virology, 3000 Leuven, Belgium.

8 ²KU Leuven, Department of Public health and Primary Care, Academic Center for General Practice,
9 Leuven, Belgium

10 ³University Hospitals Leuven, Department of Laboratory Medicine, National Reference Center for
11 Respiratory Pathogens, 3000 Leuven, Belgium.

12

13 *Corresponding author: elke.wollants@kuleuven.be ; +32 16 32 13 41; Herestraat 49 box 1040,
14 BE3000 Leuven, Belgium; ORCID: 0000-0002-4057-2236.

15

16 Key words: Wastewater based epidemiology (WBE), acute respiratory tract infection (ARI),
17 respiratory panel, influenza virus, adenovirus, RSV, entero/rhinovirus, EV-D68, coronavirus

18 Word count: 2768

19

20 **ABSTRACT**

21 ***Background***

22 Detailed information on the circulation of respiratory viruses in the community is crucial to gain
23 better understanding of the burden of respiratory infections on society.

24 **Methods**

25 By using an in-house respiratory panel for simultaneous detection of 29 respiratory pathogens (22
26 viruses and 7 bacteria/fungi), we explored the possibility to use wastewater sampling to monitor the
27 circulation of respiratory pathogens at population level.

28 **Results**

29 We were able to detect all respiratory viruses included in the panel (influenza A, respiratory syncytial
30 virus (RSV), human metapneumovirus (HMPV), parainfluenza viruses (PIV) 1-4, adenovirus (Adv),
31 human bocavirus (HBoV), enterovirus/rhinovirus (EV/RV), enterovirus D68 (EV-D68), parechovirus
32 (HPeV), human coronaviruses (HCoV)-NL63, -229E, -OC43, -HKU-1 and -SARS, cytomegalovirus (CMV)
33 and herpes simplex virus (HSV)-1 and -2), except for influenza B and HCoV MERS which were not
34 circulating in Belgium during the two year study period. An upsurge of EV-D68 infections in Europe in
35 September 2021 was clearly reflected in the wastewater samples. For the viruses where
36 epidemiological data on virus circulation in Belgium were available, these matched the wastewater
37 data. The wastewater pretreatment that was used, optimized for viral enrichment, was as such not
38 suited for the surveillance of bacteria and fungi.

39 **Conclusions**

40 Community circulation levels of respiratory viruses were well reflected in wastewater samples,
41 indicating that wastewater-based epidemiology can be a valuable tool in the epidemiology and
42 management of respiratory infections.

43

44 INTRODUCTION

45 Wastewater-based epidemiology (WBE) can be used as a non-invasive tool to monitor the circulation
46 of pathogens within a population. Since various body fluids are discharged into wastewater systems,
47 the method includes but is not limited to detection of fecally excreted viruses and bacteria. Several
48 countries use regular testing of wastewater in the surveillance of poliovirus [1]. Investigation of
49 sewage water can also be applied to monitor the circulation of other enteric viruses such as
50 norovirus, hepatitis A virus and hepatitis E virus [2], [3]. Since the start of the COVID-19 pandemic,
51 wastewater has also been used to follow circulation levels of SARS-CoV-2 and its different variants of
52 concern (VOC) [4], [5]. For this purpose, we have collected samples from a regional sewage
53 treatment plant in the region of Leuven (Belgium) since December 2020. We found wastewater
54 based surveillance to be an important objective indicator of SARS-CoV-2 circulation in the
55 community, which can prove to be highly valuable in times of reducing testing capacity or lower
56 willingness to test [6].

57 Acute respiratory tract infections (ARIs), including pneumonia, constitute a major disease burden
58 worldwide, especially in young children and the elderly. According to the WHO, lower respiratory
59 infections were the world's most deadly communicable disease, ranked as the 4th leading cause of
60 death globally in 2019 (so irrespective of the effect of the COVID-19 pandemic). Respiratory viruses,
61 including influenza virus, EV/RV, RSV, HMPV, PIV, HBoV, Adv and HCoVs, are the leading cause of
62 ARI's in children as well as in older adults [7]–[10]. A large proportion of respiratory infections do not
63 lead to hospitalization or medical care seeking, and therefore go undetected. To gain a better
64 understanding of the burden of respiratory viruses on society, it is important to obtain detailed
65 information on their circulation in the community. This knowledge could allow a better prediction
66 and management of major outbreaks, and could guide physicians in their diagnosis. The current
67 approach is usually based on reporting by a limited number of sentinel physicians and labs, which can
68 lead to bias in the available data.

69 To allow quick diagnosis of respiratory pathogens in immunocompromised and/or critically ill
70 patients with serious lower respiratory infection, a respiratory panel (RP) for simultaneous detection
71 of 22 respiratory viruses (influenza A, influenza B, RSV, HMPV, PIV-1 to -4, Adv, HBoV, RV/EV, EV-D68,
72 HPeV, HCoV-NL63, -229E, -OC43, -HKU-1, -SARS and – MERS, CMV, HSV-1 and -2) and 7
73 bacteria/fungi (*Mycoplasma pneumoniae*, *Coxiella burnetii*, *Chlamydia pneumoniae*, *Chlamydia*
74 *psittaci*, *Streptococcus pneumoniae*, *Legionella pneumophila* and *Pneumocystis jiroveci*) was
75 developed by the Department of Laboratory Medicine of the University Hospitals Leuven. Using this
76 RP, we explored whether wastewater sampling can be used to monitor the circulation of respiratory
77 pathogens at population level.

78

79 **METHODS**

80 ***Sampling***

81 Wastewater samples were collected on average weekly, starting December 6th 2020, from a
82 large regional wastewater treatment plant (WWTP) in Leuven that treats municipal
83 wastewater of approximately 115000 inhabitants. Samples (500 mL) of 24-hour composite
84 influent wastewater were collected through a time-proportional automated sampler, which
85 collects 50 mL of wastewater every 10 minutes in a large container. The samples were stored
86 in a refrigerator at 4°C before transport to the laboratory.

87 ***Viral TNA extraction***

88 Virus concentration and filtration from the wastewater samples was performed as described
89 previously [6]. Total nucleic acids (TNA) were extracted from 500 µL concentrated and
90 filtered sewage samples or DNA/RNA free water (negative controls) using the MagMAX™
91 Viral/Pathogen Nucleic Acid Isolation Kit on Kingfisher Flex-96 (ThermoFisher Scientific,
92 Europe). TNA were eluted in 50 µL elution buffer. Extractions were done in duplicate to
93 obtain sufficient volume of elute for all the tests performed.

94 ***Respiratory panel (RP)***

95 An in-house developed RP consisting of 12 real-time multiplex PCRs was performed on a
96 QuantStudio 7 (Thermo Fisher Scientific, Waltham, MA, USA) in 96 well plates. The end
97 volume of each PCR reaction mix was 20 µL consisting of 5 µL of TNA, 5 µL of master mix
98 (TaqMan Fast Virus Mix) and 10 µL of primer/probe mix. Primer and probe sequences and
99 final concentrations are provided in Supplementary table 1. The temperature profile used
100 was 50°C for 10 minutes; 95°C for 20 seconds; 45 cycles composed of 95°C for 3 seconds and
101 60°C for 30 seconds.

102 Specificity of this lab-developed RP was validated in a clinical context using External Quality

103 Control (EQC) samples, virus cultures and clinical respiratory samples.

104 ***Detection of PMMoV***

105 Human fecal indicator pepper mild mottle virus (PMMoV) was analyzed as an internal

106 extraction control, and to check for extensive differences in human waste input and/or

107 rainwater infiltration, as described previously [6], [11].

108

109 **RESULTS AND DISCUSSION**

110 Full RP results are listed in Supplementary table 2. The human fecal indicator PMMoV was detected
111 in all wastewater samples with only limited variation in Ct values (Ct 22 to 26) (Fig. 1i), indicating
112 there were no extensive differences in fecal input or wastewater dilution throughout the study.
113 Although we did not determine the normalized pathogen concentrations using standard curves, the
114 stable detection of PMMoV indicates that Ct values can be used as a proxy for semi-quantitative
115 assessment of the pathogen load in our wastewater samples.

116 ***Influenzavirus (Fig. 1a)***

117 Whereas influenza B was not detected in any of the collected samples, influenza A was detected in
118 only 1 sample in 2021 (dating from October 4th 2021), and was repeatedly detected between mid
119 Februari and mid May 2022. This is in line with the influenza data collected by the network of
120 sentinel physicians and laboratories, coordinated by the national public health institute of Belgium
121 Sciensano, who reported a very mild influenza season 2020-2021, and a late influenza season 2021-
122 2022 between end of Februari 2022 and end of April 2022. Only influenza A was detected by the
123 sentinel labs during the 2021-2022 season [12].

124 ***RSV and HMPV (Fig. 1b)***

125 RSV was detected in all but one wastewater sample collected between March 8th and June 7th 2021.
126 Sporadic detection occurred during the summer of 2021. As of mid October 2021, RSV was
127 continuously detected until end of December 2021, and discontinuously in January 2022. From the
128 end of February 2022 until the end of July 2022, RSV was present in almost all wastewater samples.
129 This is in accordance with the data from the sentinel labs reported by Sciensano, which indicated a
130 late RSV season in 2019-2020 and a low, almost continuous RSV presence up to week 31 (beginning
131 of August) during 2021-2022 [12]. Since the end of August 2022, RSV was again detectable in
132 wastewater, but no infections have been reported by Sciensano yet in the weekly bulletin of week
133 40, which might be attributable to a delay in data reporting by the sentinel labs. HMPV was not

134 detected in the wastewater samples during the winter season 2020-2021, and was found in one
135 sample during the summer of 2021. From the end of September until the end of December 2021,
136 HMPV was nearly continuously present in the wastewater samples. HMPV is not included in the
137 respiratory virus report of Sciensano, so no data on the circulation of HMPV from sentinel labs in
138 Belgium are available.

139 In the past, we have studied the prevalence and seasonality of respiratory viruses in respiratory
140 samples of ARI patients in the University Hospitals of Leuven, from 2011-2016 [13]. These data
141 showed that the HMPV season had a median onset around mid-December (week 50), peaked in the
142 beginning of April (week 14.5) and lasted until May (week 21). For RSV, the median start of the
143 epidemic season was in the late fall (week 44), with a peak on average at the beginning of December
144 (week 48), and ending at the beginning of Februari (week 5). This does not match the circulation
145 patterns we now see in the wastewater samples. As for other respiratory viruses, the circulation of
146 RSV and HMPV was impacted by the nonpharmacological interventions such as extensive hand
147 hygiene, mask-wearing, and social distancing that were implemented globally in response to the
148 COVID-19 pandemic [14]–[16]. This can account for alterations in levels as well as timing of virus
149 circulation.

150 ***Parainfluenzaviruses (Fig. 1c)***

151 Of the parainfluenzaviruses, PIV-1 was not detected in 2021 and only sporadically in 2022. PIV-2 was
152 detected almost continuously during the fall of 2021, then more sporadically during the winter 2021-
153 2022, and again almost continuously as off mid March 2022 until the end of August. PIV-3 was
154 detected in all wastewater samples from the beginning of Februari 2021 until the beginning of May
155 2021. As of the beginning of August 2021, PIV-3 was again detectable in almost all wastewater
156 samples up to the beginning of September 2022. PIV-4 was detectable in almost all samples between
157 the August 2021 and December 2021, and again as of September 2022, and only sporadically in
158 samples in between. In the data on respiratory pathogens detected by sentinel labs, shared by
159 Sciensano, a peak in PIV circulation is visible between mid Februari 2021 and the beginning of April

160 2021. During the season 2021-2022, the PIV circulation was more spread out, with no clear peak
161 being present [12]. Data on circulation of different PIV subtypes is not available from Sciensano. In
162 our study on seasonality of respiratory viruses in ARI samples, we found that the 4 subtypes of PIV
163 had their seasonal peaks at different moments during the year. Subtype 3 had the tendency to have
164 its seasonal peaks around April/May, subtype 4 around December, while subtypes 1 and 2 did not
165 present with a clear seasonal pattern [13]. According to our wastewater data, this seasonality was
166 more or less maintained in 2021 (with a spring peak of PIV-3 and an (earlier) winter peak of PIV-4),
167 but in 2022 PIV-3 was more continuously present.

168 **Adenovirus and bocavirus (Fig. 1d)**

169 Both Adv and HBoV were detected consistently and in similar concentrations in all wastewater
170 samples during the entire study. This is in accordance with the continuous Adv circulation in 2020-
171 2021 and 2021-2022 reported by Sciensano, and with our previous study on ARI samples, where we
172 found Adv infections to be present throughout the year, with a slight downfall during the Belgian
173 summer season (week 25-38) [12], [13]. The circulation of HBoV is not reported by Sciensano, but its
174 global prevalence has been reported from many countries, including Belgium [17]. To date, 4
175 genotypes of HBoV have been identified, all of which can be detected by the primers and probe
176 located in the highly conserved NP-1 gene, used in our respiratory panel [18]. HBoV1 was mainly
177 identified in respiratory samples from children with ARI [19], whereas HBoV2 is associated with
178 clinical symptoms of patients with gastroenteritis [20]. All HBoV genotypes are known to be present
179 in stool, and HBoV can frequently be detected in sewage samples [21]. Adv infections can also cause
180 gastrointestinal symptoms, even when the primary site of involvement is the respiratory tract, and
181 some serotypes (notably Adv-40 and -41) have an affinity for the gastrointestinal tract, with
182 predominant symptoms of gastroenteritis or diarrhea [22]. The presence of HBoV and Adv in our
183 wastewater samples is therefore likely linked to enteric infections rather than respiratory infections.

184 **Enterovirus/rhinovirus, EV-D68 and parechovirus (Fig. 1e)**

185 Of the 15 classified species of enterovirus, EV A–D and RV A–C infect humans and are a frequent
186 cause of respiratory infections. The fecal-oral route is the most common mode of EV transmission
187 and the primary sites of replication are the gastrointestinal and respiratory tract [23]. Most infections
188 are mild with fever and/or common cold symptoms, but some types, especially those belonging to EV
189 A–D, may cause meningitis, encephalitis, paralysis, neonatal sepsis, myalgia, myocarditis, or
190 exanthema. Of note, EV-D68 has recently been shown to cause large outbreaks of more severe
191 disease with complications such as acute flaccid myelitis [24], [25]. Although EV/RV in general were
192 detected continuously in the wastewater, EV-D68 detection was limited to early September 2021
193 until early December 2021, with the highest concentrations in October 2021. These data provide
194 indications for an outbreak of EV-D68 in the region of Leuven during the fall of 2021, which is in line
195 with the upsurge of EV-D68 infections in Europe as of September 2021, reported by the ECDC [26]. A
196 comparison with the numbers of EV-D68 positive samples tested in the University Hospitals Leuven
197 showed that a total of 20 patients were diagnosed with EV-D68 in 2021, all of which between
198 October and December 2021 (unpublished data). There was a sporadic EV-D68 detection in January
199 2022, and a brief resurgence in April 2022. As of mid September 2022, EV-D68 was again detected in
200 the wastewater samples, which could indicate the start of a new EV-D68 upsurge in the autumn of
201 2022.

202 Like EV/RV, HPeV was present in all samples during the entire study period. HPeV infections without
203 specific symptoms are very common in children, but disease manifestations ranging from
204 gastroenteritis and respiratory infections to neurological disease, particularly in neonates, have been
205 described in association with HPeV [27]. Given their involvement in gastrointestinal infections, HPeVs
206 are frequently detected in sewage samples, where their presence can be an early indicator of clinical
207 cases, including CNS-infections [3], [28]. However, since HPeV were continuously detected in our
208 wastewater samples, indicating a year-round circulation of HPeV in the community, this
209 measurement does not seem useful as an early warning in our setting.

210 **Coronaviruses (Fig. 1f and 1g)**

211 With the SARS-CoV assay in the respiratory panel, targeting a conserved region in the ORF1ab
212 polyprotein gene to allow detection of both SARS-CoV-1 and -2, we did not detect SARS-CoV-2 until
213 the end of September 2021 and only discontinuously thereafter. We also used the 2019-nCoV CDC
214 EUA kit, targeting two regions of the N-gene, for detection of SARS-CoV-2 in all wastewater samples
215 and were able to detect SARS-CoV-2 viral RNA at all measured timepoints, with viral RNA levels
216 largely corresponding to the number of positive cases in Leuven [6]. This indicates that, although
217 suitable in a clinical context, the universal RP SARS-CoV assay is not sensitive enough for
218 environmental surveillance.

219 As expected, the highly pathogenic MERS-CoV was not detected at any timepoint, since MERS-CoV
220 infections have not been reported in Europe for over 4 years now [29], [30].

221 Of the 4 endemic seasonal coronaviruses infecting humans, HCoV-NL-63 was only detected in one
222 sample whereas HCoV-229E and HCoV-OC43 were present in 72% and 84% of all samples
223 respectively. HCoV-HKU-1 was only detected as off the end of November 2021. These endemic
224 coronaviruses are mainly associated with mild and self-limiting upper respiratory tract infections,
225 accounting for about 15–30% of common cold cases, and normally circulate with annual peaks of
226 infections in the winter months [31], [32]. Besides the occurrence in the respiratory tract, all endemic
227 HCoVs can also be detected in stool samples but they do not seem to be a major cause of
228 gastroenteritis, and this detection is most likely explained by the presence of ingested virus particles
229 from the respiratory tract rather than resulting from productive replication in intestinal tissue [31]. In
230 regions with a temperate climate, such as Belgium, the highest activity of endemic HCoVs is observed
231 during the winter months [32]. As is the case for other respiratory viruses, their circulation pattern
232 has been affected by contact restrictions and hygienic measures that were implemented in reaction
233 to the COVID-19 outbreak, especially during the 1st year after the emergence of SARS-CoV-2. This
234 could account for the lower levels of endemic HCoV's detected in the wastewater samples up to the
235 spring of 2021.

236 **Herpesviruses (Fig. 1h)**

237 HSV can cause influenza like illness, but also meningitis and encephalitis. CMV is a ubiquitous
238 pathogen mainly infecting children, with childhood infections usually going unnoticed whereas
239 infections in adults can be more severe, especially in the immunocompromised. There are only
240 limited data available on the presence of human herpesviruses in wastewater, especially for HSV
241 [33]. In our study, HSV-1 and CMV were only sporadically detected, each at 4 scattered timepoints,
242 and HSV-2 was only detected in a single wastewater sample.

243 **Bacteria and fungi**

244 In view of SARS-CoV-2 and polio surveillance, the current method was developed for optimal
245 retrieval of viral material, by including filtration using a 1 µm syringe filter. The drawback of this viral
246 enrichment is that the majority of bacterial and fungal material is lost prior to extraction. Although
247 we were able to detect bacteria/fungi in some of the samples, we consider this method unsuited to
248 confidently detect their presence in wastewater.

249

250 **CONCLUSION**


251 By using the RP on our sample collection spanning almost 2 years, we were able to detect the
252 presence and seasonal variation for most of the tested respiratory viruses. This indicates that
253 wastewater sampling could provide a means to monitor the circulation of respiratory pathogens at
254 population level, and provide an early warning signal in case of local upsurges. By using wastewater
255 samples, the inherent bias posed by data reporting by limited numbers of sentinel labs can be
256 circumvented.

257


258 **FIGURE LEGENDS**

259 **Figure 1. Respiratory viruses in wastewater (Aquafin, Leuven).**

260 Evolution of viruses detected by the respiratory panel from December 2020 to September 2022, as
261 captured by the Ct value of targeted PCR assays performed on wastewater. Plots were generated
262 using R and the ggplot2 package.



263



264

265 REFERENCES

- 266 [1] T. Hovi, L. M. Shulman, H. Van Der Avoort, J. Deshpande, M. Roivainen, and E. M. De
 267 Gourville, "Role of environmental poliovirus surveillance in global polio eradication and
 268 beyond," *Epidemiol. Infect.*, vol. 140, no. 1, pp. 1–13, 2012, doi:
 269 10.1017/S095026881000316X.
- 270 [2] M. Hellmér *et al.*, "Detection of pathogenic viruses in sewage provided early warnings of
 271 hepatitis A virus and norovirus outbreaks," *Appl. Environ. Microbiol.*, vol. 80, no. 21, pp. 6771–
 272 6781, 2014, doi: 10.1128/AEM.01981-14.
- 273 [3] A. Thongprachum *et al.*, "Detection of nineteen enteric viruses in raw sewage in Japan,"
 274 *Infect. Genet. Evol.*, vol. 63, no. May, pp. 17–23, 2018, doi: 10.1016/j.meegid.2018.05.006.
- 275 [4] S. Agrawal *et al.*, "Prevalence and circulation patterns of SARS-CoV-2 variants in European
 276 sewage mirror clinical data of 54 European cities," *Water Res.*, vol. 214, no. February, p.
 277 118162, 2022, doi: 10.1016/j.watres.2022.118162.

- 278 [5] G. Medema, L. Heijnen, G. Elsinga, R. Italiaander, and A. Brouwer, "Presence of SARS-
279 Coronavirus-2 RNA in Sewage and Correlation with Reported COVID-19 Prevalence in the Early
280 Stage of the Epidemic in the Netherlands," *Environ. Sci. Technol. Lett.*, vol. 7, no. 7, pp. 511–
281 516, 2020, doi: 10.1021/acs.estlett.0c00357.
- 282 [6] M. Bloemen *et al.*, "Detection Of SARS-COV-2 Variants Of Concern In Wastewater Of Leuven,
283 Belgium," *medRxiv*, p. 2022.05.12.22274823, May 2022, doi: 10.1101/2022.05.12.22274823.
- 284 [7] G. Comach *et al.*, "Sentinel Surveillance of Influenza-Like Illness in Two Hospitals in Maracay,
285 Venezuela: 2006-2010," *PLoS One*, vol. 7, no. 9, pp. 2006–2010, 2012, doi:
286 10.1371/journal.pone.0044511.
- 287 [8] D. Wang *et al.*, "Viral Etiology of Medically Attended Influenza-Like Illnesses in Children Less
288 Than 5 Years Old in Suzhou, China," *J. Med. Virol.*, vol. 88, pp. 1334–1340, 2016, doi:
289 10.1002/jmv.24480.
- 290 [9] T. Shi *et al.*, "The etiological role of common respiratory viruses in acute respiratory infections
291 in older adults: A systematic review and meta-analysis," *J. Infect. Dis.*, vol. 222, no. Xx Xxxx,
292 pp. S563–S569, 2020, doi: 10.1093/INFDIS/JIY662.
- 293 [10] K. Otani *et al.*, "Incidence of lower respiratory tract infection and associated viruses in a birth
294 cohort in the Philippines," *BMC Infect. Dis.*, vol. 22, no. 1, pp. 1–11, 2022, doi:
295 10.1186/s12879-022-07289-3.
- 296 [11] M. Kitajima, H. P. Sassi, and J. R. Torrey, "Pepper mild mottle virus as a water quality
297 indicator," vol. 1, p. 19, 2018, doi: 10.1038/s41545-018-0019-5.
- 298 [12] Bossuyt Nathalie *et al.*, "Weekly bulletin respiratory infections." <https://epidemiologie.wiv-isp.be/ID/diseases/Pages/Influenza.aspx>
299
- 300 [13] K. Ramaekers *et al.*, "Prevalence and seasonality of six respiratory viruses during five
301 consecutive epidemic seasons in Belgium," *J. Clin. Virol.*, vol. 94, no. March, pp. 72–78, 2017,
302 doi: 10.1016/j.jcv.2017.07.011.
- 303 [14] M. Stein *et al.*, "Human metapneumovirus prevalence during 2019-2021 in Israel is influenced
304 by the COVID-19 pandemic," *Int. J. Infect. Dis.*, vol. 120, pp. 205–209, Jul. 2022, doi:
305 10.1016/j.ijid.2022.04.037.
- 306 [15] H. J. Shi *et al.*, "Effects of Non-Pharmacological Interventions on Respiratory Viruses Other
307 Than SARS-CoV-2: Analysis of Laboratory Surveillance and Literature Review From 2018 to
308 2021," *J. Korean Med. Sci.*, vol. 37, no. 21, 2022, doi: 10.3346/jkms.2022.37.e172.
- 309 [16] Y. Li, X. Wang, B. Cong, S. Deng, D. R. Feikin, and H. Nair, "Understanding the Potential Drivers
310 for Respiratory Syncytial Virus Rebound during the Coronavirus Disease 2019 Pandemic," *J.*
311 *Infect. Dis.*, vol. 225, no. 6, pp. 957–964, Mar. 2022, doi: 10.1093/infdis/jiab606.
- 312 [17] D. Polo, A. Lema, E. Gándara, and J. L. Romalde, "Prevalence of human bocavirus infections in
313 Europe. A systematic review and meta-analysis," *Transbound. Emerg. Dis.*, no. July, pp. 1–11,
314 2021, doi: 10.1111/tbed.14233.
- 315 [18] F. Neske *et al.*, "Real-time PCR for diagnosis of human bocavirus infections and phylogenetic
316 analysis," *J. Clin. Microbiol.*, vol. 45, no. 7, pp. 2116–2122, 2007, doi: 10.1128/JCM.00027-07.
- 317 [19] Q. B. Lu *et al.*, "Epidemic and molecular evolution of human bocavirus in hospitalized children
318 with acute respiratory tract infection," *Eur. J. Clin. Microbiol. Infect. Dis.*, vol. 34, no. 1, pp. 75–
319 81, 2015, doi: 10.1007/s10096-014-2215-7.
- 320 [20] R. De *et al.*, "Risk of acute gastroenteritis associated with human bocavirus infection in

- 321 children: A systematic review and meta-analysis,” *PLoS One*, vol. 12, no. 9, pp. 1–11, 2017,
322 doi: 10.1371/journal.pone.0184833.
- 323 [21] T. Booranathawornsom, K. Pombubpa, M. Tipayamongkholgul, and L. Kittigul, “Molecular
324 characterization of human bocavirus in recycled water and sewage sludge in Thailand,” *Infect.*
325 *Genet. Evol.*, vol. 100, no. March, p. 105276, 2022, doi: 10.1016/j.meegid.2022.105276.
- 326 [22] J. P. Lynch Iii, A. E. Kajon, and S. K. Singh, “Adenovirus: Epidemiology, Global Spread of Novel
327 Serotypes, and Advances in Treatment and Prevention,” *Semin. Respir. Crit. Care Med.*, vol.
328 37, no. 4, pp. 586–602, 2016, doi: 10.1055/s-0036-1584923.
- 329 [23] S. Cordey *et al.*, “Comparative analysis of viral shedding in pediatric and adult subjects with
330 central nervous system-associated enterovirus infections from 2013 to 2015 in Switzerland,”
331 *J. Clin. Virol.*, vol. 89, pp. 22–29, 2017, doi: 10.1016/j.jcv.2017.01.008.
- 332 [24] K. Messacar, M. J. Abzug, and S. R. Dominguez, “2014 outbreak of enterovirus D68 in North
333 America,” *J. Med. Virol.*, vol. 88, no. 5, pp. 739–745, May 2016, doi: 10.1002/jmv.24410.
- 334 [25] S. B. Larsson, D. Vracar, M. Karlsson, J. Ringlander, and H. Norder, “Epidemiology and clinical
335 manifestations of different enterovirus and rhinovirus types show that EV-D68 may still have
336 an impact on severity of respiratory infections,” *J. Med. Virol.*, vol. 94, no. 8, pp. 3829–3839,
337 Aug. 2022, doi: 10.1002/jmv.27767.
- 338 [26] K. S. M. Benschop *et al.*, “Re-emergence of enterovirus d68 in europe after easing the covid-
339 19 lockdown, september 2021,” *Eurosurveillance*, vol. 26, no. 45, 2021, doi: 10.2807/1560-
340 7917.ES.2021.26.45.2100998.
- 341 [27] H. Harvala and P. Simmonds, “Human parechoviruses: Biology, epidemiology and clinical
342 significance,” *Journal of Clinical Virology*, vol. 45, no. 1. pp. 1–9, May 2009. doi:
343 10.1016/j.jcv.2009.03.009.
- 344 [28] H. Harvala *et al.*, “Comparison of diagnostic clinical samples and environmental sampling for
345 enterovirus and parechovirus surveillance in Scotland, 2010 to 2012,” *Eurosurveillance*, vol.
346 19, no. 15, pp. 1–9, 2014, doi: 10.2807/1560-7917.ES2014.19.15.20772.
- 347 [29] ECDC, “Communicable disease threats report, Week 27, 3-9 July 2022”.
- 348 [30] ECDC, “Rapid risk assessment: Severe respiratory disease associated with Middle East
349 respiratory syndrome coronavirus (MERS-CoV), 22nd update,” 2018.
350 [https://www.ecdc.europa.eu/en/publications-data/rapid-risk-assessment-severe-respiratory-](https://www.ecdc.europa.eu/en/publications-data/rapid-risk-assessment-severe-respiratory-disease-associated-middle-east-11)
351 [disease-associated-middle-east-11](https://www.ecdc.europa.eu/en/publications-data/rapid-risk-assessment-severe-respiratory-disease-associated-middle-east-11)
- 352 [31] V. M. Corman, D. Muth, D. Niemeyer, and C. Drosten, “Hosts and Sources of Endemic Human
353 Coronaviruses,” *Adv. Virus Res.*, vol. 100, pp. 163–188, Jan. 2018, doi:
354 10.1016/BS.AIVIR.2018.01.001.
- 355 [32] Y. Li, X. Wang, and H. Nair, “Global seasonality of human seasonal coronaviruses: A clue for
356 postpandemic circulating season of severe acute respiratory syndrome coronavirus 2?,” *J.*
357 *Infect. Dis.*, vol. 222, no. 7, pp. 1090–1097, 2020, doi: 10.1093/infdis/jjaa436.
- 358 [33] C. McCall, H. Wu, B. Miyani, and I. Xagorarakis, “Identification of multiple potential viral
359 diseases in a large urban center using wastewater surveillance,” *Water Res.*, vol. 184, p.
360 116160, 2020, doi: 10.1016/j.watres.2020.116160.
- 361 [34] A. Clancy, B. Crowley, H. Niesters, and C. Herra, “The development of a qualitative real-time
362 RT-PCR assay for the detection of hepatitis C virus,” *Eur. J. Clin. Microbiol. Infect. Dis.*, vol. 27,
363 no. 12, pp. 1177–1182, Dec. 2008, doi: 10.1007/S10096-008-0556-9.

- 364 [35] J. Kuypers, N. Wright, and R. Morrow, "Evaluation of quantitative and type-specific real-time
365 RT-PCR assays for detection of respiratory syncytial virus in respiratory specimens from
366 children," *J. Clin. Virol.*, vol. 31, pp. 123–129, 2004, doi: 10.1016/j.jcv.2004.03.018.
- 367 [36] W. A. Nix *et al.*, "Detection of all known parechoviruses by real-time PCR," *J. Clin. Microbiol.*,
368 vol. 46, no. 8, pp. 2519–2524, Aug. 2008, doi: 10.1128/JCM.00277-08.
- 369 [37] H. G. M. Niesters, "Molecular and diagnostic clinical virology in real time," *Clin. Microbiol.*
370 *Infect.*, vol. 10, no. 1, pp. 5–11, 2004, doi: 10.1111/j.1469-0691.2004.00699.x.
- 371 [38] A. Tiveljung-Lindell *et al.*, "Development and implementation of a molecular diagnostic
372 platform for daily rapid detection of 15 respiratory viruses," *J. Med. Virol.*, vol. 81, no. 1, pp.
373 167–175, Jan. 2009, doi: 10.1002/JMV.21368.
- 374 [39] L. J. R. Van Elden *et al.*, "Frequent Detection of Human Coronaviruses in Clinical Specimens
375 from Patients with Respiratory Tract Infection by Use of a Novel Real-Time Reverse-
376 Transcriptase Polymerase Chain Reaction," *J. Infect. Dis.*, vol. 189, pp. 652–659, 2004.
- 377 [40] A. Piralla, A. Girello, M. Premoli, and F. Baldanti, "A new real-time reverse transcription-PCR
378 assay for detection of human enterovirus 68 in respiratory samples," *J. Clin. Microbiol.*, vol.
379 53, no. 5, pp. 1725–1726, May 2015, doi: 10.1128/JCM.03691-14/FORMAT/EPUB.
- 380 [41] M. Panning *et al.*, "High throughput detection of *Coxiella burnetii* by real-time PCR with
381 internal control system and automated DNA preparation," *BMC Microbiol.*, vol. 19, no. 8,
382 2008, doi: 10.1186/1471-2180-8-77.
- 383 [42] V. M. Corman *et al.*, "Detection of a novel human coronavirus by real-time reverse-
384 transcription polymerase chain reaction," *Eurosurveillance*, vol. 17, no. 39, Sep. 2012, doi:
385 10.2807/ese.17.39.20285-en.
- 386 [43] J. Hardick *et al.*, "Real-Time PCR for *Chlamydia pneumoniae* Utilizing the Roche Lightcycler
387 and a 16S rRNA Gene Target," *J. Mol. Diagnostics*, vol. 6, no. 2, 2004.
- 388 [44] A. Ménard, M. Clerc, A. Subtil, F. Mégraud, C. Bébéar, and B. De Barbeyrac, "Development of a
389 real-time PCR for the detection of *Chlamydia psittaci*," *J. Med. Microbiol.*, vol. 55, no. Pt 4, pp.
390 471–473, 2006, doi: 10.1099/JMM.0.46335-0.
- 391 [45] M. Da *et al.*, "Evaluation and Improvement of Real-Time PCR Assays Targeting *lytA*, *ply*, and
392 *psaA* Genes for Detection of Pneumococcal DNA," *J. Clin. Microbiol.*, vol. 45, no. 8, pp. 2460–
393 2466, 2007, doi: 10.1128/JCM.02498-06.

394

395 **Supplementary table 1:** Primers and probes used in the 12 multiplex PCR reactions of the respiratory panel.

Multiplex PCR	Pathogen	Primer/probe	Primer/probe sequence (5' → 3')	Concentration (nM)	Target gene	Amplicon size (bp)
RP1	Influenza A	Forward primer	TTCATGGAATGGCTAAAGACAAG	300	Matrix protein gene	116
		Reverse primer	CAAAGCGTCTACGCTGCAGT	300		
		Probe	FAM-TTCACGCTCACCGTGC-MGB	200		
	Influenza B	Forward primer	AATACGGTGGATTAACAAAAGCA	300	Hemagglutinin gene	170
		Reverse primer	ACCAGCAATAGCTCCGAAGAA	300		
		Probe	NED-TATATTTGGTTCCATTGGC-MGB	200		
	PDV (RNA IC)	Forward primer	GGTGGTGCCTTTTACAAGAAC [34]	250	Hemagglutinin gene	83
		Reverse primer	ATCTTCTTCTCAACCTCGTCC [34]	250		
		Probe	VIC-ATGCAAGGGCCAATT-MGB [34]	100		
RP2	RSV A + RSV B	Forward primer	AATACGCCAAATCTAACCACTTACA [35]	300	RNA polymerase gene	94
		Reverse primer	GCCAAGGAAGCATGCAATAAA [35]	300		
		Probe A	FAM-TGATGTGCTATTGTGCACTA-MGB [35]	100		
		Probe B	FAM-CACTATTCCTACTAAAGATGTC-MGB [35]	100		
	HMPV	Forward primer	GCGTYAGCTTCAGTCAATTCAA	500	Fusion protein gene	75
		Reverse primer	TGTTATYCCWGCATTGTCTGA	500		
		Probe	NED-CTAAATGTTGTGCGGCAAT-MGB	200		
RP3	PIV-1	Forward primer	CCTTCATTATCAATTGGTGATGCA	500	Hemagglutinin-neuraminidase gene	180
		Reverse primer	CCTGTTGTCGTTGATGTCATAGGT	500		
		Probe	FAM-TCAAACCTAATCACTCAAGGAT-MGB	100		
	PIV-2	Forward primer	CATTTACCTAAGTGATGGAATCAATCG	500	Hemagglutinin-neuraminidase gene	141
		Reverse primer	GCAAGTCTCAGTTCAGCTAGATCAGT	500		
		Probe	NED-AAAGCTGTTCACTACTG-MGB	150		
RP4	PIV-3	Forward primer	CAAAGTTGATGAAAGATCAGATTATGC	500	Hemagglutinin-neuraminidase gene	169
		Reverse primer	GTAGTATATCCCTGGTCCAACAGATG	500		
		Probe	FAM-CAATCTCRACAACAAGATT-MGB	100		
	PIV-4	Forward primer	GGAGCAAAAGAYTCATACACAATAACTACT	500	Hemagglutinin-neuraminidase gene	116
		Reverse primer	CTAGGAA TTAARATTCACAATCTTCAGAA	500		
		Probe	NED-AAATGCACTCTGTATAAGTC-MGB	150		
RP5	Adenovirus	Forward primer	TYGARGTGGAYCCCATGGAYGAG	800	Hexon gene	112
		Reverse primer	CGCAGGTAGACBGCYTCRATGA	800		

		Probe 1	FAM-ACGTCGAAGACTTC-MGB	100	Polyprotein gene	150-154
		Probe 2	FAM-ACGTCGAAAACCTTC-MGB	100		
		Probe 3	FAM-CACGTCAAAGACTTC-MGB	100		
	Rhino- enterovirus	Forward primer 1	CAWGGTGYGAAGAGYCTATTGAGCT	500		
		Forward primer 2	GTGTGAA GASCCSMGTGYGCT	500		
		Reverse primer	GAAACACG GACACCCAAAGTAGT	500		
		Probe	VIC-TCCGGCCCTGAATGYGGCTAA-TAMRA	200		
RP6	CMV	Forward primer	CGTAACGTGGACCTGACGTTT	250	Major capsid protein gene	148
		Reverse primer	CACGGTCCCGTTTAGCA	250		
		Probe	FAM-TATCTGCCCGAGATCGCGTTACA-TAMRA	200		
	Parechovirus	Forward primer	CASWWGCCTCTGGGCCAAAAAG [36]	500	Polyprotein gene	189
		Reverse primer	GGCCCCWGRTCAGATCCAYAGT [36]	500		
		Probe	Cy5-CCTRYGGGTA CCTYCWGGGCATCCTTC-BHQ3 [36]	200		
	PhHV-1 (DNA IC)	Forward primer	GGGCGAATCACAGATTGAATC [37]	100	Glycoprotein B gene	89
		Reverse primer	GCGGTTCCAAACGTACCAA [37]	100		
		Probe	VIC-TCCGCCACCATCTG-MGB	100		
RP7	HCoV NL63	Forward primer	CAGGGCTGACAAGCCTTCTCA [38]	500	Nucleocapsid protein gene	144
		Reverse primer	GCATCAACACCAATTCTGAACAAGA [38]	500		
		Probe	FAM-CGTTGGAAGCGTGTTCCTACCAGAGAGG-BHQ-1 [38]	200		
	HCoV 229E	Forward primer	TGGAAGTGCAAGGTGTTGTGGC [38]	500	Nucleocapsid protein gene	99
		Reverse primer	TGACTATCAAACAGCATA GCAGCTG [38]	500		
		Probe	Cy5-CCACAATTTGCTGAGCTTGCCGTC-BHQ-3 [38]	200		
	HCoV OC43	Forward primer	CGATGAGGCTATTCGACTAGGT [39]	500	Nucleocapsid protein gene	75
		Reverse primer	CCTTCCTGAGCCTTCAATATAGTAACC [39]	500		
		Probe	VIC-TCCGCCTGGCACGGTACTCCCT-TAMRA [39]	200		
RP8	HSV-1 + HSV-2	Forward primer	AACCTGGGRTTCTGATGCA	250	Nonfunctional glycoprotein D gene (US6)	84
		Reverse primer	CTCCGTCCAGTCGTTTATCTTAC	250		
		Probe HSV-1	FAM-TTTGAGACCGCCGGCACGTAC-BHQ-1	100		
		Probe HSV-2	Cy5-CCTTCGAGACCGGGTACGTA-BHQ-3	100		
	Enterovirus D68	Forward primer	TGGCGGCTACTCATGG [40]	500	Polyprotein gene	64
		Reverse primer	AATAGACTCTTACACCTTGTTTATGT [40]	500		
		Probe	NED-AAAA CCATGAGACGCT-MGB [40]	200		
	HCoV SARS	Forward primer	CWGGCATAACWAAGGACATGACCTA	500	ORF1ab polyprotein gene	147
		Reverse primer	CKACATCRAAGCCAATCCA	500		
		Probe	VIC-TTTATCACCCGCGAAGAA-MGB	200		

RP9	<i>Mycoplasma pneumoniae</i>	Forward primer	AGGCTTCAAGTGACAAAAGTGAC	250	Adhesin P1 gene	76
		Reverse primer	GATTGTCTGCTGGYCCAT	250		
		Probe	FAM-ACCACACCAAGTTCA-MGB	200		
	HCoV HKU1	Forward primer	CACTTCTATCCCTCCGATGTTTC [38]	500	Nucleocapsid protein gene	129
		Reverse primer	TTAGAAGCAGACCTTCTGAGCC [38]	500		
		Probe	Cy5-CGCCTGGTACGATTTTGCCTCAAGGCT-BHQ-3 [38]	200		
RP10	<i>Pneumocystis jirovecii</i>	Forward primer	GCACTGAATACTCGAGGGAGTATG	250	Large subunit ribosomal RNA gene	145
		Reverse primer	TTGGGAGCTTTAATTACTGTTCTGG	250		
		Probe	FAM-TGTTTCCCTTTCGACTATC-MGB	200		
	<i>Coxiella burnetii</i>	Forward primer	CGATAGCCCGATAAGCATCAAC [41]	250	IS1111 transposase gene	88
		Reverse primer	TGCATTGATATCCGGCATC [41]	250		
		Probe	NED-TCATCAAGGCACCAATG-MGB [41]	200		
	HCoV MERS	Forward primer	GCAACGCGCGATTGAGTT [42]	500	Envelope gene (upE)	92
		Reverse primer	GCCTTACACGGACCCATA [42]	500		
		Probe	Cy5-CTCTTACATAATCGCCCCGAGCTCG-BHQ-3 [42]	200		
RP11	<i>Chlamydia pneumoniae</i>	Forward primer	CATCCGTGTCGGAGCTAACGT [43]	250	16S rRNA gene	184
		Reverse primer	TGCGGAAAGCTGTATTCTACAGTT [43]	250		
		Probe	FAM-ATGCCCGCTGAGGAGTACACTCG-BHQ1 [43]	200		
	<i>Chlamydia psittaci</i>	Forward primer	GCCATCATGCTTGTTCGTTT [44]	250	Inclusion membrane protein A gene	74
		Reverse primer	CGGCGTGCCACTTGAGA [44]	250		
		Probe	VIC-TTGTATTATGGTGATTCAGGA-MGB [44]	200		
	<i>Streptococcus pneumoniae</i>	Forward primer	ACGCAATCTAGCAGATGAAGCA [45]	250	Autolysin gene (lytA)	75
		Reverse primer	TCGTGCGTTTTAATCCAGCT [45]	250		
		Probe	Cy5-TGCCGAAAACGCTTGATACAGGGAG-BHQ3 [45]	200		
RP12	<i>Legionella pneumophila</i>	Forward primer	TCCGGAAGCAATGGCTAAAG	250	Macrophage infectivity potentiator gene	69
		Reverse primer	TGCTGTTCCGTTAAAGCCAAT	250		
		Probe	FAM-CATGCAAGACGCTATGAGTGCGCT-TAMRA	200		
	Bocavirus	Forward primer	GCACGCCACGTGACGAA [18]	250	Nonstructural protein 1 gene	76
		Reverse primer	TGGACTCCCTTTCTTTGTAGGA [18]	250		
		Probe	Cy5-TGAGCTCAGGGAATATGAAAGACAAGCATCG-BHQ-3 [18]	200		

396 Base ambiguity codes: R: A or G; Y: C or T; W: A or T; S: C or G; K: G or T

397 PDV: phocine distemper virus; RSV: respiratory syncytial virus; IC: internal control; HMPV: human metapneumovirus; PIV: parainfluenzavirus; CMV: cytomegalovirus; PhHV: phocine

398 herpesvirus; HCoV: human coronavirus; HSV: herpes simplex virus

399

400 **Supplementary table 2:** Ct values of pathogens detected by the respiratory panel in wastewater samples
 401 (Aquafin, Leuven) at different timepoints.

	Influenza A	Influenza B	RSV	HMPV	PIV-1	PIV-2	PIV-3	PIV-4	Adv	HBov	EV/RV	EV-D68	HPeV	SARS-CoV	MERS-CoV	HCoV-NL63	HCoV-229E	HCoV-OC43	HCoV-HKU-1	HSV-1	HSV-2	Cytomegalovirus	PMMoV
6/12/20									27	30	29		32										23
5/01/21									30	31	31		35					36					24
11/01/21									28	32	29		34										23
18/01/21									30	33	31		35					36					24
1/02/21									33	33	31		36										25
8/02/21							36		30	31	29		35										24
22/02/21							34		26	31	30		34			37		36					23
24/02/21							36		28	30	30		34				36	35					25
8/03/21			39				32		30	29	30		34				38	36					23
21/03/21			39				33		27	31	30		33					35					23
31/03/21			36				34	44	27	32	30		34				36	33					23
12/04/21			39				37		30	33	30		34				35	34					25
3/05/21			37				37		29	31	31		36					34					24
10/05/21									31	31	29		33				38	37					26
19/05/21			37						29	32	28		35					34					24
26/05/21			39						30	34	29		35					32					24
31/05/21			39						29	31	29		33				38	34					23
7/06/21			38					38	31	31	29		34				37	35					24
5/07/21							38		31	32	32		35										24
12/07/21				40				37	30	31	29		34					34					23
19/07/21									32	35	34		38					37					24
26/07/21									28	36	32		37					35					25
2/08/21							37	38	25	35	31		36					37					25
10/08/21			40				44	39	27	33	31		35				37	35					24
16/08/21							37	37	23	31	29		34					36					23
23/08/21									25	32	30		37				37						24
30/08/21			39				36	38	26	34	29		33				35	35					23
8/09/21			36				36	37	22	32	27	44	32					34					23
15/09/21								38	25	33	28	37	33				37	34					25
20/09/21							37	37	22	31	26	37	32				37	34					23
27/09/21				40			36	37	23	32	26	36	31	36				32					23
4/10/21	40					39	37	36	23	31	28	36	33	36			34	32					24
13/10/21			36	40			36	35	22	29	27	35	33				35	30				35	23
20/10/21			36			39	36	36	23	31	26	35	33	37			36	32		35			24
25/10/21			36	39		38	35	35	22	29	26	34	32				34	31					24
8/11/21			36	39		38	34	35	22	30	27	35	32	35			33	32					23
15/11/21			36	37			36	35	23	30	27	36	33	35			34	32					24
22/11/21			35			36	34	37	23	30	27	37	33	34			33	31					23
29/11/21			34	39			34	39	20	28	27	37	33				32	31	36				24
6/12/21			34			38	34	38	21	29	27	38	33				34	33	34				22
13/12/21			37	40			36		22	30	27		33	34			31	33	39				22
20/12/21			38	39			37		22	30	27		34	36			33	32	34				23
27/12/21			36				36	38	21	29	27		33	36			32	33	34				23
3/01/22						41			23	31	F	39	35	36			33	32	34				24
10/01/22							37		22	31	28		36	35			33	34	33				24

