

1
2
3
4 It's not just about pads! Adolescent reproductive health in Kenya: *A qualitative secondary analysis*.

5
6
7
8 Sylvia Ayieko^{1*}, Angela Nguku^{2,3}, Nancy Kidula⁴

9
10 ¹Department of Health Promotion and Behavioral Sciences, The University of Texas Health Science Center at
11 Houston School of Public Health, Houston, Texas, United States of America

12
13 ²White Ribbon Alliance -Kenya, Nairobi, Kenya

14 ³White Ribbon Alliance-Global, Washington DC, United States of America

15 ⁴Department of Reproductive Health and Research (RHR), World Health Organization, Genève, Switzerland
16

17
18
19 *Corresponding Author:

20 Email: sylvia.a.ayieko@uth.tmc.edu or saachieng127@gmail.com (SA)
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 **Abstract**

42 Many adolescents face barriers to accessing reproductive health care even though quality reproductive health
43 care is a fundamental human right. This qualitative analysis describes quality reproductive health requests
44 among adolescent high school girls in Kenya. We conducted a secondary analysis of qualitative data from a
45 sub-sample of adolescent girls in Kenya who participated in the *What Women Want* global campaign. The
46 campaign utilized one open-ended survey question. We also analyzed interview data from key informants
47 involved in the survey. We used pre-existing codes and current literature to design the coding framework and
48 thematic analysis to describe emerging themes. Atlas. ti 8 was used to organize and analyze codes. Over 4,500
49 high school girls, ages 12 and 19 years, were included in the analysis, with 61.6% from all-girls boarding
50 schools and 13.8% from mixed-day schools. Data from nine key informants complemented findings from the
51 survey. Emerging themes included: 1) The need for improved menstrual health and hygiene: Sanitary towels
52 and cleaner toilets; 2) Prevention of adolescent pregnancy: Access to contraception; 3) Respect and dignity:
53 Participants want privacy and confidentiality; and 4) The need to address social determinants of health:
54 Economic stability and a safe physical environment. This study indicated that adolescent high school girls have
55 varied requests for reproductive health care and services. While menstrual health and hygiene are key issues,
56 reproductive needs are beyond just sanitary products. The results suggest a need for targeted reproductive
57 health interventions using a multi-sectoral approach.

58
59
60
61
62
63
64
65
66
67

68 **Introduction**

69 Access to health care, including sexual and reproductive health care, is a fundamental human right.^{1,2}

70 However, many adolescents face barriers to accessing sexual and reproductive health services, which put them
71 at risk for teenage pregnancy, unsafe abortions, reproductive tract cancers, and sexually transmitted infections,
72 including Human Immunodeficiency Virus, Acquired Immunodeficiency Syndrome (HIV/AIDS).^{3,4} In Sub-
73 Saharan Africa, for example, due to limited and inequitable sexual and reproductive health care services,
74 adolescent girls and young women (15 to 24 years) bear a disproportionate burden (25%) of HIV infections,
75 with new HIV infections occurring at a higher rate among 15–19-year-old females compared to their male
76 counterparts.⁵

77 Pregnancy-related complications are among the leading causes of death among adolescent girls in Sub-
78 Saharan Africa⁶ due to poor health care or non-utilization of health care services. Adolescent girls underutilize
79 reproductive and maternal health services due to stigma, lack of information, weakened health systems, and
80 poor implementation of health policies.^{7,8} In addition, poverty, low decision-making power, lack of negotiation
81 skills, gender-based violence, and alcohol and drug abuse contribute to adolescents' adverse reproductive health
82 outcomes.^{8,9} In low-to-middle income countries, only about 10% of adolescent girls sought care from health
83 facilities and received contraception counseling.¹

84 In 2015, all 193 United Nations member states adopted the sustainable development goals (SDGs).
85 Within the health-related SDG 3, targets 3·7 and 3·8 seek to ensure universal access to sexual and reproductive
86 health care services, including family planning, information, and education, while striving to achieve universal
87 health coverage, including access to quality essential health care services by 2030 respectively.^{10,11}

88 *Reproductive health care for adolescents in Kenya*

89 Kenya, a lower-middle-income country in Eastern Africa, is a signatory to the SDGs and is committed to
90 ensuring Universal Health Coverage. While the country has expanded access to adolescent sexual and
91 reproductive health and rights (SRHR), impactful implementation of the SRHR interventions recommended by
92 the *Global Accelerated Action for the Health of Adolescents (AA-HA!): Guidance to Support Country*
93 *Implementation*¹² has been limited. In 2017, about 24% of adolescent women aged 15 - 19 years in Kenya had

94 an unmet need for family planning, resulting in 63% of unintended pregnancies in this population.¹³ Adolescent
95 girls in Kenya are also more susceptible to HIV due to non-consensual, unprotected sexual intercourse and
96 sexual violence.¹⁴ About 18% of teenage girls in Kenya between the ages of 15-19 years are either pregnant or
97 mothers.¹⁴ Research suggests that factors associated with adolescent pregnancy in Kenya include lack of
98 knowledge, unmet need for contraception, female genital mutilation (FGM) practices, and poverty.¹⁵ Adolescent
99 girls in Kenya who engage in transactional sex due to low socioeconomic conditions also face challenges
100 navigating through conflicting societal constructs that condemn adolescent contraceptive use and stigmatize
101 pregnant adolescents.¹⁶

102 To reduce HIV infections and address unintended pregnancy in Kenya, the Kenyan government
103 distributes free male condoms, targeting youth who may not afford or have access to condoms.¹⁷ Moreover, in
104 collaboration with policymakers and civil society organizations, the Ministry of Health in Kenya has
105 encouraged implementing the National Adolescent Sexual and Reproductive Health policy at all levels of
106 care.¹⁸ The growing body of evidence suggests that the delivery of quality adolescent reproductive health care is
107 linked to more positive health-seeking behaviors, increased adherence to birth control methods, and improved
108 overall health outcomes among young people.¹⁹ Other studies allude to inadequate coordination and policy
109 implementation that fails to incorporate pregnancy, menstruation, and sanitation into reproductive health.²⁰ In
110 addition, interventions on menstrual health hygiene programs in schools have not been well adopted because
111 adolescents and young adults are not recognized as sources of information^{21,22}. Yet, they are the key
112 stakeholders in their health.

113 Access to quality reproductive healthcare for adolescent girls in Kenya is a pressing public health issue
114 warranting further investigation. To address this gap, this study utilizes qualitative data from the *What Women*
115 *Want* survey, and interviews, conducted by the White Ribbon Alliance to explore what adolescent girls enrolled
116 in different schools across Kenya demand as quality reproductive health care. It is expected that this secondary
117 analysis of qualitative surveys and interviews will contribute to the current literature by discussing how
118 adolescent girls conceptualize quality reproductive health and how to improve care.

120 **Materials and Methods**

121 *Parent Study*

122 The White Ribbon Alliance, a non-profit organization, conducted the global *What Women Want*
123 campaign between April 2018 and March 2019 to understand what women and girls want for maternal and
124 reproductive health care.^{23,24} The organization utilized a qualitative survey with one open-ended question: *My*
125 *one request for quality reproductive and maternal health care services is _____?*

126 Although the campaign initially administered the survey online, most countries, including Kenya, later
127 distributed paper surveys and had different consenting procedures. In Kenya, data collection was via
128 convenience sampling "citizen journalists," community members, and volunteers associated with the White
129 Ribbon Alliance in Kenya. About 120,000 women and girls completed the survey, with 16% being adolescents
130 aged 15-19.²⁴

131 *Ethics statement*

132 De-identified data from the *What Women Want* results were obtained from the *What Women Want*
133 dashboard.²⁴ The current study does not involve interactions with human subjects but uses existing data. Given
134 the nature of the study, the authors did not seek Institutional Review Board approval since the data was publicly
135 available. The campaign outlined the consent processes during data collection for each country.²⁴ In Kenya,
136 verbal consent was obtained from girls and women who wished to remain anonymous and just provided their
137 response and, in some cases, their age. In addition, approval and permission was obtained from schools before
138 any data collection was done. Both verbal and written consent was obtained for adolescents aged 18 years or
139 older who had their pictures taken.²⁴

140 *Sample data source*

141 We received authorization from the White Ribbon Alliance to conduct a secondary analysis of the
142 qualitative survey from a sub-sample of adolescent girls in Kenya. The current study restricted analysis to data
143 from teenage girls enrolled in sample high schools in eight counties in Kenya. Although some survey responses
144 had missing information about the age or the type of school (due to the sensitive nature of the topic), the
145 qualitative survey responses were sufficient for this current study. The study also included interview reports

146 from nine key informants (teachers, youth advocates, and mobilizers) involved in the *What Women Want*
147 campaign in Kenya.

148 *Type of schools*

149 In Kenya, high schools are usually either day schools or boarding schools. Although few high
150 (secondary) schools have some students enrolled as day scholars and others as boarders, this is not common.
151 Moreover, boarding schools in Kenya are predominantly single-sex institutions. Thus, the survey responses
152 from boarding schools reflected perspectives from mostly all-girls schools. Most day schools, by contrast, are
153 co-ed/ mixed local schools, with many students making daily commutes from their homes.

154 *Data analysis*

155 We conducted secondary analysis by re-examining survey responses from adolescent girls and analyzing
156 transcripts from key informant interviews involved in the *What Women Want* campaign in Kenya. We
157 conducted thematic content analysis based on emerging codes and existing quality-of-care frameworks²⁵ to
158 elucidate the phenomenon of quality reproductive care among adolescents. The initial coding system utilized
159 categories and codes from the parent *What Women Want* study.²⁴ SA conducted the initial coding by reviewing
160 previously assigned codes and identifying mismatched responses. Then, three investigators (SA, AN, and NK)
161 reviewed merging and reassigning codes. All the authors discussed the inconsistencies in the coding
162 assignments and categories until the team achieved consensus. The iterative process resulted in new codes and
163 categories for the coding framework. Thematic analysis was conducted using Atlas.ti-8 computer software.²⁶

164 **Results**

165 The secondary analysis included 4,570 survey responses from adolescent girls between 12 and 19 years
166 in eight counties. Although 20% did not report their ages, their information was included for analysis. Most of
167 the adolescent girls were enrolled in all-girls boarding schools (61·6%) compared to those who attended mixed-
168 day schools (13·8%). **Table one** provides a summary of participant characteristics.

169

170

171

Table 1: Demographic characteristics of study participants

Characteristics	Number (n)	Percentage (%)
Type of school	n	%
All Girls' Boarding	2814	61.6%
Mixed-Day	633	13.9%
Other	1123	24.6%
Total	4570	100.0%
County	n	%
Busia	437	9.6%
Uasin Gishu	1053	23.0%
Elgeyo Marakwet	489	10.7%
Isiolo	405	8.9%
Kisumu	280	6.1%
Nairobi	499	10.9%
Narok	980	21.4%
Vihiga	427	9.3%
Total	4570	100.0%
Age (years)	n	%
12	26	0.6%
13	127	2.8%
14	269	5.9%
15	580	12.7%
16	769	16.8%
17	732	16.0%
18	681	14.9%
19	464	10.2%
Not disclosed	922	20.2%
Total	4570	100.0%

A few overarching themes emerged from the analysis of the findings from the qualitative survey and key informant interview data. The themes indicated the need for improved menstrual health and hygiene, adolescent pregnancy prevention, respect and dignity, and the need to address social determinants of health. **Table two** summarizes selected themes and examples of quotes from the qualitative survey.

Table 2: Sample categorization of themes, categories and codes used based on the survey responses.

Themes	Categories	Codes	Examples
Improved Menstrual Health and Hygiene Management	Period poverty	Sanitary towels	<i>"Sanitary towels"</i>
		Pain relief	<i>"Free sanitary towels for them to be comfortable during the periods and special treatment for those with cramps"</i>
		Free sanitary products	<i>"The government to distribute pads to both primary and high schools, free Education on maternal health in different schools and institutions."</i>
	Water and sanitation	Toilets/ Latrines	<i>"Improved toilets in schools that are hygienic"</i>
		Water	<i>"Water in the toilets"</i>
		Proper disposal	<i>"Teach women how to dispose of sanitary pads."</i>
	Extra clothes and underwear	Extra skirts	<i>"Be given an extra skirt in case of periods leaking, which can cause embarrassment." "I kindly request for underwear and bikers."</i>
Nutrition during menstruation		<i>"Give a list of *nutritious foods, especially when it comes to monthly periods and cramps."</i>	
Prevention of adolescent pregnancy	Risk Factors	Period poverty	<i>"Provide sanitary pads to the needy to reduce teenage pregnancy."</i>
	Access to contraception	Condom use	<i>"Girls are becoming pregnant in our county at a young age; I would request the government to issue free condoms to the youth and *educate girls about teenage pregnancy and its consequences."</i>
		Abstinence	<i>"Teach the girls about abstinence from sex to avoid early pregnancy."</i>
	Address cultural issues	Female genital mutilation	<i>"To prevent female genital mutilation and early pregnancy among girls to avoid school dropouts."</i>
	Abortion	Prevention/ Care	<i>"Serious pregnancy tests to be conducted in institutions to prevent abortions."</i>
Respect and Dignity	Friendly	Non-abusive language	<i>"When young girls end up pregnant and go to a hospital to deliver, the nurses should be kind and stop using abusive languages since they lower self-esteem, most girls try to give birth on their own due to fear of bad treatment"</i>
		Privacy/ confidentiality	<i>"A private and confidential room available concerning personal healthcare issues, provision of enough sanitary pads for quality education"</i>

181
182

Social determinants of health	Transportation		<i>"Provision of good road transport to and from school"</i>
	Infrastructure	Roads	<i>"Government to improve local transport roads to enable easy access to the hospital."</i>
	Poverty	Basic school needs	<i>"Boarding fees and uniform."</i>
		Resources	<i>"Resources are targeted towards women with extreme poverty."</i>

183
184

185 Emerging themes

186 *1. Improved Menstrual Health and Hygiene*

187 Menstrual health care encompassed the physical, emotional, and psychological elements and environmental
188 factors such as the availability of water and toilets. The issues related to menstruation, such as sanitary products,
189 safe disposal, the need for underwear, risk factors for period poverty, and pain medication, emerged across
190 different ages, schools, and regions. One key informant stated:

191 *"I engaged [with] young school-going girls above 15 years of age. Their main request was menstrual hygiene
192 management and the provision of sanitary towels. What breaks my heart is how these goods do not reach some
193 of these schools, and girls are missing school." (Female Youth Advocate)*

194 Adolescent girls also expressed the need for water and sanitation during menstruation. Adolescent girls
195 perceived clean toilets, water availability, and safe disposal of used menstrual products as quality reproductive
196 healthcare. Responses ranged from needing more restrooms and accessible water sources to cleaner toilets and
197 bathrooms. A few girls described unsanitary conditions as risk factors for reproductive health infections. One
198 adolescent girl had this to say:

199 *"To provide a private place where one can change or bath in case of messing because of her periods, have
200 other places to dispose of *sanitary (sanitary towels) other than toilets (Adolescent girl, 17 yrs-Mixed day
201 school).*

202 In the analysis, the need for extra skirts, underwear, and other requests related to menstrual hygiene
203 emerged as a significant ask among adolescent girls in mixed-day schools because they felt embarrassed when

they stained their skirts during their period. A teacher in a mixed school in western Kenya further emphasized the theme of girls' unique needs in her school.

"I buy the sanitary towels from my pocket because I care for the girls, but it is not enough. I also have extra skirts and keep a supply of pain relief medication. I have to have all these because we noticed that when the girls had their periods, they would skip school a few years back. Some were afraid of "spotting" their skirts and being laughed at by the boys, while others stated that they could not afford to buy sanitary pads." (Female teacher - Mixed-day school)

Many adolescent girls requested free or reduced costs of sanitary products and improved quality sanitary pads while directing their requests to government and non-profit organizations. There was also an emphasis of clean, safe spaces, indicating that adolescent girls in Kenya had different wants for menstrual health and hygiene.

2. Prevention of Adolescent Pregnancy

Adolescents perceived prevention of adolescent/teenage pregnancy as quality reproductive care. Some categories associated with reduced adolescent pregnancies included pregnancy prevention methods, risk factors for adolescent pregnancy, abortion and post-abortion care, and proper care for pregnant teens and adolescent mothers. One mobilizer who interacted with teenagers reported that the girls were more concerned about contraception.

"For those who never had children, they were less concerned with what maternal health entails, but rather family planning affordability and accessibility" (Mobilizer, Nairobi)

While some requested condoms and contraception pills for pregnancy prevention, others felt abstinence should be emphasized in schools. Some girls also criticized health care providers who prescribed family planning methods, stating that it promoted sexual immorality. Although the need for birth control access was prevalent, a few condemned oral contraception prescriptions, citing that it led to reproductive health issues.

A few girls mentioned the association between female genital mutilation, child marriages, and teen pregnancies. Other responses noted that the lack of sanitary pads puts girls at risk of teenage pregnancy and that providing menstrual supplies helps reduce teen pregnancies. Adolescents also had differing opinions about

230 abortion. A few girls believed that conducting pregnancy tests in institutions could deter abortions, while others
231 expressed the need for policies and legal processes around safe abortions. Some, however, felt that abortion was
232 dangerous and should be condemned at all costs.

233 3. *Respect and dignity*

234 Consistent with domains in quality-of-care frameworks, the request for respect and dignity was a key
235 demand from adolescent girls. Responses indicated that adolescents want a safe space where their reproductive
236 health needs are met without judgment. However, a few girls noted that health care workers treated pregnant
237 teenagers and adolescent mothers disrespectfully as they sought care.

238 *"When young girls end up pregnant and go to a hospital to deliver, the nurses should be kind and stop using*
239 *abusive language since they lower self-esteem. Most girls try to give birth on their own due to fear of bad*
240 *treatment" (Adolescent girl, 17 years- Boarding school)*

241 Adolescents perceived quality care as using appropriate language, with a few mentioning that they wanted
242 doctors or nurses to be kind and polite and stop using abusive language. Some requested a change in attitude
243 among providers and the elimination of harassment and criticisms that might lead to fear and low self-esteem
244 among adolescent girls. A key informant, a teacher in an all-girls school, echoed the need for respectful care.

245 *"Many girls quietly suffer because of reproductive health infections or sexually transmitted infections.*
246 *Some who have contracted sexually transmitted infections rarely seek medical care for fear of being*
247 *called out, especially when in line. Girls do not feel that the 'youth-friendly corners' meet their needs*
248 *without prejudice." (Female teacher- Boarding school)*

249 Many key informants noted that some teenage girls are reluctant to seek care because of negative past
250 experiences with nurses.

251 4. *Addressing social determinants of health*

252 Other themes emerged during the coding process that initially appeared unrelated to maternal or
253 reproductive. In the parent study, some of these requests were classified as either "other asks" or
254 "undeterminable asks." However, based on literature and further examination, the authors considered economic
255 situations, the physical environment, and social contexts as critical determinants of health that impact

adolescent reproductive health outcomes. A youth advocate's statement reinforces poverty's impact on reproductive health.

"Don't make this just a pad issue!" Girls did not want pamphlets, materials, or instructions on menstrual health either. What they want, what they need are income-generating opportunities. "Girls told us it's not just about whether they have sanitary napkins or not. It's about the difficult choices between basic needs.

"..... "If I'm comparing the price of sanitary towels or napkins to a packet of flour for dinner—I'm thinking I need to eat first." (Female mobilizer- Nairobi)

Access to food and nutrition was also linked to quality reproductive health care, with a few responses requesting balanced diets in schools and others demanding relief food in areas of famine. Some survey responses such as requests for "tuition and school uniform," "need for books," or "school fees" suggested that a few girls were from low socioeconomic households or families experiencing extreme poverty.

Participants not only expressed the need for economic empowerment but also felt that women should be actively involved in politics.

"Women should also participate actively in nation politics. This will help to fight for their rights freely and even come up with possible solutions to the problems that may be hindering them (17-year-old-Boarding school).

Adolescents also want improved physical environment for quality reproductive health care. For example, adolescents requested safe streets, proper lighting, and security in health facilities. Social inclusion emerged as a surprising sub-theme under social determinants of reproductive health. Adolescent girls conveyed the need to empower and educate women and girls to enhance their decision-making power.

Discussion

Overall, the perceptions of quality reproductive care varied across adolescent high school girls in Kenya. Although the study results were similar to the overall *What Women Want* global survey findings, this analysis added value by highlighting unique themes among school-going adolescent females. Study findings acknowledge that while the survey question focused on maternal and reproductive healthcare services, the

281 respondents also highlighted social determinants associated with the healthcare of women and girls. We
282 identified issues adolescents felt were essential for quality reproductive health care.

283 A few emerging cross-cutting themes are consistent with adolescent reproductive health literature. For
284 example, reducing teenage pregnancy rates is a health goal for many countries.^{18,27} Engaging in transactional
285 sex due to period poverty (lack of sanitary towels) is also a risk factor in other studies.^{15,16} Although the survey
286 question was about requests for quality care rather than complaints, certain reproductive health care services,
287 such as contraception, elicited negative feedback from some participants. Similar to other studies, there was
288 evidence of stigma around adolescent use of contraception among a few respondents,¹⁶ although many others
289 requested increased access and additional information on birth control methods. The Adolescent Sexual and
290 Reproductive Health policy in Kenya recognizes the need for a multi-sectoral approach in addressing teenage
291 pregnancy,¹⁸ given the economic, sociocultural, and health system factors that impact adolescent girls.¹⁵
292 Historically, sociocultural systems prepared adolescents for menstruation, marriage, and pregnancy, where
293 uncles and aunts provided sexuality education informally or during initiation ceremonies.²⁸ However, given the
294 societal shifts that emphasize nuclear families, and migration to cities for employment, adolescents have to rely
295 on other sources for sexual and reproductive health education.

296 The need for improved menstrual health and hygiene was a common thread among female adolescents
297 of all ages and schools. As such, we combined some categories (*water and sanitation* and *other determinants*
298 "skirts") into one theme, "Improved Menstrual Health and Hygiene," based on the study findings. The request
299 for additional skirts, including underwear, was more salient among girls in mixed-day schools probably
300 because, similar to other studies, menstruating girls are likely to face ridicule from boys or community members
301 when they spot during their period.²⁹ Compared to girls in boarding schools who can go to their dorms during
302 breaks and change, those in day schools are forced to continue classes the entire day with stained skirts or go
303 home to change. It is also likely that some girls in day schools who come from poor households and cannot
304 afford the cost of attending boarding schools also face challenges in obtaining sanitary products. Previous
305 studies on menstruation report that absenteeism among adolescent girls was associated with menstruation due to
306 a lack of sanitary towels, dysmenorrhea, embarrassment, or non-functional toilets.^{29,30} Although issues related

307 to period poverty were significant requests for quality reproductive and maternal care, menstrual health
308 management originated as an intervention related to education.²⁹

309 The enactment of the Kenya Menstrual Hygiene Management Policy was lauded as a step toward
310 improving educational outcomes for adolescent girls attending public schools in Kenya. However, there have
311 been gaps in proper implementation due to challenges in the narrow conceptualization of menstrual health
312 hygiene as just "pad distribution".³¹ Studies in the United States and India also share similar findings on the
313 need to frame Menstrual Health Hygiene as a comprehensive policy that impacts mental health, water and
314 sanitation, education, and financial sectors.³¹ As such implementation of menstrual health hygiene in schools not
315 only addresses sustainable development goal (SDG) 3 of improving health and well-being but also SDG 4, in
316 the provision of equitable and quality education especially for adolescent girls.^{13, 14}

317 It is important to recognize that even though investments are made to improve sanitation and increase
318 reproductive health supplies and products, adolescent girls may not fully seek care or utilize services if they feel
319 disrespected and undervalued. Health care providers and practitioners working with adolescents should seek to
320 uphold the dignity of adolescents without discrimination. Although it may be challenging, health providers
321 should incorporate respectful language, dignified care, enhance privacy and confidentiality as a strategy for
322 engaging adolescents seeking reproductive health care.

323 Overall, quality reproductive healthcare should not only be examined from the perspective of healthcare
324 providers or professionals in the adolescent reproductive healthcare space. However, incorporating adolescent
325 requests can be significant, as adolescents are experts on their health.²³ The impact of economic and social
326 determinants of health on reproductive health was also evident in the responses from adolescent girls in
327 secondary schools in Kenya. As such, reproductive health programs should consider working with individuals
328 in the ministries of Education, Finance, Food and Agriculture, Transport and Infrastructure, as well as those in
329 the Sports, Culture & Heritage sectors.

330 Strengths of the study

331 To our knowledge, this is the first study in Kenya on reproductive health that has asked women and girls
332 what they want regarding quality reproductive health. By utilizing an open-ended question, participants felt free

333 to express what they wanted and what they hoped could be implemented to improve their health. The large
334 sample size elicited diverse responses on quality reproductive health care from the perspective of adolescent
335 high school girls in Kenya. Along with these unique responses, our data highlighted the different demands by
336 type of school and region, suggesting the need to target interventions unique to specific populations. The
337 perspectives of key informants further reinforced the survey results. Reproductive health advocates are currently
338 disseminating the findings from the study to policymakers and program implementers as evidence of demands
339 from women and girls. Adolescent reproductive health care programs are likely to be more successful if they
340 target the self-articulated needs of adolescents.

341 Limitations

342 This study has limitations. First, the qualitative survey asked for only one request for quality healthcare.
343 Adolescents probably had more requests but had to pick only one. Second, while the study offered girls a
344 chance to speak up, it did not explore the reason behind the "asks" and how to navigate the issues. Additionally,
345 this sub-sample was not nationally representative of all adolescent girls in Kenya. Some girls living with
346 disabilities, in humanitarian settings, or in remote/ arid areas in Kenya may have different perceptions of quality
347 reproductive health care. Future studies should explore conducting in-depth studies such as focus groups on
348 quality reproductive healthcare among adolescent girls.

349 Acknowledgments

350 We would like to thank all the adolescent girls who completed the survey and their respective schools. We also
351 appreciate the mobilizers, data collectors, and data entry personnel for their hard work and dedication. We
352 would also like to acknowledge the White Ribbon Alliance staff who were involved in analyzing the entire
353 global *What Women Want* survey.

354

355

356

357

References

1. Woog V, Alyssa B, Jesse P. Adolescent Women's Need for and Use of Sexual and Reproductive Health Services in Developing Countries. New York: Guttmacher Institute. 2015[Cited 2022 August 9]. Available from: <https://www.guttmacher.org/report/adolescent-womens-need-and-use-sexual-and-reproductive-health-services-developing-countries> Accessed
2. World Health Organization: WHO recommendations on adolescent sexual and reproductive health and rights. [Cited 2022 August 9]. Available from: <https://www.who.int/publications/i/item/9789241514606>
3. Liang M, Simelane S, Fortuny Fillo G, et al. The State of Adolescent Sexual and Reproductive Health. *J Adolesc Health*. 2019;65(6S): S3-S15. doi:10.1016/j.jadohealth.2019.09.015
4. Ravindran TKS, Govender V. Sexual and reproductive health services in universal health coverage: a review of recent evidence from low- and middle-income countries. *Sex Reprod Health Matters*. 2020;28(2):1779632. doi:10.1080/26410397.2020.1779632
5. Murewanhema G. Adolescent girls, a forgotten population in resource-limited settings in the COVID-19 pandemic: implications for sexual and reproductive health outcomes. *Pan Afr Med J*. 2020;37(Suppl 1):41. doi:10.11604/pamj.supp.2020.37.41.26970
6. Grønvik T, Fossgard Sandøy I. Complications associated with adolescent childbearing in Sub-Saharan Africa: A systematic literature review and meta-analysis. *PLoS One*. 2018;13(9):e0204327. doi:10.1371/journal.pone.0204327
7. Hackett K, Lenters L, Vandermorris A, et al. How can engagement of adolescents in antenatal care be enhanced? Learning from the perspectives of young mothers in Ghana and Tanzania. *BMC Pregnancy Childbirth*. 2019;19(1):184. doi:10.1186/s12884-019-2326-3
8. Melesse DY, Mutua MK, Choudhury A, et al. Adolescent sexual and reproductive health in sub-Saharan Africa: who is left behind?. *BMJ Glob Health*. 2020;5(1):e002231. doi:10.1136/bmjgh-2019-002231
9. Birdthistle IJ, Floyd S, Machingura A, Mudziwapasi N, Gregson S, Glynn JR. From affected to infected? Orphanhood and HIV risk among female adolescents in urban Zimbabwe. *AIDS*. 2008;22(6):759-766. doi:10.1097/QAD.0b013e3282f4cac7
10. United Nations: Transforming our world: The 2030 agenda for sustainable development, United Nations A/RES/70/1, New York, 21 October 2015. [Cited 2022 August 9]. Available from: https://www.un.org/en/development/desa/population/migration/generalassembly/docs/globalcompact/A_RES_70_1_E.pdf
11. Waage J, Yap C, Bell S, et al. Governing the UN sustainable development goals: interactions, infrastructures, and institutions. *Lancet Glob Health*. 2015;3(5):e251-e252. doi:10.1016/S2214-109X(15)70112-9
12. World Health Organization. Global accelerated action for the health of adolescents (AA-HA!): guidance to support country implementation. Global Accelerated Action for the Health of Adolescents (AA-HA!) Guidance to Support Country Implementation. 2017. [Cited 2022 August 20]. Available from: <https://www.who.int/publications/i/item/9789241512343>
13. Darroch JE, *Adding It Up: Investing in Contraception and Maternal and Newborn Health, 2017—Estimation Methodology*, New York: Guttmacher Institute, 2018. [Cited 2022 August 20]. Available from: <https://www.guttmacher.org/report/adding-it-up-investing-in-contraception-maternal-newborn-health-2017-methodology>.
14. Kenya National Bureau of Statistics: Kenya Demographic Health Survey 2014. Nairobi, Kenya: Ministry of Health Kenya and ICF international. 2015. [Cited 2022 August 20]. Available from: <https://dhsprogram.com/pubs/pdf/FR308/FR308.pdf>.
15. Olenja J, Krugu JK, Kwaak v.d. A, Kawai D, Karanja S, Apanja M, et al., Factors influencing teenage pregnancy among Maasai girls in Kajiado West Sub-County, Kenya. An operational qualitative study report as part of the YES I DO programme implemented from 2016 to 2020. [Cited 2022 August 20]. Available from: <https://www.kit.nl/wp-content/uploads/2021/01/YID-Kenya-Report-Teenage-pregnancy-final.pdf>

- 409 16. Harrington EK, Casmir E, Kithao P, et al. "Spoiled" girls: Understanding social influences on adolescent
410 contraceptive decision-making in Kenya. *PLoS One*. 2021;16(8):e0255954.
411 doi:10.1371/journal.pone.0255954
- 412 17. Mburu A, Itsura P, Mabeya H, Kaaria A, Brown DR. Knowledge of Cervical Cancer and Acceptability
413 of Prevention Strategies Among Human Papillomavirus-Vaccinated and Human Papillomavirus-
414 Unvaccinated Adolescent Women in Eldoret, Kenya. *Biores Open Access*. 2019;8(1):139-145..
415 doi:10.1089/biores.2019.0007
- 416 18. Ministry of Health Kenya: Kenya Adolescent Sexual and Reproductive Health Policy. Nairobi, Kenya:
417 Ministry of Health, Kenya, 2015. [Cited 2022 September 3]. Available from:
418 https://www.popcouncil.org/uploads/pdfs/2015STEPUP_KenyaNationalAdolSRHPolicy.pdf
- 419 19. Mazur A, Brindis CD, Decker MJ. Assessing youth-friendly sexual and reproductive health services: a
420 systematic review. *BMC Health Serv Res*. 2018;18(1):216. doi:10.1186/s12913-018-2982-4
- 421 20. Girod C, Ellis A, Andes KL, Freeman MC, Caruso BA. Physical, social, and political inequities
422 constraining girls' menstrual management at schools in informal settlements of Nairobi, Kenya. *Journal*
423 *of Urban Health*. 2017 Dec;94(6):835-46. doi: 10.1007/s11524-017-0189-3.
- 424 21. Phillips-Howard PA, Caruso B, Torondel B, Zulaika G, Sahin M, Sommer M. Menstrual hygiene
425 management among adolescent schoolgirls in low-and middle-income countries: research priorities.
426 *Global health action*. 2016 Dec 1;9(1):33032. doi: 10.3402/gha.v9.33032.
- 427 22. Pilgrim NA, Jennings JM, Sanders R, et al. Understanding Quality of Care and Satisfaction With Sexual
428 and Reproductive Healthcare Among Young Men. *J Healthc Qual*. 2018;40(6):354-366.
429 doi:10.1097/JHQ.000000000000149
- 430 23. Ayieko S, Nguku A. Engaging Youth in Citizen-Led Advocacy and Accountability for Adolescent
431 *Sexual and Reproductive Health*. *East Afr Health Res J*. 2019;3(2):85-87. doi: 10.24248/eahrj.v3i2.603.
- 432 24. White Ribbon Alliance. What Women Want 2022. [Cited 2022 September 3]. Available from:
433 <https://whiteribbonalliance.org/resources/www-dashboard/>
- 434 25. Jain AK, Hardee K. Revising the FP Quality of Care Framework in the Context of Rights-based Family
435 Planning [published correction appears in *Stud Fam Plann*. 2018 Dec;49(4):397-401]. *Stud Fam Plann*.
436 2018;49(2):171-179. doi:10.1111/sifp.12052
- 437 26. ATLAS. ti Scientific Software Development GmbH. 2019. ATLAS. ti. [Cited 2022 September 3].
438 Available from: <https://atlasti.com>
- 439 27. Lugo-Gil J, Lee A, Vohra D, Harding J, Ochoa L, Goesling B. Updated findings from the HHS teen
440 pregnancy prevention evidence review: August 2015 through October 2016. US Department of Health
441 and Human Services. 2018. [Cited 2022 September 3]. Available from:
442 https://tppevidencereview.youth.gov/pdfs/Summary_of_findings_2016-2017.pdf
- 443 28. De Haas B, Hutter I. Teachers' conflicting cultural schemas of teaching comprehensive school-based
444 sexuality education in Kampala, Uganda. *Cult Health Sex*. 2019;21(2):233-247.
445 doi:10.1080/13691058.2018.1463455
- 446 29. Mohammed S, Larsen-Reindorf RE. Menstrual knowledge, sociocultural restrictions, and barriers to
447 menstrual hygiene management in Ghana: Evidence from a multi-method survey among adolescent
448 schoolgirls and schoolboys. *PLoS One*. 2020;15(10): e0241106. doi: 10.1371/journal.pone.0241106
- 449 30. Sommer M, Hirsch JS, Nathanson C, Parker RG. Comfortably, Safely, and Without Shame: Defining
450 Menstrual Hygiene Management as a Public Health Issue. *Am J Public Health*. 2015;105(7):1302-1311.
451 doi:10.2105/AJPH.2014.302525
- 452 31. Alhelou N, Kavattur PS, Rountree L, Winkler IT. 'We like things tangible!' A critical analysis of
453 menstrual hygiene and health policy-making in India, Kenya, Senegal and the United States [published
454 online ahead of print, 2021 Dec 9]. *Glob Public Health*. 2021;1-14.
455 doi:10.1080/17441692.2021.2011945
- 456
- 457
- 458