

1 **A DISTRIBUTIONAL REGRESSION APPROACH TO MODELING THE IMPACT**
2 **OF STRUCTURAL AND INTERMEDIARY SOCIAL DETERMINANTS ON**
3 **COMMUNITIES BURDENED BY TUBERCULOSIS IN EASTERN AMAZONIA –**
4 **BRAZIL**

5
6 Authors:

7 Clóvis Luciano Giacomet¹, Antônio Carlos Vieira Ramos², Heriederson Sávio Dias Moura²,
8 Thaís Zamboni Berra², Yan Mathias Alves², Felipe Mendes Delpino^{2,3}, Jason E. Farley⁴, Nancy
9 R. Reynolds⁴, Jonas Bodini Alonso⁵, Titilade Kehinde Ayandeyi Teibo², Ricardo Alexandre
10 Arcêncio^{6*}

11
12 ¹Interunits PhD Program in Nursing, University of São Paulo College of Nursing at Ribeirão
13 Preto, Ribeirão Preto, Brazil

14 ²Graduate Program in Public Health Nursing, University of São Paulo College of Nursing at
15 Ribeirão Preto, Ribeirão Preto, Brazil

16 ³Postgraduate Program in Nursing, Federal University of Pelotas, Pelotas, Brazil

17 ⁴The Center for Infectious Disease and Nursing Innovation, Johns Hopkins School of
18 Nursing, Baltimore, United States of America

19 ⁵Research Support Center, University of São Paulo College of Nursing at Ribeirão Preto,
20 Ribeirão Preto, Brazil

21 ⁶Department of Maternal and Child Nursing and Public Health, University of São Paulo
22 College of Nursing at Ribeirão Preto, Ribeirão Preto, Brazil

23

24 *Corresponding author

25 Email: ricardo@ernp.usp.br
Note: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

26 **ABSTRACT**

27 **Background:** TB is a disease affected by social determinants of health; however, it is unclear
28 what its structural and intermediary determinants are in Eastern Amazonia. The region contains
29 many natural resources, yet it suffers drastically from poverty, inequality, and neglected
30 diseases. Here, we aimed to employ mathematical modeling to evaluate the influence of
31 structural and intermediary determinants of health on TB in Eastern Amazonia – Brazil.

32 **Methods:** We conducted an ecological study. We considered cases diagnosed of TB and
33 collected data by census tract to measure the social determinants. We applied the *generalized*
34 *additive model for location, scale, and shape* (GAMLSS) framework to identify the effect of
35 social determinants on communities with a high prevalence of TB. The Double Poisson
36 distribution (DPO) was selected and we tested the inclusion of quadratic effects.

37 **Results:** 1,730 people were selected. The majority were female (59.3%), aged 31 to 59 years
38 (47.6%), blacks (67.9%), schooling level of 5 to 8 years (18.7%). Prevalence of alcoholism
39 was 8.6% and mental illness 0.7%. The GAMLSS analyses showed that the risk of community
40 incidence of TB is associated with the proportion of the population without basic sanitation
41 and also with the age groups 16-31 years and > 61 years.

42 **Conclusions:** The study revealed that GAMLSS is a strategic tool to identify territories at
43 greatest risk for TB. Models should have broader scope to include social determinants so as to
44 better inform policy to reduce inequality and achieving the goal of the End TB strategy.

45 INTRODUCTION

46 Brazil is among the countries with the highest tuberculosis (TB) burden[1]. According
47 to the latest report from the World Health Organization (WHO), Brazil identified more than
48 66,819 cases in 2020. This shows a decrease of 18% compared with data from the year 2019,
49 but at the same time, there was an increase in TB-related mortality[1].

50 Although Brazil has committed to ending TB by 2050, the country has faced enormous
51 challenges in achieving this ambitious goal. This is largely due to austerity measures that have
52 cut social benefits to lower socio-economic groups as well as to a serious economic crisis
53 brought on by the pandemic and government policies[2]. According to forecast models, the
54 TB targets of the UN Sustainable Development Goals (SDGs) (which aims for a 90% reduction
55 in TB deaths by 2030) are not likely to be achieved.

56 There is a broad agreement that progress in tuberculosis control will require not only
57 investment in strengthening tuberculosis control programs, diagnostics, and treatment, but also
58 action on the Social and structural Determinants of TB. The Structural factors include
59 governing action, economic and social policies, and the position of power, prestige, and
60 resources in accordance with the social level occupied by an individual, family, or
61 community[3]. Intermediary determinants consist of material circumstances, psychosocial
62 behavior, biological factors, and the health system[3]. Literature is replete with evidence of
63 their influence in the context of TB,[4-9] but are without making distinctions on the dimensions
64 of these determinants.

65 Mathematical modelling can be useful in exploring the contribution of health drivers to
66 face the epidemic and browse through evidences to End TB strategies. However, current TB
67 models have limitations, for example, a systematic review conducted by Pedrazzoli and
68 colleagues showed that that few studies employing mathematical modeling have addressed
69 underlying Social and Structural determinants, which makes this a knowledge gap[4].

70 Although many mathematical models have been used for forecasting TB, most of them tended
71 to ignore the differences in variance and its asymmetry/imbalance and they retain almost
72 exclusively the mean for explaining the outcome[10], which has revealed an important gap in
73 knowledge.

74 Studies have provided results that are more satisfactory when alternative regressions
75 techniques are applied. Generalized additive models for location, scale, and shape (GAMLSS)
76 has allowed a more expanded approach where it worked with not only the mean (or location)
77 but also all the parameters of the conditional distribution of outcome to be modeled as
78 parametric/or nonparametric additive functions of independent variables and/or random-effects
79 terms[11,12]. Therefore, here, we aimed to employ mathematical modeling to evaluate the
80 impact of structural and intermediary social determinants on TB in Eastern Amazonia – Brazil.

81

82 **METHODS**

83 **Study design**

84 We used an ecological, mathematical modeling study design.

85

86 **The setting of the study**

87 We conducted the study in Macapá, the capital of Amapá state, Eastern Amazonia (Fig
88 1). It has a population of about 398,204 inhabitants, and a demography density near 62.14
89 people per km². It is the most populous city in Northern Amazonas State.

90

91 **Fig 1. Map of the study setting.**

92

93 Regarding TB, the city presented an incidence of 17.5 cases per 100,000 in 2020, which
94 represented a decrease of 25% compared to the previous year's data[13]. A problematic

95 situation is that only 39.3% of the patients diagnosed with TB had confirmation from the
96 bacteriological analysis (microscopy or GeneXpert MTB/RIF), and 47.3% did not have
97 confirmation of the diagnosis of TB through a bacteriological test[14].

98

99 **The population of the study and criteria**

100 The population was composed of all the cases diagnosed with tuberculosis and registered
101 in the Notification of Notifiable Diseases System (SINAN) in the period from 2001 to 2017.
102 We gathered socio-demographic data (age, gender, race/ color self-reported and years of study
103 and occupation) and clinical information (type of case – new or retreatment; clinical form –
104 pulmonary or extrapulmonary; coinfection with TB-HIV; alcoholism; mental disorders;
105 comorbid TB-diabetes).

106 In accordance with the WHO guidelines, the diagnosis of TB refers to the recognition of
107 an active case when a patient with *Mycobacterium tuberculosis complex* is identified from a
108 clinical specimen, either by microscopy, culture, or a newer method such as a molecular line
109 probe assay[15]. In Brazil, a pulmonary case with one or more initial sputum smear
110 examinations positive for acid-fast bacilli (AFB) is also defined as a "case"[16]. Eventually,
111 the diagnosis can be established only through clinical examination by a physician and X-ray;
112 however, it is not recommended by the Brazilian sanitary authorities. New patients are defined
113 as those who have no history of prior TB treatment or who have been treated for at least/ less
114 one month. These cases should receive a regimen containing six months of rifampicin:
115 2HRZE/4HR[16].

116

117 **Unity of study analysis and variables**

118 The unit of study analysis was 811 Urban Census Tract (UCT) contained in Macapá,
119 collected from the Brazilian Institute of Geography and Statistics (IBGE). The UCT consists

120 of the smallest territorial unit formed by a continuous area located in an urban area, with a
121 defined size, number of households, and number of residents, used for Brazilian surveys and
122 statistical research[17]. The information regarding UCT was obtained from the 2010 Brazilian
123 Demographic Census, which collected data regarding both structural and intermediary
124 determinants expressed by household conditions and characteristics of the territories (UCT). In
125 Fig 2 is shown the variable under study.

126

127 **Fig 2. Structural and intermediary determinants selected for the study, Eastern**
128 **Amazonia – Brazil.**

129

130 **Analysis plan**

131 Initially, the variables were analyzed by descriptive statistics. Additionally, for
132 identifying the structural and intermediary determinants associated with TB, we used the model
133 GAMLSS[11]. The reason for applying this model is because the modeling of the response
134 variable (TB cases) did not follow a distribution of exponential family, and it showed
135 heterogeneity in terms of distribution scale and shape. The response variable changed with the
136 explanatory variables[19].

137 Since $y^T = (y_1, \dots, y_n)$ is a vector of size n of the response variable with density function
138 $f(y_i|\theta^i)$, where $\theta^i = (\theta_{1i}, \theta_{2i}, \theta_{3i}, \theta_{4i}) = (\mu_i, \sigma_i, \nu_i, \tau_i)$, and let $k = 1, 2, 3, 4$ e let $g_k(\cdot)$ a monotone
139 link function that relates the parameters to the independent variables from the equations:

$$\left\{ \begin{array}{l} g_1(\mu) = n_1 = X_1\beta_1 + \sum_{j=1}^{J_1} Z_{j1}\gamma_{j1}, \\ g_2(\sigma) = n_2 = X_2\beta_2 + \sum_{j=1}^{J_2} Z_{j2}\gamma_{j2}, \\ g_3(\nu) = n_3 = X_3\beta_3 + \sum_{j=1}^{J_3} Z_{j3}\gamma_{j3}, \\ g_4(\tau) = n_4 = X_4\beta_4 + \sum_{j=1}^{J_4} Z_{j4}\gamma_{j4}. \end{array} \right.$$

140

141 Where $\mu, \sigma, v \in \tau$ are vectors of length n , $\beta_k = (\beta_{1k}, \beta_{2k}, \dots, \beta_{jk})$ is a vector of length j^k
142 eX_k is the delineation matrix of order $n \times j^k$. The function h_{jk}

143 Non additive function of the independent variable X_k evaluated at x_{jk} .

144 The selection of the dependent variable distribution was performed using the
145 Generalized Akaike information criterion, defined by $GAIC = -2l(\hat{\theta}) + bdf$, where $l(\hat{\theta})$ the
146 likelihood function, b is a penalty parameter, and df denotes the degrees of freedom of the
147 model[20]. For $b = 2$ we have the original Akaike information criterion (AIC). According to
148 literature[21] all distributions falling in the GAMLSS class are presented.

149 The selection of the independent variables was made in 2 steps. In the first step, the
150 presence of multicollinearity among the independent variables was evaluated. The
151 multicollinearity assessment evaluates the entry of variables in the model that are highly
152 correlated with each other. One of the most used measures is the Variance Inflation Factor,
153 whose expression is defined by:

154
$$VIF_j = \frac{1}{1 - R_j^2}$$

155 Where R_j^2 is the multiple correlation coefficient resulting from regressing X_j on the other
156 $p - 1$ regressor. The higher the degree of dependence of X_j on the remaining regressors, the
157 stronger the dependence and the higher the value of R_j^2 . A VIF value > 5 was adopted as the
158 cutoff point[12,22]. Since the independent variables are the same for both outcomes, the result
159 of this analysis is also considered valid for both.

160 Additionally, we applied the stepwise method using the Generalized Akaike
161 Information Criterion (GAIC) with $k = 4$ [12] for the remaining variables from the first stage.
162 After this analysis, The Double Poisson distribution (DPO) was selected in accordance with
163 the AIC value (Table 3), where DPO (μ, σ) has the following probability density function[21].

164
$$f(y|\mu, \sigma) = \left(\frac{1}{\sigma}\right)^{\frac{1}{2}} e^{-\frac{\mu}{\sigma}} \left(\frac{e^{-y} y^y}{y!}\right) \left(\frac{e\mu}{y}\right)^{y/\sigma} C$$

165 Where $y = 0, 1, 2, \dots, \infty$, $\mu > 0$ and $\sigma > 0$, where C is a proportionality constant that is
 166 calculated numerically. The link function between the parameters and the independent
 167 variables is the log function, i.e., $g_1(\mu) = \log(\mu)$ and $g_2(\sigma) = \log(\sigma)$ [21].

168 We verified the model's adequacy through the model's diagnostic graphics: Fitted
 169 Values x Residuals, Order of Observations x Residuals, Distribution of Residuals, and the
 170 Quantile-Quantile plot (Q-Q plot). Additionally, the Shapiro-Wilk Normality test was applied
 171 to the model's residuals to verify if they are fit with the Standard Normal distribution. We
 172 considered the dependent variable, the number of TB cases in each UCT, and the structural and
 173 intermediary determinants indicators as the independent variables (Table 1). These variable/
 174 proxy indicators were selected in accordance with the theoretical framework defined in the
 175 study[4, 19].

176

177 **Table 1. Socio-demographic and clinical-epidemiological profile of Tuberculosis cases,**
 178 **Eastern Amazonia (Brazil).**

Variables	N=1,730 (%)
Age (years)	
0 to 14 years	74 (4.3)
15 to 30 years old	640 (37.0)
31 to 59 years	823 (47.6)
Above 60 years old	175 (10.1)
Blank/Ignored	18 (1.0)
Gender	
Male	704 (40,7)
Female	1026 (59,3)
Years of study	
Illiterate	128 (7.4)
Elementary school complete	112 (6.5)
Elementary School Incomplete	803 (46.6)
High School Complete	224 (12.9)
Higher Education Incomplete	52 (3.0)
Higher education complete	127 (7.3)
Blank/Ignored	284 (16.4)
Registration types	
New	682 (39.4)
Relapse	9 (0.5)

Retreatment after abandonment	0 (0.0)
Retreatment	1 (0.1)
Transfer	1 (0.1)
After death	26 (1.5)
Blank/Unknown	1.011 (58.4)
Clinic form	
Extrapulmonary	236 (13.6)
Pulmonary	1472 (85.1)
Pulmonary + Extrapulmonary	18 (1.0)
Blank/Ignored	4 (0.3)
Coinfection TB-HIV	
No	787 (45.5)
Yes	77 (4.5)
Blank/Ignored	866 (50.0)
Coinfection TB-Diabetes	
No	912 (52.8)
Yes	94 (5.4)
Blank/ Ignored	724 (41,8)
Alcoholism	
No	896 (51.7)
Yes	148 (8.6)
Blank/ Ignored	686 (39.7)
Mental Disorders	
No	1001 (57.8)
Yes	12 (0.7)
Blank/Ignored	717 (41.5)

179

180 We performed the GAIC selection of the independent variables considering only the
 181 linear effects. Additionally, we tested the inclusion of quadratic effects since in the scatter plots,
 182 the fitted curve (by the loess method = local polynomial regression) evidenced a possible
 183 quadratic relationship. To compare the two models, linear terms versus quadratic terms, we
 184 used the Likelihood Ratio Test (LR). Once we selected the fittest model, we estimated the
 185 Relative Increase, expressed in percentage, in the Average Number of Tuberculosis Cases
 186 through the expression $AR(\beta) = [\exp(\beta) - 1] * 100\%$. The R program version 4.1.1 through
 187 the GAMLSS library[23] was used to perform the data analysis for this part.

188

189 **Ethics approval and consent to participate**

190 In compliance with Resolution 499/2012 of the National Health Council in Brazil, the
191 study was authorized by the Municipal Health Secretariat of Macapá- Brazil and approved by
192 the Research Ethics Committee from the University of São Paulo College of Nursing at
193 Ribeirão Preto, which had the Certificate of Submission for Ethics Appreciation (CAAE)
194 Number. 23043019.2.0000.5393. Consent to participate not applicable, because we work with
195 secondary data. All participant identifiers were removed.

196

197 **RESULTS**

198 We identified 1,730 patients diagnosed with TB. Table 1 shows the main characteristics
199 of the patients, in which is observed an age ranging from 1 to 89 years (median = 44.5 years).
200 Most were female (59.3%), aged 31 to 59 years (47.6%), blacks (67.9%), and with an education
201 level of 5 to 8 incomplete years (18.7%). In terms of the clinical and epidemiological profile,
202 cases were mostly characterized as new (39.4%) and pulmonary (85.1%) TB. Regarding
203 comorbidities, we observed a TB-HIV prevalence of 4.5%; TB-Diabetes of 5.4%; Alcoholism
204 at 8.6%; and mental illness at 0.7%. We observed an excess of cases numbers with ignored or
205 blank data.

206 In Fig 3, evidences the spatial distribution of TB cases, considering the comorbid or
207 health condition: TB-HIV coinfection mental disorder, diabetes or alcoholism.

208

209 **Fig 3. Spatial distribution of tuberculosis in the study, Eastern Amazonia.**

210

211 In Table 2, we presented the main information obtained from the descriptive statistics.
212 For the dependent variable, we considered the total of cases and the independents (proxy
213 variables of structural and intermediary social determinants) the proportion of households in
214 each UCT under those conditions characterized by the variable.

215 **Table 2. Statistics descriptive of the variables defined in the study, Eastern Amazonia.**

216

Variables	Mean	Standard Deviation	Minimum	Q0.25	Q0.5	Q0.75	Maximum
Cases of TB per UCT *	3.43	3.00	0.00	1.00	3.00	5.00	23.00
v1 Proportion of households with 1 to 3 people	42.32	7.56	19.15	37.77	42.21	47.19	76.81
v2 Proportion of households with 4 to 6 people	43.79	5.42	15.94	40.70	43.81	47.34	56.52
v3 Proportion of households with 7 to 9 people	10.82	4.07	0.00	8.28	10.17	12.86	29.79
v4 Proportion of households with higher than 10 people	3.08	2.19	0.00	1.60	2.70	4.04	14.89
v5 Proportion of males without schooling	18.33	7.28	3.28	13.09	17.60	22.73	44.06
v6 Proportion of females without schooling	17.14	6.82	2.33	12.33	16.12	21.38	43.90
v7 Proportion of own and paid-up domiciles	76.05	12.20	17.55	69.28	77.94	84.40	100.00
v8 Proportion of rented dwellings	16.70	9.72	0.00	9.45	15.32	22.65	67.11
v9 Proportion of domiciles in another property' condition (not owned, rented, and not granted)	0.19	0.67	0.00	0.00	0.00	0.00	10.20
v10 Proportion of households with a bathroom for exclusive use by residents or a toilet and drainage in a general sewage or rainwater network	65.63	14.40	9.09	56.64	65.60	75.00	100.00
v11 Proportion of houses without a bathroom for the exclusive use of residents or toilet	1.11	1.86	0.00	0.00	0.36	1.48	12.00
v12 Permanent particular residences with water supply from the general network	59.39	36.91	0.00	22.02	70.27	94.16	100.00
v13 Permanent particular residences with water supply from wells or springs on the property	36.73	34.79	0.00	4.49	25.65	70.83	100.00
v14 Permanent Particular Domiciles with Another Form of Water Supply	3.28	7.78	0.00	0.00	0.80	2.88	82.50
v15 Permanent private households with a bathroom exclusively for residents or sanitary	98.29	4.36	22.33	97.89	99.32	100.00	100.00
v16 Permanent private households with exclusive use of bathroom or toilet and sanitary sewage via the general sewage network or rainwater drainage	10.06	21.83	0.00	0.00	1.06	5.27	98.90

v17 Permanent private households with a bathroom for exclusive use by the residents or a toilet and sanitary sewage via a septic tank	18.73	26.59	0.00	0.88	6.02	25.09	100.00
v18 Permanent private households with a bathroom for exclusive use by the residents or a toilet and sanitary sewage via a rudimentary septic tank	49.23	33.94	0.00	14.18	54.88	81.91	99.36
v19 Permanent private households with a bathroom for exclusive use by the residents or a toilet and sewage via river, lake, or sea	17.28	28.16	0.00	0.00	1.09	21.06	100.00
v20 Proportion of households with no nominal monthly income per capita	4.07	4.83	0.00	1.31	2.68	5.06	38.64
v21 Proportion of households with per capita nominal monthly income of up to 1/8 to 1 minimum wage	58.02	20.22	7.63	43.61	60.00	73.71	96.09
v22 Proportion of households with a nominal monthly per capita income of more than 1 to 2 minimum wages	18.82	7.02	2.61	13.28	20.19	24.24	38.94
v23 Proportion of domiciles with nominal monthly income per capita of more than 2 to 3 minimum wages	7.75	5.63	0.00	3.14	6.85	11.41	36.36
v24 Proportion of households with a nominal monthly income per capita of more than 3 to 5 minimum wages	6.47	5.70	0.00	1.64	5.00	10.21	24.29
v25 Proportion of households with a nominal monthly income per capita of more than 5 to 10 minimum wages	3.93	4.57	0.00	0.71	2.24	5.55	23.25
v26 Proportion of households with nominal monthly income per capita higher than 10 minimum wages	1.20	2.05	0.00	0.00	0.42	1.42	
v27 Proportion of resident people of white race or ethnicity	27.10	6.51	5.88	23.31	26.96	30.70	14.12
v28 Proportion of black residents	9.18	5.49	0.00	5.63	8.57	11.57	52.63
v29 Proportion of resident people of yellow race or ethnicity	1.19	1.34	0.00	0.26	0.80	1.80	39.60
v30 Proportion of resident people of mixed race or ethnicity	62.34	8.35	31.15	57.59	61.96	67.50	94.12
v31 Proportion of resident people of indigenous race or ethnicity	0.20	0.51	0.00	0.00	0.00	0.16	

v32 Proportion of people aged 0-15	32.74	7.21	9.44	27.46	32.64	38.28	6.77
v33 Proportion of people aged 16 to 30 years old	31.35	3.12	23.23	29.43	31.16	33.26	51.47
v34 Proportion of persons aged 31 to 60 years old	32.76	5.08	19.22	29.36	32.72	36.37	54.44
v35 Proportion of people aged 61 and over	5.16	2.93	0.87	3.21	4.16	6.10	47.73

217

218 In Table 3, we observed the results from the AIC criterion's application for the selection
 219 of the most appropriate probability distribution considering the total number of TB cases, by
 220 which we can identify that the Double Poisson distribution (DPO) presented the best result (the
 221 highest AIC).

222

223 **Table 3. The main distributions for fitting a GAMLSS model selected in accordance with**
 224 **the Akaike Information Criterion value, Eastern Amazonia (Brazil).**

225

Distribution for fitting a GAMLSS model	Akaike information criterion (AIC)
The Double Poisson distribution (DPO)	2004.25
Zero Inflated Negative Binomial Distribution (ZINBI)	2006.04
Zero adjusted negative binomial distribution (ZANBI)	2006.04
Zero adjusted (hurdle) of the beta negative binomial distribution (ZABNB)	2008.02
Zero-inflated of the beta negative binomial distribution (ZIBNB)	2008.02
The Zipf and zero adjusted Zipf distributions (ZINBF)	2008.04
The Sichel distribution (ZISICHEL)	2008.04
Zero-inflated Poisson inverse Gaussian distribution (ZIPIG)	2008.22
Negative Binomial type II distribution (SNBII)	2010.27
Negative Binomial type I distribution (SBNBI)	2010.27

226

227 In Fig 4, we have the superposition of the DPO density over the data distribution. The
 228 application of the Shapiro-Wilk normality test on the model's residuals shows the adequacy of
 229 the fit ($W = 0.9967$; $p\text{-value} = 0.5245$).

230

231 **Fig 4. Histograms of Tuberculosis cases with fit to the transformed Double Poisson**
 232 **distribution, Eastern Amazonia (Brazil).**

233

234 In Table 4, we included the statistics obtained from the modeling analysis, where we
 235 identify the complete model (saturated) selected through GAIC (no outlier) with quadratic

236 terms and no quadratic outlier (v33 and v19). These variables represent factors associated with
 237 territories specifically deficient or absent of a sewage disposal system and the prevalence of
 238 younger people (16 to 30 years old) or older (higher than 61 years old). After the application
 239 of LR, the model with the quadratic terms was more fit when compared to the linear model. In

μ	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	0.4331	0.2063	2.0995	0.0364**
v19^1 Permanent private households with a bathroom for exclusive use by the residents or a toilet and sewage via river, lake, or sea	-6.9247	1.9192	-3.6081	0.0003**
v19^2 Permanent private households with a bathroom for exclusive use by the residents or a toilet and sewage via river, lake, or sea	-3.4713	1.1011	-3.1526	0.0017**
v33^1 Proportion of people aged 16 to 30 years old	2.6266	0.9350	2.8091	0.0052
v33^2 Proportion of people aged 16 to 30 years old	-2.4703	1.0115	-2.4423	0.0150**
v17 Permanent private households with a bathroom for exclusive use by the residents or a toilet and sanitary sewage via a septic tank	0.0044	0.0023	1.9035	0.0577
v35 Proportion of people aged 61 and over	0.0731	0.0163	4.4791	0.0000**
v18 Permanent private households with a bathroom for exclusive use by the residents or a toilet and sanitary sewage via a rudimentary septic tank	0.0042	0.0021	2.0369	0.0423**
Σ	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	0,7195	0,0761	9,4514	0,0000**

240 the table, v19^1 is the linear term of v19 and v19^2 is the quadratic term of v19. The same
 241 explanation should be considered in v33.

242

243 **Table 4. Model of the structural and intermediary social determinants associated with**
 244 **community TB infection in Eastern Amazonia (Brazil).**

245

246 *Shapiro-Wilk normality test; data: residuals (ajuste_o2c)
 247 W = 0.99681, p-value = 0.5654; Likelihood Ratio Test for nested GAMLSS models;
 248 (No check whether the models are nested is performed). Null model: deviance= 1884.811 with
 249 7 deg. of freedom
 250 Alternative model: deviance= 1866.812 with 9 deg. of freedom; LRT = 17.99909 with 2 deg.
 251 of freedom and p-value= 0.0001234661
 252 **p<0.05
 253

254 Fig 5 presents the graph of predicted values containing the values of the variables with
255 quadratic terms v19 and v33. Considering that variables v17, v18 and v35 are equal in terms
256 of their respective medians, in other words, v17 = 18.74%, v18 = 49.35% and v35 = 5.17%).
257 In this case, if we consider that v19=10%, v33=20%, the expected average number of TB cases
258 is 0.7367; When we consider v19 = 20% and v33 = 30%, the expected average number of cases
259 is 3.7824; and if we consider v19 = 30% and v33 = 40%, the average expected number of cases
260 is 2.86. Fig 6 has been added as supplementary material for predicting the number of cases in
261 accordance with the ranged value of v19 and v33.

262

263 **Fig 5. Graph with predicted values obtained from the modeling with quadratic terms,**
264 **Eastern Amazonia (Brazil).**

265

266 **Fig 6. Predicting the number of cases according to the range value of v19 and v33.**

267

268 **DISCUSSION**

269 We found evidence that GAMLSS is superior when compared to the other techniques
270 applied for studying the social determinants[12]. The first potentiality is because the GAMLSS
271 estimates the complete conditional distribution, which means that the response variable can be
272 better estimated by a probability distribution such as the gamma or lognormal distribution[24].
273 Another potentiality or advantage is that all resources are available through R packages, where
274 it is allowed to adjust more than 50 different kinds of distributions.

275 Through our findings, we identified the structural determinants (represented by age)
276 and intermediary (expressed by sanitary conditions of the environment such as treated sewage).
277 The literature has evidenced that people of economically active age who are males were more
278 commonly affected by TB, which might be related to our findings[25].

279 According to the national data, nearly 8.5% of TB cases that were diagnosed in 2017
280 were in people 19 or younger[25]. In Brazil, TB in adolescents has long remained a hidden
281 pandemic and still continues to be neglected. TB in adolescents needs to be considered a
282 sentinel event since it is related to a recent infection through contact with a bacilliferous
283 adult[25].

284 Another issue that may be associated with our findings is a situation revealed by a recent
285 IBGE survey, which concluded that the country has 14.8 million unemployed people, which
286 represents 14.7% of the economically active population. However, this rate is even higher
287 among young people; in the 14 to 17 age group, 46% are looking for work; and from 18 to 24
288 years old, unemployment affects 31% of people, which takes these people into an extreme
289 social vulnerability zone. The study also evidenced that long-term unemployment is
290 characterized by the predominance of people between the ages of 17 and 29. This data is from
291 Brazil, but when it is considered data from Eastern Amazonia, this situation can become even
292 more serious[26].

293 We also found the association with territories with a predominance of elderly people
294 (age older than 61 years old), and the literature has shown that the elderly are more vulnerable
295 to falling into poverty when compared to the other groups[27]. This is because the elderly have
296 a lesser chance of recovering from a negative income shock, and they have difficulties
297 (re)entering the labor market because productivity and employability decline with age from
298 around the age of 60. As a result of the difficulty of recovering from negative income shocks
299 among the elderly, poverty tends to be a more permanent feature than with other groups in the
300 society[27], which makes them vulnerable to poverty as well as to the TB determinants[28].

301 A study from Eastern Amazonia has shown a growth of TB among the elderly, which
302 is related to the deterioration of social conditions because of austerity politics adopted in Brazil,

303 mainly related to the weakness in social security in ensuring social protection to the elderlies,
304 and their difficulty in getting good food, quality of life and improved house conditions[29].

305 We also observed that the context favorable for sickness due to TB is associated with
306 access to basic sanitation. This is another serious situation observed in developing countries;
307 according to literature, more than 2 billion people worldwide do not have access to basic
308 sanitation, which represents more than 25% of the world's population[30]. According to a
309 study that assessed the access to basic sanitation in Brazil, almost 90% of the residents of
310 Macapá do not have access to the sewage network, and as in the whole country, Macapá invests
311 less than 30% of their financial resources picked up with taxes[31].

312 Investments in terms of better housing conditions and basic sanity is a condition
313 essential for the quality of life of people as well as for avoiding neglected diseases such as TB.
314 The Sustainable Development Goals, also known as the Global Goals, seek to reconcile
315 economic growth, environmental balance, and social progress, ensuring that all people have
316 the same opportunities and can lead a better life without compromising the planet[32]. It is
317 among its goals by 2030 to achieve universal and equitable access to basic sanitation and clean
318 water; however, achieving these goals is a challenge in Brazil, mainly in Eastern Amazonia.

319 Eastern Amazonia has the largest percentage of its territory set aside for integral natural
320 protection through the so-called Legally Allocated Areas, and contradictorily, has almost half
321 of its population living below the poverty line, with 45.9% of people having one full meal every
322 three days. A large part living in subhuman conditions in hangover areas, worsening the
323 pictures of violence, suicide, and public health, while leaving 20.2% of its entire labor force
324 unemployed. Therefore, all these aspects contributed a lot to the sickness of the community
325 due to TB. It is important to find strategies not only for prevention, diagnosis, treatment, and
326 recovery / rehabilitation, but to fundamentally find solutions sustainable and balanced with the
327 environment in Amazonia.

328 Therefore, politics addressed to ensure sustainability is quite important for bringing
329 better conditions of life to the people who live in the region. Historically, the poverty in the
330 region became evident from the decolonization process with the formation of a dependent labor
331 market in extremely precarious conditions. New strategies are important and effective public
332 policies aimed at social justice to the specificities of the Amazon region and not just an attempt
333 to solve social inequalities with compensatory policies such as the *Bolsa Familia*[33].

334 The study advances knowledge by evidencing the structural and intermediary social
335 determinants of TB in Eastern Amazonia. However, the model only included cases of TB. It
336 would be interesting to advance with other aspects of the disease through mathematical models
337 such as infection and confirm if they are the same as the disease[4]. We used the GAMLSS
338 that evidenced important aspects for understanding the context of TB and how the structural
339 and intermediary determinants impacted on communities burdened by TB, where it was
340 possible to estimate the number of cases for each territory or UCT under study.

341 We used a more usual approach, which did not assume the existence of the constraint,
342 which may be a potential limitation of the study. This is the first study applied in Eastern
343 Amazonia that confirms its novelty and originality. Unfortunately, Brazil has not carried out
344 its Demographic Census yet, mainly because of the budget shortfall. Therefore, the social
345 situation identified may have worsened, mainly due to the COVID-19 pandemic, which means
346 that the goal End TB by 2050 seems more distant than we think.

347

348 **Competing interests**

349 The authors declare that they have no competing interests.

350

351 **Acknowledgements**

352 The authors greatly appreciate the outstanding and committed study assistance of Health
353 Bureau of Mapacá – AP Brazil. We thank Jason Farley and Nancy Reynolds for critically
354 reading the manuscript. Moreover, we thank CAPES and CNPq for contributing to the study.

355

356 **Author Contributions**

357 **Conceptualization:** Clóvis Luciano Giacomet, Antônio Carlos Vieira Ramos, Heriederson
358 Sávio Dias Moura, Thaís Zamboni Berra, Ricardo Alexandre Arcêncio.

359 **Data curation:** Clóvis Luciano Giacomet, Antônio Carlos Vieira Ramos, Thaís Zamboni
360 Berra, Jonas Bodini Alonso.

361 **Formal analysis:** Clóvis Luciano Giacomet, Antônio Carlos Vieira Ramos, Thaís Zamboni
362 Berra, Jonas Bodini Alonso.

363 **Funding acquisition:** Clóvis Luciano Giacomet, Heriederson Sávio Dias Moura, Ricardo
364 Alexandre Arcêncio.

365 **Methodology:** Clóvis Luciano Giacomet, Antônio Carlos Vieira Ramos, Thaís Zamboni
366 Berra, Felipe Mendes Delpino, Jonas Bodini Alonso, Ricardo Alexandre Arcêncio.

367 **Software:** Jonas Bodini Alonso, Ricardo Alexandre Arcêncio.

368 **Supervision:** Clóvis Luciano Giacomet, Ricardo Alexandre Arcêncio.

369 **Validation:** Clóvis Luciano Giacomet, Antônio Carlos Vieira Ramos, Thaís Zamboni Berra,
370 Jason E. Farley, Nancy R. Reynolds, Jonas Bodini Alonso, Ricardo Alexandre Arcêncio.

371 **Writing – original draft:** Clóvis Luciano Giacomet, Antônio Carlos Vieira Ramos,
372 Heriederson Sávio Dias Moura, Thaís Zamboni Berra, Yan Mathias Alves, Felipe Mendes
373 Delpino, Jonas Bodini Alonso, Ricardo Alexandre Arcêncio

374 **Writing – review & editing:** Clóvis Luciano Giacomet, Antônio Carlos Vieira Ramos,
375 Heriederson Sávio Dias Moura, Thaís Zamboni Berra, Yan Mathias Alves, Felipe Mendes
376 Delpino, Jason E. Farley, Nancy R. Reynolds, Jonas Bodini Alonso, Titilade Kehinde
377 Ayandeyi Teibo, Ricardo Alexandre Arcêncio.

378

379 **Funding**

380 CAPES (code 001) and CNPq (scholarship Research productivity fellowship at the 1C level –
381 process 304483/2018-4).

382

383 **REFERENCES**

384 1. World Health Organization. Global tuberculosis report 2021, 2021. Accessed: Dec 10,
385 2021. Available from: <https://apps.who.int/iris/handle/10665/346387>.

386 2. United Nations. World Social Report 2020: Inequality in a Rapidly Changing World,

- 387 2020. Accessed Dec 10, 2021. Available from: [https://www.un-ilibrary.org/economic-](https://www.un-ilibrary.org/economic-and-social-development/world-social-report-2020_7f5d0efc-en)
388 [and-social-development/world-social-report-2020_7f5d0efc-en](https://www.un-ilibrary.org/economic-and-social-development/world-social-report-2020_7f5d0efc-en).
- 389 3. Silva S, Arinaminpathy N, Atun R, Goosby E, Reid M. Economic impact of
390 tuberculosis mortality in 120 countries and the cost of not achieving the Sustainable
391 Development Goals tuberculosis targets: a full-income analysis. *Lancet Glob Health*.
392 2021;9(10): e1372–9. doi: 10.1016/S2214-109X(21)00299-0.
- 393 4. Pedrazzoli D, Boccia D, Dodd PJ, Lönnroth K, Dowdy DW, Siroka A, et al.
394 Modelling the social and structural determinants of tuberculosis: opportunities and
395 challenges. *Int J Tuberc Lung Dis Off J Int Union Tuberc Lung Dis*. 2017;21(9): 957–
396 64. doi: 10.5588/ijtld.16.0906.
- 397 5. Moreira ASR, Kritski AL, Carvalho ACC. Social determinants of health and
398 catastrophic costs associated with the diagnosis and treatment of tuberculosis. *J Bras*
399 *Pneumol*. 2020;46(05): e20200015. doi: 10.36416/1806-3756/e20200015.
- 400 6. Duarte R, Lönnroth K, Carvalho C, Lima F, Carvalho ACC, Muñoz-Torrico M, et al.
401 Tuberculosis, social determinants and co-morbidities (including HIV). *Pulmonology*
402 2018;24(2): 115-119. doi: 10.1016/j.rppnen.2017.11.003.
- 403 7. Zille AI, Werneck GL, Luiz RR, Conde MB. Social determinants of pulmonary
404 tuberculosis in Brazil: an ecological study. *BMC Pulm Med*. 19(1): 87. doi:
405 10.1186/s12890-019-0855-1.
- 406 8. Kilabuk E, Momoli F, Mallick R, Van Dyk D, Pease C, Zwerling A, et al. Social
407 determinants of health among residential areas with a high tuberculosis incidence in a
408 remote Inuit community. *J Epidemiol Community Health*. 73(5): 401-406. doi:
409 10.1136/jech-2018-211261.
- 410 9. Wingfield T, Tovar MA, Datta S, Saunders MJ, Evans CA. Addressing social
411 determinants to end tuberculosis. *Lancet*. 391(10126): 1129-1132. doi:

- 412 10.1016/S0140-6736(18)30484-7.
- 413 10. Rigby RA, Stasinopoulos DM. Generalized additive models for location, scale and
414 shape. *J R Stat Soc Ser C Appl Stat.* 2005;54(3): 507–54. doi: 10.1111/j.1467-
415 9876.2005.00510.x.
- 416 11. De Bastiani F, Rigby RA, Stasinopoulos DM, Cysneiros AHMA, Uribe-Opazo MA.
417 Gaussian Markov random field spatial models in GAMLSS. *J Appl Stat.* 45(2018):
418 168–86. doi: 10.1080/02664763.2016.1269728.
- 419 12. Giacomet CL, Santos MS, Berra TZ, Alves YM, Alves LS, Costa FBP, et al.
420 Temporal trend of tuberculosis incidence and its spatial distribution in Macapá –
421 Amapá. *Rev Saude Publica.* 55: 96. doi: 10.11606/s1518-8787.2021055003431.
- 422 13. Santana JLA, Magrini AT, Ribeiro LA, Rigelli RH, Araujo MHM. Adesão ao
423 tratamento da tuberculose no Amapá: Um quinquênio de análise epidemiológica. *Rev*
424 *Científica Multidiscip Núcleo Conhecimento.* 10: 69–87.
425 doi: 10.32749/nucleodoconhecimento.com.br/saude/tratamento-da-tuberculose.
- 426 14. World Health Organization. Meeting report of the WHO expert consultation on drug-
427 resistant tuberculosis treatment outcome definitions, 17-19 November 2020.
428 Accessed: Dec 13, 2021. Available from:
429 <https://apps.who.int/iris/handle/10665/340284>.
- 430 15. Brasil. Ministério da Saúde. Secretaria de Vigilância em Saúde. Departamento de
431 Vigilância das Doenças Transmissíveis. Manual de Recomendações para o Controle
432 da Tuberculose no Brasil, 2019; 2a ed. atualizada, 364. Accessed: Dec 13, 2021.
433 Available from:
434 https://bvsms.saude.gov.br/bvs/publicacoes/manual_recomendacoes_controle_tubercu
435 [lose_brasil_2_ed.pdf](https://bvsms.saude.gov.br/bvs/publicacoes/manual_recomendacoes_controle_tubercu).
- 436 16. Boing AF, Boing AC, Subramanian SV. Inequalities in the access to healthy urban

- 437 structure and housing: an analysis of the Brazilian census data. *Cad Saúde Pública*.
438 37(6): e00233119. doi: 10.1590/0102-311X00233119.
- 439 17. Solar O, Irwin A. A conceptual framework for action on the social determinants of
440 health, *Social Determinants of Health Discussion Paper 2 (Policy and Practice)*.
441 Geneva, Switzerland: World Health Organization. 2010. Accessed: Dec 10, 2021.
442 Available from: <https://apps.who.int/iris/handle/10665/44489>.
- 443 18. Florencio LA. Engenharia de avaliações com base nos modelos GAMLSS. Master's
444 Thesis, Federal University of Pernambuco, 2010. Available from:
445 <https://repositorio.ufpe.br/handle/123456789/6227>.
- 446 19. Akaike H. A new look at the statistical model identification. *IEEE Trans Autom*
447 *Control*. 19(6): 716–23. doi: 10.1109/TAC.1974.1100705.
- 448 20. Stasinopoulos M, Rigby B, Akantziliotou C. Instructions on how to use the *gamlss*
449 package in R, 2008. Accessed: Dec 14, 2021. Available from:
450 <http://www.gamlss.com/wp-content/uploads/2013/01/gamlss-manual.pdf>.
- 451 21. Myers RH, Montgomery DC, Anderson-Cook CM. *Response surface methodology:*
452 *process and product optimization using designed experiments*. Vanderbilt University,
453 Hoboken, New Jersey: Wiley, 2016. Accessed: Dec 21, 2021. Available from:
454 <https://catalog.library.vanderbilt.edu/discovery/fulldisplay/alma991043342547203276>
455 [/01VAN_INST:vanui](#).
- 456 22. Stasinopoulos DM, Rigby RA. Generalized Additive Models for Location Scale and
457 Shape (GAMLSS) in R. *Journal of Statistical Software*. 23(7): 1–46.
458 <https://doi.org/10.18637/jss.v023.i07>.
- 459 23. Jesus FB, Lima FCA, Martins CBG, Matos KF, Souza SPS. Vulnerabilidade na
460 adolescência: a experiência e expressão do adolescente. *Rev Gaúcha Enferm*. 32(2):
461 359-67. doi: 10.1590/S1983-14472011000200021.

- 462 24. Shelby T, Meyer AJ, Ochom E, Turimumahoro P, Babirye D, Katamba A, et al.
463 Social determinants of tuberculosis evaluation among household contacts: a
464 secondary analysis. *Public Health Action*. 8(3): 118-123. doi: 10.5588/fa.18.0025.
- 465 25. Santos BA, Cruz RPS, Lima SVMA, Santos AD, Duque M, Araújo KCGM, et al.
466 Tuberculose em crianças e adolescentes: uma análise epidemiológica e espacial no
467 estado de Sergipe, Brasil, 2001-2017. *Ciênc Saúde Coletiva*. 25(8): 2939–48. doi:
468 10.1590/1413-81232020258.25692018.
- 469 26. Brasil. Instituto Brasileiro de Geografia e Estatísticas. Pesquisa aponta que os jovens
470 são os mais afetados pelo desemprego, 2021. Accessed: Jan 21, 2021. Available from:
471 [https://agenciabrasil.ebc.com.br/radioagencia-nacional/economia/audio/2021-](https://agenciabrasil.ebc.com.br/radioagencia-nacional/economia/audio/2021-08/pesquisa-aponta-que-os-jovens-sao-os-mais-afetados-pelo-desempregohtml)
472 [08/pesquisa-aponta-que-os-jovens-sao-os-mais-afetados-pelo-desempregohtml](https://agenciabrasil.ebc.com.br/radioagencia-nacional/economia/audio/2021-08/pesquisa-aponta-que-os-jovens-sao-os-mais-afetados-pelo-desempregohtml).
- 473 27. Travassos GF, Coelho AB, Arends-Kuenning MP. The elderly in Brazil: demographic
474 transition, profile, and socioeconomic condition. *Rev Bras Estud Popul*. 37: 1–27. doi:
475 10.20947/S0102-3098a0129.
- 476 28. Negin J, Abimbola S, Marais BJ. Tuberculosis among older adults – time to take
477 notice. *Int J Infect Dis*. 32: 135–7. doi: 10.1016/j.ijid.2014.11.018.
- 478 29. Mesquita CR, Lima KVB, Souza e Guimarães RJP, Santos BO, Rodrigues LHA,
479 Costa RJF, et al. Análise retrospectiva de casos de tuberculose em idosos. *Rev Bras*
480 *Em Promoção Saúde* 2021;34: 1117. doi: 10.5020/18061230.2021.11117.
- 481 30. Swe KT, Rahman MM, Rahman MS, Teng Y, Abe SK, Hashizume M, et al. Impact of
482 poverty reduction on access to water and sanitation in low- and lower-middle-income
483 countries: country-specific Bayesian projections to 2030. *Trop Med Int Health*. 26(7):
484 760-774. doi: 10.1111/tmi.13580.
- 485 31. Instituto Trata Brasil. Ranking do Saneamento 2021, 2021. Accessed: Jan 19, 2022.
486 Available from:

487 https://tratabrasil.org.br/images/estudos/Ranking_saneamento_2021/Ranking_do_San
488 [eamento_2021_-_tabela_das_100_maiores_cidades_do_Brasil_.pdf](https://tratabrasil.org.br/images/estudos/Ranking_saneamento_2021/Ranking_do_Saneamento_2021_-_tabela_das_100_maiores_cidades_do_Brasil_.pdf).


489 32. Duran DC, Artene A, Gogan LM, Duran V. The Objectives of Sustainable
490 Development - Ways to Achieve Welfare. *Procedia Econ Finance*. 26(2015): 812–7.
491 doi: 10.1016/S2212-5671(15)00852-7.

492 33. Rolim DC. A pobreza e a riqueza na região amazônica e a contribuição da política de
493 assistência social: o Estado do Amazonas em foco. VII Jornada Internacional Políticas
494 Públicas. Universidade Federal do Maranhão. Centro de Ciências Sociais. Programa
495 de Pós-Graduação em Políticas Públicas 2015:13. Accessed: Jan 19, 2022. Available
496 from: [http://www.joinpp.ufma.br/jornadas/joinpp2015/pdfs/eixo4/a-pobreza-e-a-](http://www.joinpp.ufma.br/jornadas/joinpp2015/pdfs/eixo4/a-pobreza-e-a-riqueza-na-regiao-amazonica-e-a-contribuicao-da-politica-de-assistencia-social-o-estado-do-amazonas-em-foco.pdf)
497 [riqueza-na-regiao-amazonica-e-a-contribuicao-da-politica-de-assistencia-social-o-](http://www.joinpp.ufma.br/jornadas/joinpp2015/pdfs/eixo4/a-pobreza-e-a-riqueza-na-regiao-amazonica-e-a-contribuicao-da-politica-de-assistencia-social-o-estado-do-amazonas-em-foco.pdf)
498 [estado-do-amazonas-em-foco.pdf](http://www.joinpp.ufma.br/jornadas/joinpp2015/pdfs/eixo4/a-pobreza-e-a-riqueza-na-regiao-amazonica-e-a-contribuicao-da-politica-de-assistencia-social-o-estado-do-amazonas-em-foco.pdf).

499

500 **Supporting information**

501 S1 – Dataset





medRxiv preprint doi: <https://doi.org/10.1101/2022.11.03.22281901>; this version posted November 4, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](#).

medRxiv preprint doi: <https://doi.org/10.1101/2022.11.03.22281901>; this version posted November 1, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).




Social Determinants	Definition	proxy indicators	Variables	Code
Structural	The structural determinant are those that interplay between context and socioeconomic position, generating and reinforcing class divisions, define the socio-economical position within hierarchies of power, prestige, and access to resources[18].	Income	- Proportion of households with no nominal monthly income per capita	v20
			- Proportion of households with per capita nominal monthly income of up to 1/8 to 1 minimum wage	v21
			- Proportion of households with a nominal monthly per capita income of more than 1 to 2 minimum wages	v22
			- Proportion of domiciles with nominal monthly income per capita of more than 2 to 3 minimum wages	v23
			- Proportion of households with a nominal monthly income per capita of more than 3 to 5 minimum wages	v24
		- Proportion of households with a nominal monthly income per capita of more than 5 to 10 minimum wages	v25	
		Education and gender	- Proportion of males without schooling - Proportion of females without schooling	v05 v06
		Race/ ethnicity	- Proportion of resident people of white race or ethnicity	v26
			- Proportion of black residents	v27
			- Proportion of resident people of yellow race or ethnicity	v28
			- Proportion of resident people of mixed race or ethnicity	v29
			- Proportion of resident people of indigenous race or ethnicity	v30
Intermediary	Structural determinants operate through a series of factors that will influence the health condition of a person, which WHO[18] named as the intermediary social factors. These factors are accumulative in a person's life course and flow from the social stratification, determining differences in exposure and vulnerability and	Agglomeration (quantitative of people who live in the domiciles)	- Proportion of households with 1 to 3 people - The proportion of households with 4 to 6 people - The proportion of households with 7 to 9 people - The proportion of households with higher than 10 people	v01 v02 v03 v04
		Age	- Proportion of people aged 0-15 - Proportion of people aged 16 to 30 years old - Proportion of persons aged 31 to 60 years old - Proportion of people aged 61 and over	v31 v32 v33 v34
		Conditions of domiciles' property	- Proportion of own and paid-up domiciles - Proportion of rented dwellings - Proportion of domiciles in another property condition (not owned, rented, and not granted)	v07 v08 v09
		Sanitation conditions	- Proportion of households with a bathroom for exclusive use by residents or a toilet and drainage in a general sewage or rainwater network	v10
			- Proportion of houses without a bathroom for the exclusive use of residents or toilet	v11
			- Permanent particular residences with water supply from the general network	v12


	<p>are therefore, health-compromising conditions. These factors are material circumstances; psychosocial circumstances; behavioral and/or biological factors, and the health system</p>		<ul style="list-style-type: none"> - Permanent particular residences with water supply from wells or springs on the property - Permanent Particular Domiciles with Another Form of Water Supply - Permanent private households with a bathroom exclusively for residents or a toilet and sanitary sewage via the general sewage system or rainwater drainage system - Permanent private households with a bathroom for exclusive use by the residents or a toilet and sanitary sewage via a septic tank - Permanent private households with a bathroom for exclusive use by the residents or a toilet and sanitary sewage via a rudimentary septic tank - Permanent private households with a bathroom for exclusive use by the residents or a toilet and sewage via river, lake, or sea 	<p>v13</p> <p>v14</p> <p>v15</p> <p>v16</p> <p>v17</p> <p>v18</p> <p>v19</p>
--	---	--	--	--


medRxiv preprint doi: <https://doi.org/10.1101/2022.11.03.22281901>; this version posted November 4, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).



medRxiv preprint doi: <https://doi.org/10.1101/2022.11.03.22281901>; this version posted November 4, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).







medRxiv preprint doi: <https://doi.org/10.1101/2022.11.03.22281901>; this version posted November 4, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .

v19	v33	v17	v18	v35	Predicted
0	0	18.74	49.35	5.17	0.0001
1	0	18.74	49.35	5.17	0.0001
2	0	18.74	49.35	5.17	0.0001
3	0	18.74	49.35	5.17	0.0001
4	0	18.74	49.35	5.17	0.0001
5	0	18.74	49.35	5.17	0.0001
6	0	18.74	49.35	5.17	0.0001
7	0	18.74	49.35	5.17	0.0001
8	0	18.74	49.35	5.17	0.0001
9	0	18.74	49.35	5.17	0.0001
10	0	18.74	49.35	5.17	0.0001
11	0	18.74	49.35	5.17	0.0001
12	0	18.74	49.35	5.17	0.0001
13	0	18.74	49.35	5.17	0.0001
14	0	18.74	49.35	5.17	0.0001
15	0	18.74	49.35	5.17	0.0001
16	0	18.74	49.35	5.17	0.0001
17	0	18.74	49.35	5.17	0.0001
18	0	18.74	49.35	5.17	0.0001
19	0	18.74	49.35	5.17	0.0001
20	0	18.74	49.35	5.17	0.0001
21	0	18.74	49.35	5.17	0.0001
22	0	18.74	49.35	5.17	0.0001
23	0	18.74	49.35	5.17	0.0001
24	0	18.74	49.35	5.17	0.0001
25	0	18.74	49.35	5.17	0.0001
26	0	18.74	49.35	5.17	0.0001
27	0	18.74	49.35	5.17	0.0001
28	0	18.74	49.35	5.17	0.0001
29	0	18.74	49.35	5.17	0.0001
30	0	18.74	49.35	5.17	0.0001
31	0	18.74	49.35	5.17	0.0001
32	0	18.74	49.35	5.17	0.0001
33	0	18.74	49.35	5.17	0.0001
34	0	18.74	49.35	5.17	0.0001
35	0	18.74	49.35	5.17	0.0001
36	0	18.74	49.35	5.17	0.0001
37	0	18.74	49.35	5.17	0.0001
38	0	18.74	49.35	5.17	0.0001
39	0	18.74	49.35	5.17	0.0001
40	0	18.74	49.35	5.17	0.0001
41	0	18.74	49.35	5.17	0.0001
42	0	18.74	49.35	5.17	0.0001

medRxiv preprint doi: <https://doi.org/10.1101/2022.11.03.22281901>; this version posted November 4, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

43	0	18.74	49.35	5.17	0.0001
44	0	18.74	49.35	5.17	0.0001
45	0	18.74	49.35	5.17	0.0001
46	0	18.74	49.35	5.17	0.0001
47	0	18.74	49.35	5.17	0.0001
48	0	18.74	49.35	5.17	0.0001
49	0	18.74	49.35	5.17	0.0001
50	0	18.74	49.35	5.17	0.0001
51	0	18.74	49.35	5.17	0.0001
52	0	18.74	49.35	5.17	0.0001
53	0	18.74	49.35	5.17	0.0001
54	0	18.74	49.35	5.17	0.0001
55	0	18.74	49.35	5.17	0.0001
56	0	18.74	49.35	5.17	0.0001
57	0	18.74	49.35	5.17	0.0001
58	0	18.74	49.35	5.17	0.0001
59	0	18.74	49.35	5.17	0.0001
60	0	18.74	49.35	5.17	0.0001
61	0	18.74	49.35	5.17	0.0001
62	0	18.74	49.35	5.17	0.0001
63	0	18.74	49.35	5.17	0.0001
64	0	18.74	49.35	5.17	0.0001
65	0	18.74	49.35	5.17	0.0001
66	0	18.74	49.35	5.17	0.0001
67	0	18.74	49.35	5.17	0.0001
68	0	18.74	49.35	5.17	0.0001
69	0	18.74	49.35	5.17	0.0001
70	0	18.74	49.35	5.17	0.0001
71	0	18.74	49.35	5.17	0.0001
72	0	18.74	49.35	5.17	0.0000
73	0	18.74	49.35	5.17	0.0000
74	0	18.74	49.35	5.17	0.0000
75	0	18.74	49.35	5.17	0.0000
76	0	18.74	49.35	5.17	0.0000
77	0	18.74	49.35	5.17	0.0000
78	0	18.74	49.35	5.17	0.0000
79	0	18.74	49.35	5.17	0.0000
80	0	18.74	49.35	5.17	0.0000
81	0	18.74	49.35	5.17	0.0000
82	0	18.74	49.35	5.17	0.0000
83	0	18.74	49.35	5.17	0.0000
84	0	18.74	49.35	5.17	0.0000
85	0	18.74	49.35	5.17	0.0000
86	0	18.74	49.35	5.17	0.0000
87	0	18.74	49.35	5.17	0.0000

medRxiv preprint doi: <https://doi.org/10.1101/2022.11.03.22281901>; this version posted November 4, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

88	0	18.74	49.35	5.17	0.0000
89	0	18.74	49.35	5.17	0.0000
90	0	18.74	49.35	5.17	0.0000
91	0	18.74	49.35	5.17	0.0000
92	0	18.74	49.35	5.17	0.0000
93	0	18.74	49.35	5.17	0.0000
94	0	18.74	49.35	5.17	0.0000
95	0	18.74	49.35	5.17	0.0000
96	0	18.74	49.35	5.17	0.0000
97	0	18.74	49.35	5.17	0.0000
98	0	18.74	49.35	5.17	0.0000

medRxiv preprint doi: <https://doi.org/10.1101/2022.11.03.22281901>; this version posted November 4, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/) .